

Sarringhaus Lauren (Orcid ID: 0000-0002-8819-5118)
Mitani John C. (Orcid ID: 0000-0001-7042-5854)

Title: The ontogeny of knuckle-walking and dorsal metacarpal ridge prominence in chimpanzees.

Authors: Sarringhaus, L.^{1,2}, Mitani, J. C.², MacLatchy, L.M.^{2*}

Affiliation: ¹Duke University, Department of Evolutionary Anthropology, ²University of Michigan, Department of Anthropology

***Corresponding author maclatch@umich.edu**

Abstract:

Objectives: This study examines the sensitivity of the development of the dorsal metacarpal ridge (DMR) to the frequency of knuckle-walking during ontogeny in chimpanzees, and compares the prevalence and prominence of the DMR in hominoids and baboons.

Materials and Methods: We characterized the type and frequency of quadrupedalism of wild chimpanzees of different ages from the Ngogo community, Uganda. Using museum specimens, we quantified third metacarpal DMR angle and height in *Pan* individuals of different developmental stages, and in adult *Gorilla*, *Pongo*, *Hylobates* and *Papio*.

Results: When terrestrial, all quadrupedalism consisted of knuckle-walking, although older individuals knuckle-walked more than younger. The total amount of quadrupedalism engaged in by different age groups while arboreal was consistent; however, knuckle usage while arboreally quadrupedal increased with age. The DMR ridge did not emerge in *Pan* until older infancy, after the onset of knuckle-walking, and reached prominence by the juvenile stage. The DMR angle increased significantly after juvenility but relative height did not. Adult *Pan* and *Gorilla* have more prevalent and prominent DMRs than *Papio*; the DMR was rare in *Pongo* and absent in *Hylobates*.

Discussion: The timing of development of the DMR in chimpanzees supports it as a knuckle walking character, as does its occurrence in gorillas, and virtual absence in suspensory *Pongo* and *Hylobates*. The DMR in baboons differs in appearance and frequency from DMRs in African apes, and so differentiates knuckle-walking from digitigrady in catarrhines. The relationship between the DMR and loading of a vertical hand while terrestrial deserves further study.

Keywords: locomotor development, *Gorilla*, *Papio*, digitigrade, quadrupedalism

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/ajpa.24477

This article is protected by copyright. All rights reserved.

Introduction

Bipedality is a hallmark of humankind and is frequently used as the *sine qua non* to recognize our earliest human ancestors (Haile-Selassie, 2001; MacLatchy, DeSilva, Sanders, & Wood, 2010; Zollikofer et al., 2005). Due to significant gaps in the fossil record, how our ancestors moved before they became bipedal is unknown, yet multiple hypotheses have been proposed for over a century (Böhme et al., 2019; Fleagle et al., 1981; Keith, 1923; Kimura, 2019; Morton, 1922 & 1924; Prost, 1980; Schultz 1930; Stern, 1975; Stern & Susman, 1981; Straus, 1949; Tuttle, 1967; Tuttle & Basmajian, 1974; Wood Jones, 1929 to reference a subset see Rose, 1991; Richmond, Begun, & Strait, 2001; Pilbeam & Lieberman, 2017 for reviews). One of the most intensely debated hypotheses is whether early humans evolved from a chimpanzee-like, knuckle-walking ancestor that moved quadrupedally on the ground (Begun, 1992; Dainton & Macho, 1999; Gregory, 1916; Kivell & Schmitt, 2009; Richmond & Strait, 2000; Washburn, 1967; Williams, 2010). Other hypotheses, such as an orangutan-like ancestor that moved upright using orthograde clambering and hand-assisted bipedality in the trees (Crompton, Sellers, & Thorpe, 2010; Crompton, Vereecke, & Thorpe, 2008; Thorpe, Holder, & Crompton, 2007; Thorpe, McClymount, & Crompton, 2014; see Böhme et al., 2019 for a variant) or a *Proconsul*-like arboreal ancestor that walked on the palms of its hands (Lovejoy, Suwa, Simpson, Matternes, & White, 2009; Lovejoy, Suwa, Spurlock, Asfaw, & White, 2009; Lovejoy & McCollum, 2010; White, Lovejoy, Asfaw, Carlson, & Suwa, 2015) have gained traction recently, but the knuckle-walking hypothesis continues to remain influential (Pilbeam & Lieberman, 2017).

Testing the knuckle-walking hypothesis will depend, in part, on determining if there are skeletal correlates unique to this hand posture, and then looking for these features in the fossil record. Resolving such form-function relationships would be aided by data on the actual hand postures used during locomotion in the wild and further quantification of hypothetically distinctive bony features. Prior studies indicate that knuckle-walking is a rare form of quadrupedalism performed by only two genera of primates, *Pan* and *Gorilla*. During knuckle-walking, the fingers of these apes are tucked under the hand, the metacarpophalangeal (MCP) joints are extended, and the interphalangeal joints are flexed so that the middle phalanges contact the substrate. Additional research shows that quadrupedal walking increases as chimpanzees age (Doran, 1992; Sarringhaus, MacLatchy, & Mitani, 2014), with infants spending less than a quarter of locomotor time, juveniles spending nearly half, and adolescents and adults devoting the majority of their time moving this way (Sarringhaus et al., 2014). The increase in quadrupedal locomotion from infancy to juvenility is even more pronounced when distance traveled is taken into consideration (Doran, 1992).

Although the approximate timeline of the transition to quadrupedalism has been characterized, the development of knuckle-walking *per se* has yet to be determined. Adult chimpanzees primarily knuckle-walk when moving quadrupedally, but prior studies indicate that knuckle-walking develops gradually (Kimura, 1986; Doran, 1992; Sarringhaus, 2013). Doran (1992) observed one infant chimpanzee (4-6 months) using its palms during a bout of quadrupedal locomotion. She observed older infants (6 months - 2 years) knuckle-walking in 2 of 8 quadrupedal bouts (*ibid.*). Additional qualitative observations suggested to Doran that the degree to which chimpanzees knuckle-walk during ontogeny depends in part on whether they are arboreal or terrestrial (*ibid.*). These observations provide a valuable framework for tracking the development of knuckle-walking, but small samples and the lack of quantitative data on the behavior preclude making any general conclusions.

Given the hypothesis that the evolution of bipedalism involved a knuckle-walking phase, there has been considerable scrutiny of possible anatomical correlates of knuckle-walking hand postures. The wrist has been a region of interest in this regard, and several anatomical features have been proposed to be related to knuckle-walking behavior, including articular facets in the capitate (Corruccini, 1978; Wuthrich, MacLatchy, & Nengo, 2019), distal projection of the dorsal radius (Richmond et al., 2001; Tuttle, 1967), and os central fusion (Kivell & Begun, 2007; Orr, 2018). However, no single wrist feature has emerged as unequivocally predictive of knuckle-walking behavior, as some features such as os central fusion and distal radial projection are not unique to knuckle-walking apes (see Kivell, 2016 for a review). Moreover, some features of the wrist, such as degree of capitate waisting, manifest along a continuum among genera (Kivell & Schmitt, 2009). A recent multivariate study successfully partitioned *Pan* and *Gorilla* from Asian apes and other catarrhines using anatomical landmarks associated with capitate surface architecture (Wuthrich et al., 2019). However, this study also noted that individual knuckle-walking traits are not exclusively associated with knuckle-walking, and that inferring this positional behavior from wrist morphology remains challenging.

The metacarpal (MC) represents another promising bone for detecting anatomical signatures of knuckle-walking, as the hyperextension that occurs at the metacarpal phalangeal (MCP) joint during hand contact with the substrate is predicted to direct loads to the dorsal aspect of the MC head (Chirchir, Zeininger, Nakatsukasa, Ketchman, & Richmond, 2017; Zeininger, Richmond, & Hartman, 2011;). In contrast, suspension and climbing are predicted to preferentially load the distal and palmar aspects of the MC heads, respectively (*ibid.*). Some support for this prediction comes from studies of the internal trabecular distribution in the third MC head of chimpanzees and other hominoids. Although imaging, sampling and analytical methods have varied, multiple studies have found that trabecular bone in the dorsal aspect of the metacarpal head has a relatively high bone volume fraction in knuckle-walkers compared to other hominoids (Chirchir et al. 2017; Dunmore, Kivell, Bardo, & Skinner, 2019; Tsegei et al. 2013). In contrast, Barak, Sherratt, & Lieberman (2017) did not find a relatively high bone volume fraction in the dorsal metacarpal head of chimpanzees, but did detect an architectural difference between humans and chimpanzees, with the 3D principal trabecular orientation aligning with the long axis of the metacarpal shaft in humans, but deviating either medially or laterally in chimpanzees. The authors argued this orientation possibly reflects directional loading on the MC head that results from subtle responses by the hands in balancing more lateral or medial hindlimb postures. Matarazzo (2015) also found that trabecular alignment in the metacarpal head of knuckle-walkers is distinguishable from that of suspensors and generalized quadrupeds, while Zeininger et al. (2011) has demonstrated lower mineralization in the dorsal aspect of the third MC head of chimpanzees compared to humans, possibly reflecting higher rates of bone remodeling in this region due to the joint reaction forces generated by knuckle-walking.

The dorsal aspect of the MC head has also been shown to have a dorsal metacarpal ridge (DMR). This raised ridge of bone on the distal end of the MC has been hypothesized to be a knuckle-walking feature that acts to stabilize the metacarpophalangeal joint during hyperextension (Tuttle, 1967; Lewis, 1977; Sarmiento, 1994; Williams, 2010). Furthermore, the DMR is purportedly found in knuckle-walking gorillas and chimpanzees but is absent in non-knuckle-walking humans, gibbons, and orangutans (Lewis, 1977; Preuschoft, 1973; Richmond et al., 2001; Susman, 1979; Tuttle, 1967). The DMR is present most often on the third MC in African apes (79%-100%) and displays variation in its incidence and degree on other digits (Inouye, 1994; Inouye & Shea, 2004; Susman, 1979). The third MC is also the digit loaded most consistently by captive chimpanzees, while there is less variation in gorilla loading patterns (Matarazzo, 2008; Wunderlich & Jungers, 2009). Adult *Pan troglodytes*, *Gorilla gorilla* and *Gorilla beringei* display the DMR more frequently than do subadults, and the height of the feature scales with body size (Inouye & Shea, 2004). Taken together, the preceding observations suggest that the DMR is a plastic feature that forms during ontogeny, a period when bones are most sensitive to repeated locomotor activity (Lieberman, Pearson, Polk, Demes, & Crompton, 2003; Pontzer et al., 2006; Turner, Takano, & Owan, 1995)¹. Nonetheless, the precise time the DMR starts to emerge during development in the African apes remains unclear.

The DMR has anecdotally been noted to be present in large-bodied digitigrade primates such as *Papio* (Richmond et al., 2001), a taxon that aligns the manus vertically with hyperextended metacarpophalangeal joints (Patel, 2010). During digitigrade locomotion, the MC head is load bearing and projected to experience the greatest strain on the dorsal and central sides (Chirchir et al., 2017). The presence of the DMR in African apes and baboons suggests that the DMR may not represent a unique knuckle-walking feature. Instead, the DMR may be a MCP joint stabilizer during MCP hyperextension for individuals above a certain body size that engage in habitual terrestriality, whether through digitigrady or knuckle-walking (Richmond et al., 2001; Orr, 2005; Patel & Maiolino, 2016.). Testing this hypothesis requires comparing DMR prevalence and prominence in digitigrade *Papio* and knuckle-walking *Pan* and *Gorilla*.

While multiple researchers have attributed a stabilizing, functional role to the DMR during knuckle-walking and/or digitigrady (Tuttle, 1967; Lewis, 1977; Sarmiento, 1994; Williams, 2010; Richmond *et al.*, 2001; Orr, 2005; Patel and Maiolino, 2016), Simpson, Latimer, and Lovejoy (2018) have challenged this interpretation on two grounds. First, they suggest that the ridge cannot act as a stabilizer because regular loading of the adult bone in this fashion would lead to osteoarthritis, and second, they suggest that the DMR is an incidental byproduct of loads experienced by the subadult cartilaginous epiphysis during knuckle-walking. This caution is warranted. However, addressing whether the DMR is a direct stabilizer or a byproduct of habitual use would require clinical and/or experimental approaches beyond the scope of this study.

There are three goals of this study. First, we assess the onset of knuckle-walking in chimpanzees, a behavior characterized by alterations in the hand- and wrist-loading environment. We document developmental changes in the loading environment by differentiating hand contact and degree of arboreality used by chimpanzees of different ages. We then test the

¹ These references do not explicitly examine bony ridge formation near the epiphyseal plate, which may utilize different developmental pathways.

hypothesis that hand posture changes during ontogeny, with knuckle-walking increasing with increasing quadrupedalism and terrestriality. Our second goal is to assess changes in chimpanzee DMR features over the same developmental timeline through an analysis of an ontogenetic series of aged MCs from museum collections. Here we investigate whether the DMR changes during development in tandem with the change in loading environment experienced by the wrist and hand with the onset of knuckle-walking. Specifically, we seek to determine whether DMR development tracks knuckle-walking. Such tracking would be consistent with a phenotypic plasticity explanation, although it does not unequivocally demonstrate it. Third, we attempt to ascertain whether DMR morphology is unique to knuckle-walkers. To do so, we conduct comparisons between chimpanzees and other primate species, who move in different ways, and test three hypotheses:

The null hypothesis (H0) proposes that the DMR is not affected by hand usage. H0 predicts that the presence and prominence of the DMR will fail to track the development of knuckle-walking in chimpanzees (i.e. it will not increase with increasing knuckle usage). It also predicts that the prevalence and prominence of the DMR will not vary among genera (*Pan*, *Gorilla*, *Papio*, *Pongo* and *Hylobates*) according to hand usage.

The first alternate hypothesis (H1) proposes the DMR reflects knuckle-walking. This hypothesis predicts that the DMR will develop after subadult *Pan* start to knuckle-walk and that the DMR will be more prevalent and prominent in *Pan* and *Gorilla* compared to *Papio*, *Pongo* and *Hylobates*.

The second alternate hypothesis (H2) proposes that the DMR reflects vertical manus terrestrial quadrupedalism, including both knuckle-walking and digitigrady. Like H1, H2 predicts that the DMR will develop after knuckle-walking in subadult *Pan*. Additional predictions of H2 are that the DMR will be prevalent in adult *Pan*, *Papio*, and *Gorilla* but not in *Pongo* and *Hylobates*, and furthermore, that the prevalence and prominence of the DMR will be similar among *Pan*, *Gorilla*, and *Papio*.

Materials and Methods

Behavioral Methods

Data on the locomotor behavior of chimpanzees were collected by L.S. at Ngogo, Kibale National Park, Uganda. The unusually large size of the Ngogo community (approximately 160 individuals) provided a rare opportunity to sample the locomotor behavior of many chimpanzees who varied in age. Chimpanzees were divided into four age categories: infants (0.1 – 5 years), juveniles (5.1 – 10 years), adolescents (10.1 – 14.9 years), and adults (15 + years; Table 1). When examining only infants, the largest increase in quadrupedal locomotion occurs when individuals reach three years of age (Sarringhaus et al., 2014; Table 1). The infant category therefore was further divided into *young infant* (≤ 3 years) and *old infant* (3 - 5 years) groups. Ages of subadults were determined based on their birth dates known to within a few days or a calendar year (Wood, Watts, Mitani, & Langergraber, 2017). The birth dates of some adults were estimated, as these individuals were all over 20 years of age. Their ages were assigned using genealogical information, derived from genetic data and behavioral observations, and information regarding their physical appearance (Wood et al., 2017). Sarringhaus et al. (2014) provides additional details and a rationale for how age groups were defined.

Rates of arboreal and terrestrial quadrupedal locomotion were calculated using observations of 53 chimpanzees, including 9 young infants, 11 old infants, 11 juveniles, 11 adolescents, and 11 adults. Each of these chimpanzees was sampled during five observation sessions, each of which was one hour long. During observation sessions, the positional behavior of focal individuals was recorded every two minutes during instantaneous scan samples. Chimpanzee positional behavior was classified according to body parts that bear the individual's weight, using categories defined in prior studies (Hunt et al., 1996; Thorpe & Crompton, 2006; Sarringhaus et al., 2014). Quadrupedal locomotion included both quadrupedal walking and running. The percentage of time members of each age class spent in quadrupedal locomotion was calculated for arboreal and terrestrial substrates. Rates of arboreal and terrestrial quadrupedal locomotion were normally distributed and analyzed using analysis of variance (ANOVA) with Tamhan T's posthoc comparisons as equal variance was not assumed (Sarringhaus et al. 2014).

The hand placements of chimpanzees during quadrupedal locomotion were recorded and included three categories: knuckle, palmigrade grasp, and palmigrade non-grasp¹. In knuckle, the intermediate phalanges on digits II-V contact the substrate. In palmigrade grasp, the palm and at least two digits are in contact with the substrate, and the digits are used to grasp the substrate. In palmigrade non-grasp, the palm contacts the substrate but the fingers do not grasp the substrate. Quadrupedal locomotion was also coded as occurring on terrestrial or arboreal substrates. The preceding data were collected over 340 hours observing chimpanzees with hand contact recorded in 789 of the 960 instances of quadrupedal locomotion (Sarringhaus et al., 2014). Because quadrupedal locomotion occurs infrequently in infants, especially young infants, 32 hours of locomotor video footage were also analyzed to determine hand usage during quadrupedal locomotion. Videos were recorded by LS on two

cameras, a Canon 2GL and a Canon XHAI HDV 3CCD, and analyzed frame-by-frame. For each locomotor bout, the hand postures of individuals were recorded from the second visible forearm step. Video data were recorded in the same manner as in focal observations, resulting in 563 samples of chimpanzees moving quadrupedally. There was no overlap in data derived from video bouts and focal observation sessions. Hand contact data were derived from 119 individuals, including 10 young infants, 11 old infants, 16 juveniles, 21 adolescents, and 61 adults.

Hand contact data during arboreal locomotion were analyzed using a negative binomial regression with data aggregated by individual ID (Hilbe, 2011; SPSS v.26). Terrestrial quadrupedal walking and usage was too homogenous to include in the model (see results). The dependent variable was the number of times individuals used their knuckles while moving. The predictor variable was age coded as a continuous variable in years. We included the natural log of the number of times individuals engaged in arboreal quadrupedal walking as an offset in the model.

Skeletal Methods

The MCs of 323 wild-caught individual primates from 7 museums were assessed (see Table 2). The 173 chimpanzee skeletal specimens were partitioned into age categories based on tooth eruption (see Table 1, Smith, Crummett, & Brandt, 1994; Smith et al., 2010; Smith & Boesch, 2011, Smith et al., 2013; Machanda et al., 2015). Specifically, we used maxilla tooth alveolar eruption to separate individuals into age categories of young infant (no first molar (M1) erupted), old infant (M1 but no second molar (M2) erupted), juvenile (M2 but no canine (C) erupted), and adolescent (C erupted but epiphyses not fused, Table 1). Adults had C and third molar (M3) erupted and all postcranial epiphyseal sutures fused. These tooth eruption patterns were translated into age estimates in years (see Table 1) so that the timing of the onset of skeletal features can be directly compared to the timing of the behavioral transitions we document among individuals of known age in our locomotor study.

The MCs of *Gorilla*, *Papio*, *Hylobates*, and *Pongo* were also analyzed to incorporate comparisons to species who moved in similar and different ways from chimpanzees (Table 2). The third MC was chosen for examination because the third ray has been reported to be used consistently during knuckle-walking by captive African apes while the use of the other digits is more variable (Inouye, 1994). Even when digits II-V are used during knuckle-walking, the third middle phalanx consistently bears the greatest weight in captive chimpanzees (Matarazzo, 2008; Wunderlich & Jungers, 2009), suggesting that MC III will display morphological changes related to the inception of knuckle-walking, if such changes exist.

The DMR was measured using ImageJ software from photographs of the bone taken in medial view. MCs were aligned (using clay) so that the ventral and dorsal anterior-posterior diaphyseal surface midpoints, just proximal to the metaphysis, were equidistant from the supporting surface. The distal and proximal surfaces were in anatomical position. Measurements were standardized using a 1 cm scale, positioned in the same plane as the bone, in each image. The presence and degree of the DMR was assessed using Inouye's MC torus measurement method (Inouye, 1994; Inouye & Shea, 2004), which involves measuring the ridge angle (formed by drawing lines between the three points labeled A, B, and C, Figure 1) and ridge height (the distance from point C perpendicular to the 180° line (Figure 1)). If the angle formed by ABC was approximately a straight line (180° or greater), a DMR was considered absent. If the ridge angle was less than 180°, a DMR was considered present. All angle measurements were capped at 180°; while this method detects a dorsal ridge, it is not ideal for assessing the extent of roundedness of the MC head, however, we are not assessing this condition in this study. The DMR angle was used for comparative analysis because it captures ridge prominence in specimens more precisely than simply recording presence or absence of the trait using the 180° threshold. Angle was used as it is a dimensionless unit and even the absence of a ridge produces a numeric value. Angle degrees were converted to radians ($1 \text{ rad} = 180/\pi$) for analysis.

Inouye has demonstrated that MC length does not scale with body size in *Gorilla* and *Pan* (1992 & 1994). Long bone midshaft diaphyseal measurements and joint dimensions are generally better at inferring body size than lengths (Burgess, McFarlin, Mudakikwa, Cranfield, & Ruff, 2018; Ruff, 2002). Therefore, we measured metacarpal midshaft anterior-posterior diameter from the medial-view photographs as a proxy for body size. We performed a linear regression for LN(ridge height) on LN(AP diameter) and found $R^2=0.225$ and $R^2=0.286$ for chimpanzees and gorillas respectively. As ridge height data were not normally distributed (at least one group test $p<0.05$), the regression p values cannot be used to assess significance of the relationship between ridge height and AP diameter. Previous work has shown that ridge height scales with body size (Inouye & Shea, 2004). We therefore use the natural log of the ratio of ridge height/AP diameter in analyses to account for body size differences when making comparisons among individuals in the same and different genera (see Jungers, Falsetti, & Wall, 1995 for the merits of using ratios).

Because the data were not normally distributed, we used nonparametric statistics in all of the analyses. We used Kruskal-Wallis tests to examine heterogeneity among multiple groups, and conducted pairwise posthoc comparisons with Bonferroni corrections to determine whether differences existed between specific groups (Sokal & Rohlf, 1995). In these analyses, individuals were categorized into the five age groups corresponding to dental eruption categories described above. We analyzed angle differences using all individuals. Our analyses of height differences included only those individuals who displayed a ridge. We used Mann-Whitney U tests to compare sex differences in the DMR heights and angles of adult chimpanzees and gorillas (Siegel & Castellen, 1988).

Results

Locomotor Development in Chimpanzees

As chimpanzees developed, they spent an increasing amount of time moving quadrupedally, and also shifted to proportionally more terrestrial as opposed to arboreal quadrupedalism (Figure 2). Young and old infants spent only a small percentage of their time moving quadrupedally (11% and 24% of locomotor behavior respectively, Figure 2), but the latter did so significantly more than the former (Mann-Whitney $p = 0.046$; see Sarringhaus et al., 2014). When moving quadrupedally, both young and old infants did so more often in the trees than the ground (Figure 2). In contrast, quadrupedalism was primarily a terrestrial activity for juveniles, adolescents, and adults (Figure 2). Individuals in different age groups did not vary in the amount of locomotor time spent moving quadrupedally in the trees (ANOVA $F = 1.510$, $p = 0.213$; Figure 2). The amount of locomotor time spent moving quadrupedally on the ground, however, displayed heterogeneity among the five different age classes (ANOVA $F = 35.887$, $p < 0.001$; Figure 2). Pairwise posthoc tests revealed that members of older groups engaged in more terrestrial quadrupedalism than individuals in younger groups ($p < 0.05$ for all comparisons), except old vs. young infants and adolescents vs. adults ($p = 0.926$ and 0.878 , respectively; Figure 2).

We examined the way chimpanzees used their hands during terrestrial and arboreal quadrupedalism. During 1,008 instances of terrestrial quadrupedal locomotion, chimpanzees of all ages engaged in knuckle-walking 100% of the time (Figure 3). By contrast, chimpanzee hand use during arboreal quadrupedalism varied as chimpanzees aged, with individuals employing three distinct types of hand positions: knuckle, palm grasp, and palm non-grasp (Figure 3). Individuals in both young and old infant groups employed all three of these hand postures but predominantly used palm grasp when moving arboreally (Figure 3). As chimpanzees grew older, their hand use during arboreal quadrupedalism changed. For every one-year increase in age, knuckle-usage increased by 33% (Exp(B) = 1.033, CI = 2.3 - 4.4%, $p < 0.001$). Including age as a predictor in a negative binomial regression analysis improved the fit of the model compared to the null model that excluded predictors ($X^2(1) = 8.135$, $p < 0.004$).

Dorsal Metacarpal Ridge

Dorsal Metacarpal Ridge Development in Chimpanzees

The DMR was present to variable degrees in chimpanzees of different ages. The DMR was absent in all young infants with no M1 but had developed (angle $< 180^\circ$) in 41% of the old infants, whose M1s had erupted (Table 2, Figure 4). The DMR was present in 93% of the juveniles, defined as individuals whose M2s had erupted, and in 98-100% of the adults and adolescents (Table 2, Figure 4). In individuals with a ridge present, LN (DMR height) and angle (radians) were inversely correlated; those with large DMR heights possessed small angles ($r = -0.837$, $p < 0.05$).

The DMR angle varied among chimpanzees in the different age classes (Kruskal-Wallis $H = 97.041$, $p < 0.001$; Figure 4A). Pairwise posthoc tests revealed that the DMR became more prominent as chimpanzees aged, i.e., the DMR angle decreased with age. Young and old infants possessed DMRs with relatively larger angles. They did not differ from each other ($p = 0.755$), but their DMR angles were much larger than those of members in all other age groups ($p < 0.001$ for all comparisons). The DMR angle continued to remain relatively large during juvenility but differed from the smaller angle in adults ($p < 0.001$). Finally, the DMR angle by adolescence converged on those found in adults, as the values did not differ significantly ($p = 1.00$, Figure 4A blue lines).

DMR height varied among chimpanzees of different ages (Kruskal-Wallis $H = 12.10$, $p = 0.007$; Figure 4B). While mean height increased as chimpanzees aged (Table 2 & Figure 4B), pairwise posthoc tests indicated that only old infants differed from adults in this regard ($p = 0.011$, all other comparisons $p > 0.05$). Ridge height cannot be analyzed for young infants because these individuals lacked ridges (Figure 4B).

Dorsal Metacarpal Ridge Comparisons Among Genera

While almost all adult *Pan* (98%; n=43/44) and *Gorilla* (97%; n=28/29) MCs possessed ridges, they were rare or absent in other taxa. None of the *Hylobates* specimens displayed ridges (n=41); however, ridges were present in a few *Pongo* individuals (n=3/16) and about half of the *Papio* (n=11/20) specimens. The presence of the ridge does not seem to be a function of size. The *Pongo* and *Papio* individuals with a ridge present were not those with the largest MC midshaft AP diameter. For example the *Pongo* individual with largest DMR had a MC diameter = 6.714mm, below the *Pongo* mean of 8.93mm, but not the smallest individual (range = 6.1-14.25 mm). Moreover, lack of a ridge in *Gorilla* and *Pan* did not occur in the smallest individuals (as measured by diameter). For example, the one *Gorilla* adult who lacked a DMR had an MC diameter = 11.8 mm, just below the *Gorilla* adult mean of 12.40 mm (range = 9.47-15.15 mm). The one *Pan* adult who lacked a ridge had a MC diameter = 8.95 mm, close to the adult mean of 9.03 mm (range = 7.3-11.2 mm).

The angle of the DMR varied among subadult and adult *Pan*, subadult and adult *Gorilla*, adult *Pongo*, adult *Hylobates*, and adult *Papio* (Kruskal-Wallis H 144.181, df = 6, p < 0.001, Figure 5A). Post hoc tests indicated that the DMR angles of adult *Pan* and *Gorilla* did not differ from one another, but members of both species had lower angles (i.e., more pronounced ridges) compared to all other groups (p<0.001, see Table 3). Similarly, the DMR angles of subadult *Pan* and *Gorilla* did not differ from one another nor did those between *Pongo*, *Hylobates*, and *Papio*. *Hylobates*, whose specimens displayed relatively large angles, was the only group that differed from the subadult *Gorilla* and subadult *Pan* (Table 3).

Relative DMR height also varied among genera (Kruskal-Wallis H = 32.725, p < 0.001, Figure 5B). Pairwise posthoc tests indicated that adult *Gorilla* differed from all other genera with the exception of adult *Pan* (Table 3). The DMR heights of adult *Pan*, subadult *Pan*, and subadult *Gorilla* did not differ from one another. The DMR heights of *Papio* specimens were significantly lower than those of adult and subadult *Pan* members but were not statistically different from the heights observed in subadult *Gorilla* (Table 3).

Dorsal Metacarpal Ridge Comparisons Between Sexes

The DMR angles of males and females did not differ in *Pan* or *Gorilla* (*Pan*: Mann-Whitney U z = -1.217, p = 0.240, n = 20 females and 7 males; *Gorilla*: Mann-Whitney U z = -1.20, p = 0.244, n = 11 females and 17 males). There was also no difference between the sexes with regard to DMR height in either genus (*Pan*: Mann-Whitney U z = -0.885, p = 0.400; *Gorilla*: Mann-Whitney U z = -1.341, p = 0.191). There were an insufficient number of *Papio* specimens for sex comparison analysis; however, ridges were present in approximately 60% of the male and female specimens whose sex was known (3/5 and 7/12, respectively).

Discussion

Documenting ontogenetic transitions in knuckle-walking requires taking two variables into consideration: 1) the amount of time individuals engage in terrestrial and arboreal quadrupedalism and 2) the breakdown of hand postures within those types of quadrupedalism. Sarringhaus et al. (2014) and Doran (1992) found that as chimpanzees aged, the amount of time they spent moving quadrupedally increased. However, neither study documented the ontogeny of hand usage with regard to substrate usage. This study analyzed arboreal and terrestrial quadrupedalism separately. We found that terrestrial quadrupedalism increased with increasing age, with significant increases in the transitions from old infancy to juvenility and from juvenility to adolescence. While moving quadrupedally on the ground, chimpanzees of all ages did so only using knuckle hand postures. In terms of frequency, arboreal quadrupedalism did not vary between age groups. However, chimpanzee hand postures did vary across development during arboreal quadrupedal locomotion, with knuckle-usage increasing with increasing age. Young infants, old infants, and juveniles primarily used palmigrade postures during arboreal quadrupedalism, while adolescents and adult chimpanzees typically knuckle-walked while arboreal. Notably, infants engaged in knuckle-walking only a small part of the time. The most pronounced shifts involving increased knuckle-usage took place at juvenility and at adolescence.

After clarifying how wild chimpanzees use their hands during locomotion and how both quadrupedal frequency and quadrupedal hand usage change with age, we proceeded to investigate the relationship between knuckle-walking and the prevalence and prominence of the DMR. We measured the DMR angle in addition to height to examine the sharpness of the incline of the ridge. Relatively larger ridges potentially provide greater resistance against dislocation of the proximal end of the phalanx during hyperextension at the metacarpophalangeal joint. We partitioned the subadult group into four different age

categories and found significant differences among groups of subadults. The DMR ridge is completely absent in young infants, and does not emerge until older infancy (M1 dental stage) in chimpanzees. It then reaches adult prominence in terms of DMR height during juvenility (M2 dental stage) when individuals begin to knuckle-walk frequently. A difference in relative height appears only between infants and adults. The mean DMR angle, however, does continue to increase after juvenility.

If knuckle-walking generates ridge formation (a component of hypotheses 1 and 2), the feature should *not* appear before the behavior is introduced into the locomotor repertoire, but instead after some delay. This is what occurs in chimpanzees, namely, there is some knuckle-walking during the young infant stage that precedes ridge development. The DMR does not appear until old infancy (between M1 and M2 emergence), *after* the onset of knuckle-walking. This finding fails to support a central tenet of the null hypothesis (H0), that the DMR is not at all behaviorally induced and will fail to track the onset of knuckle-walking in chimpanzees.

Although the prominence of the DMR angle continues to increase steadily over the course of development, the adult relative ridge height is established in juvenility when individuals begin to knuckle-walk as the predominant mode of locomotion. While the relationship between strain components and cortical bone modeling/remodeling is complex and poorly understood (Huiskes & van Reitbergen, 2005; Judex & Rubin, 2010), cortical bone plasticity, when present, does not typically display a linear relationship with loading frequency. Instead it responds to a minimal number of strains, after which increased loading cycles produce no additional bony response until the event of fracture (Umemura, Ishiko, Yamauchi, Kurono, & Mashiko, 1997; Rubin & Lanyon, 1984 & 1985). We are unaware of similar experiments investigating bone modeling/remodeling in the metaphyseal regions of the long bone shaft, and it is possible that distal metacarpal bone ridge formation may be dissimilar to patterns of appositional bone growth shown in prior studies. Articular surface shape and size have been shown to remain relatively constant over development, and are not strongly influenced by behavior (Lieberman, Devlin, Pearson, 2001; Ruff & Runestad, 1992; but see Plochocki, 2004). Pathological ridge formation has been documented on the dorsal surface of the distal hallux in humans (Bingold & Collins, 1950; Chou, 2000), but the pathological remodeling seen in such examples of osteoarthritis (Mueller & Tuan, 2011; Martel-Pelletier, Battista, Lajuenesse, & Pelletier, 1998) is likely not applicable to the modeling/remodeling that occurs in nonpathological dorsal metacarpal ridge formation in nonhuman primates. Regardless of the process of bone adaptation, our findings leave open the possibility that infant levels of terrestrial knuckle-walking may induce ridge formation on the dorsal MC.

Subadult *Gorilla* followed a similar pattern to subadult *Pan* in terms of DMR emergence with no individuals displaying ridge formation before M1 development and most individuals displaying ridges by M2 development. Our *Gorilla* sample is composed of *Gorilla gorilla* and *Gorilla beringei*, as the sample was small, and we elected to make comparisons at the level of genus. Adult *Gorilla* of both species are understood to be less arboreal compared to *Pan* (Doran, 1997; Doran & Hunt, 1994; Remis, 1998), but studies of locomotor development exist for only one population of gorillas, Virunga mountain gorillas (*Gorilla beringei beringei*; Doran, 1997). That study found that knuckle-walking predominates in the Virunga gorilla behavioral repertoire by one year of age (*ibid.*), much earlier than in *Pan*, for whom knuckle-walking gradually develops and does not become the predominant locomotor mode until juvenility. Given that no infant gorillas (i.e., no M1 present) measured in our study had a DMR (N=5) but about half of those at the later development stage (M1 emerged, but without M2 emerged) did have an incipient DMR (N= 8 of 17 total), it is possible that gorilla DMR development lags further behind their locomotor development, and is distinct from *Pan*. It is worth noting that while many developmental and life history milestones have been found to overlap between *Gorilla* and *Pan* (Robbins & Robbins, 2021; Smith et al., 1994; Stoinski, Perdue, Breuer, & Hoff, 2013), Virunga mountain gorillas may have faster life history trajectories and development compared to other gorillas, and even other populations of mountain gorillas (e.g. see body size discussion in Galbany et al., 2017; weaning in Robbins & Robbins, 2017; skeletal maturation in Taylor, 1997; but also see Kralick et al., 2017 for a review of dental development). Additional study of locomotor development in lowland and mountain gorillas, and a larger subadult morphological sample that is specific to different populations of *Gorilla* will be needed to clarify the factors contributing to these patterns of DMR emergence in *Gorilla*.

Previous research has suggested that because the ridge is not always found on the third MC of all individuals, it is not necessary for knuckle-walking performance (i.e. it is not a critical stabilizer at the MCP joint) but rather a plastic bony response to hyperextension at the MCP joint (Inouye & Shea, 2004). Our data indicate this feature was present in most (97-98%) but not *all* African ape individuals, as shown by Inouye & Shea (2004). Furthermore, we provide evidence of within-genus variability with a slight ridge occurring in a few *Pongo* specimens (N=3). However, the level of ridge height in these *Pongo* specimens is lower in relative and absolute terms compared to *Pan* old infants (Table 2 & Figures 4B & 5B). Both types of rare occurrences (absence in African apes, presence in *Pongo*) appear independent of body size. Our finding that there are no sex differences

present in DMR angle and height in sexually dimorphic *Gorilla* also supports the contention that this feature is not tightly driven by differences in body size. In addition, Inouye & Shea (2004) found that the presence or absence of the DMR did not correlate with body size in chimpanzees and gorillas.

Variability in locomotor performance is understudied in wild ape populations. For example, no information is currently available regarding inter-individual variability in the degree of hyperextension at the MC joint in great apes as they move on different substrates in the wild. However, there are some indications of variable hand usage in *Gorilla* and *Pongo*. Thompson et al. (2017) found that knuckle is the most common but not the only hand posture used during terrestrial quadrupedalism in mountain gorillas; several individuals displayed other hand postures frequently, including fist walking with metacarpal heads in direct contact with the substrate. This variation does not occur in chimpanzees as shown here. Nevertheless, there may be variation among chimpanzees (and other apes) in the digit that bears the most weight. It is also important to note that while *Pongo* are primarily arboreal (Manduell et al. 2012, Thorpe & Crompton, 2006), it is not uncharacteristic for them to be terrestrial in the wild (Mitani, pers. obs., Ancrenaz et. al., 2014; Ashbury et al. 2015; Loken, Spehar, Rayadin, 2013), and observations in captivity have documented individuals utilizing different terrestrial hand postures, including knuckle-walking (Susman, 1974; Tuttle, 1967). Such data suggest that variation in morphology does not preclude a behavioral influence on the formation of the DMR in great apes.

Despite the occasional absence of the DMR in knuckle-walking chimpanzees, our findings indicate the DMR is more prevalent and prominent in knuckle-walkers (*Gorilla* and *Pan*) than in suspensory taxa (*Pongo* and *Hylobates*). This is consistent with the hypothesis (H1) that the DMR reflects knuckle-walking. The results presented here also provide partial support for the hypothesis (H2) that the DMR reflects terrestrial vertical manus terrestrial postures with hyperextended MCP joints (including both knuckle-walking and digitigrady). Our analysis revealed that about a half of the baboons sampled in this study display a DMR. These animals move quadrupedally using their digits and occasionally their palms to contact the substrate (Napier & Napier, 1967; Patel, 2010; Patel & Wunderlich, 2010; Zeininger et. al., 2017). Further examination of *Papio* DMR formation in baboons during development and in relation to ontogenetic changes in locomotion (as performed in this study for *Pan*), will be needed to determine why ridges fail to appear about half the time in this species. The predictions of the null hypothesis (H0), that the DMR will fail to track knuckle-walking and does not vary among genera according to hand usage, were not upheld by our findings.

While the DMR appears in knuckle-walking and digitigrade individuals, it is possible to distinguish the two, as the DMR in African ape knuckle-walkers is more prominent than that in digitigrade *Papio*. The former display more acute angles and larger relative heights (LN height/MC diameter) than do the latter. This observation leads us to provisionally suggest that a ridge can be diagnostic of knuckle-walking if it exceeds a threshold ratio of 0.13 (DMR height/MC midshaft AP diameter; Figures 4 & 5). We also found a qualitative difference in ridge morphology between knuckle-walking African apes and digitigrade *Papio*. The DMR in *Pan* and *Gorilla* spans the dorsal surface of the third MC, while the *Papio* DMR is typically composed of two ridges, one angled medially and one laterally, that do not connect on the dorsal midline (see Figure 6). These preliminary qualitative observations call for further examination of the dorsal MC morphology of habitually terrestrial digitigrade primates.

In sum, results of this study provide the first quantitative analysis of hand contact and substrate use during quadrupedal locomotion in wild chimpanzees of different ages. These data are required to understand the biomechanical signals experienced by the hands and wrists of these animals. Increases in chimpanzee knuckle-walking mirror increases in quadrupedal locomotion during different periods of development. Changes in the DMR angle and height correspond to developmentally mediated shifts in the frequency of knuckle-walking behavior. The DMR appears to be a terrestrial quadrupedal walking feature that is most prevalent and prominent in adult chimpanzees and gorillas. It is less prevalent and prominent in baboons. By contrast, it occurs rarely and is not very prominent in orangutans, and it is entirely absent in gibbons. Provisionally, we propose this feature is diagnostic of knuckle-walking when it exceeds a DMR height/MC diameter threshold of ~ 0.13 (see Figure 5). Because the timing of the development of the DMR occurs after the initial transition to knuckle-walking, the loads produced by knuckle-walking likely influence the development of this trait. This study does not resolve whether the DMR is a functionally salient trait that acts as a buttress to prevent hyperextension rather than a byproduct of knuckle-walking that reflects dorsal loading but does not contribute to joint stability. To date, only a handful of epigenetic features have been identified and used to reconstruct the locomotor behavior of fossil hominoids (Ward, 2002). These include the femoral bicondylar angle (e.g. Tardieu & Trinkaus, 1994; Tardieu, 1999) and lumbar lordosis (e.g. Nakatsukasa, 2004; Nakatsukasa & Hayama, 1996;). Results from this study furnish provisional support for one additional plastic feature indicative of a particular locomotor mode during the course of ontogeny, namely that the DMR reflects a vertical manus terrestrial quadrupedal posture. However, the quantitatively lower expression and qualitative differences in appearance between the DMR in adult African apes and *Papio* augur well for being able

to differentiate between a DMR that reflects knuckle-walking vs. digitigrady in a fossil taxon. Pending the discovery of relevant hominoid fossils, the results of such analyses could help resolve whether a terrestrial quadrupedal phase preceded human bipedal evolution.

References

- Ancrenaz, M. et al. (2014) Coming down from the trees: Is terrestrial activity in Bornean orangutans natural or disturbance driven?. *Scientific Reports*, 4, 4024; DOI:10.1038/srep04024 (2014).
- Ashbury, A. M., Posa, M. R. C., Dunkel, L. P., Spillmann, B., Atmoko S. S. U., van Schaik C. P., & van Noordwijk, M. A. (2015). Why do orangutans leave the trees? Terrestrial behavior among wild Bornean orangutans (*Pongo pygmaeus wurmbii*) at Tuanan, Central Kalimantan. *American Journal of Physical Anthropology* 77:1216–122 doi:10.1002/ajp.22460.
- Barak, M. M., Sherratt, E., & Lieberman, D. E. (2017). Using principal trabecular orientation to differentiate joint loading orientation in the 3rd metacarpal heads of humans and chimpanzees. *Journal of Human Evolution*, 113, 73- 182. <http://dx.doi.org/10.1016/j.jhevol.2017.08.018>
- Begun, D. R. (1992). Miocene fossil hominids and the chimp-human clade. *Science* 257, 1929-1933.
- Bingold, A. C. and Collins, D. H. 1950. Hallux Rigidus. *Journal of Bone and Joint Surgery* 32, 214-222 DOI: [10.1302/0301-620X.32B2.214](https://doi.org/10.1302/0301-620X.32B2.214)
- Böhme, M., Spassov, N., Fuss, J., Tröscher, A., Deane, A. S., Prieto, J., Kirscher, U., Lechner, T., & Begun, D. R. (2019). A new Miocene ape and locomotion in the ancestor of great apes and humans. *Nature* <https://doi.org/10.1038/s41586-019-1731-0>
- Burgess, L. M., McFarlin, S. C., Mudakikwa, A., Cranfield, M. R., Ruff, C. B. (2018). Body mass estimation in hominoids: Age and locomotor. *Journal of Human Evolution*, 115, 36-46. <http://dx.doi.org/10.1016/j.jhevol.2017.07.004>
- Chirchir, H., Zeininger, A., Nakatsukasa, M., Ketchman, R. A., & Richmond, B. G. (2017). Does trabecular bone structure within the metacarpal heads of primates vary with hand posture? *Comptes Rendus Palevol* 16, 533–544. <http://dx.doi.org/10.1016/j.crpv.2016.10.002>
- Chou L. B. 2000. Disorders of the first metatarsophalangeal joint: diagnosis of great-toe pain. *Phys Sportsmed* 28, 32-45. doi: 10.3810/psm.2000.07.1075.
- Corruccini, R. S. (1978). Comparative osteometrics of the hominoid wrist joint, with special reference to knuckle-walking. *Journal of Human Evolution*, 7, 307–321.
- Crompton, R. H., Sellers, W. I., & Thorpe, S. K. S. (2010). Arboreality, terrestriality and bipedalism. *Philosophical Transactions of the Royal Society B*, 365, 3301–3314. doi: 10.1098/rstb.2010.0035
- Crompton, R. H., Vereecke, E. E., & Thorpe, S. K. S. (2008). Locomotion and posture from the common hominoid ancestor to fully modern hominins, with special reference to the last common panin/hominin ancestor. *Journal of Anatomy*, 212, 501–543. doi: 10.1111doi: 10.1111/j.1469-7580.2008.00870.x
- Dainton, M., Macho, G. A. (1999). Did knuckle-walking evolve twice? *Journal of Human Evolution*, 36, 171-194.
- Doran, D. M. (1992). The ontogeny of chimpanzee and pygmy chimpanzee locomotor behavior: a case study of paedomorphism and its behavioral correlates. *Journal of Human Evolution*, 23, 139-157.
- Doran, D. M. (1997). Ontogeny of locomotion in mountain gorillas and chimpanzees. *Journal of Human Evolution*, 32, 323-344.

- Doran, D. M., Hunt, K. D. (1994). Comparative locomotor behavior of chimpanzees and bonobos: Species and Habitat Differences. In: R.W. Wrangham, B. C. McGrew, F. B. M. De Waal, P. G. Heltne (Eds.). *Chimpanzee Cultures* (pp. 93 - 108). Cambridge: Harvard University Press.
- Dunmore, C. J., Kivell, T. L., Bardo, A., & Skinner, M. M. (2019). Metacarpal trabecular bone varies with distinct hand-positions used in hominid locomotion. *Journal of Anatomy*, 235, 45-66. Doi: 10.1111/joa.12966
- Fleagle, J. G., Stern, J. T., Jungers, W. L., Susman, R. L., Vangor, A. K., & Wells, J. P. (1981). Climbing: a biomechanical link with brachiation and bipedalism. *Symposia of Zoological Society of London*, 48, 359-373.
- Galbany J., Abavandimwe D., Vakiener M., Eckardt, W., Mudakikwa, A., Ndagijimana, F., Stoinski, T., McFarlin, S. C. (2017). Body growth and life history in wild mountain gorillas (*Gorilla beringei beringei*) from Volcanoes National Park, Rwanda. *American Journal of Physical Anthropology*, 163, 570 - 590. <https://doi.org/10.1002/ajpa.23232>
- Gregory, W. K. (1916). Studies on the evolution of primates. Part II. Phylogeny of recent and extinct anthropoids, with special reference to the origin of man. *Bulletin of the American Museum of Natural History*, 35, 258-355.
- Haile-Selassie, Y. (2001). Late Miocene hominids from the Middle Awash, Ethiopia. *Nature* 412, 178-191.
- Hilbe, J. M. (2011). *Negative binomial regression*. Cambridge University Press.
- Huiskes, R., & van Reitbergen, B. (2005). Mechanics of Bone. In: V. Mow & R Huiskes (Eds.), *Basic Orthopedic Biomechanics and Mechano-Biology*. Third Edition. (pp. 123-177). Philadelphia, PA: Lippincott Williams and Wilkins.
- Hunt, K. D., Cant, J. G. H., Gebo, D. L., Rose, M. D., Walker, S. E., & Youlatos, D. (1996). Standardized descriptions of primate locomotor and postural modes. *Primates*, 37, 363-387.
- Inouye, S.E. (1992). Ontogeny and allometry of African ape manual rays. *Journal of Human Evolution*, 23, 107-138.
- Inouye, S. E. (1994). Ontogeny of knuckle-walking hand postures in African apes. *Journal of Human Evolution*, 26, 459-485.
- Inouye, S. E., & Shea, B. T. (2004). The implications of variation in knuckle-walking features for models of African hominoid locomotor evolution. In: E. Bruner (Ed.) *Journal of Anthropological Sciences*, Torino, 67-88.
- Jenkins, F. A. J., & Fleagle, J. G. (1975). Knuckle-walking and the functional anatomy of the wrist. In: R.H. Tuttle, *Primate Functional Morphology and Evolution* (pp. 213-231). The Hague: De Gruyter.
- Judex, S., & Rubin, C. (2010). Mechanical Influences on Bone Mass and Morphology. In: R.A. Adler (Ed.). *Contemporary Endocrinology: Osteoporosis: Pathophysiology and Clinical Management*. New York: Humana Press. DOI 10.1007/978-1-59745-459-9_8,
- Jungers, W.L., Falsetti, A.B., & Wall, C.E. (1995). Shape, relative size, and size-adjustments in morphometrics. *Yearbook of Physical Anthropology*, 38, 137-161.
- Keith, A. (1923). Hunterian Lectures On Man's Posture: Its Evolution And Disorders. Given At The Royal College Of Surgeons Of England. *British Medical Journal*, 1, 624-626.
- Kimura, T. (1986). Kinesiology of quadrupedal locomotion in infant chimpanzees. *Baiomekanizumu*, 8, 77-85.
- Kimura, T. (2019). How did humans acquire erect bipedal walking? *Anthropological Science*, 127, 1-12. DOI: 10.1537/ase.190219
- Kivell, T. L. (2016). The Primate Wrist. In: T. L. Kivell, P. Lemelin, B. G. Richmon, D. Schmitt (Eds.). *Evolution of the Primate Hand: Anatomical, Developmental, Functional, and Paleontological Evidence* (pp. 17-54). New York: NY Springer Science+Business Media. DOI 10.1007/978-1-4939-3646-5_1

Kivell, T. L., & Begun, D. R. (2007). Frequency and timing of scaphoid-centrale fusion in hominoids. *Journal of Human Evolution* 52, 321 - 340. doi:10.1016/j.jhevol.2006.10.002

Kivell, T. L., & Schmitt, D. (2009). Independent evolution of knuckle-walking in African apes shows that humans did not evolve from a knuckle-walking ancestor. *Proceedings of the National Academy of Science* 106, 14241-14246. doi:10.1073/pnas.0901280106.

Kralick, A. E., Burgess M. L., Glowacka, H., Arbenz-Smith, K., McGrath, K., Ruff, C. B., Chan, K. C., Cranfield, M. R., Stoinski, T. S., Bromage, T. G., Mudakikwa, A., McFarlin, S. C. (2017). A radiographic study of permanent molar development in wild Virunga mountain gorillas of known chronological age from Rwanda. *American Journal of Physical Anthropology*, 163, 129–147. <https://doi.org/10.1002/ajpa.23192>

Lewis, O. J. (1977). Joint remodeling and the evolution of the human hand. *Journal of Anatomy* 123, 157-201.

Lieberman, D. E., Pearson, O. M., Polk, J. D., Demes, B., & Crompton, A. W. (2003). Optimization of bone growth and remodeling in response to loading in tapered mammalian limbs. *Journal of Experimental Biology*, 206, 3125–3138. doi:10.1242/jeb.00514

Loken, B., Spehar, S., Rayadin, Y. (2013). Terrestriality in the Bornean Orangutan (*Pongo pygmaeus morio*) and Implications for Their Ecology and Conservation. *American Journal of Physical Anthropology* 75, 1129-1138.

Lovejoy, C. O., Suwa, G., Simpson, S. W., Matternes, J. H., & White, T. D. (2009a). The great divides: *Ardipithecus ramidus* reveals the postcrania of our last common ancestors with African apes. *Science*, 326, 100 – 106.

Lovejoy, O. C., Suwa, G., Spurlock, L., Asfaw, B., & White, T.D. (2009b). The pelvis and femur of *Ardipithecus ramidus*: the emergence of upright walking. *Science*, 326, 71. DOI: 10.1126/science.1175831

Lovejoy, C.O., & McCollum, M.A. (2010). Spinopelvic pathways to bipedality: why no hominids ever relied on a bent-hip–bent-knee gait. *Philosophical Transactions of the Royal Society B*, 365, 3289–3299. doi:10.1098/rstb.2010.0112

Machanda, Z., Brazeau, N. F., Bernard, A. B., Donovan, R. M., Papakyrikos, A. M., Wrangham, R., & Smith, T. M. (2015). Dental eruption in East African wild chimpanzees. *Journal of Human Evolution* 82, 137-144. <http://dx.doi.org/10.1016/j.jhevol.2015.02.010>

MacLatchy, L., DeSilva, J., Sanders, W., & Wood, B. (2010). The Hominins. In: L. Werdelin, W. J. Sanders (Eds.), *Cenozoic Mammals of Africa* (pp. 471-540). Berkeley: University of California Press.

Manduell, K. L., Harrison, M. E., & Thorpe, S. K. (2012). Forest structure and support availability influence orangutan locomotion in Sumatra and Borneo. *American Journal of Primatology*, 74, 1128-1142. doi: 10.1002/ajp.22072.

Martel-Pelletier, J., Di Battista, J. A., Lajeunesse, D & Pelletier, J.P. (1998). IGF/IGFBP axis in cartilage and bone in osteoarthritis pathogenesis. *Inflammation Research* 47, 90-100. doi:1023-3830/98/030090-11

Matarazzo, S. (2008). Knuckle walking signal in the manual digits of *Pan* and *Gorilla*. *American Journal of Physical Anthropology*, 135, 27–33. DOI 10.1002/ajpa.20701

Matarazzo, S. A. (2015) Trabecular architecture of the manual elements reflects locomotor patterns in primates. *PLoS ONE*, 10, e0120436. doi:10.1371/journal.pone.0120436

Morton, D. J. (1922). Evolution of the human foot. *American Journal of Physical Anthropology*, 4, 305-366

Morton, D. J. (1924). Evolution of the human foot II. *American Journal of Physical Anthropology*, 7, 1-47.

Mueller M. B., Tuan, R. S. (2011). Anabolic/Catabolic Balance in Pathogenesis of Osteoarthritis: Identifying Molecular Targets. *RM&R* 3, S3-S11. doi.org/10.1016/j.pmrj.2011.05.009

Nakatsukasa, M. (2004). Acquisition of bipedalism: the Miocene hominoid record and modern analogues for bipedal protohominids. *Journal of Anatomy*, 204, 385-402.

Nakatsukasa, M., & Hayama, S. (1996). Development of bipedal standing posture in trained Suo monkeys. *Primate Research*, 12, 263.

Napier, J. R., & Napier, R. H. (1967). A handbook of living primates. London: Academic Press.

Orr, C. M. (2005). Knuckle-walking anteater: A convergence test of adaptation for purported knuckle-walking features of African Hominidae. *American Journal of Physical Anthropology*, 128, 639–658. DOI 10.1002/ajpa.20192

Orr, C. M. (2018). Kinematics of the anthropoid os centrale and the functional consequences of scaphoid-centrale fusion in African apes and hominins. *Journal of Human Evolution*, 114, 102 - 117. <https://doi.org/10.1016/j.jhevol.2017.10.002>

Patel, B. A., & Maiolino, S. A. (2016). Morphological diversity in the digital rays of primate hands. In: T. L. Kivell, P. Lemelin, B. G. Richmon, D. Schmitt (Eds.). *Evolution of the Primate Hand: Anatomical, Developmental, Functional, and Paleontological Evidence* (pp. 55-100). New York: NY Springer Science+Business Media. DOI 10.1007/978-1-4939-3646-5_1

Patel, B. A. (2010). Functional morphology of cercopithecoid primate metacarpals. *Journal of Human Evolution*, 58, 320-337. doi:10.1016/j.jhevol.2010.01.001

Patel, B. A., & Wunderlich, R. E. (2010). Dynamic pressure patterns in the hands of olive baboons (*Papio anubis*) during terrestrial locomotion: implications for cercopithecoid primate hand morphology. *Anatomical Record*, 293, 710 – 718. DOI 10.1002/ar.21128

Pilbeam, D. R., & Lieberman, D. E. (2017). Reconstructing the last common ancestor of chimpanzees and humans. In: M. Muller (Ed.). *Chimpanzees and Human Evolution* (pp 22-98). Boston: Harvard University Press.

Pontzer H, Lieberman DE, Momin E, Devlin MJ, Polk JD, Hallgrímsson B, Cooper DML. 2006. Trabecular bone in the bird knee responds with high sensitivity to changes in load orientation. *J. Exp. Biol.* 209: 57-65.

Preuschoft, H. (1973). Functional anatomy of the upper extremity. In: G. H. Bourne (Ed.) *The Chimpanzee*, 6, 34-120. Basel: Karger.

Prost, J.H. (1980). Origin of bipedalism. *American Journal of Physical Anthropology* 52:175-189.

Remis, M. J. (1998). The gorilla paradox. In: E. Strasser, J. Fleagle, A. Rosenberger, & H. M. McHegry (Eds.). *Primate Locomotion* (pp. 95-106). New York: Plenum Press.

Richmond, B. G., Begun, D. R., & Strait, D. S. (2001). Origin of human bipedalism: the knuckle-walking hypothesis revisited. *Yearbook of Physical Anthropology*, 44, 70-105.

Richmond, B. G., & Strait, D. S. (2000). Evidence that humans evolved from a knuckle-walking ancestor. *Nature*, 404, 382-385.

Robbins, M. M., Robbins, A. M. (2021). Variability of weaning age in mountain gorillas (*Gorilla beringei beringei*). *American Journal of Physical Anthropology*, 174, 776-174. <https://doi.org/10.1002/ajpa.24237>

Rose, M.D. (1991). The process of bipedalization in hominids. In: Y. Coppens, & B. Senut (Eds.). *Origine(s) de la Bipédie Chez les Hominidés*, pp. 37–48. Paris: Editions du CNRS.

Rubin, C. T. & Lanyon, L. E. (1984). Regulation of bone formation by applied dynamic loads. *Journal of Bone and Joint Surgery* 66A, 397-402.

Rubin, C. T. & Lanyon, L. E. (1985). Regulation of Bone Mass by Mechanical Strain Magnitude. *Calcified Tissue International*, 37, 411- 417.

- Ruff, C. B. (2002). Long bone articular and diaphyseal structure in old world monkeys and apes. I: Locomotor effects. *American Journal of Physical Anthropology*, 119, 305-342.
- Sarmiento, E. E. (1994). Terrestrial traits in the hands and feet of gorillas. *American Museum Novitates*, 309, 1-56.
- Sarringhaus, L. (2013). Changes in long bone strength correspond to shifts in locomotor behavior during development in chimpanzees (*Pan troglodytes*). *American Journal of Physical Anthropology*, 150 S5, 241.
- Sarringhaus, L., MacLatchy L. M., & Mitani, J. C. (2014). The locomotor and postural development of wild chimpanzees. *Journal of Human Evolution*, 66, 29 – 38. <http://dx.doi.org/10.1016/j.jhevol.2013.09.006>
- Schultz, A. (1930). The skeleton of the trunk and limbs of higher primates. *Human Biology*, 2, 303-438.
- Siegel, S., Castellen, N.J. (1988). Nonparametric Statistics for the Behavioral Sciences, Second edition. New York: McGraw-Hill.
- Simpson, S. W., Latimer, B., & Lovejoy, C. O. (2018). Why do knuckle-walking African apes knuckle-walk? *The Anatomical Record*, 301, 496–514. DOI 10.1002/ar.23743
- Smith, B. H., Crummett, T.L., & Brandt, K. L. (1994). Ages of eruption on primate teeth: a compendium for aging individuals and comparing life histories. *Yearbook of Physical Anthropology*, 37, 177–231.
- Smith, B. H., & Boesch, C. (2011). Mortality and the magnitude of the “wild effect” in chimpanzee tooth emergence. *Journal of Human Evolution*, 60, 34–46. doi:10.1016/j.jhevol.2010.08.006
- Smith, T. M., Smith, B. H., Reid, D. J., Siedel, H., Vigilant, L., Hublin, J. J., & Boesch, C. (2010). Dental development of the Tai Forest chimpanzees revisited. *Journal of Human Evolution*, 58, 363-373. doi:10.1016/j.jhevol.2010.02.008
- Smith, T. M., Machanda, Z., Bernard, A. B., Donovan, R. M., Papakyrikos, A. M., Muller, M. N., & Wrangham, R. (2013). First molar eruption, weaning, and life history in living wild chimpanzees. *Proceedings of the National Academy of Science*, 110, 2787-2791.
- Sokal, R.R., & Rohlf, F.J. (1995). *Biometry*. New York: Freeman and Company.
- Stern, J. T. (1975) Before bipedality. *Yearbook of Physical Anthropology*, 19, 59–68.
- Stern, J.T. & Susman, R. (1981). Electromyography of the Gluteal Muscles in *Hylobates*, *Pongo*, and *Pan*: Implications for the Evolution of Hominid Bipedality. *American Journal of Physical Anthropology*, 55, 153-166.
- Stoinski, T. S., Perdue, B., Breuer, T., & Hoff, M. P. (2013). Variability in the developmental life history of the genus *Gorilla*. *American Journal of Physical Anthropology* 152, 165 - 172. DOI: 10.1002/ajpa.22301
- Straus, W. L. (1949). The riddle of man's ancestry. *Q. Rev. Biol*, 24, 200–223.
- Susman R.L. (1974). Facultative terrestrial hand postures in an orangutan (*Pongo pygmaeus*) and pongid evolution. *American Journal of Physical Anthropology*, 40, 27-37.
- Susman, R.L. (1979). Comparative and functional morphology of hominoid fingers. *American Journal of Physical Anthropology*, 50, 215-236.
- Tardieu, C. (1999). Ontogeny and phylogeny of femoro-tibial characters in humans and hominid fossils: Functional influence and genetic determinism. *American Journal of Physical Anthropology*, 110, 365-377.
- Tardieu, C., & Trinkaus, E. (1994). Early ontogeny of the human femoral bicondylar angle. *American Journal of Physical Anthropology*, 95, 183-195.
- Taylor, A. B. (1997). Relative Growth, Ontogeny, and Sexual Dimorphism in Gorilla (*Gorilla gorilla gorilla* and *G. g. beringei*): Evolutionary and Ecological Considerations. *American Journal of Primatology* 43, 1 - 31.

- Thompson, N. E., Ostrofsky, K. R., McFarlin, S. C., Robbins, M. M., Stoinski, T. S., & Alméjida, S. (2017). Unexpected terrestrial hand posture diversity in wild mountain gorillas. *American Journal of Physical Anthropology*, 166, 84-94. DOI: 10.1002/ajpa.23404
- Thompson, N.E. (2020). The biomechanics of knuckle-walking: 3-D kinematics of the chimpanzee and macaque wrist, hand and fingers. *Journal of Experimental Biology*, 223. doi:10.1242/jeb.224360
- Thorpe, S. K., & Crompton, R. H. (2006). Orangutan positional behavior and the nature of arboreal locomotion in Hominoidea. *American Journal of Physical Anthropology*, 131, 384-401.
- Thorpe, S. K. S., Holder, R. L., & Crompton, R.H. (2007). Origin of human bipedalism as an adaptation for locomotion on flexible branches. *Science*, 316, 1328-1331.
- Thorpe, S. K. S., McClymount, J. M., & Crompton, R.H. (2014). The arboreal origins of human bipedalism. *Antiquity*, 88, 906–926.
- Turner, C. H., Takano, Y., Owan, I. (1995). Aging changes mechanical loading thresholds for bone formation in rats. *Journal for Bone and Mineral Research*, 10, 1544-1549.
- Tuttle, R.H. (1967). Knuckle-walking and the evolution of hominoid hands. *American Journal of Physical Anthropology*, 26, 171-206.
- Tuttle, R., Basmajian, J. V. (1974). Electromyography of brachial muscles in *Pan gorilla* and hominoid evolution. *American Journal of Physical Anthropology*, 41, 71-90.
- Tsegai, Z. J., Kivell, T. L., Gross, T., Nguyen, N. H., Pahr, D. H., Smaers, J. B., & Skinner, M. M. (2013). Trabecular Bone Structure Correlates with Hand Posture and Use in Hominoids. *PLoS ONE*, 8(11), 78781. <https://doi.org/10.1371/journal.pone.0078781>
- Umemura, Y., Ishiko, T., Yamauchi, T., Kurono, M., Mashiko, S. (1997). Five jumps per day increase bone mass and breaking force in rats. *Journal of Bone and Mineral Research* 12, 1480-1485. doi:10.1359/jbmr.1997.12.9.1480
- Ward, C.V. (2002). Interpreting the Posture and Locomotion of *Australopithecus afarensis*: Where Do We Stand? *Yearbook of Physical Anthropology*, 45, 185 – 215.
- Washburn, S.L. (1967). Behaviour and the origin of man. *Proceedings of the Royal Anthropological Institute of Great Britain and Ireland*, 1967, 21–27.
- White, T. D., Lovejoy, C. O., Asfaw, B., Carlson, J. P., & Suwa, G. (2015). Neither chimpanzee nor human, *Ardipithecus* reveals the surprising ancestry of both. *Proceedings of the National Academy of Sciences*, 112, 4877–4884. doi: 10.1073/pnas.1403659111
- Williams, S.A. (2010). Morphological integration and the evolution of knuckle-walking. *Journal of Human Evolution*, 58, 432-440. doi:10.1016/j.jhevol.2010.03.005
- Wood, B. M., Watts, D. P., Mitani, J. C., Langergraber, K. E. (2017). Favorable ecological circumstances promote life expectancy in chimpanzees similar to that of human hunter-gatherers. *Journal of Human Evolution*, 105, 41-56. <https://doi.org/10.1016/j.jhevol.2017.01.003>
- Wood Jones, F. (1929). *Man's place among the mammals*. New York: Longman Green.
- Wunderlich, R. E., & Jungers, W. L. (2009). Manual digital pressures during knuckle-walking in chimpanzees (*Pan troglodytes*). *American Journal of Physical Anthropology*, 139, 394-403. DOI 10.1002/ajpa.20994

Wuthrich, C., MacLatchy, L. M, Nengo, I. O. (2019). Wrist morphology reveals substantial locomotor diversity among early catarrhines: an analysis of capitates from the early Miocene of Tinderet (Kenya). *Scientific Reports*, 9,3728. <https://doi.org/10.1038/s41598-019-39800-3>

Zeininger, A., Richmond, B. G., & Hartman, G. (2011). Metacarpal head biomechanics: A comparative backscattered electron image analysis of trabecular bone mineral density in *Pan troglodytes*, *Pongo pygmaeus*, and *Homo sapiens*. *Journal of Human Evolution*, 60, 703-710.

Zeininger, A., Shapiro, L. J., & Raichlen, D. A. (2017). Ontogenetic changes in limb postures and their impact on effective limb length in baboons (*Papio cynocephalus*). *American Journal of Physical Anthropology*, 163, 231–241. <https://doi.org/10.1002/ajpa.23201>

Zihlman, A. L., Bolter, D. R., Boesch, C. (2007). Skeletal and dental growth and development in chimpanzees of the Tai National Park, Cote D'Ivoire. *Journal Zoology*, 273, 63-73.

Zollikofer, C. P. E., Ponce de Leon, M., Lieberman, D. E, Guy, F., Pilbeam, D., Likius, A., Mackaye, H.T., Vignaud, P., & Brunet, M. (2005). Virtual cranial reconstruction of *Sahelanthropus tchadensis*. *Nature*, 434, 755 -759.

Acknowledgements

The authors thank Jacinta Beehner, Maureen Devlin, and B. Holly Smith for providing useful feedback on earlier versions of the paper. Thanks are also due to Daniel Schmitt, Angel Zeininger, and Gabe Yapuncich. for helpful comments and discussions. The authors also thank Eileen Westwig and Darrel Frost (AMNH), Yohannes Haile-Selassie and Lyman Jellema (CMNH), Judy Chupasko (HMCZ), Paula Jenkins (LMNH), Marcia Ponce de León (UZ), Stefan Merker (SMNS) and Angela Gill and Malcolm Marman (QuexM). Field research was sponsored by the Uganda Wildlife Authority, Uganda National Council for Science and Technology, and the Makerere University Biological Field Station. This research was funded by grants from the L.S.B. Leakey Foundation, the National Science Foundation (IOB-0516644 and BCS-0850951), and the University of Michigan International Institute.

Conflict of Interest

The authors declare no competing interests.

Data

Data that supports these findings are available upon request from authors.

Figure Legends

Figure 1. Measurement of the dorsal metacarpal ridge. Angle $\angle ABC$ (measured in radians) is used to quantify the presence of the DMR. DMR height is measured as the length of the line perpendicular to the 180 degree line connecting points A and B, that intersects with point C. Line D represents the midshaft diameter.

Figure 2. Quadrupedal locomotion by chimpanzees in different age classes. The proportion of time spent moving quadrupedally arboreally (orange) and terrestrially (blue) is shown relative to all locomotor behavior.



Figure 3. Quadrupedal locomotion using different types of hand contact. Terrestrial and arboreal locomotion are plotted separately. Red line = Palm grasp, Green = Palm non-grasp, Blue = Knuckle. Results show combined video and frequency data. Image A is an example of palmigrade grasp hand posture, image B is an example of knuckle hand posture.



Figure 4A. Distal metacarpal ridge angles (in radians) in chimpanzees of different age classes. Means and 95% confidence intervals are shown. The dashed line at 3.14 radians indicates that the epiphyseal area is straight or rounded with no distal metacarpal ridge present. Below 3.14 indicates a distal metacarpal ridge is present. B. Chimpanzee distal metacarpal ridge height/MC diameter in chimpanzees of different age classes. Means and 95% confidence intervals are shown. Blue data are all

individuals. Red data are only individuals with the feature present. Blue data young infant N = 17, old infant N = 51, juvenile N = 30, adolescent N = 25, adult N = 45; red data old infant N = 21, juvenile N = 28, adolescent N = 25, adult N = 44.


Figure 5A. Distal metacarpal ridge angles (in radians) in *Pan*, *Gorilla*, *Pongo*, *Hylobates*, and *Papio*. Means and 95% confidence intervals are shown. The dashed line at 3.14 radians indicates that the epiphyseal area is straight or rounded with no distal metacarpal ridge present. Below 3.14 indicates a distal metacarpal ridge is present. B. Distal metacarpal ridge height/MC diameter in *Pan*, *Gorilla*, *Pongo*, *Hylobates*, and *Papio*. Means and 95% confidence intervals are shown. Blue data are all individuals. Red data are only individuals with the feature present. Blue data *Pan* subadult N = 125, *Pan* adult N = 45, *Gorilla* subadult N = 43, *Gorilla* adult N = 29, *Pongo* adult N = 16, *Hylobates* adult N = 41, *Papio* adult N = 20. Red data *Pan* subadult N = 75, *Pan* adult N = 44, *Gorilla* subadult N = 28, *Gorilla* adult N = 28, *Pongo* adult N = 3, *Papio* adult N = 11.

Figure 6. The DMR of *Papio*, *Pan*, and *Gorilla* in medial (top) and dorsal (bottom) view. When present in *Papio*, the DMR is two separate ridges angled to the lateral and medial side (blue arrows). The DMR of African apes (*Pan* and *Gorilla*) presents as a continuous ridge from the medial to lateral edge of the distal dorsal metacarpal (blue arrows).






A



B






Figure 7.png




Table 1. Morphological and behavioral markers of chimpanzee age categories.

Category	Age ^a	Locomotor Behavior ^b	Dental eruption markers ^c
Young Infant	0 - 3	Predominantly suspensory with the highest levels of suspensory behavior for any age group.	deciduous dentition M1 not erupted
Old Infant	3.1 - 5	Still predominantly suspensory but higher levels of independence from mother and higher levels of quadrupedal locomotion compared to young infants.	M1 erupts I1 and M2 not erupted
Juvenile	5-10	Completely independent from mother, a substantial decrease in suspensory locomotion and increase in quadrupedal walking and running compared to older infants.	I1 & M2 erupted, C frequently erupts at this time M3 not erupted
Adolescent	10--15	A substantial decrease in suspensory locomotion and vertical climb and increase in quadrupedal walking compared to juveniles.	C fully erupted M3 erupts at this time postcrania not all fused
Adult	~15+	No significant change in locomotion compared to adolescents.	Some wear on teeth, postcrania all fused ^d

a. Absolute ages used for behavioral research. b. Sarringhaus et al., 2014; Doran, 1992 c. Dental age markers used to create age categories for anatomical specimens. Dental markers were used to classify museum specimens according to age, as they correspond to chronological and behavioral age categories (Smith, Crummett, & Brandt, 1994; Smith et al., 2010, Smith & Boesch, 2011, Smith et al., 2013, Machanda et al., 2015; D. postcrania from Zihlman, Bolter, & Boesch, 2007.

Table 2. Dorsal metacarpal ridge angle and height for different age classes and genera of primates.

Category	DMR angle (all individuals)		DMR angle <180 (only DMR present)		DMR height in mm (all individuals)		DMR height <180 (only DMR present)	
	N	Mean (SD)	N	Subset Mean (SD)	N	Mean (SD)	N	mean (SD)
<i>Pan</i> adult	45	158.91 (8.46)	44/45	158.43 (7.91)	45	1.47 ^a (0.87)	44	1.50 (0.85)
<i>Pan</i> subadult	125 ^a	171.30 (10.14)	75/125	165.27 (9.18)	125	0.58 (0.71)	74	0.96 (0.65)
Young Inf. ^b (Pre M1)	17	180 (0)	0/17	NA	17	0	0	NA
Old Inf. (M1 pre M2)	51	176.09 (6.30)	21/51	170.50 (6.61)	49	0.24 (0.44)	21	0.59 (0.51)
Juvenile (M2 pre C)	30	167.04 (8.33)	28/30	166.10 (7.82)	31	0.93 (0.70)	28	0.99 (0.67)
Adolescent (C)	25	160.02 (10.04)	25/25	160.02 (10.05)	25	1.29 (0.65)	25	1.29 (0.65)
<i>Gorilla</i> adult	29	151.63 (11.76)	28/29	150.62 (10.62)	29	2.55 ^c (1.35)	28	2.64 (1.28)
<i>Gorilla</i> subadult	44	170.27 (9.62)	28/43	164.71 (7.70)	44	0.77 (0.87)	28	1.20 (0.80)
Pre M1	5	180.00 (0)	0/5	NA	5	0	0	NA
M1 pre M2	17	174.65 (7.18)	8/17	168.64 (6.30)	17	0.36 (0.48)	8	0.77 (0.42)
M2 pre C	17	167.96 (6.60)	15/17	166.36 (5.13)	17	0.93 (0.60)	15	1.06 (0.52)
C not fused	4	151.66 (5.56)	4/4	151.66 (5.56)	4	2.61 (0.96)	4	2.61 (0.96)
<i>Pongo</i> adult	16	178.38 (3.56)	3/16	171.37 (2.02)	16	0.04 (0.11)	3	0.22 (0.17)
<i>Hylobates</i> adult	41	180 (0)	0/41	NA	41	0.00	NA	0
<i>Papio</i> adult ^c	20	174.11 (7.30)	11/20	169.29 (6.68)	20	0.17 (0.24)	11	0.31 (0.24)

Primate specimens were from the American Museum of Natural History, Cleveland Museum of Natural History, Harvard Museum of Comparative Zoology, Natural History Museum London, Naturkundemuseum Stuttgart, the University of Zurich, and the Quex Museum (Kent, U.K.). a. 2 chimpanzee subadult individuals are subadult but have no reliable age estimates; b. inf = infant; c. 9 baboon individuals were measured from the lateral side. No

difference between samples measured from the lateral and medial side for baboons $F=0.864$, $df=1,21$, $p=0.363$); d. Compared to Inouye height measurement of *Pan troglodytes* for 1.42 (0.96); e. compared to Inouye *Gorilla beringei* 1.84 (1.53) and *Gorilla gorilla* 2.41 (1.77).

Table 3. Kruskal-Wallis post hoc results for DMR angle (green) and height (purple).

	Pan SA	Pan	Gorilla SA	Gorilla	Pongo	Hylobates	Papio
<i>Pan SA</i>	X	0.536	1.000	0.001	NA	NA	0.049
<i>Pan</i>	<0.001	X	1.000	0.792	NA	NA	0.001
<i>Gorilla SA</i>	1.000	<0.001	X	0.020	NA	NA	0.115
<i>Gorilla</i>	<0.001	1.000	<0.001	X	NA	NA	<0.001
<i>Pongo</i>	0.118	<0.001	0.104	<0.001	X	NA	NA
<i>Hylobates</i>	<0.001	<0.001	<0.001	<0.001	1.000	X	NA
<i>Papio</i>	1.000	<0.001	1.000	<0.001	1.000	0.148	X

Pairwise comparison results shown with Bonferroni correction for multiple tests. SA indicates subadults, all other groups comprised of adult specimens.

Angle (using radians) for ALL individuals were compared and height was compared ONLY between individuals with ridge PRESENT using LN DMR height/MC midshaft AP diameter (as LN transformation produces no value for ratios equivalent to zero)

