

Systematic review of family engagement interventions in neonatal, paediatric, and adult ICUs

Natalie S. McAndrew PhD, RN, ACNS-BC, CCRN-K, Assistant Professor and Nurse Scientist^{1,2}  | Teresa Jerofke-Owen PhD, RN, Assistant Professor³ | Christine A. Fortney PhD, RN, Assistant Professor⁴ | Deena K. Costa PhD, RN, FAAN, Assistant Professor⁵ | Breanna Hetland PhD, RN, CCRN-K, Assistant Professor⁶ | Jill Guttormson PhD, RN, Associate Professor³ | Eric Harding MLS, Clinical Services Librarian⁷

¹College of Nursing, University of Wisconsin-Milwaukee, Milwaukee, Wisconsin

²Froedtert & the Medical College of Wisconsin Froedtert Hospital, Milwaukee, Wisconsin

³College of Nursing, Marquette University, Milwaukee, Wisconsin

⁴College of Nursing, The Ohio State University, Martha S. Pitzer Center for Women, Children & Youth, Columbus, Ohio

⁵School of Nursing, University of Michigan, Ann Arbor, Michigan

⁶Omaha Division, College of Nursing, University of Nebraska Medical Center, Omaha, Nebraska

⁷Medical College of Wisconsin Libraries, Medical College of Wisconsin, Milwaukee, Wisconsin

Correspondence

Natalie S. McAndrew, Assistant Professor and Nurse Scientist, College of Nursing, University of Wisconsin-Milwaukee, 1921 East Harford Avenue, Milwaukee, WI 53211.

Email: mcandre3@uwm.edu

Funding information

National Institute of Nursing Research, Grant/Award Number: P20-NR015331; Center for Complexity and Self-management of Chronic Disease at the University of Michigan School of Nursing

Abstract

Aims and objectives: The purpose of this systematic review was to evaluate interventions that have been used to engage families in direct care activities (active family engagement) in adult, paediatric, and neonatal intensive care unit (ICU) settings.

Background: Family engagement is universally advocated across ICU populations and practice settings; however, appraisal of the active family engagement intervention literature remains limited.

Search strategy: Ovid Medline, PsycArticles & PsycInfo, Scopus, and CINAHL were searched for family interventions that involved direct care of the patient to enhance the psychological, physical, or emotional well-being of the patient or family in neonatal, paediatric, or adult ICUs.

Inclusion/exclusion criteria: Studies were included if an active family engagement intervention was evaluated. Studies were excluded if they were not published in English or reported non-interventional research.

Results: A total of 6210 abstracts were screened and 19 studies were included. Most studies were of low to moderate quality and were conducted in neonatal ICUs within the United States. Intervention dosage and frequency varied widely across studies. The interventions focused on developmental care (neonatal ICU) and involved families in basic patient care. Family member outcomes measured included satisfaction, stress, family-centred care, confidence, anxiety, and depression. Most studies found improvements in one or more outcomes.

Conclusions: There is a paucity of literature about active family engagement interventions, especially in adult and paediatric populations. The optimal dosage and frequency of family engagement interventions remains unknown. Our systematic review found that data are limited on the relationship between family engagement and patient outcomes, and provides a timely appraisal to guide future research.

Relevance to Clinical Practice: Further research on the efficacy of family engagement interventions is warranted. The translation of active family engagement interventions into clinical practice should also be supported.

KEY WORDS

family engagement, family involvement, family care in critical care, family-centred care, intensive care, role of family in ICU

1 | INTRODUCTION

Family engagement is an *active* partnership among health professionals, patients, and their families that can improve individual health and well-being and health care quality and safety.¹⁻³ Current family-centred care (FCC) guidelines⁴ stress the importance of family involvement, particularly in decision-making. However, the evidence base for active family engagement, in which family members contribute to aspects of direct patient care,⁵⁻⁷ is limited. Involving family in the delivery of care is advocated across ICU populations and practice settings, but to our knowledge, an appraisal of active family engagement interventions has not been conducted. Thus, we undertook a systematic review of active family engagement interventions in neonatal, paediatric, and adult ICUs.

2 | BACKGROUND

Family engagement has become an important concept in the critical care literature,^{2,5-8} with numerous calls to improve family engagement in the ICU.^{2,5,6,9} In a recent scoping review of family involvement interventions in adult ICUs, family engagement is described on a continuum moving from passive (eg, physical presence at the bedside and receiving and having needs met) to more active activities (eg, sharing and receiving information, involvement in decision-making, and making contributions to the care of the patient).³ Empirical evidence supports the value of family presence, communication with families, and the importance of decision-making support in the ICU.^{3,4,10,11} However, the evidence on interventions that directly involve families in the care of the critically ill patient and the effect of such interventions on patient and family outcomes has not been well described.

Active family engagement is an important element of and vehicle to achieving FCC,¹ and relevant to health care delivery across the lifespan.⁴ Although paediatrics has embraced an FCC philosophy longer than adults specialties, universal barriers to FCC in neonatal, paediatric, and adult ICUs include: inadequate guidance and support for families; lack of guidelines and policies for family engagement; inadequate time to engage with families; and a lack of unit and organizational support for

What is known about this topic

- Family engagement is universally advocated across ICU populations and practice settings; however, appraisal of the active family engagement intervention literature remains limited

What this paper adds

- There is a paucity of literature about active family engagement interventions, especially in adult and paediatric populations.
- The optimal dosage and frequency of family engagement interventions remains unknown.
- Our systematic review found that data are limited on the relationship between family engagement and patient outcomes and provides a timely appraisal to guide future research.

FCC.¹²⁻¹⁶ Hence, there is a need for evaluation of the existing evidence to identify effective strategies for promoting active family engagement in ICU environments across the lifespan.^{3,5,13}

As more researchers begin to test active family engagement interventions in paediatric and adult settings, a detailed review of existing interventions and their impact on patient and family outcomes is needed. The purpose of this systematic review is to describe and evaluate interventions that have been used to actively engage families in neonatal, paediatric, and adult ICUs.

3 | METHODS

3.1 | Design

A systematic review following the preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines¹⁷ was

conducted. The review was registered in the International Prospective Register of Systematic Reviews (National Institute for Health Research)—(CRD42018109259).

3.2 | Search strategy

The literature search strategies were designed by a medical librarian. Ovid Medline, PsycARTICLES, PsycINFO, SCOPUS, and Cumulative Index of Nursing and Allied Health Literature (CINAHL) databases were searched using a combination of standardized terms and keywords including: empower, involve, activate, engage, participate, collaborate, FCC, and patient participation. The full search strategy for Ovid Medline is shown in Figure 1.

3.3 | Study inclusion criteria

To be included for review, articles had to: (a) include family (defined as family member, family caregiver, partner, significant other, relative, parent, spouse, or children) (b) be conducted in an ICU setting (neonatal, paediatric, or adult), and (c) report quantitative or qualitative outcomes of a family-focused intervention that actively engaged family in some aspect of patient care defined as—doing something with/for the

1. intensive care units.mp. or exp Intensive Care Units/
2. critical care.mp. or exp Critical Care/
3. critical care nursing.mp. or exp Critical Care Nursing/
4. ICU.mp.
5. 1 or 2 or 3 or 4
6. exp FAMILY/ or exp NUCLEAR FAMILY/ or exp FAMILY RELATIONS/ or family.mp.
7. family caregiver.mp. or exp Caregivers/
8. partner*.mp.
9. significant other.mp.
10. relative*.mp.
11. exp Parents/ or exp PARENT-CHILD RELATIONS/ or parent*.mp.
12. family member*.mp.
13. 6 or 7 or 8 or 9 or 10 or 11 or 12
14. patient participation.mp. or exp Patient Participation/
15. professional-family relations.mp. or exp Professional-Family Relations/
16. patient preference.mp. or exp Patient Preference/
17. patient-centered care.mp. or exp Patient-Centered Care/
18. nurse-patient relations.mp. or exp Nurse-Patient Relations/
19. patient satisfaction.mp. or exp Patient Satisfaction/
20. physician-patient relations.mp. or exp Physician-Patient Relations/
21. decision making.mp. or exp Decision Making/
22. personal satisfaction.mp. or exp Personal Satisfaction/
23. 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21 or 22
24. empower*.mp.
25. engage*.mp.
26. participat*.mp.
27. involve*.mp.
28. perception*.mp.
29. collaborate.mp.
30. collaboration.mp.
31. family-centered care.mp.
32. patient activation.mp.
33. family activation.mp.
34. activation.mp.
35. experience.mp.
36. 24 or 25 or 26 or 27 or 28 or 29 or 30 or 31 or 32 or 33 or 34 or 35
37. 5 and 13 and 23 and 3

FIGURE 1 Ovid Medline search strategy

patient to enhance their psychological, physical, or emotional well-being. We excluded: (a) non-interventional study designs, (b) quality improvement and dissertation studies, and (c) studies not published in English.

3.4 | Review process

Literature searches were completed in September 2018 and updated in May 2019. Results were exported into EndNote, and uploaded into an open source software to manage systematic reviews (Rayyan QCRI).¹⁸ Six reviewers independently screened all of the abstracts. If at least two reviewers included an abstract, a full text review was completed. Discussions were held among all six reviewers until consensus was reached about study inclusion.

3.5 | Data extraction

A comprehensive data extraction form was developed based on the Cochrane data collection for intervention reviews¹⁹ and the Grading of Recommendations, Assessment, Development, and Evaluations (GRADE).²⁰ Data extraction focused on the population, setting, sample sizes, intervention descriptions, measures, and intervention

outcomes. Two team members were assigned to each study and independently extracted data and graded the quality of the study using the GRADE criteria (quality of evidence based on confidence that the true effect is close to the estimate of the effect and rated as very low, low, moderate, or high).²⁰ The GRADE criteria examines factors such as limitations in study design and other risks of bias. Differences were resolved through collaborative review and discussion until consensus was achieved among the two reviewers. Data extraction results were entered into a table format to synthesize the findings.

4 | RESULTS/FINDINGS

A total of 6210 records were identified. With automated duplicate finding, 663 duplicates were removed for a total of 5547 citations. After abstract screening, 147 full text articles were reviewed for inclusion. Nineteen articles met all inclusion criteria and were included in the review (Figure 2). The summative table of the interventions and study characteristics can be found in Table 1. Given the heterogeneity across the studies, a narrative approach was used to describe the study findings.⁴⁰ Intervention descriptions are grouped by patient population (neonates, children, or adults).

4.1 | Study characteristics

4.1.1 | Design

There were eight reports of randomized controlled trials (RCT),^{24-27,29,30,32,33} and seven quasi-experimental studies.^{21,22,28,31,35,36,39} Four studies were pilots and/or feasibility/acceptability studies.^{23,34,37,38} All of the RCTs were conducted in the NICU. Two of the RCTs reported on different elements of the same longitudinal study.^{26,27} A quasi-experimental design was the strongest design used in studies conducted in the PICU or adult ICU.

4.1.2 | Study quality

Study quality varied, with only one study rated as high quality.³⁰ Seven studies were of moderate quality,^{21,25-27,32,33,36} four were low quality,^{22,24,28,29} three were very low quality,^{31,35,39} and four studies could not be assigned a quality grade because of design (pilot/feasibility/acceptability).^{23,34,37,38} Only one study included any element of blinding³⁰; however, given the types of multifaceted interventions blinding was not feasible in most studies.




FIGURE 2 Preferred reporting items for systematic reviews and meta-analyses (PRISMA) diagram

TABLE 1 Summative table of interventions and study characteristics

Author/Year	Location	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
NICU studies								
Byers et al. ²¹ 2006 United States	Setting: neonatal ICU; single centre Sample: Parents of premature infants 32 weeks or less	Evaluate the impact of individualized developmental supportive family care on infant and family outcomes	Developmentally supportive care (based on Synactive Theory of Development)	Quasi experimental 114 parent/infant dyads Specific number in control and intervention group not reported	Intervention Individualized, developmentally supportive family-centred care, which included a specialized section of NICU and staff education 1. Specialized Section: open visitation, acoustic panels, privacy curtains (reduce noise and light) 2. Staff Education: developmentally supportive family care course included family centred care, ethics, communication, neonatal developmental NICU experiences; also newborn assessment training (Brazeltton, oral/motor; NIDCAP). Full NIDCAP completed within first 7 days of admission.	Physiologic (vitals) Medical and development progress (feeding, growth, behavioural stress cues) Return to sleep Length of stay Complication rates (ventilation days, days to open crib) Parent satisfaction and perceptions Resource utilization	Lower behavioural stress cues intervention group No statistical difference in other measures including cost, physiologic (with exception of lower activity respiratory rate in intervention group), growth, feeding, return to sleep, achievement of infant progress, complication rates, parental perceptions/satisfaction	Moderate Different inclusion/exclusion intervention and control No comparison of parents on demographics Data collection not blinded to study group No intervention fidelity monitoring reported Exclusion criteria was 32 weeks or less, yet demographics indicate infants more than 32 weeks at enrolment

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Cooper et al ²² 2007 United States	Setting: 8 Neonatal ICUs Sample: Hospital administrators, staff, parents/ family	Evaluate the impact of March of Dimes NICU Family Support Program on overall care and family centred practices	None	Quasi experimental Three groups NICU with full (four NICUs), partial (three NICUs), and no implementation of program (one NICU) Total Sample includes: 11 NICU administrators, 502 NICU staff members, 216 NICU families	<u>Intervention</u> March of Dimes Family Support Program (NFS): national program designed to promote family-centred care in NICUs. Includes: <ul style="list-style-type: none">• Family support specialist• Baby photos and scrapbooking• Parent to parent support• Education for parent and siblings• Staff education <u>Control</u> Usual care	Presence of intervention components (yes/no) Parental knowledge, comfort, confidence, connectedness Parental behaviours (asking questions of HCPs, ability to describe infant condition, level of involvement in care) Parent self-efficacy Parent report of programs and policies that endorsed intervention components	Parents/family reported: <ul style="list-style-type: none">• increased family receipt• increase parent comfort after education• increased parent-parent support• NFS specialist helped decrease stress and increase confidence for parentsAt partially and fully implemented sites, parents were more comfortable knowing what to expect for baby's medical condition and baby's growth and development Parents partially and fully implemented sites felt their opinions were taken seriously "often" or "a lot"	Low Measures post implementation only No data given on comparison sites to determine equivalence No statistical control for unit factors Data self-report from surveys and interviews No intervention fidelity monitoring/ reporting

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
De Bernardo et al ²³ 2017 Italy	Setting: NICU, single centre Sample: Parents of NICU infants at least 30 days post single surgery	Compare satisfaction and stress between parents in a family centred care group vs non-family centred care	Synactive Theory of Development	Non-randomized, prospective cohort pilot 144	Implementation of the Family Centred Care (FCC) Model • Physical changes to the NICU: addition of kitchen and family rooms	Infant weight Parent satisfaction (receipt of information, health care team communication and collaboration; privacy)	Higher infant weight at day 60 day intervention group Parent increased satisfaction with communication/ collaboration; (Parent Stress Scale)	Low Small sample size One unit; pre/post intervention (different groups control/ intervention) No intervention fidelity monitoring

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Francik et al. ²⁴ 2011 England	Setting: 4 NICUs Sample: Parents of NICU babies	Feasibility and effect of an intervention to increase parental involvement in pain management for NICU infants in relation to parents' stress and post discharge parenting	None	Randomized controlled trial (RCT) 169 (84 intervention; 85 control)	Information received; privacy Lower stress for parental roles, baby appearance/treatment; and NICU environment	Scale: Neonatal Intensive Care [PSS;NICU])	Information received; privacy Lower stress for parental roles, baby appearance/treatment; and NICU environment	No control variables when evaluating parental stress (eg, socio-economic status, lack of sleep, etc.)

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Hoffenkamp et al. ²⁵ 2015 Netherlands	Setting: NICU (2) and maternity ward (7); 7 hospitals Sample: Parents with infants born at <37 weeks	Competence and confidence		it may affect infants • How pain is assessed and managed in the NICU • The important role parents can play • Specific instructions on how to comfort infants • Advice on how parents can work in partnership with NICU staff. Parents also received two visits (~45 minutes each from a research nurse) to show them how to apply the comforting techniques and to answer questions	Parental confidence in infant care- giving (Self- Efficacy in Infant Care Scale [SICS]) Parental perceptions of role attainment (What Being a Parent of a New Baby Is Like [WPBBL-R])	Parental confidence in infant care- giving (Self- Efficacy in Infant Care Scale [SICS]) Parental perceptions of role attainment (What Being a Parent of a New Baby Is Like [WPBBL-R])	No differences in parental stress No group differences satisfaction with infant pain care or confidence in ability of staff to manage infant pain and support parents	No differences in parental stress No group differences satisfaction with infant pain care or confidence in ability of staff to manage infant pain and support parents
								Moderate Interrater agreement for observational coding has some limitations Not clear which part of the intervention actually

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Holditch-Davis et al. ^a 2013 United States	Setting: 4 NICUs; 2 academic medical centres, 2 community-hospitals Sample: Mothers of preterm infants	Examine mothers' satisfaction with auditory-tactile-visual-vestibular (ATVV) intervention and kangaroo care.	None	Longitudinal 3-group experimental 249 (208 mothers completed satisfaction survey reported)	Intervention (two groups) 1. ATVV: perform message that involves moderate stroking, eye	Satisfaction with intervention Helpfulness of nurses who taught them intervention	All groups were satisfied with the intervention and helpfulness of the nurses. Lower satisfaction at 2 months	Moderate Same nurse performed all interventions Investigator-developed instrument
				Control Usual care				(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Holditch-Davis et al. ^a	Setting: Four NICUs: two academic medical	Explore whether mother and infant characteristics affected maternal satisfaction ratings	here) (73 control, 67 ATVV, 68 KCI)	2. Kangaroo care: skin-to-skin contact and holding Instructed to provide interventions for at least 15 minutes at least once a day, 3 times a week until infant was 2 months corrected age. Were able to provide whichever care they wanted to their infants Control ^b	contact, talking to, and rocking infant	RCT-3-group longitudinal study	Infant sleep-wake responses to the intervention (two groups)	Moderate ATVV infants had significantly more alert periods,
							Infant sleep-wake responses to the intervention	Limited sample size

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
2014 United States	centres; two community hospitals Sample: Mothers of preterm infants	vestibular (ATVV) intervention and kangaroo care (KC)		240 (81 in control, 78 ATVV, and 81 KC)	<p>1. Auditory-tactile-visual-vestibular intervention (ATVV): stimulation in a gradual progression over 15 minutes beginning with voice only then auditory and tactile, with visual stimulation as infant becomes alert. Horizontal rocking added and tactile component withdrawn for final 5 minutes.</p> <p>2. Kangaroo Care (KC): holding infant skin to skin in upright position between mother's breasts. As long as wanted for at least 15 minutes.</p>	<p>Arousal was scored for 5 minutes from the videotapes. The predominant sleep-wake state (alertness, drowsiness, active waking, sleep-wake transition, active sleep, and quiet sleep was recorded once a minute following ATVV and beginning following the start of KC.</p>	<p>total waking, and total alertness. No significant differences in infant responsiveness to stimulation.</p> <p>No difference maternal anxiety, depressive symptoms, or Posttraumatic stress symptoms, or parenting stress.</p> <p>KC mothers more rapid decline in worry than the other groups.</p> <p>Maternal positive involvement, developmental stimulation, and HOME total score did not differ.</p>	<p>Crossover between groups (20%-30% of KC and ATVV moms and 58% of control moms engaged in a non-assigned intervention)</p> <p>High attrition rate (21%)</p>

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Lee et al ²⁸ 2013 Taiwan	Setting: NICU, single centre Sample: Fathers of infants <37 weeks gestation with expected LOS of at least 2 weeks	Evaluate the effectiveness of an intervention on fathering ability, perceived nurse's support and parental stress after a preterm infant's admission to a NICU	Four components of support 1. information 2. emotional 3. instrumental 4. esteem	Historical comparison group 69 (34 control group; 35 intervention group)	Intervention Booklet distributed to fathers providing information about premature babies and NICU (equipment, developmental care, nutrition, infant appearance and behaviour, ways to interact with infant, and relaxation tips). A nurse present during visits to answer questions and encourage use what was in booklet, and	Parental stress: (Parent Stress Scale; Neonatal Intensive Care [PSS:NICU]) Fathering ability (FA:NICU; author developed) Father perceived nursing support (Nurse-Parent Support Tool [NPST])	Low Intervention fathers had significantly higher fathering ability and perceived nursing support Fathers receiving the intervention program had greater reduction in stress than the comparison group Fathers rated booklet as being helpful	No randomization Historical control Only one NICU Self-developed tools

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Matricardi et al. ²⁹ 2013 Italy	Setting: NICU, single centre academic hospital Sample: Mothers and fathers of preterm infants <32 weeks hospitalization	Determine the effects of a parental intervention in reducing parental stress levels during hospitalization	None	RCT 42 mother and father dyads (21 dyads control group; 21 dyads intervention group)	Intervention Each couple met with the unit physical therapist for eight sessions lasting 1 hour each (from 31 to 36 weeks post-menstrual age of infant). Goal of sessions were to increase parental ability to recognize signs of infant stress and well-being, help them to soothe their infant, improve physical contact with appropriate stimulation with infant.	Parental stress (Parent Stress Scale: Neonatal Intensive Care [PSS:NICU])	Mothers (not fathers) reported lower role-stress at time 2 in the intervention group compared with the standard care group Mothers spent more time in NICU than fathers in both the intervention and control groups	Low Unclear how close the sample represented the population Limited description of what occurred with each session of the intervention

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Melnyk et al. ³⁰ 2006 United States	Setting: NICU, multicentre, two academic hospitals Sample: Mothers and fathers of infants with gestational age of 26 to 34 weeks, a birth weight of less than 2500 g, and anticipated to survive, had no severe deficits	Evaluate the efficacy of Creating Opportunities for Parent Empowerment [COPE] program (educational- behavioural intervention to enhance parent- infant interactions) and determine effects on parental	Self-regulation and control theories	RCT 260 73 fathers control group; 147 mothers and 81 fathers intervention group)	Intervention COPE; provides information (written and audiotape) to parents in four phases from time of admission to after discharge.	Infant length of stay (LOS) Anxiety (State-Trait [STA]) Depression (Becks Depression Inventory [BDI]) Phase I content: information provided on infant-behaviour and parent-role;	Shorter infant LOS in NICU and hospital Mothers in COPE had significantly less parental stress Parents in COPE has more positive parenting interactions with their infant Fathers in COPE more involved in infant care and	High Blinded study Active control group Analysis controlled for infant characteristics
						Parental Stress (Parent Stress Scale: Neonatal Intensive Care [PSS: NICU]) Parent-infant interaction		

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Noergaard et al. ³¹ 2018 Denmark	Setting: NICU, single centre regional hospital	Investigate the impact of a more father-friendly NICU on paternal stress and	None	Quasi-experimental 109 (55 control group; 54 intervention group)	parents asked to track milestones Phase II content:— recognizing stress cue and readiness for interaction Phase III content(prior to discharge) smoothing transition from hospital to home; recognizing cues and helping stressed infant Phase IV content (1 week post discharge); Information preterm infant development, suggestions for positive parent-infant relationships and activities to foster cognitive development <u>Control</u> Audiotapes and written information on hospital services, discharge, and immunization provided at same timepoint as COPE phases.	ratings (Index of Parental Behavior in the NICU) Parents' beliefs about their infants and their parental role during hospitalization (Parental Belief Scale-NICU)	more sensitive to the babies needs At 2 months, mothers in COPE significantly less anxiety and depressive symptoms No Fathers had no difference between groups for in anxiety and depression at 2 months	Higher stress for father's in intervention group in all domains with

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
	Sample: Fathers of infants admitted to NICU	participation in newborn care		friendly NICU including: <ul style="list-style-type: none">• encouragement of fathers' participation (skin-to-skin contact and routine care of newborn)• direct communication with fathers about newborns status and development• counselling from a social worker on maternity leave• provision of support groups for fathers• inclusion of other family members in care of newborn• opportunity for older siblings to stay overnight at the hospital	childcare at the time of discharge for fathers	exception of parental roles subscale	than 60% of the items. No data reporting extent to which the intervention was implemented	
Van der Pal et al ³² 2007 Netherlands	Setting: NICU; multicentre, academic hospitals	Two studies reported: #1: Compare the effect of basic developmental care (incubator covers and nests) to usual care #2: Compare the effect of basic developmental	Synactive Theory of Infant Development	RCT (Two studies) RCT #1: 192 infant/parent dyads (94 control; 98 intervention) RCT #2: 168 infant/parent dyads (84 control; 84 intervention)	RCT #1 <u>Intervention</u> Basic developmental care which included reduction of light and sound using incubator covers and nests to	1. Confidence (Two Scales from the Mothers and Baby Scale): (1) Confidence in Caregiving (CC) (2) Global Confidence Scale (GCS)	No difference confidence, perceived nurse support or parental stress between groups in either RCT	Moderate No blinding of intervention No report of intervention fidelity

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
care to Newborn Individualized Developmental Care and Assessment Program (NIDCAP)								
Welch et al. ³³ 2016 United States	Setting: NICU, single academic centre Sample: Mothers who had delivered a singleton or set of twins between 26 and 34 weeks gestational age	Evaluate the effects of the Family Nurture Intervention (FNI) on mothers and infants	Calming cycle	RCT 115 (56 control group; 59 intervention group)	support motor development <u>RCT #1 Control</u> Usual care <u>RCT #2</u> <u>Intervention</u> Newborn Individualized Developmental Care and Assessment Program (NIDCAP); observations of the infant before, during and after caregiving every 7 to 10 days by NIDCAP trained developmental specialist First observation within 48 hours of birth. Only nurses trained in NIDCAP cared for infants	Parental perception of nurse support (Nurse Parent Support Tool (NPST)) Parental Stress (Parent Stress Scale: Neonatal Intensive Care [PSS;NICU])		Moderate No differences in FNI and standard care (SC) groups at baseline or near term age for depression and anxiety At 4 months for FNI mothers vs SC:

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Sarin et al. ³⁴ 2019 India	Setting: NICU, single academic centre	Understand the acceptability of Family centred Care (FCC) from providers' and parents'	Philosophy of Family centred Care	Qualitative 12 family members (5 mothers, 5 fathers, 2 grandparents), 6 health care providers (3 paediatric residents; 3 nurses)	Inhibition/ Behavioral Activation Scales [BISBAS])	• Lower depression and anxiety scores • Higher percentage of mothers breastfeeding • More frequent and longer skin to skin sessions for FNI mothers vs mothers receiving standard care.	• Lower depression and anxiety scores • Higher percentage of mothers breastfeeding • More frequent and longer skin to skin sessions for FNI mothers vs mothers receiving standard care.	Unable to grade (qualitative data only) Bias: did not include inter-rater coding and interpretation: data were coded and analysed by the lead researcher only. Site: hospital had FCC in place prior the study and, reported greater
	Sample: Family of neonates and NICU staff	Explore the integration of providers' and clients' activities in NICU Examine the continuing care			Intervention At admission, family members received an introduction to FCC in a face to face session with doctor or nurse. Session included information on the training process required to become a parent-attendant	Acceptability and perceived benefits and challenges of the FCC intervention	Family members reported FCC increased knowledge, promoted adaptation to NICU, increased access to their infant and increased infant well-being parents/ caregivers reported greater	

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Kuntaros et al. ³⁵ 2007 Thailand	Setting: PICU, single centre Sample: Mothers of children admitted to PICU in the last 24 hours	Competencies of parents after discharge			and the parental role in care provision Four education sessions were offered. Session 1 and 2 topics included: infection prevention, breastfeeding, breast milk expression, assisted feeding Session 3 for parents of low weight newborns- addressed kangaroo care. Session 4 focused on preparation for discharge and care of newborn at home.		capability and more empowerment to care for newborn Health care providers had a favourable perception of FCC because of benefit of parental presence for well-being of the newborn	therefore, may have had additional support available in other hospital settings. One group design
PICU	Kuntaros et al. ³⁵ 2007 Thailand	Setting: PICU, single centre Sample: Mothers of children admitted to PICU in the last 24 hours	Compare maternal self-efficacy in participatory involvement in child care and satisfaction with nursing care between control group and experimental group that received family-centred care approach	Effective Helpgiving Practices 1. technical quality 2. helpgiver traits/ attribution 3. help receiver participatory involvement	Quasi-experimental 32 (16 control 16 intervention) Mothers in both control and experimental groups were matched with their years of education.	Intervention Mothers allowed to visit at any time and were more involved in care than control group mothers. Over 4 days, the investigator accompanied mom and family to bedside visit and provided information about unit, assessed	Higher self-efficacy and satisfaction in intervention group. Higher scores on satisfaction survey also	Very low It is not clear how mothers were placed into groups. Intervention not well described Small sample size; only one site Investigator developed instruments; not previously tested Because many patients were

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
					perceptions of the child's illness and expectations for nursing services, clarified/ corrected misperceptions, gave updates about child's condition, encouraged mental and emotional support and sharing of concerns, needs, and ideas, provided a chance for joint decision-making, and allowed them to provide care to the child by themselves.			under sedation some activities in the experimental intervention could not be performed.
Adult ICU	Black et al³⁶ 2011 Northern Ireland	Setting: adult ICU; single centre Sample: Patient/ family dyads	Examine the effects of nurse- facilitated family participation in psychological care (operationalized as emotional support and presence) on the extent of patient delirium and psychological recovery following critical illness	Neuman's Systems Model	Time Series (pre/post) 170 (83 control group; 87 intervention group)	Intervention Information booklet provided to family focused on how to provide psychological support through communication with patient; included suggestions on how to interact and topics/type of information to share with	Length of Stay Physiologic (labs; vitals), Medications (sedatives, analgesics, muscle relaxants) Delirium (Intensive Care Delirium Screening Checklist) Impact of illness (Sickness Impact Profile [SIP])	Moderate Pre/post design; unclear if groups equivalent at baseline Single setting (Seven Bed ICU in one hospital) No intervention fidelity remained when controlling for severity of illness, length of ICU stay, and delirium

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Davidson et al. ³⁷ 2010 United States	Setting: adult ICU; single centre Sample: Family members or significant support person of adult mechanically ventilated ICU patients	Evaluate the feasibility of implementing interventions from the Facilitated Sensemaking Theory into the ICU	Facilitated Sensemaking	Pilot/feasibility 30 family members or significant support person	Intervention Two main components: personalized instruction and provision of family visiting kits 1. Personalized instruction: introduction and explanation of the project, decoding of the environment at the bedside (done by the nurse), instructions on helpful visiting activities, coaching on how to ask questions of the doctor and identify unmet needs,	Feasibility and family evaluation of the program Family report of needs (importance and if met)	All items offered within the intervention were found useful to some family members. All proposed family interventions (eg, bedside and cognitive recovery activities) were used and found helpful. The journal was least useful and personal care supplies were most useful. Observation of families; most engagement when receiving information on how to participate at bedside and how to decode/ debriefing at the	Unable to grade difference delirium or length of stay between groups

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Marshall et al. ³⁸ 2016 Australia	Setting: Adult ICU, multicentre, academic hospitals Sample: Patients on mechanical ventilation for 48 hours and more	Evaluate the feasibility and acceptability of an intervention that aimed to educate families about the importance of breathing	Family-centred Theory	Pilot/feasibility 126/49 patients, 51 family members, 4 physicians, 20 nurses, 2 dieticians 30 family members received low intensity	Intervention (two groups) 1. Low intensity: patient nutritional history acquired from family shared with	Feasibility and acceptability Recruitment and retention rates	Recruitment/ consent rate of 26% Retention rate of 67% Families found both the low and moderate intensity	Unable to grade Small sample Only two ICUs No control group Only qualitative data obtained

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
	their families; health care professionals	nutrition for recovery from critical illness		intensity intervention 19 family members received moderate intensity intervention	health care team; short education session supplemented with a printed resource emphasizing importance of nutrition and explaining therapies during and after critical illness. Guided prompts given to families to promote discussion with health care team		interventions acceptable The intervention prompted conversations with family, friends, and health care professionals, and supported planning nutrition support post ICU Some limitations with food diary, participants unclear about how it should be used	This intervention also prompted families to ask questions related to other aspects of care
				2. Moderate intensity: low intensity	components plus a daily nutritional diary that was completed by the family during the two-week period following extubation of the patient. Diary used to promote conversation between the family and the patient about nutrition and to encourage discussion with health care team			Control None

(Continues)

TABLE 1 (Continued)

Author Year Location	Setting and sample	Aims/research questions	Theoretical model	Design/sample size	Study conditions	Outcome measures	Results	Quality appraisal
Mitchell et al. ³⁹ 2009 Australia	Setting: Medical/Surgical Adult ICU, multicentre two academic hospitals Sample: Families of patients predicted to be in the ICU more than 2 days	Determine the effect of a family-centred nursing intervention on family members' perceptions of family-centred care	Family-centred care philosophy/ model	Pragmatic clinical trial; non-equivalent control group, pretest-posttest design 174/75 control group; 99 intervention group)	Intervention Families participated in fundamental care activities at the discretion of the nurse including: hair combing, hand massage and bathing Control Usual care	Family member perception of collaboration, respect, and support from ICU staff (Family centred Care Survey)	Those in the intervention were more likely to perceive higher FCC (respect, collaboration and support) than those in the control group.	Very low No control for unit culture and other variables in analysis Adapted outcome measure was not validated

^aHolditch-Davis et al 2013 and 2014 report results from the same study.^bPublication does not report satisfaction for control group, satisfaction only reported for intervention groups.

4.1.3 | Family sample

Sample sizes ranged from 12 (acceptability study)³⁴ to 414 (RCT)³⁰ family members. Of the studies that reported family member demographics, there were larger percentages of female than male participants, with female (mother or other family member) participation ranging from 50% to 95%.^{24,25,29,30,32,34,37-39} Four studies included only mothers,^{26,27,33,35} and two studies included only fathers.^{28,31} Of the studies reporting race, samples ranged from 2.5% to 84% White,^{24,26,27,30,32,33} with the most diverse sample reported as 77% Black and 16% Hispanic.²⁷

4.1.4 | Study setting

Seven studies were conducted in the United States,^{21,22,26,27,30,33} two in Italy,^{23,29} two in the Netherlands,^{25,32} and two in Australia.^{38,39} Other locations included India,³⁴ Denmark,³¹ China,²⁸ Thailand,³⁵ and Ireland.³⁶ The majority of the studies were conducted in the neonatal ICU (NICU) ($n = 14$),²¹⁻³⁴ followed by the adult ICU ($n = 4$).³⁶⁻³⁹ Only one study was conducted in a paediatric ICU.³⁵ More than half were single centre studies ($n = 11$), and many were conducted in large academic medical centres ($n = 8$). Most NICU settings were designated as Level III care centres (*prompt and readily available access to a full range of paediatric medical subspecialties*), with one Level IV NICU (*highest level of neonatal care for complex and critically ill infants*).³³

4.2 | Intervention characteristics

4.2.1 | Theoretical framework

Thirteen studies included a theoretical model/framework that guided the intervention.^{21,23,25,28,30,32,33,35-37,39} NICU frameworks included developmentally supportive care and the Synactive Theory of Infant Development,⁴¹ self-regulation and control theories, mediated learning based on social learning theory, and the philosophy of FCC.^{34,38,39} The PICU study utilized helpgiving practices.⁴² Adult ICU frameworks included Neuman's System's Model, facilitated sense-making,⁴³ and FCC.^{38,39}

4.2.2 | Types of interventions

NICU interventions were focused on involving a parent or parents in some aspect of developmental supportive care for the infant. Intervention components included: education (informational packets and conversations with health care professionals) about the NICU environment and behaviours and characteristics commonly exhibited by NICU babies, and structured guidance from health care professionals on how to care for the baby using techniques such as kangaroo care, auditory-tactile-visual-vestibular stimulation, infant massage, and calming touch. Almost all of the NICU interventions reviewed included

an educational component, and many utilized nurses or other staff (family support specialist or physical therapist) to help parents to master the knowledge and skills required for infant care. Two studies used video-recorded sessions of parents interacting with their infant to provide behaviour-based feedback to parents about how to bond and engage with the infant in a developmentally appropriate way.^{25,30} Two studies used the Newborn Individualized Developmental Care and Assessment Program (NIDCAP) as a component of an intervention or as an intervention.^{21,32} The Family Nurture Intervention (FNI) used in one study involved calming activities for the mother to perform with the infant³³ including the use of scent cloths for both the infant and the mother to encourage bonding. The PICU intervention involved flexible visiting with structured support from nursing staff to encourage the mother to talk to, touch, and hug the child, and participate in daily care.³⁵ These paediatric-based interventions used behaviour change strategies such as modelling, feedback, and increasing parent self-efficacy through mastery of skills.^{23-25,28-31,33-35} Standard care was most commonly used in the control groups with the exception of one study that used an attention control.³⁰

Similar to NICU interventions, adult ICU interventions often included a mix of education, information packets, and nurse facilitation of family involvement in care.³⁶⁻³⁹ Specific strategies included nurse-facilitated family communication and interaction with the patient,³⁶ personalized nurse instruction and family visitation kits,³⁷ family involvement in basic patient care,³⁹ and family involvement in the assessment and management of the critically ill patient's nutritional status.³⁸ These interventions included educational components (providing information via written material or with a health care professional on the ICU environment and how to deliver patient care activities), as well as behavioural change strategies such as feedback, modelling, and reinforcement of skills in patient care, and teaching families what to expect to guide goal setting. There was only one study that took place in the adult ICU that had a control group receiving standard care.³⁶

4.2.3 | Frequency and duration of intervention (dosage)

The majority of the studies required multiple sessions for the delivery of all of the intervention components. NICU intervention sessions ranged from 15 to 60 minutes in length^{24-27,29} and occurred from three to four times per week^{26,27,33} to as frequently as daily.²³ In other NICU studies there was a specified number of sessions ranging from two to eight total sessions.^{24,29,30,34} There was an adult ICU study that included both a low and a moderate intensity intervention.³⁸ In the paediatric and adult ICU studies dosage was difficult to discern.

4.3 | Effects of interventions

4.3.1 | Infant outcomes

There were no significant differences in physiologic outcomes of the newborn such as HR, O₂ saturations, growth, feeding, complication

rates, and time to return to sleep when comparing infants in control and intervention groups.²¹ Pain levels were actually perceived to be higher by parents in the intervention group²⁴ than parents in a control group. Infant weights were significantly higher,²³ length of stay in the NICU and hospital were shorter³⁰ and infants had more alert periods and total waking time in the intervention groups compared with the control groups.²⁷ Infants whose parents were instructed on developmentally supportive care demonstrated lower behavioural stress cues and lower respiratory rates during activity than infants in the control group who received usual care with restricted visitation.²¹

4.3.2 | Adult patient outcomes

There were lower physical and psychosocial impact of illness scores for patients in the intervention vs the control; however, there were no differences in delirium or length of stay.³⁶

4.3.3 | Parental psychological factors

Ten of the NICU studies examined stress as an outcome. Results varied: Six studies demonstrated a significant association between the engagement intervention and decreased parental stress levels related to parental role, NICU environment, or appearance of the infant,^{22,23,27-30} three studies showed no significant relationship between parental stress level and participation in an engagement intervention.^{24,25,32} Two studies were conducted with just the fathers of infants hospitalized in a NICU,^{28,31} with conflicting outcome results for stress levels. In one study there was reduction in stress in the intervention group vs the control²⁸; in the other study the intervention group experienced more stress than the control.³¹

Nine of the NICU studies examined outcomes in both mothers and fathers. In some of the studies, mothers were more likely to experience a decrease in parental role stress and overall stress than fathers.^{29,30} While both parents demonstrated more sensitive interactions with infants (increase in sensitivity and positive regard) and fewer withdrawn behaviours (detachment and flat affect) following video-interactive guidance, fathers demonstrated greater increases in parental bonding.²⁵

Two studies examined more distal parental psychological outcomes. Melnyk et al³⁰ demonstrated a reduction in symptoms of anxiety and depression in mothers after participation in the Creating Opportunities of Parent Empowerment program, whereas there were no differences in fathers' symptoms. The study conducted by Welch et al³³ demonstrated a decrease in depressive and anxiety symptoms in mothers following participation in the FNI.

4.3.4 | Parental satisfaction, knowledge and adaptation

In studies with parental satisfaction measures, parent satisfaction with the interventions were generally favourable.^{23,26,34} Outcomes

included improved communication, collaboration, information,²³ helpfulness of the nursing staff,²⁶ knowledge acquisition,²² adaption to the NICU environment, and increased closeness to the infant.³⁴ Mothers of children hospitalized in the PICU who were part of the intervention group reported higher self-efficacy levels in participatory involvement and overall satisfaction with nursing care.³⁵ Parents who received education about infant pain and comforting infants expressed a stronger preference to be present or involved than parents in the control group.²⁴ Two studies demonstrated no differences in: parental perceptions of care delivered,²¹ confidence levels for caring for their infant, or perceived nursing staff support.³²

4.3.5 | NICU staff perspectives

Nurses, physicians, and other interprofessional team members had positive perceptions of the engagement interventions.^{22,34} They reported more informed parents, increased bonding, open and honest communication, more involvement in decision-making,²² and increased parental presence that enhances the well-being of the newborn.³⁴

4.3.6 | Adult ICUs—Family member feedback on feasibility and acceptability of interventions

Outcomes assessed from the four adult ICU studies mainly focused on feasibility of the interventions. Family members reported higher FCC (collaboration, support, respect) when they were engaged in fundamental care activities such as hair combing, hand massage, or bathing than those who did not.³⁹ Educating families on how to use nutrition diaries prompted them to ask other relevant questions about the care and health of their loved ones.³⁸ Family members found personalized instructions by the nursing staff and family visiting kits to be useful in helping them make sense of the situation and their new role as caregiver.³⁷ Family members were most engaged in the learning process when given information about what to expect in the ICU environment (monitors, alarms, surroundings, etc.) and how to participate in care at the bedside (personal care for the patient such as manicure, lip balm application, or passive range of motion).³⁷ Family maintained journals summarizing patient progress and daily activities were not as helpful as verbal instructions from nurses during interactive care activities.³⁷

5 | DISCUSSION

All except one study³² included in the review described one or more benefits of family engagement interventions. Positive outcomes for family members included increased satisfaction, self-efficacy, empowerment, and desire to be involved in the care of an infant, as well as reduced stress, anxiety, and depression. Positive infant outcomes included shorter length of stay, increased weight.^{23,30} Adult patients

had a lower impact of illness in one study.³⁶ Although the studies were heterogeneous in terms of the intervention studied, measures, and outcomes, overall there is evidence that active family engagement results in positive outcomes for patients and families. The evidence base is stronger in NICU studies than adult ICU studies, and research is lacking in PICUs.

The quality of the evidence for the included studies was predominately moderate to low. Three studies were very low quality^{31,35,39} and three could not be graded because of design.^{34,37,38} Single site study location, lack of intervention fidelity monitoring, inadequate reporting of participant demographics, attrition, lack of statistical control for demographics or unit factors, small sample sizes, lack of randomization, possible intervention contamination, and investigator developed instruments were possible sources of bias in some of the studies. We found that the family samples were predominately female with limited diversity with the exception of two NICU studies. In addition, the majority of the studies were conducted in Western countries, an important consideration, as health care delivery systems, and family definitions, composition, and function may differ in other areas of the world.

There was considerable variability in study design and interventions, making it difficult to compare outcomes. Although more than half the studies included a theoretical framework, few studies made a clear linkage between the theoretical underpinnings and the components of the intervention and expected outcomes. Intervention fidelity was rarely discussed. Family engagement interventions included multiple components, some of which were nurse-led. It was not clear in most studies how nurses enacted intervention delivery. Active family engagement may require more complex interventions to be successful, therefore, clear descriptions of intervention components are needed for replication. Multi-site RCTs of family engagement interventions are a priority for future research to increase the strength of the evidence base for active family engagement interventions, and to support their translation into clinical practice.

Reporting of patient-related outcomes of family engagement interventions was limited.^{21,24,30,36} Satisfaction was used as an outcome in many of the reviewed studies, and while important, it may not fully capture the benefits of engaging in care. Studying outcomes beyond stress, anxiety, and depression, such as resilience, adaptation, well-being, empowerment, and confidence, could yield important data about other potential benefits of engagement. Only two studies looked at the long term outcomes of interventions.^{27,30} There is a need for more research in this area as psychological symptoms for patients and their family members persist after hospitalization.⁴⁴⁻⁴⁶ Understanding the reach of family engagement beyond the ICU may provide important insights.

Exploring the impact of active family engagement based on family role may be important across ICU settings. As parenting roles differ, it is not unexpected that outcomes of family engagement interventions differed for mothers and fathers.^{25,29,30} Fathers may experience greater stress because of demands outside of the NICU environment such as employment, juggling home demands, and caring for other children.^{31,47} Some studies have found that mothers and fathers have

different perspectives, with mothers tending to be more detailed oriented and fathers preferring a global picture of their infant's condition and care.⁴⁸ To our knowledge there are no known studies addressing differences among family members in response to active family engagement interventions in the adult ICU. Further, understanding the effects of family engagement beyond individual patients and family members—the impact on families, remains an important focus for future research.⁴⁹ Studying families presents methodological challenges; however, family-based analysis could result in new opportunities to promote active family engagement in more targeted ways.

Family engagement is posited to occur on a continuum,³ with the current review focused on interventions that involve the family in direct care of the patient. A question remains about how to prepare families to be involved in care. Future research should focus on how family members move along the care continuum and specific ways to increase their motivation and confidence to be part of direct care. The interactive care model⁵⁰ includes an assessment of a family member's capacity to be engaged, and may serve as an important framework to guide the development of individualized family engagement interventions and the implementation of active family engagement in the ICU.

The current review offers a lifespan perspective on active family engagement. In the NICU the practice of FCC has long been supported as a philosophy of care that provides optimal support to infants during hospitalization.⁵¹ Beginning in the 1950s, Bowlby's work highlighted the emotional, psychological, and developmental consequences of keeping mothers and infants apart.⁵² Additional research has elucidated the importance of including all caregivers in infant bonding, expanding earlier maternal-based models to the entire family.⁵³ The opportunity to learn about engagement from other developmental stages of life has been described by others.⁵⁴ Physical touch is highlighted as an important aspect of developmental care in the NICU^{51,52,55}; however, less is known about the importance of touch for adults. Similarly, in adult ICUs emphasis is placed on family engagement in terms of communication and decision-making^{6,14,56} and less is known about communication and decision-making aspects of family engagement in the NICU. A theoretical framework for family engagement requires further development, particularly in the PICU and adult ICU settings. A stronger connection between theory and engagement interventions is needed in future studies. Conceptualizing engagement as a fluid and dynamic phenomenon may lead to the development of family engagement interventions that can be tailored to families in different phases of readiness for involvement in ICU care.

6 | LIMITATIONS

Our review was limited to the English language, and thus, we may not have included papers that contributed to this area of science. However, an important strength of this review is the comprehensiveness of the search with assistance of a health librarian and inclusion of a wide range of databases. We included pilot/feasibility and acceptability studies to add a rich description of engagement interventions. However, these studies are only preliminary and cannot provide any

evidence of outcomes associated with the interventions. There was also a wide variation in the methodology and quality of the intervention studies reviewed. There is the potential risk of reporting bias—negative or non-significant findings may have not been reported in the literature and therefore are not included in this review.

7 | IMPLICATIONS AND RECOMMENDATIONS FOR PRACTICE

In this systematic review, four studies were focused on feasibility indicating that 20% of the literature was early stage intervention work. This suggests that substantial work is needed in the development and evaluation of family engagement interventions, particularly in the PICU and adult ICU settings. Without clear data on the safety and efficacy of involving families in care of the patient it is difficult to develop policies and procedures for practice. More research on who should be engaged, how to engage families, and the outcomes for patients and their families should be the focus of future research.

8 | CONCLUSION

There remains a limited evidence base for active family engagement in the PICU and adult ICU populations. More high quality family engagement interventional studies are needed. This review highlights important directions for active family engagement in ICUs across the lifespan.

9 | IMPACTS

Family engagement is theorized to be an important part of high quality critical care but little is known about how to best engage families in direct patient care in the ICU, particularly in the adult and paediatric practice settings. This review summarizes the existing evidence base of active family engagement interventions in the ICU and highlights heterogeneity in interventions and outcomes. There are multiple opportunities to enhance research on active family engagement in the ICU to improve care for patients and their families.

ORCID

Natalie S. McAndrew <https://orcid.org/0000-0002-8572-6761>

REFERENCES

- Carman KL, Dardess P, Maurer M, et al. Patient and family engagement: a framework for understanding the elements and developing interventions and policies. *Health Aff.* 2013;32:223-231.
- Brown SM, Rozenblum R, Aboumatar H, et al. Defining patient and family engagement in the intensive care unit. *Am J Respir Crit Care Med.* 2015;191:358-360.
- Olding M, McMillan SE, Reeves S, Schmitt MH, Puntillo K, Kitto S. Patient and family involvement in adult critical and intensive care settings: a scoping review. *Health Expect.* 2016;19:1183-1202.

4. Davidson JE, Aslakson R, Long AC, et al. Guidelines for family-centered care in the neonatal, pediatric, and adult ICU. *Crit Care Med.* 2017;45:103-128.
5. Kleinpell R, Zimmerman J, Vermoch KL, et al. Promoting family engagement in the ICU: experience from a national collaborative of 63 ICUs. *Crit Care Med.* 2019;47:1692-1698.
6. Kleinpell R, Heyland DK, Lipman J, et al. Patient and family engagement in the ICU: report from the task force of the World Federation of Societies of Intensive and Critical Care Medicine. *J Crit Care.* 2018; 48:251-256.
7. Ågård AS, Hofhuis JGM, Koopmans M, et al. Identifying improvement opportunities for patient- and family-centered care in the ICU: using qualitative methods to understand family perspectives. *J Crit Care.* 2019;49:33-37.
8. Agency for Healthcare Research and Quality. *Strategies for Patient, Family, and Caregiver Engagement.* Technical Brief. Rockville, MD: Agency for Healthcare Research and Quality; 2020. <https://effectivehealthcare.ahrq.gov/sites/default/files/pdf/patient-engagement-draft-technical-brief.pdf>.
9. Haines KJ, Kelly P, Fitzgerald P, Skinner EH, Iwashyna TJ. The untapped potential of patient and family engagement in the organization of critical care. *Crit Care Med.* 2017;45:899-906.
10. Goldfarb MJ, Bibas L, Bartlett V, Jones H, Khan N. Outcomes of patient- and family-centered care interventions in the ICU: a systematic review and meta-analysis. *Crit Care Med.* 2017;45:1751-1761.
11. Davidson JE, Powers K, Hedayat KM, et al. Clinical practice guidelines for support of the family in the patient-centered intensive care unit: American College of Critical Care Medicine Task Force 2004-2005. *Crit Care Med.* 2007;35:605-622.
12. Mirlashari J, Brown H, Fomani FK, de Salaberry J, Zadeh TK, Khoshkhou F. The challenges of implementing family-centered care in NICU from the perspectives of physicians and nurses. *J Pediatr Nurs.* 2020;50:e91-e98.
13. Umberger E, Canvasser J, Hall SL. Enhancing NICU parent engagement and empowerment. *Semin Pediatr Surg.* 2018;27:19-24.
14. Hamilton R, Kleinpell R, Lipman J, Davidson JE. International facilitators and barriers to family engagement in the ICU: results of a qualitative analysis. *J Crit Care.* 2020;58:72-77.
15. Kiwanuka F, Shayan SJ, Tolulope AA. Barriers to patient and family-centred care in adult intensive care units: a systematic review. *Nurs Open.* 2019; 6, 676-684. <https://doi.org/10.1002/nop2.253>.
16. Hetland B, McAndrew N, Perazzo J, Hickman R. A qualitative study of factors that influence active family involvement with patient care in the ICU: survey of critical care nurses. *Intens Crit Care Nur.* 2018;44:67-75.
17. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 2009;6:e1000100.
18. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan—a web and mobile app for systematic reviews. *Syst Rev.* 2016;5:210. <https://doi.org/10.1186/s13643-016-0384-4>.
19. Cochrane. Data extraction forms. Data-extraction-forms; 2020. Accessed February 13, 2020. <https://dplp.cochrane.org/data-extraction-forms>.
20. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ.* 2008;336:924-926.
21. Byers JF, Lowman LB, Jennifer F, et al. A quasi-experimental trial on individualized, developmentally supportive family-centered care. *J Obstet Gynecol Neonatal Nurs.* 2006;35:105-115.
22. Cooper LG, Gooding JS, Gallagher J, Sternesky L, Ledsky R, Berns SD. Impact of a family-centered care initiative on NICU care, staff and families. *J Perinatol.* 2007;27:S32-S37.
23. De Bernardo G, Svelto M, Giordano M, Sordino D, Riccitelli M. Supporting parents in taking care of their infants admitted to a neonatal intensive care unit: a prospective cohort pilot study. *Ital J Pediatr.* 2017;43:36. <https://doi.org/10.1186/s13052-017-0352-1>.
24. Franck LS, Oulton K, Nderitu S, Lim M, Fang S, Kaiser A. Parent involvement in pain management for NICU infants: a randomized controlled trial. *Pediatrics.* 2011; 128(3), 510-518. <https://doi.org/10.1542/peds.2011-0272>.
25. Hoffenkamp HN, Tooten A, Hall RAS, et al. Effectiveness of hospital-based video interaction guidance on parental interactive behavior, bonding, and stress after preterm birth: a randomized controlled trial. *J Consult Clin Psychol.* 2015;83:416-429.
26. Holditch-Davis D, White-Traut R, Levy J, Williams KL, Ryan D, Vonderheid S. Maternal satisfaction with administering infant interventions in the neonatal intensive care unit. *J Obstet Gynecol Neonatal Nurs.* 2013;42:641-654.
27. Holditch-Davis D, White-Traut RC, Levy JA, O'Shea TM, Geraldo V, David RJ. Maternally administered interventions for preterm infants in the NICU: effects on maternal psychological distress and mother-infant relationship. *Infant Behav Dev.* 2014;37:695-710.
28. Lee T-Y, Wang M-M, Lin K-C, Kao C-H. The effectiveness of early intervention on paternal stress for fathers of premature infants admitted to a neonatal intensive care unit. *J Adv Nurs.* 2013;69:1085-1095.
29. Matricardi S, Agostino R, Fedeli C, Montiroso R. Mothers are not fathers: differences between parents in the reduction of stress levels after a parental intervention in a NICU. *Acta Paediatr.* 2013;102:8-14.
30. Melnyk BM, Feinstein NF, Alpert-Gillis L, et al. Reducing premature infants' length of stay and improving parents' mental health outcomes with the creating opportunities for parent empowerment (COPE) neonatal intensive care unit program: a randomized, controlled trial. *Pediatrics.* 2006;118:e1414-e1427.
31. Noergaard B, Ammentorp J, Garne E, Fenger-Gron J, Kofoed P-E. Fathers' stress in a neonatal intensive care unit. *Adv Neonatal Care.* 2018; 18(5), 413-422.
32. van der Pal S, Maguire C, le Cessie S, Wit J, Walther F, Bruij J. Parental experiences during the first period at the neonatal unit after two developmental care interventions. *Acta Paediatr.* 2007;96:1611-1616.
33. Welch MG, Halperin MS, Austin J, et al. Depression and anxiety symptoms of mothers of preterm infants are decreased at 4 months corrected age with Family Nurture Intervention in the NICU. *Arch Womens Ment Health.* 2016;19:51-61.
34. Sarin E, Maria A. Acceptability of a family-centered newborn care model among providers and receivers of care in a public health setting: a qualitative study from India. *BMC Health Serv Res.* 2019;19: 184. <https://doi.org/10.1186/s12913-019-4017-1>.
35. Kuntaros S, Wichiencharoen K, Prasopkittikun T, Stavorn D. Effects of family-centered care on self-efficacy in participatory involvement in child care and satisfaction of mothers in PICU. *Thai J Nurs Res.* 2007;11:203-213.
36. Black P, Boore JRP, Parahoo K. The effect of nurse-facilitated family participation in the psychological care of the critically ill patient: nurse-facilitated family participation in psychological care. *J Adv Nurs.* 2011;67:1091-1101.
37. Davidson JE, Daly BJ, Agan D, Brady NR, Higgins PA. Facilitated sensemaking: a feasibility study for the provision of a family support program in the intensive care unit. *Crit Care Nurs Q.* 2010;33:177-189.
38. Marshall AP, Wake E, Weisbrodt L, Dhaliwal R, Spencer A, Heyland DK. A multi-faceted, family-centred nutrition intervention to optimise nutrition intake of critically ill patients: the OPTICS feasibility study. *Aust Crit Care.* 2016;29:68-76.
39. Mitchell M, Chaboyer W, Burmeister E, Foster M. Positive effects of a nursing intervention on family-centered care in adult critical care. *Am J Crit Care.* 2009;18:543-552.
40. Cochrane Consumers and Communication Review Group. Cochrane Consumers and Communication Review Group: data synthesis and analysis; 2013. <http://ccrcrg.cochrane.org>. Accessed January 4, 2018.

41. Als H. Toward a synactive theory of development: promise for the assessment and support of infant individuality. *Infant Ment Health J.* 1982;3:229-243.
42. Dunst CJ, Trivette CM. Empowerment, effective helpgiving practices and family-centered care. *Pediatr Nurs.* 1996;22:334-337. 343.
43. Davidson JE. Facilitated sensemaking: a strategy and new middle-range theory to support families of intensive care unit patients. *Crit Care Nurse.* 2010;30:28-39.
44. Davidson JE, Harvey MA. Patient and family post-intensive care syndrome. *AACN Adv Crit Care.* 2016;27:184-186.
45. Fernandes A, Jaeger MS, Chudow M. Post-intensive care syndrome: a review of preventive strategies and follow-up care. *Am J Health Syst Pharm.* 2019;76:119-122.
46. Needham DM, Davidson J, Cohen H, et al. Improving long-term outcomes after discharge from intensive care unit: report from a stakeholders' conference*. *Crit Care Med.* 2012;40:502-509.
47. Ireland J, Khashu M, Cescutti-Butler L, van Teijlingen E, Hewitt-Taylor J. Experiences of fathers with babies admitted to neonatal care units: a review of the literature. *J Neonatal Nurs.* 2016;22:171-176.
48. Reid S, Bredemeyer S, Chiarella M. Integrative review of parents' perspectives of the nursing role in neonatal family-centered care. *J Obstet Gynecol Neonatal Nurs.* 2019;48:408-417.
49. Grzywacz JG, Ganong L. Issues in families and health research. *Fam Relat.* 2009;58:373-378.
50. Drenkard K, Swartwout E, Deyo P, O'Neil MB. Interactive care model: a framework for more fully engaging people in their healthcare. *J Nurs Adm.* 2015;45:503-510.
51. Craig JW, Glick C, Phillips R, Hall SL, Smith J, Browne J. Recommendations for involving the family in developmental care of the NICU baby. *J Perinatol.* 2015;35:S5-S8.
52. van der Horst FCP, van der Veer R. Loneliness in infancy: Harry Harlow, John Bowlby and issues of separation. *Integr Psychol Behav Sci.* 2008;42:325-335.
53. Harrison TM. Family-centered pediatric nursing care: state of the science. *J Pediatr Nurs.* 2010;25:335-343.
54. Davidson JE, Hudson CA. Family-centered care: a reflection. *Crit Care Nurs Clin North Am.* 2020;32:xv-xx.
55. Feeley N, Genest C, Niela-Vilén H, Charbonneau L, Axelin A. Parents and nurses balancing parent-infant closeness and separation: a qualitative study of NICU nurses' perceptions. *BMC Pediatr.* 2016;16:134. <https://doi.org/10.1186/s12887-016-0663-1>.
56. Doucette E, Sanzone L, Albahouth A, De Luca W, Santella G, Wang K. The role of technology in enhancing a family-centred approach to care: navigating nurse-family communication in the ICU. *Can Crit Care Nurs J.* 2019;30:29-34.

How to cite this article: McAndrew NS, Jerofke-Owen T, Fortney CA, et al. Systematic review of family engagement interventions in neonatal, paediatric, and adult ICUs. *Nurs Crit Care.* 2022;27:296-325. <https://doi.org/10.1111/nicc.12564>