

**ARTICLE TYPE****Constrained Data Smoothing via Optimal Control**Assen Dontchev<sup>1</sup> | Ilya V. Kolmanovsky<sup>1</sup> | Trung B. Tran<sup>2</sup><sup>1</sup>Department of Aerospace Engineering, The University of Michigan, Ann Arbor, MI, USA<sup>2</sup>Department of Mechanical Engineering, The University of Michigan, Ann Arbor, MI, USA**Correspondence**

Ilya V. Kolmanovsky - Email: ilya@umich.edu

**Present Address**

1320 Beal Avenue, François-Xavier Bagnoud Aerospace Building, Ann Arbor, MI 48109-2140

**Summary**

The paper considers a problem of best smoothing in a strip, where the objective is to find a function  $f : [0, 1] \rightarrow \mathbb{R}$  that satisfies bilateral constraints on its values,  $d(t) \leq f(t) \leq e(t)$  for all  $0 \leq t \leq 1$  and minimizes a weighted sum of the  $L_2$ -norm of the second derivative and squared deviations from specified values,  $y_i$ , at discrete points  $0 = t_1 < t_2 < \dots < t_{N+2}$ . We assume that constraints  $d(t)$  and  $e(t)$  are continuous functions that are linear in each interval  $[t_i, t_{i+1}]$ ,  $i = 1, \dots, N + 1$ . We connect this problem to a state-constrained optimal control problem for the double integrator, and give conditions for the existence and uniqueness of the solution under which we also show that the solution is a cubic spline with knots at  $t_i$  and no more than two additional knots in each interval  $(t_i, t_{i+1})$ . We propose a numerical algorithm for solving this problem based on a two stage minimization, where the outer loop optimization problem is finite-dimensional and convex, while the inner loop optimization problem admits a solution which is easy to compute. Numerical results that show the efficacy of the proposed approach are reported.

**KEYWORDS:**

Data Smoothing, Optimal Control, State Constraints, Smoothing Spline, Optimal Smoothing

**1 | INTRODUCTION**

Given data  $(t_i, y_i)$  that represent points on the graph of an unknown function, the classical best interpolation problem is to find a function  $f$  with minimal  $L_2$  norm of the second derivative whose values at  $t_i$  equal  $y_i$ . As is well known<sup>1</sup>, the solution of this problem is a natural cubic spline interpolating the data. Dontchev<sup>2</sup> has considered a version of this problem subject to bilateral constraints; namely, the graph of the interpolating function is restricted to a strip defined by two piecewise linear functions across the mesh  $\{t_i\}$ ; it was shown that the solution of this problem is a cubic spline with not more than two additional knots in each interpolation interval  $(t_i, t_{i+1})$ . Later<sup>3</sup> the structure of the solution was further characterized as follows. On each interpolation interval one of the four cases may appear: (i) the solution is in the interior of the strip and no additional knots occur; (ii) there is a single point on one of the constraints where the constraint is active (one additional knot); (iii) there is an interval on one of the constraints where this constraint is active (two additional knots on the same constraint); (iv) there are two single points on each of the constraints where the respective constraints are active (two additional knots on different constraints).

In this paper we consider a problem of *data smoothing* subject to bilateral constraints representing a strip. By applying optimality conditions and duality in optimal control, we obtain a characterization of the solution which is somewhat similar to that of the best interpolation in a strip<sup>2,3</sup>, but there are important differences. Based on that characterization, we propose an algorithm for data smoothing in a strip and illustrate its effectiveness through computer experiments.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/oca.2890

Constrained interpolation and smoothing have been studied from various viewpoints in the past<sup>4,5,6,7</sup> and more recently<sup>8</sup>. Specifically, for constrained smoothing, Irvine et. al.<sup>9</sup> considered the smoothing problem as an extension of constrained interpolation, where the inequality constraint is imposed on the  $k$ th derivative being non-negative, e.g.,  $f^{(k)} \geq 0$ . Elfving and Andersson<sup>10</sup> have studied the problem of constrained smoothing splines, where some inequality constraints were imposed on the first derivative  $f'$  or second derivative  $f''$ . Mammen et. al.<sup>11</sup> proposed a unifying framework in a normed vector space for treating constrained smoothing problems; their work also contained an overview of the progress on constrained data smoothing up to the year 2001. Mammen and Thomas-Agnan<sup>12</sup> then employed the framework of Mammen et. al.<sup>11</sup> to study the problem of constrained smoothing spline interpolation with monotone shape restriction, i.e.,  $f^{(k)} \geq 0$ . Turlach<sup>13</sup> also studied the problem of constrained smoothing spline interpolation with monotone shape restriction and proposed the method which is extendable to multiple simultaneous shape constraints. Kano and Martin<sup>14</sup> studied constrained and optimal smoothing and interpolating splines; the examined constraint types include point, interval, and integrated values. In their work they assumed that the solution function uses the normalized uniform B-splines of some fixed degree  $k$  as the basis functions; our work differs by considering  $f$  being in a Sobolev space.

It has been known for quite awhile that problems in approximation theory have important connections to optimal control<sup>15</sup>, and this connection is also exploited in the present work. In particular, Shen and Wang<sup>16</sup> studied the constrained spline smoothing problem through optimal control; but unlike our work, which deals with the constraints on the state of the reformulated optimal control problem, the work of Shen and Wang deals with the constraint on the control  $u$ . Ikeda et. al.<sup>17</sup> also addressed the problem of constrained spline smoothing using optimal control; however, the constraints were imposed only at the knots, whereas in this paper, we study the case where the bilateral constraints on the function are piecewise linear and imposed pointwise over an interval.

## 2 | PROBLEM SETTING

We consider the following problem:

$$\text{Minimize } 0.5 \left( \|f''\|_2^2 + \sum_{i,j=1}^{N+2} q_{ij}(f(t_i) - y_i)(f(t_j) - y_j) \right) \quad (1)$$

$$\text{subject to } e(t) \leq f(t) \leq d(t) \text{ for all } t \in [0, 1], \text{ and } f \in W^{2,2}[0, 1], \quad (2)$$

where  $0 = t_1 < t_2 < \dots < t_{N+2} = 1$ ,  $N$  is a natural number,  $y_i, i \in \{1, 2, \dots, N+2\}$  are given real numbers,  $\|\cdot\|_2$  is the usual  $L^2$  norm of a function on the interval  $[0, 1]$ , and  $W^{2,2}[0, 1]$  denotes the Sobolev space of functions on  $[0, 1]$  with absolutely continuous first derivatives and second derivative in  $L^2$ . In the problem described by (1)-(2), the first term is the standard in best interpolation problems squared  $L^2$  norm of the second derivative of  $f$ , while the second term represents least squares with weights  $q_{ij}$  of the deviation of values of  $f$  at  $t_j$  from given points  $y_j$ . The functions  $d$  and  $e$  describe a strip where the values of the function  $f$  should belong.

We denote by  $Q \in \mathbb{R}^{(N+2) \times (N+2)}$  the square matrix  $[q_{ij}]$  and we assume that  $Q$  is symmetric and positive semi-definite  $Q = Q^T \geq 0$ . Furthermore, there exist  $i, j, i < j$ , such that

$$\tilde{Q} = \begin{bmatrix} q_{ii} & q_{ij} \\ q_{ji} & q_{jj} \end{bmatrix} > 0. \quad (3)$$

Condition (3) could be satisfied, for instance, by choosing weights  $q_{ii}$  and  $q_{jj}$  at some two points  $t_i$  and  $t_j$  to be positive while selecting the corresponding cross-weights  $q_{ij}$  and  $q_{ji}$  to be zero.

We assume that the functions  $e$  and  $d$  are piecewise linear and continuous across the knots  $t_i$  and such that  $e(t) < d(t)$  for all  $t \in [0, 1]$ . Furthermore, we assume that

$$e(t_j) < y_j < d(t_j), \quad j = 1, \dots, N+2. \quad (4)$$

Indeed, we need to find a best fit of the points  $y_i$  with a function having values in the strip, hence it should be expected that the points  $(t_i, y_i)$  are inside the strip. Without the strip constraint, the solution to this problem is the well-known smoothing cubic spline<sup>1</sup>.

We will now prove the existence and uniqueness of solution by reformulating the problem as an optimal control problem. Let  $x_1 = f, x_2 = f', u = f''$ . Then problem (1)-(2) can be written in the following form:

$$\text{Minimize } J = 0.5 \left( \|u\|_2^2 + \sum_{i,j=1}^{N+2} q_{ij}(x_1(t_i) - y_i)(x_1(t_j) - y_j) \right) \quad (5)$$

$$\text{subject to } \dot{x}_1 = x_2,$$

$$\dot{x}_2 = u,$$

$$\text{and } u \in \mathcal{U} = \{u \in L^2[0, 1] \mid \exists x_1 \text{ with } \ddot{x}_1(t) = u(t) \text{ a.e. } 0 \leq t \leq 1 \text{ and } e(t) \leq x_1(t) \leq d(t) \text{ for all } t \in [0, 1]\}. \quad (6)$$

Note that in the optimal control problem we interpret  $t$  as time and we use dots to designate time derivatives. This is a (non-standard) optimal control problem for the double integrator, where  $u$  is the control and  $x = (x_1, x_2)$  is the state. Note that the inequality state constraints appear in the definition of the feasible set  $\mathcal{U}$  for the control.

The set  $\mathcal{U}$  is clearly convex. It is also closed. Indeed, suppose  $u^n \in \mathcal{U}, u^n \rightarrow \bar{u}$  in  $L_2$  and let  $x_1^n$  be the corresponding  $x_1$ . Note that

$$x_1^n(t) = x_1^n(0) + t \left( x_1^n(1) - x_1^n(0) - \int_0^1 (t - \sigma) u^n(\sigma) d\sigma \right) + \int_0^t (t - \sigma) u^n(\sigma) d\sigma. \quad (7)$$

The sequences of real numbers  $x_1^n(0)$  and  $x_1^n(1)$  are bounded between  $e(0)$  and  $d(0)$  and between  $e(1)$  and  $d(1)$ , respectively; hence we can extract their convergent subsequences to some  $\bar{x}_1(0)$  and  $\bar{x}_1(1)$ , respectively. Without loss of generality, we can assume that the original sequences  $x_1^n(0)$  and  $x_1^n(1)$  converge to  $\bar{x}_1(0)$  and  $\bar{x}_1(1)$  while  $u^n \rightarrow \bar{u}$  in  $L_2$  as  $n \rightarrow \infty$ . Define  $\bar{x}_1$  by boundary conditions  $\bar{x}_1(0)$  and  $\bar{x}_1(1)$  and  $\ddot{\bar{x}}_1 = \bar{u}$ , i.e.,

$$\bar{x}_1(t) = \bar{x}_1(0) + t \left( \bar{x}_1(1) - \bar{x}_1(0) - \int_0^1 (t - \sigma) \bar{u}(\sigma) d\sigma \right) + \int_0^t (t - \sigma) \bar{u}(\sigma) d\sigma. \quad (8)$$

Exploiting Cauchy-Schwartz inequality in the expression for  $|x_1^n(t) - \bar{x}_1(t)|$  formed using (7) and (8), we obtain that  $L^2$  convergence of controls and convergence of boundary conditions implies uniform ( $C^0$ ) convergence of the corresponding state trajectories, i.e.,  $x_1^n \rightarrow \bar{x}_1$  as  $n \rightarrow \infty$ . Thus  $e(t) \leq \bar{x}_1(t) \leq d(t)$  for all  $0 \leq t \leq 1$  and  $\bar{u} \in \mathcal{U}$ .

Based on condition (4), we will now show that there exists a feasible trajectory of the state  $x_1$  whose values at  $t_j$  are in the interior of the strip. Let  $f_0$  be the piecewise linear function whose graph connects the points  $(t_i, y_i)$ . Clearly,  $e(t) < f_0(t) < d(t)$  for all  $t \in [0, 1]$ . For any natural  $m$ , let  $t_{i,j} = t_i + j(t_{i+1} - t_i)/m$ . It is well known that there exists a sequence  $S_m$  of cubic splines interpolating points  $(t_{i,j}, f_0(t_{i,j}))$  that converges to  $f_0$  uniformly in  $[0, 1]$ . Then for  $m$  sufficiently large we will have  $e(t) < S_m(t) < d(t)$  for all  $t \in [0, 1]$ . Thus,  $S_m$  is a feasible trajectory for the state  $x_1$  with values in the interior of the strip, and the second time-derivative of  $S_m$  is a feasible control for which the value of the objective function is finite. This yields that there exists a sublevel set of  $J$  in (5), viewed as a function of  $u$ , which is nonempty; it is obviously closed and convex in  $L^2$ . Since

$$0.5\|u\|_2^2 \leq J(u) \text{ for all } u \in L^2,$$

the sublevel sets are furthermore bounded, hence weakly compact.

The rest of the proof of the existence is standard. The intersections of the nonempty sublevel set with the feasible set  $\mathcal{U}$  is weakly compact too and the minimization over this intersection is equivalent to the minimization over  $\mathcal{U}$ . The objective function is convex and continuous, hence weakly lower semicontinuous. Thus, under the assumption made, problem (5)-(6) has a solution and hence problem (1)-(2) has a solution.

The condition (3) ensures that  $J$  in (5), viewed as a function of  $x_1$ , is strictly convex<sup>1</sup>. This gives us uniqueness of the solution.

We denote the solution by  $(\bar{x}, \bar{u}, \bar{z})$ . In the following section we will show that the solution  $\bar{x}_1$  is in fact a cubic spline with no more than 2 additional knots in each interval  $[t_i, t_{i+1}]$ . We also give a detailed description of the solution together with an algorithm to compute it.

<sup>1</sup>This follows since  $L_2$ -squared norm is strictly convex,  $\bar{Q} > 0, Q \geq 0$  and noting that  $u = \ddot{x}_1^{(1)} = \ddot{x}_1^{(2)}, x_1^{(1)}(t_i) = x_1^{(2)}(t_i), x_1^{(1)}(t_j) = x_1^{(2)}(t_j)$  imply that  $x_1^{(1)}(t) = x_1^{(2)}(t)$  for all  $0 \leq t \leq 1$ . The full details of the proof are analogous to the ones of strict convexity of the function  $\varphi$  in Section 4; we chose not to repeat them here.

### 3 | CHARACTERIZING THE SOLUTION

Suppose  $f \in W^{2,2}[0, 1]$  and  $0 = t_1 < t_2 < \dots < t_{N+2} = 1$  are given. The  $i$ th normalized linear B-spline is defined by<sup>1</sup>:

$$B_i(t) = \begin{cases} 0 & \text{if } t_{i+2} < t, \\ \frac{t_{i+2} - t}{t_{i+2} - t_{i+1}} & \text{if } t_{i+1} < t \leq t_{i+2}, \\ \frac{t_{i+2} - t_{i+1}}{t_{i+2} - t_{i+1}} - \frac{t_{i+1} - t}{t_{i+2} - t_{i+1}} - \frac{t_{i+1} - t}{t_{i+1} - t_i} & \text{if } t_i < t \leq t_{i+1}, \\ 0 & \text{if } t \leq t_i. \end{cases}$$

Integrating by parts, it follows that

$$\begin{aligned} \int_0^1 f''(t)B_i(t)dt &= \int_{t_i}^{t_{i+1}} f''(t)B_i(t)dt + \int_{t_{i+1}}^{t_{i+2}} f''(t)B_i(t)dt \\ &= \left( -f'(t_{i+1}) + \frac{f(t_{i+2}) - f(t_{i+1})}{t_{i+2} - t_{i+1}} \right) + \left( f'(t_{i+1}) - \frac{f(t_{i+1}) - f(t_i)}{t_{i+1} - t_i} \right) \\ &= \frac{f(t_{i+2}) - f(t_{i+1})}{t_{i+2} - t_{i+1}} - \frac{f(t_{i+1}) - f(t_i)}{t_{i+1} - t_i}. \end{aligned}$$

Defining the vector function  $B(t) = (B_i)_{i=1}^N = (B_1(t), \dots, B_N(t))^T$ , where  $B_i$ 's are as above, we have

$$\int_0^1 f''(t)B(t)dt = \begin{pmatrix} \frac{f(t_3) - f(t_2)}{t_3 - t_2} - \frac{f(t_2) - f(t_1)}{t_2 - t_1} \\ \frac{f(t_4) - f(t_3)}{t_4 - t_3} - \frac{f(t_3) - f(t_2)}{t_3 - t_2} \\ \vdots \\ \frac{f(t_{N+2}) - f(t_{N+1})}{t_{N+2} - t_{N+1}} - \frac{f(t_{N+1}) - f(t_N)}{t_{N+1} - t_N} \end{pmatrix} = Kz,$$

where

$$K = \begin{bmatrix} \frac{1}{t_2 - t_1} - \frac{t_3 - t_1}{(t_3 - t_2)(t_2 - t_1)} & \frac{1}{t_3 - t_2} & 0 & \dots & 0 & 0 & 0 \\ 0 & \frac{1}{t_3 - t_2} & -\frac{1}{(t_4 - t_3)(t_3 - t_2)} & \frac{1}{t_2 - t_1} & \dots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \dots & \ddots & 0 \\ 0 & 0 & 0 & 0 & \dots & \frac{1}{t_{N+1} - t_N} - \frac{t_{N+2} - t_N}{(t_{N+2} - t_{N+1})(t_{N+1} - t_N)} & \frac{1}{t_{N+2} - t_{N+1}} \end{bmatrix},$$

$$z = \begin{pmatrix} f(t_1) \\ f(t_2) \\ f(t_3) \\ \vdots \\ f(t_N) \\ f(t_{N+1}) \\ f(t_{N+2}) \end{pmatrix}.$$

Because  $t_1 < t_2 < \dots < t_{N+2}$ , the rows of  $K$  must be linearly independent, i.e.,  $K \in \mathbb{R}^{N \times (N+2)}$  is full row rank.

Let  $y = (y_1, y_2, \dots, y_{N+2})^T$ . As in Section 2, denoting  $x_1 = f, x_2 = f', u = f''$ , problem (1)-(2) can be written in the following form:

$$\begin{aligned} &\text{Minimize with respect to } (x, u, z) \in W^{2,2} \times L^2 \times \mathbb{R}^{N+2} && (9) \\ &\text{the objective function } J(x, z, u) = \frac{1}{2} (\|u\|_2^2 + (z - y)^T Q (z - y)) \end{aligned}$$

$$\begin{aligned}
&\text{subject to } \dot{x}_1 = x_2, \quad x_1(t_i) = z_i, \text{ for } i \in \{1, \dots, N+2\} \\
&\dot{x}_2 = u, \\
&e(t) \leq x_1(t) \leq d(t) \text{ for all } t \in [0, 1], \\
&\int_0^1 u(t)B(t)dt = Kz.
\end{aligned}$$

In this section we employ the duality theory and optimality conditions for state and control constrained optimal control problems developed by Hagger and Mitter<sup>18</sup> and previously applied by Dontchev<sup>2</sup> to the best interpolation in a strip problem. First, observe that, as shown in the preceding section, condition (4) implies that there exists a feasible control  $u$  such that the corresponding trajectory of the state  $x_1$  satisfies  $e(t) < x_1(t) < d(t)$  for all  $t \in [0, 1]$ . This, combined with the surjectivity of the matrix  $K$ , implies that Slater's constraint qualification holds, hence we can apply the Lagrange multiplier rule.

To do that, we introduce the Lagrange functional

$$\begin{aligned}
\mathcal{L}(x, x(0), z, u, \lambda, \nu, \mu) = & J(x, z, u) + \lambda^T \left( \int_0^1 u(t)B(t)dt - Kz \right) + \int_0^1 \left[ x_1(t) - x_1(0) - \int_0^t x_2(s)ds \right] d\nu_1 \\
& + \int_0^1 \left[ x_2(t) - x_2(0) - \int_0^t u(s)ds \right] d\nu_2 + \int_0^1 (e - x_1)d\mu_1 + \int_0^1 (x_1 - d)d\mu_2,
\end{aligned}$$

where  $\lambda = (\lambda_1, \dots, \lambda_N)^T \in \mathbb{R}^N$ ,  $\nu = (\nu_1, \nu_2)$  is a regular Borel measure, and  $\mu = (\mu_1, \mu_2)$  is a nonnegative regular Borel measure such that  $\mu_1$  is supported on the set  $T_1 = \{t \in [0, 1], x_1(t) = e(t)\}$  and  $\mu_2$  is supported on the set  $T_2 = \{t \in [0, 1], x_1(t) = d(t)\}$ . Based on separation of convex sets by a hyperplane, one can show, following similar arguments as by Dontchev<sup>2</sup>, that there exist optimal Lagrange multipliers  $(\bar{\lambda}, \bar{\nu}, \bar{\mu})$  such that

$$\mathcal{L}(\bar{x}, \bar{x}(0), \bar{z}, \bar{u}, \bar{\lambda}, \bar{\nu}, \bar{\mu}) \leq \mathcal{L}(x, \alpha, z, u, \bar{\lambda}, \bar{\nu}, \bar{\mu}) \quad (10)$$

for every continuous function  $x$  with  $x(t) \in \mathbb{R}^2$ ,  $t \in [0, 1]$ , every  $\alpha \in \mathbb{R}^2$ , every  $z \in \mathbb{R}^{N+2}$ , and every  $u \in L^2$ .

Define

$$\bar{p}_1(t) = \int_t^1 d\bar{\nu}_1, \quad \bar{p}_2(t) = \int_t^1 d\bar{\nu}_2. \quad (11)$$

Then  $\bar{p}_1$  and  $\bar{p}_2$  are functions of bounded variations that are continuous from the left. Since the Lagrange functional is convex and Fréchet differentiable, the condition (10) is equivalent to the corresponding optimality condition based on the Fermat rule "derivative equals zero." Differentiating  $\mathcal{L}$  with respect to  $z$  and evaluating at the optimal variables results in

$$Q(\bar{z} - y) + K^T \bar{\lambda} = 0. \quad (12)$$

Next, differentiating  $\mathcal{L}$  with respect to  $x(0) = (x_1(0), x_2(0))$  and evaluating at the optimal variables gives

$$\bar{p}_1(0) = \bar{p}_2(0) = 0.$$

Since

$$\int_0^t \int_1^t \bar{u}(s)dsd\bar{\nu}_2 = \int_0^t \bar{u}(t)\bar{p}_2(t)dt,$$

from the equality  $\partial\mathcal{L}/\partial u = 0$  evaluated at the optimal variables we get

$$\bar{u}(t) + B(t)^T \bar{\lambda} - \bar{p}_2(t) = 0. \quad (13)$$

Furthermore, utilizing the equality

$$\int_0^1 \bar{x}_2 d\bar{\nu}_2 - \int_0^1 \int_0^t dsd\bar{\nu}_2 = \int_0^1 \bar{x}_2 d(-\bar{p}_2 + \int_s^1 \bar{p}_1)$$

and differentiating  $\mathcal{L}$  with respect to  $x_2$  gives

$$\dot{\bar{p}}_2 = -\bar{p}_1, \quad \bar{p}_2(0) = 0. \quad (14)$$

Note that  $\bar{p}_1$  and  $\bar{p}_2$  are the familiar from Pontryagin's Maximum Principle adjoint variables. Finally, differentiating  $\mathcal{L}$  with respect to  $x_1$  results in

$$\bar{p}_1(t) = - \int_t^1 d(\bar{\mu}_2 - \bar{\mu}_1). \quad (15)$$

We conclude from (13) that the optimal control  $\bar{u}$  is an absolutely continuous function which is the sum of a piecewise linear and continuous function across the knots  $t_i$  and an absolutely continuous function whose derivative, as given in (14), is of the form (15) with  $\bar{\mu}_i$ ,  $i = 1, 2$ , being nonnegative regular measures supported on the disjoint subsets  $T_i$  of  $[0, 1]$  where the state constraints are active. In addition, (13),  $\bar{p}_2(0) = 0$ , (11) (implying  $\bar{p}_2(1) = 0$ ), and  $B(0) = B(1) = 0$  give that the optimal control satisfies  $\bar{u}(0) = \bar{u}(1) = 0$ .

Let  $\tau_1, \tau_2$  be such that for the optimal solution  $\bar{x}_1$  satisfies  $e(t) < \bar{x}_1(t) < d(t)$  for all  $t \in (\tau_1, \tau_2)$ . From (15) we have that  $\bar{p}_1(t) = 0$ , hence from (14) the adjoint variable  $\bar{p}_2$  is constant. Thus, from (13), on such an interval  $(\tau_1, \tau_2)$  the optimal control  $\bar{u}(t)$  is a piecewise linear continuous function. Furthermore, in the case when the interval  $(\tau_1, \tau_2)$  is contained in some of the intervals  $(t_j, t_{j+1})$ , the optimal  $\bar{u}$  is a linear function inasmuch  $B_j$  is linear there. Then the optimal  $\bar{x}_2$  is a quadratic polynomial there while the optimal  $\bar{x}_1$  is a cubic polynomial.

Assume that in some interval  $(t_j, t_{j+1})$  there are two points  $\tau_1$  and  $\tau_2$ ,  $t_i \leq \tau_1 < \tau_2 \leq t_{i+1}$  such that  $\bar{x}_1(\tau_1) = e(\tau_1)$  and  $\bar{x}_1(\tau_2) = e(\tau_2)$ . Let  $\zeta(t) = 0$  for  $t \in (\tau_1, \tau_2)$  and  $\zeta(t) = \bar{u}(t)$  for  $t \notin (\tau_1, \tau_2)$ . Note that  $\bar{x}_1(\tau_2) = e(\tau_2)$  and  $\bar{x}_2(\tau_2) = \dot{e}(\tau_2)$  since  $\bar{x}_2 - e$  is maximized at  $\tau_2$ . Apply the control  $\zeta$  to the double integrator with the optimal initial conditions  $\bar{x}_1(0)$  and  $\bar{x}_2(0)$  obtaining trajectories  $\xi = (\xi_1, \xi_2)$ . Clearly,  $\xi_i(t) = \bar{x}_i(t)$  for  $t \in [0, \tau_1]$ . Since  $\zeta(t) = 0$  on  $[\tau_1, \tau_2]$ , we have that  $\xi_i(t) = e(t)$  on  $[\tau_1, \tau_2]$ . Then we also have  $\xi_1(\tau_2) = e(\tau_2) = \bar{x}_1(\tau_2)$  and  $\xi_2(\tau_2) = \dot{e}(\tau_2) = \bar{x}_2(\tau_2)$ . But then  $\xi_i(t) = \bar{x}_i(t)$  for  $t \in [\tau_2, 1]$ ,  $i \in \{1, 2\}$ . Thus, we found a function  $\xi_1(t)$  such that  $\xi_1(t_i) = \bar{x}_1(t_i)$  and also  $\xi_1(t) = 0$  on  $[\tau_1, \tau_2]$ . If we assume that  $\bar{u}(v) = \ddot{x}_1(v) \neq 0$  for some  $v \in (\tau_1, \tau_2)$ , since  $\bar{u}$  is a continuous function, there exists  $\varepsilon > 0$  such that  $\bar{u}(t) = \ddot{x}_1(t) \neq 0$  in  $(v - \varepsilon, v + \varepsilon)$  and then the value of the objective function of (9) will be greater than the value obtained for the function  $\zeta(t) = \xi_1(t)$  and  $\xi_1$ . This contradicts the optimality of  $\bar{u}$ . We conclude that when  $\bar{x}_1(\tau_1) = e(\tau_1)$  and  $\bar{x}_1(\tau_2) = e(\tau_2)$  for  $t_j \leq \tau_1 < \tau_2 \leq t_{j+1}$  then  $\bar{x}_1(t) = e(t)$  for all  $t \in [\tau_1, \tau_2]$ ; moreover,  $\bar{u}(t) = 0$  for all  $t \in [\tau_1, \tau_2]$  as well. Clearly, the same argument works for the lower constraint  $d$ .

The property of the solution  $\bar{x}_1$  we found means that each of the constraints cannot be active in two separate (isolated) points in  $(t_i, t_{i+1})$ . That is, in every  $(t_i, t_{i+1})$  each of the constraints is active either at one point, or on an interval. We will also show that if one of the constraints is active on an interval with positive length in  $[t_i, t_{i+1}]$  the other constraint cannot be active in  $(t_i, t_{i+1})$ . We summarize and complement our findings in the following theorem:

**Theorem 1.** If one of the constraints is active on an interval with positive length in  $[t_i, t_{i+1}]$  the other constraint cannot be active in  $(t_i, t_{i+1})$ . This implies that the solution  $\bar{x}_1$  is a cubic spline with knots at  $t_i$ ,  $i = 1, \dots, N + 2$ , and at most two additional knots in each  $(t_i, t_{i+1})$ ,  $i = 1, \dots, N + 1$ .

*Proof.* Suppose that both constraints are active somewhere in  $[t_i, t_{i+1}]$  and such that one of the constraints is active on an interval with positive length. Let  $\tau_1, \tau_2$  be two additional consecutive knots in  $(t_i, t_{i+1})$  that belong to two different constraints; assume also  $\tau_1 < \tau_2$ . There are four cases to be considered:

1.  $\tau_1$  is the right end of a proper interval where the lower constraint  $e$  is active.
2.  $\tau_1$  is the right end of a proper interval where the upper constraint  $d$  is active.
3.  $\tau_1$  is a single (isolated) point where lower constraint  $e$  is active, meaning that  $\tau_2$  is the left end of a proper interval where the upper constraint  $d$  is active.
4.  $\tau_1$  is a single (isolated) point where the upper constraint  $d$  is active, meaning that  $\tau_2$  is the left end of a proper interval where the lower constraint  $e$  is active.

Consider the first case and let  $\bar{x}_1$  be the solution. Note that  $e(t) < \bar{x}_1(t) < d(t)$  for  $t \in (\tau_1, \tau_2)$ . Since  $\ddot{x}_1(t) = \ddot{e}(t) = 0$  for  $t < \tau_1$  and close to  $\tau_1$ , and since  $\ddot{x}_1$  is continuous on  $(0, 1)$ , it follows that  $\ddot{x}_1(\tau_1) = 0$ .

From (13), (12) and the fact that  $K$  is full rank, we have that

$$\bar{\lambda} = -(KK^T)^{-1}KQ(\bar{z} - y),$$

and for  $t \in (\tau_1, \tau_2)$ ,

$$\ddot{\bar{x}}_1(t) = \dot{B}(t)^T (KK^T)^{-1} KQ(\bar{z} - y) + \int_t^1 d(\bar{\mu}_1 - \bar{\mu}_2). \quad (16)$$

Since  $\dot{B}(t)$  is a constant for  $t \in (t_i, t_{i+1})$ ,  $\bar{\mu}_2(t) = 0$  and  $\bar{\mu}_1(t) \geq 0$  for  $t \in (t_i, \tau_1)$  and  $\bar{\mu}_1(t) = 0$  for  $t \in (\tau_1, \tau_2)$ , it follows that  $\ddot{\bar{x}}_1(t) \geq 0$  for  $t \in (\tau_1, \tau_2)$ . Note that  $\bar{x}_1$  is also a constant on  $(\tau_1, \tau_2)$ . Thus  $\ddot{\bar{x}}_1(t)$  is non-decreasing for  $t \in (\tau_1, \tau_2)$ ,  $\ddot{\bar{x}}_1(\tau_1) = 0$ , and hence  $\ddot{\bar{x}}_1(t) \geq 0$  for all  $t < \tau_2$  that are close to  $\tau_2$ . On the other hand, since  $\ddot{\bar{x}}_1$  is continuous in  $(0, 1)$ , by the Taylor theorem, for every  $t \in (\tau_1, \tau_2)$ , we have

$$\begin{aligned} \bar{x}_1(t) &= \bar{x}_1(\tau_2) + \dot{\bar{x}}_1(\tau_2)(t - \tau_2) + \frac{1}{2} \ddot{\bar{x}}_1(\bar{\tau})(t - \tau_2)^2 \\ &= d(t) + \frac{1}{2} \ddot{\bar{x}}_1(\bar{\tau})(t - \tau_2)^2, \end{aligned}$$

for some  $\bar{\tau} \in [t, \tau_2]$ . This follows since  $\bar{x}_1(\tau_2) = d(\tau_2)$ , and  $\dot{\bar{x}}_1(\tau_2) = \dot{d}(\tau_2)$  as  $\bar{x}_1 - d$  has a maximum at  $t = \tau_2$ , while  $d(t) = d(\tau_2) + \dot{d}(\tau_2)(t - \tau_2)$  for  $t \in [\tau_1, \tau_2]$  since  $d(t)$  is linear. Since  $\bar{x}_1(t) < d(t)$  for  $t \in (\tau_1, \tau_2)$ , there exist time instants  $t < \tau_2$  arbitrary close to  $\tau_2$  where  $\ddot{\bar{x}}_1(t) < 0$ . Consequently, Case 1 cannot occur.

The rest of the cases is analyzed similarly. Therefore, the solution can only have at most two additional knots in  $[t_i, t_{i+1}]$ .  $\square$

#### 4 | REDUCTION TO TWO STAGE MINIMIZATION

Given any  $f \in W^{2,2}[0, 1]$ , let  $\eta(f)$  be a vector with components  $f(t_i)$  and  $\theta(f)$  be a vector with components  $f'(t_i)$ ,  $i = 1, 2, \dots, N + 2$ . Let

$$J(f) = 0.5 \left( \|f''\|_2^2 + (\eta(f) - y)^T Q(\eta(f) - y) \right)$$

and consider the problem (1)-(2) which can be restated as

$$\text{Minimize } J(f) \text{ subject to } f \in W^{2,2}[0, 1], e(t) \leq f(t) \leq d(t). \quad (17)$$

Define  $\varphi(\eta, \theta)$  as the value function of the following optimization problem:

$$\begin{aligned} &\text{Minimize } 0.5 \left( \|f''\|_2^2 + (\eta - y)^T Q(\eta - y) \right) \\ &\text{subject to } f(t_i) = \eta_i, f'(t_i) = \theta_i, i = 1, \dots, N + 2, \\ &\quad e(t) \leq f(t) \leq d(t), 0 \leq t \leq 1, \\ &\quad f \in W^{2,2}[0, 1]. \end{aligned} \quad (18)$$

We let  $\varphi(\eta, \theta) = +\infty$  if a feasible solution to (18) does not exist. Consider the problem:

$$\begin{aligned} &\text{Minimize } \varphi(\eta, \theta) \\ &\text{subject to } \eta \in \mathcal{E} := \{\eta \in \mathbb{R}^{N+2} : e(t_i) \leq \eta_i \leq d(t_i), i = 1, \dots, N + 2\} \text{ and } \theta \in \mathbb{R}^{N+2}. \end{aligned} \quad (19)$$

The solution to (19) exists and is unique. Indeed, let  $f^*$  be the unique solution to (1)-(2), and define  $\eta^* = \eta(f^*)$ ,  $\theta^* = \theta(f^*)$ . If there exists  $f^{**}$  which is feasible for (18) for some  $\eta^{**}, \theta^{**}$  and  $\varphi(\eta^{**}, \theta^{**}) \leq \varphi(\eta^*, \theta^*)$  then

$$J(f^{**}) = \varphi(\eta^{**}, \theta^{**}) \leq \varphi(\eta^*, \theta^*) = J(f^*).$$

Since  $f^{**}$  is feasible for (1)-(2), this is only possible if  $f^{**} = f^*$  and  $\eta^{**} = \eta^*$ ,  $\theta^{**} = \theta^*$ . By an analogous argument, solving problem (18)-(19) should lead to the same solution as solving the original problem (1)-(2).

The problem (19) is a convex finite-dimensional optimization problem. Furthermore, the solution to problem (18) can be easily computed.

For completeness, we demonstrate that the function  $\varphi$  is strictly convex. Let  $\eta^1, \eta^2 \in [e(t_1), d(t_1)] \times \dots \times [e(t_{N+2}), d(t_{N+2})]$  and  $\theta^1, \theta^2 \in \mathbb{R}^{N+2}$ . Let  $f^1$  and  $f^2$  be the solutions to (18) which correspond to  $\eta^1, \theta^1$  and  $\eta^2, \theta^2$ , respectively. Consider  $\lambda \in (0, 1)$

and  $\eta^0 = \lambda\eta^1 + (1-\lambda)\eta^2$ ,  $\theta^0 = \lambda\theta^1 + (1-\lambda)\theta^2$ ,  $f^\lambda = \lambda f^1 + (1-\lambda)f^2$ . Then  $f^\lambda$  is feasible for (18) with  $\eta = \eta^0$ ,  $\theta = \theta^0$ . Let  $f^0$  denote the solution to (18) corresponding to  $\eta^0$ ,  $\theta^0$ . Note that  $2\varphi(\eta, \theta)$  can be decomposed into a sum of a function: A function

$$2\tilde{\varphi}(\eta, \theta) = \|(f)''\|_2^2 + (\tilde{\eta} - \tilde{y})^T \tilde{Q}(\tilde{\eta} - \tilde{y}), \quad (20)$$

and another convex function, where  $\tilde{\eta}$ ,  $\tilde{y}$  are  $2 \times 1$  vectors consisting of  $i$ th and  $j$ th components of  $\eta$  and  $y$ , respectively, and  $i$ ,  $j$  and  $\tilde{Q}$  are identified in (3). Hence it is sufficient to show strict convexity of  $2\tilde{\varphi}(\eta, \theta)$ . From  $\tilde{Q} > 0$ , optimality of  $f^0$  for  $\eta^0$ ,  $\theta^0$  and strict convexity of the  $L_2$  norm squared, we have

$$\begin{aligned} 2\tilde{\varphi}(\eta^0, \theta^0) &= \|(f^0)''\|_2^2 + (\tilde{\eta}^0 - \tilde{y})^T \tilde{Q}(\tilde{\eta}^0 - \tilde{y}) \\ &\leq \|(f^\lambda)''\|_2^2 \\ &\quad + (\lambda(\tilde{\eta}^1 - \tilde{y}) + (1-\lambda)(\tilde{\eta}^2 - \tilde{y}))^T \tilde{Q}(\lambda(\tilde{\eta}^1 - \tilde{y}) + (1-\lambda)(\tilde{\eta}^2 - \tilde{y})) \\ &\leq \lambda\|(f^1)''\|_2^2 + (1-\lambda)\|(f^2)''\|_2^2 \\ &\quad + \lambda(\tilde{\eta}^1 - \tilde{y})^T \tilde{Q}(\tilde{\eta}^1 - \tilde{y}) + (1-\lambda)(\tilde{\eta}^2 - \tilde{y})^T \tilde{Q}(\tilde{\eta}^2 - \tilde{y}), \end{aligned}$$

where the equality is only possible if  $(f^1)''(t) = (f^2)''(t)$  for almost all  $0 \leq t \leq 1$  and  $\tilde{\eta}^1 = \tilde{\eta}^2$ , i.e.,  $f^1(t_i) = f^2(t_i)$ ,  $f^1(t_j) = f^2(t_j)$ . But this implies  $f^1 = f^2$  and hence  $\eta^1 = \eta^2$  and  $\theta^1 = \theta^2$ . This proves strict convexity.

Given a set of values  $\{\eta_i\}_{i=1}^{N+2}$ ,  $\{\theta_i\}_{i=1}^{N+2}$ ,  $\eta = (\eta_1, \dots, \eta_{N+2})^T$  and  $Q \geq 0$ ,  $\varphi(\eta, \theta)$  can be computed as

$$\varphi(\eta, \theta) = 0.5 \left[ \sum_{i=1}^{N+1} \varphi_i(\eta_i, \theta_i, \eta_{i+1}, \theta_{i+1}) \right] + 0.5(\eta - y)^T Q(\eta - y),$$

where  $\varphi_i$  is the value function of the following optimization problem for the interval  $[t_i, t_{i+1}]$ :

$$\text{Minimize } \|f''\|_{L^2[t_i, t_{i+1}]}^2 \quad (21)$$

$$\text{subject to } f(t_i) = \eta_i, f'(t_i) = \theta_i, f(t_{i+1}) = \eta_{i+1}, f'(t_{i+1}) = \theta_{i+1}, \quad (22)$$

$$e(t) \leq f(t) \leq d(t) \text{ for } t_i \leq t \leq t_{i+1}, \text{ and } f \in W^{2,2}[t_i, t_{i+1}].$$

Thus the inner-loop optimization problem (18) can be broken down into  $N + 1$  independent problems. Note that each of these optimization problems is convex in  $W^{2,2}[t_i, t_{i+1}]$  but infinite-dimensional. By the same arguments as in the proof of Theorem 1 (see also a similar treatment in<sup>7</sup>), the solution to (21)-(22) is a cubic spline for which one of the following holds in each interval  $(t_i, t_{i+1})$ :

1. The constraints are not active (Case 0).
2. A constraint is active at a single (isolated) point either on the lower constraint  $e$  (Case 1) or on the upper constraint  $d$  (Case 2). The point where the constraint is active is called a *touching point*.
3. A constraint is active on a proper interval; the segment on the constraint that is active is referred to as the *subarc*. The subarc can be on a lower constraint (Case 3) or on the upper constraint (Case 4).
4. The solution touches lower constraint at a single (isolated) point first and then the upper constraint at a single (isolated) point (Case 5) or the upper constraint first at a single (isolated) point and then the lower constraint at a single (isolated) point (Case 6). Such a solution is said to have a *touching pair* in  $[t_i, t_{i+1}]$

As it will become clear from Section 5, the numerical complexity of solving (21)-(22) primarily stems from Cases 5 and 6 that need to be handled numerically (Cases 0-4 can be solved analytically), and from the objective function of (19) being differentiable almost everywhere yet potentially not everywhere. Strategies for addressing Cases 5 and 6 are further discussed in Section 5 while a plethora of algorithms for nondifferentiable optimization<sup>19</sup> is available for nonsmooth problems.



## 5 | NUMERICAL IMPLEMENTATION AND EXAMPLES

The numerical implementation has been carried out in MATLAB 2021a using the Optimization Toolbox and the Curve Fitting Toolbox. The codebase written for this paper is uploaded to GitLab; its URL is included at the end of this paper.




FIGURE 1 Top: The solution as  $q_{5,5}$  varies in the first example. Bottom: The second derivative of the solution.

The process of numerically solving the problem involves a two stage optimization. In the inner loop, independent problems (21)-(22) are solved by constructing solution candidates for each of 7 cases (if exist) and selecting the one that has the smallest  $L_2$ -norm. In the outer loop, the function  $\varphi$  is minimized in its domain. To reliably distinguish between the touching point and subarc in the numerical implementation, the domain is slightly reduced to  $e(t_i) + \epsilon \leq \eta_i \leq d(t_i) - \epsilon$  where  $\epsilon > 0$  is small.

For the inner loop optimization in Case 0, the solution is a cubic polynomial and its coefficients are determined from the boundary conditions by solving a system of four linear algebraic equations. In Cases 1 and 2, the cubic spline in  $[t_i, t_{i+1}]$  consist of two pieces adjoined at the touching point  $t_i < \tau_{i,1} < t_{i+1}$ . The boundary conditions of (21)-(22), the conditions that  $f(\tau_{i,1})$ ,  $f'(\tau_{i,1})$  match the corresponding values of the constraint and its derivative, and the condition for the continuity of the second derivative of the solution yield a cubic equation for the location of the touching point  $\tau_{i,1}$  and eight linear algebraic equations that determine the coefficients of the two cubic polynomials which constitute the cubic spline. Thus in Cases 1 and 2, up to three solution candidates corresponding to the roots of the cubic polynomial need to be considered. The cubic spline with the subarc in Cases 3 and 4 consist of two cubic polynomials adjoining a linear function from both sides. The equations for the ends of the subarc  $\tau_{i,1}$ ,  $\tau_{i,2}$  and for the coefficients of two cubic polynomials are constructed similarly; they reduce to a system of linear algebraic equations. For Cases 5 and 6, determining the location of the two additional knots ( $\tau_{i,1}$  and  $\tau_{i,2}$ ) reduces to solving two (coupled) multivariate polynomial equations of order five; then the coefficients of the three cubic polynomials are determined by solving a system of linear algebraic equations. For the former, numerical methods are used. In MATLAB, the options are either `fsolve` that uses Newton's method or `vpasolve` which uses both symbolic and numerical manipulations. In the former case, multiple starting points may need to be used to reliably determine all solution candidates (this is implemented in the function `ic_search` of our codebase). While `vpasolve` is able to reliably find solutions, its run time is about 50 times slower than `fsolve` without `ic_search`. A potential alternative solution, that we leave to future work, is to train a neural network offline to compute the locations of additional knots in the touching pair given problem data in  $(t_i, t_{i+1})$  as inputs. Typically, many of the solution candidates are discarded as they do not satisfy the constraints (this check reduces to finding minima of a cubic function) or the conditions informing each Case.

In the outer loop the function  $\varphi$  is minimized. This function is strictly convex and Lipschitz continuous but may be nonsmooth. In MATLAB, `fminsearch` function is available which implements the Nelder–Mead's method which does not rely on the use of the gradient information. The alternative is the use of the function `fmincon` which is intended for smooth problems, but as it relies only on function values for the computations, it can be applied to minimizing our function. Our numerical experiments indicated that `fminsearch` gives a larger cost solution as compared to `fmincon` and, furthermore, with `fminsearch` the second derivative of the solution is not continuous. In addition, `fminsearch` is slower than `fmincon`. Hence our final implementation uses `fmincon`. A plethora of computational methods for nonsmooth optimization exist<sup>19</sup> which could be alternatively used.

Two numerical examples are reported in Figure 1 and Figure 2. In these examples, the starting point  $(\eta^0, \theta^0)$  for for the outer-loop optimization was set so that

$$\eta_i^0 = 0.5(e(t_i) + d(t_i)), \quad \theta_i^0 = 0, \quad i = 1, \dots, N + 2.$$

Both examples were defined on an interval different from  $[0, 1]$  (originally assumed in our analysis) in order to illustrate a more general situation. In the first example, the interval is  $[t_1, t_{N+2}] = [0, 3]$  and  $N = 5$  (7 data points). In the second example, the interval is  $[t_1, t_{N+2}] = [13, 25]$  and  $N = 4$  (6 data points). The constraints  $e$  and  $d$  are indicated by the dashed lines in Figures 1 and Figure 2.

In the examples the matrix  $Q$  coincides with the identity matrix except for one diagonal element which is varied to illustrate the effects of increasing/decreasing the weight corresponding to a given data point; when this weight is set to  $\infty$  it means that  $\eta_j$  is fixed at  $y_j$  and not adjusted in the outer loop optimization (i.e., we replace smoothing by an interpolation constraint at that specific point). Such  $\eta_j$  is indicated by 'o' in Figures 1 and Figure 2. In the first example  $q_{5,5}$  is varied and in the second example  $q_{3,3}$  is varied. The increase in the weight causes the solution to approach closer the specified data point.

The locations of  $y_i$  are indicated by 'x' in Figures 1 and Figure 2 and the additional knots in each interval  $(t_i, t_{i+1})$  are indicated by '\*'. To avoid excessive annotation, the knots located at the intervals' starting and ending points ( $t_i$ 's) are not marked.

Note that the second derivative of the constructed solution is continuous and piecewise linear in both examples, and that the second derivative being zero at  $t_1$  and  $t_{N+2}$  is consistent with our theoretical results.

The computations were carried out on a Lenovo Legion Y520 computer with 16GB of RAM and an Intel Core i7-7700HQ 2.80GHz processor. When `fsolve` without `ic_search` was used, the runtime was several minutes of calculation. This run time could be improved by warm starting (e.g., from an unconstrained interpolating cubic spline).

*Remark 1:* As an alternative approach to our problem, we could consider a discretized approximation of the original problem. Specifically, suppose we discretize the interval  $[t_0, t_{N+2}]$  into  $m$  equally spaced sub-intervals, where  $m \gg N + 2$ , with the sub-interval end points being  $\{\tau_k\}_{k=0}^m$ . Then we could consider two strategies.




FIGURE 2 Top: The solution as  $q_{3,3}$  varies in the second example. Bottom: The corresponding second derivative.

The first and fairly standard strategy would be to treat  $u$  as a constant in interval  $[\tau_k, \tau_{k+1}[$  and only impose constraints on  $x_1$  at  $\tau_k$ 's. The problem with this formulation is that  $u$  is piecewise constant and may not be continuous, which is not consistent with our theoretical results derived for the original infinite-dimensional optimization problem. Specifically, we know that the optimal  $u$  must be continuous on  $[t_0, t_{N+2}]$  and satisfy  $u(t_0) = u(t_{N+2}) = 0$ . The second formulation tries to address this by treating the third derivative  $v = \ddot{x}_1$  of  $x_1$  as constant in each sub-interval  $[\tau_k, \tau_{k+1}[$ ; similarly, we only impose the constraints on  $x_1$  at  $\tau_k$ 's.

It can be easily shown that both discretized approximation approaches reduce to quadratic programming (QP) problems. Such problems can then be solved using QP solvers.

The main issue with the above discretized approximation approaches is that the constraints are only imposed and enforced at the points,  $\tau_k$ . Hence, unlike with the approach proposed in this paper, “intersample” constraint violations could occur. Additionally, the QP problem can be large-dimensional and not trivial to solve if fine discretization is used. At the same time, if the

QP problem could be solved cheaply (e.g., for not very fine discretization) then an approximate solution through the discretization and quadratic programming could be of use for warm-starting, i.e., for providing an initial guess to both the outer loop and the inner loop optimizers described in this paper. We leave the comprehensive investigation of potential synergies between the approach in this paper and alternative approaches based on discretization and quadratic programming to a future publication.

## 6 | CONCLUSIONS

In this paper we considered a problem of best smoothing in a strip and connected this problem to a state-constrained optimal control problem for the double integrator. Conditions for the existence and uniqueness of the solution were given under which the solution was shown to be a cubic spline with no more than two additional knots in each interval between the given data points. A numerical algorithm for solving this problem based on a two stage minimization was proposed; its efficacy was illustrated using numerical examples.

## ACKNOWLEDGMENTS

Assen Dontchev and Ilya Kolmanovsky acknowledge the support of AFOSR under the grant number FA9550-20-1-0385 to the University of Michigan. Trung Tran acknowledges the support of the Vingroup Science and Technology Scholarship Program.

## DATA SHARING


The data that support the findings of this study are openly available in the *Best Interpolation and Data Smoothing in a Strip* repository at <https://gitlab.eecs.umich.edu/trungbt/best-interpolation-in-a-strip>, reference number<sup>20</sup>. The numerical examples can be found in the \plots\Case Study II folder in the codebase.

## References

1. de Boor C. *A Practical Guide to Splines*. Applied Mathematical Sciences, Springer, New York . 2001.
2. Dontchev AL. Best interpolation in a strip. *Journal of approximation theory* 1993; 73(3): 334–342.
3. Dontchev AL, Kolmanovsky IV. Best interpolation in a strip II: Reduction to unconstrained convex optimization. *Computational Optimization and Applications* 1996; 5(3): 233–251.
4. Dontchev AL, Qi H, Qi L. Quadratic Convergence of Newton’s Method for Convex Interpolation and Smoothing.. *Constructive Approximation* 2003; 19(1).
5. Micchelli CA, Utreras FI. Smoothing and interpolation in a convex subset of a Hilbert space: II. The semi-norm case. *ESAIM: Mathematical Modelling and Numerical Analysis-Modélisation Mathématique et Analyse Numérique* 1991; 25(4): 425–440.
6. Schmidt JW. A Dual Method for Smoothing Histograms Using Nonnegative C1-Splines. In: Springer. 1992 (pp. 317–329).
7. Dontchev A, Kolmanovsky I. State constraints in the linear regulator problem: a case study. In: . 1. IEEE. ; 1994: 949-950 vol.1
8. Rozhenko AI, Fedorov EA. On an algorithm of bilateral restrictions smoothing with spline. *Sibirskii Zhurnal Vychislitel’noi Matematiki* 2016; 19(3): 331–342.
9. Irvine LD, Marin SP, Smith PW. Constrained interpolation and smoothing. *Constructive approximation* 1986; 2(1): 129–151.

10. Elfving T, Andersson LE. An algorithm for computing constrained smoothing spline functions. *Numerische Mathematik* 1987; 52(5): 583–595.
11. Mammen E, Marron J, Turlach B, Wand M. A general projection framework for constrained smoothing. *Statistical Science* 2001; 16(3): 232–248.
12. Mammen E, Thomas-Agnan C. Smoothing splines and shape restrictions. *Scandinavian Journal of Statistics* 1999; 26(2): 239–252.
13. Turlach BA. Shape constrained smoothing using smoothing splines. *Computational Statistics* 2005; 20(1): 81–104.
14. Kano H, Fujioka H, Martin CF. Optimal smoothing and interpolating splines with constraints. *Applied Mathematics and Computation* 2011; 218(5): 1831–1844.
15. Egerstedt M, Martin C. *Control Theoretic Splines: Optimal Control, Statistics, and Path Planning*. Princeton Series in Applied Mathematics Princeton University Press . 2009.
16. Shen J, Wang X. A constrained optimal control approach to smoothing splines. In: IEEE. ; 2011: 1729–1734.
17. Ikeda T, Nagahara M, Chatterjee D, Srikant S. Constrained Smoothing Splines by Optimal Control. *IEEE Control Systems Letters* 2021; 6: 1298–1303.
18. Hager WW, Mitter SK. Lagrange duality theory for convex control problems. *SIAM Journal on Control and Optimization* 1976; 14(5): 843–856.
19. Bagirov A, Gaudioso M, Karmitsa N, Mäkelä M, Taheri S. *Numerical Nonsmooth Optimization: State of the Art Algorithms*. Springer International Publishing . 2020.
20. Dontchev A, Kolmanovsky I, Tran T. Best Interpolation and Data Smoothing in a Strip. <https://gitlab.eecs.umich.edu/trungbt/best-interpolation-in-a-strip>; .

## AUTHOR BIOGRAPHY



### **Asen L. Dontchev.**

Asen L. Dontchev received the Ph.D. degree in control from the Warsaw University of Technology, Warsaw, Poland, in 1974, and the D.Sc. degree in mathematical sciences from the Bulgarian Academy of Sciences, Sofia, Bulgaria, in 1987. He was an Associate Editor for Mathematical Reviews from 1990 to 2019. He was also an Adjunct Associate Research Scientist in the Department of Mathematics and the Department of Aerospace Engineering at the University of Michigan, Ann Arbor, MI, USA. He was the author of the books *Perturbations, Approximations and Sensitivity Analysis of Optimal Control Systems* (Springer-Verlag, 1983), *Well-Posed Optimization Problems* (Springer-Verlag, 1993) with T. Zolezzi, and

*Implicit Functions and Solution Mappings* (Springer, 2014) with R. T. Rockafellar. His research interests included estimating the effect of perturbations and approximations on solutions of variational problems, in particular problems in optimization, control, differential equations, numerical analysis, and approximation theory. Dr. Dontchev had served as a Program Director at the Division of Mathematical Sciences of NSF and in the Boards of several journals, including the *Journal of Mathematical Analysis and Applications*, *SIAM Journal on Control and Optimization*, *SIAM Journal on Optimization*, *Computational Optimization and Applications*, and *Journal of Optimization Theory and Applications*.





**Ilya V. Kolmanovsky.** Ilya V. Kolmanovsky is a professor in the department of aerospace engineering at the University of Michigan, with research interests in control theory for systems with state and control constraints, and in control applications to aerospace and automotive systems. He received his Ph.D. degree in aerospace engineering from the University of Michigan in 1995. Prof. Kolmanovsky is a Fellow of IEEE and is named as an inventor on 104 United States patents.



**Trung B. Tran** Trung B. Tran received his B.S. in Mechanical Engineering and B.A. in Mathematics from Bucknell University, Lewisburg, PA, USA in 2020. For five years during his time at Bucknell, he was a student research assistant for the Department of Mechanical Engineering and the Department of Mathematics. He is currently pursuing a master's degree in mechanical engineering at the University of Michigan, Ann Arbor, MI, USA with a focus in control engineering.


**How to cite this article:** Dontchev AL, Kolmanovsky IV, and Tran TB (2022), Constrained Data Smoothing via Optimal Control, *Optimal Control Applications and Methods*, 2022: volume....









Example 1: Double Derivative of Smooth Spline




## Example 2: Smooth Spline




Example 2: Double Derivative of Smooth Spline




oca\_2890\_example\_2\_mini\_3\_second\_der.eps



Example 1: Double Derivative of Smooth Spline



# Example 2: Smooth Spline



Example 2: Double Derivative of Smooth Spline

