

Hill Sophie F. (Orcid ID: 0000-0002-8331-9812)

Revised July 12, 2022.

Genetic interaction between *Scn8a* and potassium channel genes *Kcna1* and *Kcnq2*

Sophie F. Hill^{1,2}, Julie Ziobro³, Paymaan Jafar-Nejad⁴, Frank Rigo⁴, and Miriam H. Meisler^{1,2,5}

Author affiliations:

¹Neuroscience Graduate Program, University of Michigan, Ann Arbor MI 48109, USA

sfhill@umich.edu Orcid 0000-0002-8331-9812

²Department of Human Genetics, University of Michigan, Ann Arbor MI 48109, USA

³Department of Pediatrics, University of Michigan, Ann Arbor MI 48109, USA

ziobroj@med.umich.edu

⁴Ionis Pharmaceuticals, Carlsbad CA 92010, USA

pjafarne@ionisph.com and FRigo@ionisph.com

⁵Department of Neurology, University of Michigan, Ann Arbor MI 48109, USA

meislerm@umich.edu Orcid 0000-0001-9107-706X

Correspondence to: Miriam Meisler, PhD

Department of Human Genetics

4909 Buhl Box 5618

University of Michigan Medical School

Ann Arbor MI 48109-5618

meislerm@umich.edu

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1111/epi.17374](https://doi.org/10.1111/epi.17374)

This article is protected by copyright. All rights reserved.

ABSTRACT

Voltage-gated sodium and potassium channels regulate the initiation and termination of neuronal action potentials. Gain-of-function mutations of sodium channel *Scn8a* and loss-of-function mutations of potassium channels *Kcna1* and *Kcnq2* increase neuronal activity and lead to seizure disorders. We tested the hypothesis that reducing expression of *Scn8a* would compensate for loss-of-function mutations of *Kcna1* or *Kcnq2*. *Scn8a* expression was reduced by administration of an antisense oligonucleotide (ASO). This treatment lengthened survival of the *Kcn1a* and *Kcnq2* mutants, and reduced seizure frequency in the *Kcnq2* mutant mice. These observations suggest that reduction of *SCN8A* may be therapeutic for genetic epilepsies resulting from potassium channel mutations.

Keywords: epilepsy; sodium channel; potassium channel; ASO; therapy

INTRODUCTION

Sodium and potassium channels concentrated at the axon initial segment (AIS) regulate the generation of neuronal action potentials.^{1,2} Action potentials are initiated by excitatory stimuli that activate voltage-gated sodium channels, permitting influx of sodium ions.² Subsequent activation of voltage-gated potassium channels permits the exit of potassium ions and repolarizes the neuron.² Potassium channels encoded by *KCNQ2* and *KCNQ3* modulate subthreshold changes in membrane potential that influence neuronal excitability.^{3,4} In genetic epilepsies, elevated sodium channel activity or reduced potassium channel activity can result in excess neuronal firing.^{5,6} Emerging genetic therapies for seizure disorders include reduction of sodium channel expression⁷ or elevation of potassium channel expression.^{8,9} Here we test the hypothesis that reduced expression of an AIS-concentrated sodium channel can compensate for loss of potassium channel activity.

Loss-of-function mutations of the potassium channel gene *KCNQ2* are a major cause of developmental and epileptic encephalopathy (DEE),^{3,4} and loss of function mutations of potassium channel *KCNA1* are responsible for episodic ataxia type 1 and rare cases of DEE.⁶ We used an antisense oligonucleotide (ASO) to reduce expression of the sodium channel gene *Scn8a* in mouse models of *Kcnq2* and *Kcna1* epilepsy. *Scn8a* transcripts were reduced to 50% of wildtype level by administration of the ASO. Transcript levels return to normal 6 weeks after a single injection.⁷ Our observations suggest that specific reduction of *SCN8A* expression may be a useful therapy for disorders of these potassium channels.

MATERIAL AND METHODS.

Mice *Kcnq2^{fl/fl}* mice¹⁰ on strain C57BL/6J were provided by Dr. Anastasios Tzingounis, University of Connecticut. *Kcna1^{+/-}* mice¹¹ on a mixed Black Swiss genetic background (Tac:N:NIHS-BC) were provided by Dr. Edward Glasscock, Southern Methodist University. *Emx1-Cre* mice on strain C57BL/6J were purchased from the Jackson Laboratory (Jax #005628). Both male and female mice were used for the experiments. Mice were housed and cared for in accordance with NIH guidelines in a 12/12-hour light/dark cycle with standard mouse chow and water available ad libitum. All experiments were approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Michigan.

Antisense oligonucleotides

The twenty bp *Scn8a* gapmer ASO (5' GACGA TTAGT GACAT AGGCT 3'), synthesized by Ionis Pharmaceuticals, is complementary to the 3' UTR of the mouse *Scn8a* transcript.⁷ The control ASO (5' CCTAT AGGAC TATCC AGGAA 3') does not match any mouse transcript and is well tolerated in vivo.⁷ ASOs were diluted in phosphate-buffered saline (PBS) for injection. At postnatal day 2 (P2), mutant mice received 45 ug ASO in 2 uL by intracerebroventricular (ICV) injection into the left lateral ventricle. Adult mice were anesthetized with isoflurane and received 100 ug ASO in a 3 uL manual ICV injection into the left lateral ventricle without a guide cannula, as described.¹²

EEG recording

Screw electrodes were implanted in *Kcnq2^{fl/fl},Emx1-Cre* mice at postnatal week 6 (n=2) and in *Kcna1^{+/-}* mice at postnatal week 20 (n=2). For surgery, mice were anesthetized with isoflurane and placed in a stereotaxic adapter. Bilateral screw electrodes were placed in the skull at approximately AP=-2.1, ML= +/-1.7 and a common reference electrode was placed over the cerebellum (approximately AP=-6.0, ML=0). The electrodes were connected to a 6-pin electrode pedestal and the headcap was secured using dental cement. After 1-7 days of recovery, simultaneous EEG recording and video monitoring were performed with a Natus recording system continuously for a minimum of 24 hours and a maximum of 14 days. Signals were acquired at 256 Hz. Data were filtered with a 70Hz low pass filter and 1 Hz high pass

filter. Seizures and interictal background were assessed manually by an experienced reader. Seizures were defined as a sudden burst of electrographic activity consisting of rhythmic spike-and-wave discharges lasting >10 seconds and evolving in frequency and amplitude. Interictal epileptiform discharges were not quantified.

Phenotypes

Hindlimb claspings in *Kcnq2* mutant mice was video recorded at P21. Mice were suspended by the tail for one minute on three consecutive days. Videos were analyzed frame by frame to quantify the percent of time spent in hindlimb claspings. Myoclonic jerks in *Kcna1* mice were counted during a 5 minute observation period at 11 AM on three successive days by two independent observers blinded to genotype.

Data availability

The authors confirm that the data supporting the findings of this study are available within the article.

RESULTS

***Kcnq2* mutant mice**

KCNQ2 encodes the AIS-localized potassium channel Kv7.2 which regulates sub-threshold neuronal excitability and resting membrane potential.^{3,4} Loss-of-function mutations of human *KCNQ2* cause a spectrum of disorders ranging from benign familial neonatal seizures to DEE, and it is the second most common gene mutated in DEE.^{3,4}

Deletion of *Kcnq2* in forebrain excitatory neurons in *Kcnq2^{fl/fl},Emx1-Cre* mice results in neuronal hyperexcitability and spontaneous seizures, as seen in human *KCNQ2* disorders.¹⁰ The seizures and premature death in these mice provide a useful endpoint for therapeutic intervention. We treated *Kcnq2^{fl/fl},Emx1-Cre* mice with 45 ug of *Scn8a* ASO or control ASO by intracerebroventricular (ICV) injection at postnatal day 2 (P2) (Fig 1A). After treatment with control ASO ($n = 12$), the first death was observed at 3 weeks of age, and 50% lethality was reached by 8 weeks of age (Fig 1B-C). In mice receiving the *Scn8a* ASO, the first death was

delayed to 11 weeks of age, and 50% lethality was not reached until 15 weeks. A second dose of 100 ug ASO at 8 weeks of age further delayed the age of 50% lethality to 19 weeks (Fig 1B-C). Overall, 65% to 75% of mice exhibited premature lethality in mice receiving the *Scn8a* ASO and the control ASO (Chi-square test of proportions, $\chi^2=0.36$, $df=2$, $p=0.83$). Reduction of *Scn8a* expression thus extended the lifespan of mice with *Kcnq2* deficiency but did not prevent premature lethality.

The frequency of spontaneous seizures in untreated *Kcnq2^{fl/fl},Emx1-Cre* mice was 40 seizures in 67 days of observation^{10,13}. After treatment with *Scn8a* ASO, we did not observe any seizures or abnormal EEG activity in 16 days of observation ($p<0.0001$, χ^2 test) (Fig 1D). However, the treated mice continued to exhibit hunched posture, scruffy appearance and hindlimb clasp (Fig 1E). Reduction of *Scn8a* expression thus prevented seizures and delays premature death but does not correct the other neurological phenotypes in *Kcnq2* mutant mice. Postnatal growth of *Kcnq2^{fl/fl},Emx1-Cre* mice did not differ from wildtype littermates.

***Kcna1* mutant mice**

KCNA1 encodes Kv1.1, an inward rectifier potassium channel concentrated at the AIS.⁶ Heterozygous loss-of-function mutations of *KCNA1* cause severe epilepsy and episodic ataxia type 1 that can be accompanied by seizures.⁶ Homozygous *Kcna1^{-/-}* null mice are a model of sudden unexpected death in epilepsy (SUDEP).¹¹

Kcna1^{-/-} mice were treated with 45 ug *Scn8a* ASO or control ASO by ICV injection at P2 (Fig 2A). Mice that received control ASO at P2 ($n = 10$) died between 2 and 6 weeks of age with 50% lethality by 1 month of age (Fig 2 B,C). A single dose of *Scn8a* ASO extended median lifespan to 3 months ($n = 9$) (Fig 2 B,C). To evaluate repeated treatments, additional doses of 100 ug ASO were administered at monthly intervals between 1 and 4 months of age (Fig 2A). The repeated treatments delayed the earliest death to 13 weeks ($n = 9$), with less than 50% lethality at 6 months of age (Fig 2 B,C). The penetrance of premature lethality was

reduced from 70% (control ASO) to 20% (*Scn8a* ASO) (Chi-square test of proportions, $\chi^2=5.05$, $df=2$, $p=0.025$).

Spontaneous seizures in untreated *Kcna1*^{-/-} mice were reported to begin at 3 weeks of age¹¹ and occur 5-20 times per day with diurnal variation in frequency.^{11,14} We implanted electrodes at 20 weeks of age in *Kcna1* null mice that had been treated with five doses of ASO starting at P2 (Fig 2A). Electrographic and electro-clinical seizures were detected with an average frequency of 13 seizures per day (Fig 2 D).

Kcna1^{-/-} mice also exhibit hunched posture, scruffy appearance, and impaired growth that was not corrected by the ASO (Figure 2E). *Kcna1*^{-/-} mice exhibit myoclonic jerks that were not corrected by ASO treatment (Figure 2F). Thus, reduction of *Scn8a* expression prolonged survival of *Kcna1*^{-/-} mice but did not completely rescue the neurological abnormalities.

DISCUSSION

Reduction of *Scn8a* expression had therapeutic benefits in two models of epilepsy caused by mutation of potassium channels localized at the AIS. *KCNQ2* is one of the most commonly mutated genes in childhood-onset epilepsy.^{3,4} Reduced *Scn8a* extended the lifespan of *Kcnq2* mutant mice by 11 weeks and restored normal EEG activity but did not eliminate other neurological abnormalities. In the *Kcn1a* null mice, reduction of *Scn8a* expression significantly prolonged lifespan. The positive effect of reducing *Scn8a* is consistent with the therapeutic effectiveness of sodium channel blockers in some individuals with *KCNQ2* mutations³ and in improving the symptoms of episodic ataxia in many individuals with null mutations of *KCNA1*.¹⁵ Nav1.6 is a major source of persistent current in mammalian brain, and reduction of persistent current led to reduced seizures in an *in silico* model of Kv1 deficiency.¹⁶ The effectiveness of the ASO demonstrates that specific reduction of *SCN8A* without changes in other sodium channels can be therapeutic for potassium channel deficiency.

Our results add to a growing body of evidence demonstrating genetic interactions between genes encoding ion channels.^{5,17} The calcium channel genes *Cacna1a* and *Cacna1g*

have been identified as modifiers of *Kcna1* epilepsy¹⁸ and Dravet syndrome.¹⁹ Quantitative variation in the GABA receptor subunit *Gabra2* modifies the severity of *Scn8a* epilepsy and Dravet syndrome.⁵ The current work demonstrates that quantitative variation in *Scn8a* modifies the severity of *Kcna1* and *Kcnq2* epilepsy in mouse models. *Scn8a* is also a modifier of Dravet syndrome.⁷ *Scn2a* modifies the severity of *Kcna1*²⁰ and *Kcnq2* epilepsy.²¹ In a human pedigree, an allelic variant of *SCN1A* was recently found to modify the severity of a null allele.²² Neuronal excitability is evidently influenced by overall ion channel composition, and compensatory modulation of channel genes can restore the balance of excitation and inhibition that is disrupted in epilepsy.

The effects of *Scn8a* modulation shown here suggest that specific reduction of *Scn8a* might replace non-specific sodium channel blockers in treating epilepsies caused by *KCNQ2* and *KCNA1*. In contrast to heterozygous affected patients, the mouse models studied here were homozygous for the potassium channel deficits, and the *Kcnq2* deficiency was expressed only in forebrain excitatory neurons. The side effects associated with non-specific sodium channel blockers are likely to be reduced by specific reduction of *SCN8A*, for example with an Nav1.6-specific drug such as NBI-921352,²³ which is currently in phase 1 clinical trial for treatment of patients with gain-of-function mutations of *Scn8a* (<https://clinicaltrials.gov/ct2/show/NCT03467100>). Genetic therapies such as viral delivery of shRNA or CRISPR/Cas9 knockout may also be applicable to the potassium channel disorders.

ACKNOWLEDGEMENTS

We are grateful to Dr. Anastasios Tzingounis for providing *Kcnq2^{fl/fl}* mice and Dr. Edward Glasscock for providing *Kcna1^{+/-}* mice. We thank Sanjna Chalasani and Aparna Sumanth for technical assistance.

FUNDING

Supported by NINDS Grant RO1 NS34509 from the National Institutes of Health and a grant from the Rackham School of Graduate Studies, University of Michigan.

Conflict of interest statement

P.J-N. and F. R. are employed by Ionis Pharmaceuticals, a for-profit company that develops ASO therapies. The remaining authors have no conflict of interest to disclose.

Ethics Statement: We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

Author contributions: S.H., M.M., P.J-N., and F.R. contributed to the conception and design of the study; S.H. and J.Z. contributed to acquisition and analysis of the data; S.H. and J.Z. contributed to preparation of figures; S.H. and M.M. drafted the manuscript.

References

1. Huang CY, Rasband MN. Axon initial segments: structure, function, and disease. *Ann NY Acad. Sci.* 2018;1420(1):46-61. doi:10.1111/nyas.13718
2. Bean BP. The action potential in mammalian central neurons. *Nat Rev Neurosci.* 2007;8(6):451-65. doi:10.1038/nrn2148
3. Miceli F, Soldovieri MV, Joshi N, Weckhuysen S, Cooper E, Tagliatela M. KCNQ2-Related Disorders. *GeneReviews®*. Published online September 27, 2018. <https://www.ncbi.nlm.nih.gov/books/NBK32534/>
4. Weckhuysen S, Mandelstam S, Suls A, et al. KCNQ2 encephalopathy: emerging phenotype of a neonatal epileptic encephalopathy. *Ann Neurol.* 2012;71(1):15-25. doi:10.1002/ana.22644
5. Meisler MH, Hill SF, Yu W. Sodium channelopathies in neurodevelopmental disorders. *Nat Rev Neurosci.* 2021;22(3):152-166. doi:10.1038/s41583-020-00418-4
6. D'adamo MC, Liantonio A, Rolland JF, Pessia M, Imbrici P. Kv1.1 Channelopathies: Pathophysiological Mechanisms and Therapeutic Approaches. *Int J Mol Sci.* 2020;21(8):2935. doi:10.3390/IJMS21082935
7. Lenk GM, Jafar-Nejad P, Hill SF, et al. Scn8a Antisense Oligonucleotide Is Protective in Mouse Models of SCN8A Encephalopathy and Dravet Syndrome. *Ann Neurol.* 2020;87(3):339-346. doi:10.1002/ana.25676
8. Colasante G, Qiu Y, Massimino L, et al. In vivo CRISPRa decreases seizures and rescues cognitive deficits in a rodent model of epilepsy. *Brain.* 2020;143(3):891-905. doi:10.1093/brain/awaa045
9. Snowball A, Chabrol E, Wykes RC, et al. Epilepsy gene therapy using an engineered potassium channel. *J Neurosci.* 2019;39(16):3159-3169. doi:10.1523/JNEUROSCI.1143-18.2019
10. Soh H, Pant R, LoTurco JJ, Tzingounis A V. Conditional deletions of epilepsy-associated KCNQ2 and KCNQ3 channels from cerebral cortex cause differential effects on neuronal excitability. *J Neurosci.* 2014;34(15):5311-5321. doi:10.1523/JNEUROSCI.3919-

13.2014

11. Smart SL, Lopantsev V, Zhang CL, et al. Deletion of the K(V)1.1 potassium channel causes epilepsy in mice. *Neuron*. 1998;20(4):809-819. doi:10.1016/s0896-6273(00)81018-1
12. Kim HY, Lee DK, Chung BR, Kim H V., Kim Y. Intracerebroventricular Injection of Amyloid- β Peptides in Normal Mice to Acutely Induce Alzheimer-like Cognitive Deficits. *J Vis Exp*. 2016;(109):53308. doi:10.3791/53308
13. Aiba I, Noebels JL. Kcnq2/Kv7.2 controls the threshold and bi-hemispheric symmetry of cortical spreading depolarization. *Brain*. 2021;144(9):2863-2878. doi:10.1093/BRAIN/AWAB141
14. Fenoglio-Simeone KA, Wilke JC, Milligan HL, Allen CN, Rho JM, Maganti RK. Ketogenic diet treatment abolishes seizure periodicity and improves diurnal rhythmicity in epileptic Kcna1-null mice. *Epilepsia*. 2009;50(9):2027-2034. doi:10.1111/J.1528-1167.2009.02163.X
15. Lauxmann S, Sonnenberg L, Koch NA, et al. Therapeutic Potential of Sodium Channel Blockers as a Targeted Therapy Approach in KCNA1-Associated Episodic Ataxia and a Comprehensive Review of the Literature. *Front Neurol*. 2021;12:703970. doi:10.3389/FNEUR.2021.703970/FULL
16. Du J, Vegh V, Reutens DC. Persistent sodium current blockers can suppress seizures caused by loss of low-threshold D-type potassium currents: Predictions from an in silico study of Kv1 channel disorders. *Epilepsia Open*. 2020;5(1):86. doi:10.1002/EPI4.12379
17. Meisler MH, O'Brien JE, Sharkey LM. Sodium channel gene family: epilepsy mutations, gene interactions and modifier effects. *J Physiol*. 2010;588(Pt 11):1841. doi:10.1113/JPHYSIOL.2010.188482
18. Glasscock E, Qian J, Yoo JW, Noebels JL. Masking epilepsy by combining two epilepsy genes. *Nat Neurosci*. 2007;10(12). doi:10.1038/nn1999
19. Calhoun JD, Hawkins NA, Zachwieja NJ, Kearney JA. Cacna1g is a genetic modifier of epilepsy in a mouse model of Dravet syndrome. *Epilepsia*. 2017;58(8):e111-e115.


doi:10.1111/EPI.13811

20. Mishra V, Karumuri BK, Gautier NM, et al. Scn2a deletion improves survival and brain–heart dynamics in the Kcna1-null mouse model of sudden unexpected death in epilepsy (SUDEP). *Hum Mol Genet.* 2017;26(11):2091-2103. doi:10.1093/HMG/DDX104
21. Kearney JA, Yang Y, Beyer B, et al. Severe epilepsy resulting from genetic interaction between Scn2a and Kcnq2. *Hum Mol Genet.* 2006;15(6):1043-1048. doi:10.1093/HMG/DDL019
22. Johnson J, Focken T, Khakh K, et al. NBI-921352, a first-in-class, Na V 1.6 selective, sodium channel inhibitor that prevents seizures in Scn8a gain-of-function mice, and wild-type mice and rats. *Elife.* 2022;11:e72468. doi:10.7554/ELIFE.72468
23. Hammer MF, Pan Y, Cumbay M, Pendziwiat M, Afawi Z, Goldberg-Stern H, Johnstone L, Helbig I, Cummins TR. Whole exome sequencing and co-expression analysis identify an *SCN1A* variant that modifies pathogenicity in a family with genetic epilepsies and febrile seizures plus. *Epilepsia.* 2022. doi: 10.1111/epi.17296

Figure legends

Figure 1. An ASO to *Scn8a* prolongs survival of *Kcnq2* mutant mice. (A) Timeline for treatment. (B, C) *Kcnq2^{fl/fl},Emx1^{Cre/+}* mice treated with *Scn8a* ASO survive longer than mice treated with control ASO. A second treatment prolonged the effect. Asterisks indicate significance of Bonferroni's multiple comparisons test: *, $p < 0.05$; ****, $p < 0.0001$. (D) Sample EEG recordings demonstrating normal EEG background in two mutant mice. (E) Abnormal hindlimb postures in *Kcnq2^{fl/fl},Emx1^{Cre/+}* mice after treatment with *Scn8a* ASO at P2.

Figure 2. *Scn8a* ASO prolongs survival of *Kcna1* mutant mice. (A) Timeline for treatment. (B, C) *Kcna1^{-/-}* mice treated with *Scn8a* ASO survive longer than mice treated with control ASO. Asterisks indicate significance of Bonferroni's multiple comparisons test: *, $p < 0.05$; ***, $p < 0.0005$. (D) EEG recordings showing abnormal interictal discharges (arrows) and electrographic seizure. (E) *Scn8a* ASO does not correct the growth deficit in *Kcna1^{-/-}* mice. (F) *Scn8a* ASO does not prevent myoclonic jerks in *Kcna1^{-/-}* mice.



EPI_17374_1_Kcnq2_survival_fonts.tif

