

ORIGINAL ARTICLE

International controlled study of revascularization and outcomes following COVID-positive mechanical thrombectomy

Adam A. Dmytriw^{1,2}  | Sherief Ghozy²  | Ahmad Sweid³ | Michel Piotin⁴ | Kimon Bekelis⁵ | Nader Sourour⁶ | Eytan Raz⁷ | Daniel Vela-Duarte⁸ | Italo Linfante⁸ | Guilherme Dabus⁸ | Max Kole⁹ | Mario Martínez-Galdámez¹⁰ | Shahid M. Nimjee¹¹ | Demetrius K. Lopes¹² | Ameer E. Hassan¹³ | Peter Kan¹⁴ | Mohammad Ghorbani¹⁵ | Michael R. Levitt¹⁶ | Simon Escalard⁴  | Symeon Missios⁵ | Maksim Shapiro⁷ | Frédéric Clarençon⁶ | Mahmoud Elhorany⁶  | Rizwan A. Tahir⁹ | Patrick P. Youssef¹¹ | Aditya S. Pandey¹⁷ | Robert M. Starke¹⁸ | Kareem El Naamani³ | Rawad Abbas³ | Ossama Y. Mansour¹⁹ | Jorge Galvan¹⁰ | Joshua T. Billingsley¹² | Abolghasem Mortazavi¹³ | Melanie Walker¹⁶ | Mahmoud Dibas² | Fabio Settecase²⁰ | Manraj K. S. Heran²⁰ | Anna L. Kuhn²¹ | Ajit S. Puri²¹ | Bijoy K. Menon²² | Sanjeev Sivakumar²³ | Ashkan Mowla²⁴ | Salvatore D'Amato¹ | Alicia M. Zha²⁵ | Daniel Cooke²⁶ | Justin E. Vranic¹ | Robert W. Regenhardt¹ | James D. Rabinov¹ | Christopher J. Stapleton¹ | Mayank Goyal²² | Hannah Wu^{27,28,29} | Jake Cohen^{27,28,29} | David Turkel-Parella^{27,28,29} | Andrew Xavier^{30,31} | Muhammad Waqas³² | Vincent Tutino³² | Adnan Siddiqui³² | Gaurav Gupta³³ | Anil Nanda³³ | Priyank Khandelwal³³ | Cristina Tiu³⁴ | Pere C. Portela³⁵ | Natalia Perez de la Ossa³⁶ | Xabier Urrea³⁷ | Mercedes de Lera³⁸ | Juan F. Arenillas³⁸ | Marc Ribo^{39,40} | Manuel Requena^{39,40}  | Mariangela Piano⁴¹ | Guglielmo Pero⁴¹ | Keith De Sousa⁴² | Fawaz Al-Mufti⁴³ | Zafar Hashim⁴⁴ | Sanjeev Nayak⁴⁴ | Leonardo Renieri⁴⁵ | Rose Du² | Mohamed A. Aziz-Sultan² | David Liebeskind⁴⁶ | Raul G. Nogueira⁴⁷ | Mohamad Abdalkader⁴⁸ | Thanh N. Nguyen⁴⁸ | Nicholas Vigilante⁴⁹ | James E. Siegler⁴⁹ | Jonathan A. Grossberg⁵⁰ | Hassan Saad⁵⁰ | Michael R. Gooch³ | Nabeel A. Herial³ | Robert H. Rosenwasser³ | Stavropoula Tjournakaris³ | Aman B. Patel¹ | Ambooj Tiwari^{27,28,29} | Pascal Jabbour³ | North American Neurovascular COVID-19 (NAN-C) Consortium & Society of Vascular and Interventional Neurology (SVIN) Investigators

¹Neuroendovascular Program, Mass General Brigham Partners, Harvard Medical School, Boston, Massachusetts, USA

²Neuroradiology & Neurosurgery Services, Brigham & Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

³Department of Neurological Surgery, Thomas Jefferson University Hospital, Philadelphia, Pennsylvania, USA

⁴Department of Interventional Neuroradiology, Rothschild Foundation Hospital, Paris, France

⁵Department of Neurosurgery, Good Samaritan Hospital Medical Center, New York, USA

⁶Department of Interventional Neuroradiology, Pitié-Salpêtrière Hospital, Paris, France

- ⁷Department of Radiology, New York University Langone Medical Center, New York, New York, USA
- ⁸Department of Interventional Neuroradiology & Neuroendovascular Surgery, Miami Cardiac and Vascular Institute, Baptist Hospital of Miami, Miami, Florida, USA
- ⁹Department of Neurosurgery, Henry Ford Hospital, Detroit, Michigan, USA
- ¹⁰Department of Interventional Neuroradiology, Hospital Clinico Universitario de Valladolid, Valladolid, Spain
- ¹¹Department of Neurosurgery, The Ohio State University Wexner Medical Center, Columbus, Ohio, USA
- ¹²Department of Neurosurgery, Advocate Aurora Health, Chicago, Illinois, USA
- ¹³Department of Neuroscience, Valley Baptist Medical Center/University of Texas Rio Grande Valley, Harlingen, Texas, USA
- ¹⁴Department of Neurosurgery, UTMB, Houston, Texas, USA
- ¹⁵Department of Neurosurgery, Firoozgar Hospital, Tehran, Iran
- ¹⁶Departments of Neurological Surgery, Radiology, Mechanical Engineering, and Stroke & Applied Neuroscience Center, University of Washington, Seattle, Washington, USA
- ¹⁷Department of Neurosurgery, University of Michigan, Ann Arbor, Michigan, USA
- ¹⁸Department of Neurosurgery & Neuroradiology, University of Miami & Jackson Memorial Hospital, Miami, Florida, USA
- ¹⁹Department of Neurology, Alexandria University Hospital, Alexandria, Egypt
- ²⁰Division of Neuroradiology, Vancouver General Hospital, University of British Columbia, Vancouver, British Columbia, Canada
- ²¹Division of Neurointerventional Radiology, Department of Radiology, UMass Memorial Medical Center, Worcester, Massachusetts, USA
- ²²Calgary Stroke Program, Cumming School of Medicine, Calgary, Alberta, Canada
- ²³Department of Medicine (Neurology), Prisma Health Upstate, USC, Greenville, South Carolina, USA
- ²⁴Department of Neurological Surgery, University of Southern California, Los Angeles, California, USA
- ²⁵Department of Neurology, UT Health Science Center, Houston, Texas, USA
- ²⁶Department of Neurointerventional Radiology, San Francisco General Hospital, San Francisco, California, USA
- ²⁷Department of Neurology, Brookdale University Hospital, Brooklyn, New York, USA
- ²⁸Department of Neurology, Jamaica Medical Center, Richmond Hill, New York, USA
- ²⁹Department of Neurology, NYU Lutheran Hospital, Brooklyn, New York, USA
- ³⁰Department of Neurology, Sinai Grace Hospital, Detroit, Michigan, USA
- ³¹Department of Neurology, St. Joseph Mercy Health, Ann Arbor, Michigan, USA
- ³²Department of Neurosurgery, University of New York at Buffalo, Buffalo, New York, USA
- ³³Department of Neurology, Robert Wood Johnson University Hospital, New Brunswick, New Jersey, USA
- ³⁴Department of Neurology, University Emergency Hospital Bucharest, Bucharest, Romania; "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania
- ³⁵Department of Neurology, Hospital Universitari, Bellvitge, Barcelona, Spain
- ³⁶Stroke Unit, Neuroscience Department, Hospital Universitari Germans Trias i Pujol, Carretera Canyet s/n, Barcelona, Spain
- ³⁷Department of Neurology, Hospital Clínic, Barcelona, Spain
- ³⁸Department of Neurology, Hospital Clínic Universitario, Valladolid, Spain
- ³⁹Stroke Unit, Department of Neurology, Vall d'Hebron Research Institute, Barcelona, Spain
- ⁴⁰Departament de Medicina, Universitat Autònoma de Barcelona, Barcelona, Spain
- ⁴¹Department of Neuroradiology, ASST Grande Ospedale Metropolitano Niguarda, Milan, Italy
- ⁴²Department of Neurology, Eastern Region, Northwell Health, New York, New York, USA
- ⁴³Department of Neurology, Radiology, and Neurosurgery, Westchester Medical Center at NY Medical College, Valhalla, New York, USA
- ⁴⁴Department of Radiology, University Hospital of North Midlands, Stoke-on-Trent, UK
- ⁴⁵Department of Radiology, Neurovascular Unit, Careggi University Hospital, Florence, Italy
- ⁴⁶Department of Neurology, Ronald Reagan UCLA Medical Center, Los Angeles, USA
- ⁴⁷Department of Neurology, Marcus Stroke and Neuroscience Center, Grady Memorial Hospital, Emory University School of Medicine, Atlanta, Georgia, USA
- ⁴⁸Departments of Neurology and Radiology, Boston Medical Center, Boston University School of Medicine, Boston, Massachusetts, USA
- ⁴⁹Cooper Neurological Institute, Cooper University Hospital, Camden, New Jersey, USA
- ⁵⁰Department of Neurosurgery, Emory University, Atlanta, Georgia, USA

Correspondence

Adam A. Dmytriw, Neuroendovascular Program, Massachusetts General Hospital, Harvard Medical School, 55 Fruit St, Boston, MA 02114, USA.
Email: admytriw@mgh.harvard.edu

Abstract

Background and purpose: Previous studies suggest that mechanisms and outcomes in patients with COVID-19-associated stroke differ from those in patients with non-COVID-19-associated strokes, but there is limited comparative evidence focusing on these populations. The aim of this study, therefore, was to determine if a significant association exists between COVID-19 status with revascularization and functional outcomes

following thrombectomy for large vessel occlusion (LVO), after adjustment for potential confounding factors.

Methods: A cross-sectional, international multicenter retrospective study was conducted in consecutively admitted COVID-19 patients with concomitant acute LVO, compared to a control group without COVID-19. Data collected included age, gender, comorbidities, clinical characteristics, details of the involved vessels, procedural technique, and various outcomes. A multivariable-adjusted analysis was conducted.

Results: In this cohort of 697 patients with acute LVO, 302 had COVID-19 while 395 patients did not. There was a significant difference ($p < 0.001$) in the mean age (in years) and gender of patients, with younger patients and more males in the COVID-19 group. In terms of favorable revascularization (modified Thrombolysis in Cerebral Infarction [mTICI] grade 3), COVID-19 was associated with lower odds of complete revascularization (odds ratio 0.33, 95% confidence interval [CI] 0.23–0.48; $p < 0.001$), which persisted on multivariable modeling with adjustment for other predictors (adjusted odds ratio 0.30, 95% CI 0.12–0.77; $p = 0.012$). Moreover, endovascular complications, in-hospital mortality, and length of hospital stay were significantly higher among COVID-19 patients ($p < 0.001$).

Conclusion: COVID-19 was an independent predictor of incomplete revascularization and poor functional outcome in patients with stroke due to LVO. Furthermore, COVID-19 patients with LVO were more often younger and had higher morbidity/mortality rates.

KEYWORDS

COVID-19, large vessel occlusion, morbidity, mortality, stroke

INTRODUCTION

Since the first reported case in Wuhan, China, coronavirus disease 2019 (COVID-19) has been prevalent globally, with approximately 45 million cases and 730,000 deaths in the United States, up to October 8, 2021 [1]. According to the World Health Organization (WHO), there have been over 241 million cases and almost 5 million deaths worldwide [2]. Patients with COVID-19 are at higher risk of thrombotic events, with an estimated prevalence rate of 22%, which can further increase to up to 43% after admission to the intensive care unit [3]. Hypercoagulability in COVID-19 patients may be caused by disease-associated stasis, cytokine storm, dysfunctional endothelium, and platelet activation [3–5].

One form of thrombosis is acute ischemic stroke (AIS). The risk of AIS in COVID-19 patients may be elevated approximately three- to eightfold in the first 3 days of respiratory symptoms [6,7]. The reported prevalence of stroke among COVID-19 patients varies from 1.3% up to 4.9% and the rate of SARS CoV-2 infection among stroke admissions is estimated to be 3.3% [8–16]. While there have been a number of reports indicating an excess number of large vessel occlusion (LVO) strokes in patients with COVID-19 [8,17,18], there are limited data on the safety and outcomes of acute revascularization of LVO in COVID-19 patients [19,20]. To address current limitations, an international multicenter study was conducted to identify differences in the demographics and stroke characteristics of LVO patients with COVID-19 versus without COVID-19. Furthermore, we

sought to evaluate whether COVID-19 has an independent association with revascularization and functional outcomes following the endovascular treatment.

MATERIALS AND METHODS

An international multicenter retrospective study of consecutively admitted COVID-19 patients with concomitant acute LVOs was performed between February 25 and December 30, 2020, across 50 international comprehensive stroke centers, from North America, Europe and the Middle East.

The institutional review boards of participating institutions reviewed and approved the study, and patient consent was waived based on the de-identified retrospective protocol with minimal risk. Diagnosis of COVID-19 was established using reverse-transcriptase PCR assays of nasopharyngeal samples for identification of SARS-CoV-2. Data will be made available by the corresponding author to qualified investigators upon reasonable request.

Data collection

Data collected included age, gender, comorbidities, clinical characteristics of the included COVID-19 patients, details of the involved vessels, procedural technique, and selected outcome measures (e.g.,

symptomatic intracerebral hemorrhage [sICH]). Onset to admission time was defined as the time from stroke onset to hospital arrival. Procedure duration was calculated as the time difference between arterial access and sheath removal. COVID-19 severity was determined based on the score and classification provided by the World Health Organization (WHO) [21]. In addition, classification of AIS subtype was performed according to the Trial of Org 10,172 in Acute Stroke Treatment (TOAST) criteria [22]. For the involved vessels, we used the same classification provided by our participating centers based on what was reported as first, second, third, and fourth.

Study endpoints

Co-primary outcomes were: (i) optimal revascularization, defined as modified Thrombolysis in Cerebral Infarction (mTICI) grade 3; (ii) unfavorable functional outcome, at discharge and at 90 days, defined as modified Rankin Scale (mRS) score 3–6; and (iii) mortality at 90 days. Recanalization scoring was estimated without central imaging adjudication. Secondary outcomes were: (i) sICH, defined as reduction of four points on the National Institute of Health Stroke Scale (NIHSS) in association with any hemorrhage, at the judgment of the treating clinician; (ii) NIHSS score 24 h following mechanical thrombectomy.

Statistical analysis

All data were analyzed using R software version 4.1.1 using the packages “Rcmdr” and “glm2” [23]. Continuous variables are shown as means and standard deviation, with skewness and kurtosis tests used to evaluate the normal distribution, with comparisons made according to COVID-19 status (COVID-19 vs. non-COVID-19) using independent *t*-tests or Mann–Whitney *U*-tests, as appropriate. Categorical variables are reported as frequencies and percentages, with the chi-squared test or Fisher's exact test used for comparisons, as appropriate. Finally, univariable logistic regression was used to test covariates predictive of revascularization (mTICI grade 2b or 3), and unfavorable outcome (mRS score 3–6). Interaction and confounding were assessed through stratification and relevant expansion covariates. Whenever possible, factors predictive on univariable analysis ($p < 0.05$) were entered into a backward multivariable logistic regression analysis, and the effect of COVID-19 was assessed to be clinically relevant in all models. Regression results are expressed as odds ratios (ORs) and 95% confidence intervals (95% CIs). A *p* value < 0.05 was taken to indicate significance for all statistical tests. The results presented here are reported in accordance with the Strengthening the Observational Reporting of Observational Studies guidelines. Missing data were not imputed.

Data availability

The relevant anonymized patient-level data are available on reasonable request from the authors.

RESULTS

The total cohort comprised 697 patients with LVO, 302 of whom had concomitant COVID-19 (43.3%). Patients with COVID-19 had a younger mean age than those without COVID-19 (61.1 ± 15.7 years vs. 71.0 ± 15.8 years; $p < 0.001$), with a lower proportion of female patients in the COVID-19 group (41.1% vs. 80.5%; $p < 0.001$). Functional status prior to stroke onset was significantly different, with a lower proportion of functional independence in the COVID-19 group (mRS score 0–2: 65.6% vs. 96.2%; $p < 0.001$ [Table 1]).

Comorbidities

Chronic heart disease (18.9% vs. 34.4%; $p < 0.001$) and atrial fibrillation (21.2% vs. 38.2%; $p < 0.001$) were less common in patients with COVID-19, while chronic liver disease (6.0% vs. 2.5%; $p = 0.022$) and type 2 diabetes mellitus (29.5% vs. 22.3%; $p = 0.014$) were more common. Hypertension, chronic lung disease and chronic kidney disease frequency did not differ between the groups (Table 1).

COVID-19 characteristics

The severity of COVID-19 at stroke onset was moderate in 73.3% of cases, severe in 14.8% and critical in 11.9%. Cough was the most frequent presenting symptom (48.7%), followed by fever (45.9%), pneumonia (14.4%), and acute respiratory distress syndrome (20.0%). COVID-19 diagnosis was established in 66.1% of the patients prior to stroke onset. The mean duration between COVID-19 symptoms and stroke onset was 8.8 ± 11.4 days, and in 33.9% of the COVID-19 group, stroke was the initial manifestation of the COVID-19 disease.

Stroke characteristics

Cardioembolic etiology represented a lower proportion in the COVID-19 group (12.9% vs. 49.6%), while large vessel atherosclerosis (37.6% vs. 16.5%) and cryptogenic stroke (16.8% vs. 0.0%) were observed at a higher proportion in the COVID-19 group ($p < 0.001$). The mean Albert Stroke Program Early Computed Tomography Score (ASPECTS) at admission was lower in the COVID-19 group (7.8 ± 2.4 vs. 8.9 ± 1.5 ; $p < 0.001$). The mean number of involved vessels was comparable between the COVID-19 and non-COVID-19 groups (1.4 ± 0.9 vs. 1.3 ± 0.5 ; $p = 0.236$). The detailed distribution of affected vessels is shown in Table 2.

Stroke treatment

The mean duration of time from last known normal to access was shorter in the COVID-19 group (357.4 ± 513.3 min vs. 474.4 ± 365.1 min; $p = 0.009$), while the mean door-to-arterial access duration

TABLE 1 Baseline characteristics of the patients with large vessel occlusion with and without COVID-19

Variables	Non-COVID-19 (N = 395)	COVID-19 (N = 302)	Total (N = 697)	<i>p</i>
Mean (SD) age, years	71.0 (15.8)	61.1 (15.7)	67.1 (16.5)	<0.001*
Gender: female, <i>n</i> (%)	318 (80.5)	124 (41.1)	442 (63.4)	<0.001*
Pre-admission mRS score 0–2, <i>n</i> (%)	380 (96.2)	198 (65.6)	578 (82.9)	<0.001*
Mean (SD) NIHSS score at admission	14.3 (7.5)	17.2 (8.5)	15.4 (8.0)	<0.001*
Hypertension, <i>n</i> (%)	271 (68.8)	174 (57.6)	445 (63.8)	0.104
Chronic heart disease, <i>n</i> (%)	136 (34.4)	57 (18.9)	193 (27.7)	<0.001*
Chronic lung disease, <i>n</i> (%)	76 (19.2)	60 (19.9)	136 (19.5)	0.575
Chronic kidney disease, <i>n</i> (%)	43 (10.9)	28 (9.3)	71 (10.2)	0.791
Chronic liver disease, <i>n</i> (%)	10 (2.5)	18 (6.0)	28 (4.0)	0.022*
Diabetes mellitus (type 2), <i>n</i> (%)	88 (22.3)	89 (29.5)	177 (25.4)	0.014*
Atrial fibrillation, <i>n</i> (%)	151 (38.2)	64 (21.2)	215 (30.8)	<0.001*
New onset atrial fibrillation, <i>n</i> (%)	0 (0.0)	21 (7.0)	21 (3.0)	0.238

Note: Missing data count per variable: age, *n* = 53; gender *n* = 17; pre-admission mRS score, *n* = 70; NIHSS at admission, *n* = 72; hypertension, *n* = 40; chronic heart disease, *n* = 24; chronic lung disease, *n* = 22; chronic kidney disease, *n* = 24; chronic liver disease, *n* = 22; diabetes mellitus (type 2), *n* = 31; atrial fibrillation, *n* = 31; new onset atrial fibrillation, *n* = 417.

Abbreviations: mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; SD, standard deviation.

*Statistically significant.

was longer in the COVID-19 group (87.5 ± 63.5 min vs. 71.6 ± 80.0 min; $p = 0.043$). Intravenous tissue plasminogen activator (tPA) administration was less common in the COVID-19 group (23.5% vs. 33.4%; $p < 0.001$). A higher proportion of mechanical thrombectomy procedures was performed under general anesthesia in the COVID-19 group (31.5% vs. 19.1%; $p < 0.001$), while mean thrombectomy pass number (1.8 ± 1.5 vs. 1.9 ± 1.2 ; $p = 0.520$), first pass effect (53.3% vs. 49.1%; $p = 0.395$), and stenting rates (7.3% vs. 8.6%; $p = 0.584$) were comparable among the two groups (Table 3). The mean duration of the procedure to achieve revascularization was prolonged in the COVID-19 group (62.2 ± 47.3 vs. 51.9 ± 31.9 ; $p = 0.002$). Moreover, there was a higher proportion of mTICI grade 3 outcomes in the control group (66.6% vs. 25.5%; $p < 0.001$ [Table 3]).

Complications, functional outcomes, and mortality

The procedure-related complication rate was higher among COVID-19 patients (26.6% vs. 10.0%; $p < 0.001$), with 19.7% of the complications in the COVID-19 group being symptomatic. There was no significant difference in sICH (6.6% vs. 5.6%; $p = 0.683$) nor in NIHSS score at 24 h after thrombectomy (11.9 ± 10.8 vs. 12.2 ± 8.2 ; $p = 0.807$) between the COVID-19 and non-COVID-19 groups.

The length of hospital stay was longer in the COVID-19 group (15.5 ± 17.6 days vs. 8.4 ± 8.5 days; $p < 0.001$). Poor functional outcome (mRS score 3–6) at discharge was observed in a significantly higher proportion of patients in the COVID-19 group (37.1% vs. 9.6%; $p < 0.001$), and favorable functional outcome (mRS 0–2) at 90-day follow-up was observed in a lower proportion of COVID-19 patients

versus non-COVID-19 patients (10.6% vs 59.0%; $p < 0.001$). Similarly, the mortality rate was more than twofold higher in the COVID-19 group (42.0% vs. 19.1%; $p \leq 0.001$ [Table 3]).

Predictors of revascularization mTICI grade 3

In the univariable model, absence of COVID-19 infection (OR 0.33, 95% CI 0.23–0.48; $p < 0.001$), female gender (OR 2.56, 95% CI 1.78–3.69; $p < 0.001$), and higher ASPECTS (OR 1.15, 95% CI 1.04–1.27; $p = 0.007$) were predictors of better revascularization (mTICI grade 3). Accounting for other possible cofounders, only COVID-19 status (OR 0.30, 95% CI 0.12–0.77; $p = 0.012$) and female sex (OR 2.83, 95% CI 1.31–6.15; $p = 0.008$) were independent predictors (Table 4).

Predictors of unfavorable outcomes (mRS score 3–6)

Unfavorable outcomes were higher with increasing age (OR 1.04, 95% CI 1.02–1.06; $p < 0.001$) and NIHSS score at admission (OR 1.10, 95% CI 1.06–1.15; $p < 0.001$). Accounting for possible confounders (including pre-admission mRS score), age (OR 1.04, 95% CI 1.02–1.06; $p < 0.001$) and NIHSS score (OR 1.10, 95% CI 1.06–1.15; $p < 0.001$) persisted as independent risk factors for unfavorable outcomes (Table 5). A sensitivity analysis including only patients with a pre-admission mRS score of 0–2, found higher rates of unfavorable outcome among COVID-19 patients at 3 months (OR 2.10, 95% CI 1.08–3.99; $p = 0.025$); however, this association was not significant after controlling for other variables (OR 0.81, 95% CI 0.22–2.59; $p = 0.731$ [Table S1]). A summary of predictors of mTICI grade 3 and mRS score 3–6 is presented in Figure S1.

TABLE 2 Summary of the included large vessel occlusions

Variables	Non-COVID-19 (N = 395)	COVID-19 (N = 302)	Total (N = 697)	<i>p</i>
Stroke classification, <i>n</i> (%)				
Large vessel atherosclerosis (50% or more narrowing in an artery)	64 (16.5)	76 (37.6)	140 (23.8)	<0.001*
Cardioembolic	192 (49.6)	26 (12.9)	218 (37.0)	
Other (dissection, hypercoagulable)	131 (33.8)	66 (32.7)	197 (33.5)	
Cryptogenic	0 (0.0)	34 (16.8)	34 (5.8)	
Mean (SD) number of vessels involved	1.3 (0.5)	1.4 (0.9)	1.4 (0.8)	0.236
Vessel involved, <i>n</i> (%)				
ICA	35 (9.0)	9 (25.7)	44 (10.4)	<0.001*
MCA (M1 segment)	156 (40.2)	3 (8.6)	159 (37.6)	
MCA (M2 segment)	89 (22.9)	3 (8.6)	92 (21.7)	
ACA (A1 segment)	4 (1.0)	3 (8.6)	7 (1.7)	
ACA (A2 segment)	4 (1.0)	15 (42.9)	19 (4.5)	
Extracranial carotid	62 (16.0)	0 (0.0)	62 (14.7)	
Basilar	24 (6.2)	2 (5.7)	26 (6.1)	
MCA (M3/4 segments)	10 (2.6)	0 (0.0)	10 (2.4)	
Vertebral	4 (1.0)	0 (0.0)	4 (0.9)	
Second vessel involved (if more than one), <i>n</i> (%)				
ICA	7 (8.5)	4 (7.4)	11 (8.1)	0.002*
MCA (M1 segment)	39 (47.6)	10 (18.5)	49 (36.0)	
MCA (M2 segment)	18 (22.0)	8 (14.8)	26 (19.1)	
ACA (A1 segment)	5 (6.1)	13 (24.1)	18 (13.2)	
ACA (A2 segment)	3 (3.7)	4 (7.4)	7 (5.1)	
Extracranial carotid	0 (0.0)	3 (5.6)	3 (2.2)	
Basilar	1 (1.2)	1 (1.9)	2 (1.5)	
PCA	2 (2.4)	6 (11.1)	8 (5.9)	
MCA (M3/4 segments)	6 (7.3)	4 (7.4)	10 (7.4)	
Vertebral	1 (1.2)	1 (1.9)	2 (1.4)	
Third vessel involved (if more than two), <i>n</i> (%)				
MCA (M1 segment)	6 (31.6)	2 (50.0)	8 (34.8)	0.149
MCA (M2 segment)	5 (26.3)	1 (25.0)	6 (26.1)	
ACA (A1 segment)	7 (36.8)	0 (0.0)	7 (30.4)	
ACA (A2 segment)	1 (5.3)	0 (0.0)	1 (4.3)	
PCA	0 (0.0)	1 (25.0)	1 (4.3)	
Fourth vessel involved (if more than three), <i>n</i> (%)				
MCA (M1 segment)	0 (0.0)	8 (42.1)	8 (36.4)	0.244
MCA (M2 segment)	3 (100.0)	7 (36.8)	10 (45.5)	
ACA (A1 segment)	0 (0.0)	3 (15.8)	3 (13.6)	
PCA	0 (0.0)	1 (5.3)	1 (4.5)	

Note: Missing data count per variable: stroke classification, *n* = 108; ASPECTS, *n* = 152; vessel involved, *n* = 273.

Abbreviations: ACA, anterior cerebral artery; ASPECTS, Alberta Stroke Program Early Computed Tomography Score; COVID-19, coronavirus disease 2019; ICA, internal carotid artery; MCA, middle cerebral artery; PCA, posterior cerebral artery; SD, standard deviation.

*Statistically significant.

DISCUSSION

Our global collaborative effort to pool data on patients with LVO during the COVID-19 pandemic provided us with the opportunity

to explore the impact of COVID-19 on interventional outcomes in this population. From analyses of compiled cases from 50 institutions worldwide, our data suggest that COVID-19 patients with concomitant LVO have poorer functional outcome and rates of

TABLE 3 Summary of the procedures performed and outcomes

Variables	Non-COVID-19 (N = 395)	COVID-19 (N = 302)	Total (N = 697)	p
Mean (SD) door to groin time, min	71.6 (80.0)	87.5 (63.5)	76.7 (75.4)	0.043*
Mean (SD) LKN to groin time, min	474.4 (365.1)	357.4 (513.3)	442.8 (413.1)	0.009*
tPA given, n (%)	132 (33.4)	71 (23.5)	203 (29.1)	<0.001*
Mean (SD) number of passes during thrombectomy	1.9 (1.2)	1.8 (1.5)	1.8 (1.3)	0.520
First pass effect, n (%)	194 (49.1)	161 (53.3)	279 (40.0)	0.395
Stenting, n (%)	34 (8.6)	22 (7.3)	56 (8.0)	0.584
Mean (SD) procedure duration, min	51.9 (31.9)	62.2 (47.3)	56.4 (39.6)	0.002*
Favorable revascularization (mTICI 2b-3)	337 (85.3)	154 (50.9)	491 (86.3)	<0.001*
Favorable revascularization (mTICI 3)	263 (66.6)	77 (25.5)	340 (48.8)	<0.001*
Complications during or after the procedure, n (%)				
None	352 (90.0)	168 (73.4)	521 (83.9)	<0.001*
Asymptomatic	39 (10.0)	16 (6.9)	55 (8.9)	
Symptomatic	0 (0.0)	45 (19.7)	45 (7.2)	
sICH, n (%)	22 (5.6)	20 (6.6)	42 (6.0)	0.683
Mean (SD) NIHSS score 24h post thrombectomy	12.2 (8.2)	11.9 (10.8)	12.1 (9.1)	0.807
mRS score at discharge, n (%)				
0-2	131 (33.2)	149 (49.3)	280 (40.2)	<0.001*
3-6	38 (9.6)	112 (37.1)	150 (21.5)	
NA	226 (57.2)	41 (13.6)	267 (38.3)	
mRS score at 3 months follow-up, n (%)				
0-2	233 (59.0)	32 (10.6)	265 (38.0)	<0.001*
3-6	73 (18.5)	18 (6.0)	91 (13.1)	
NA	89 (22.5)	252 (83.4)	341 (48.9)	
Mean (SD) length of hospital stay, days	8.4 (8.5)	15.5 (17.6)	15.5 (17.5)	<0.001*
Discharge, n (%)				
Home	81 (20.5)	41 (13.6)	122 (17.5)	<0.001*
Rehabilitation	173 (43.8)	56 (18.5)	229 (32.9)	
Hospice	23 (5.8)	5 (1.7)	28 (4.0)	
Nursing facility	57 (14.4)	17 (5.6)	74 (10.6)	
Not reported/deceased	61 (15.4)	183 (60.6)	244 (35.0)	
In-hospital mortality	74 (19.1)	111 (42.0)	185 (28.4)	<0.001*

Note: Missing data count per variable: door to groin, $n = 271$; LKN to groin, $n = 264$; tPA given, $n = 80$; number of passes, $n = 112$; first pass effect, $n = 144$; stenting, $n = 65$; procedure duration, $n = 102$; favorable revascularization, $n = 127$; complications, $n = 75$; sICH, $n = 373$; 24 h NIHSS, $n = 161$; mRS at discharge, $n = 267$; mRS at 3 months; $n = 341$; length of hospital stay, $n = 92$; discharge, $n = 191$.

Abbreviations: LKN, last known normal; mRS, modified Rankin Scale; mTICI, modified Treatment in Cerebral Infarction; NA, not available; NIHSS, National Institutes of Health Stroke Scale; SD, standard deviation; sICH, symptomatic intracerebral hemorrhage; tPA, tissue plasminogen activator.

*Statistically significant.

revascularization compared to non-COVID-19 patients, with a mortality rate reaching up to 42%. The likelihood of achieving complete revascularization was reduced by 70%, compared to patients without COVID-19, in our collaboration. Our data are supported by contemporary reports of poorer outcomes in patients with AIS in the setting of COVID-19, albeit without controls, which is a key advantage of our current report [24-30]. Similarly to the present study, other analyses also suggest that COVID-19 is an independent predictor of

LVO, less favorable functional outcome at follow-up, and increased mortality [19,30,31,32,33].

The underlying pathophysiology that drives poorer outcomes for AIS in COVID-19 patients is subject to ongoing investigation. However, it is likely that COVID-19 influences patient conditioning for post-stroke recovery. Factors known to impede recovery after stroke include patient demographic factors, baseline functional status, and comorbidities [34]. In addition, stroke characteristics including severity

Predictors	Univariable	Multivariable
	OR (95% CI)	OR (95% CI)
COVID status		
Non-COVID-19	Reference	
COVID-19	0.33 (0.23–0.48; $p < 0.001^*$)	0.30 (0.12–0.77; $p = 0.012^*$)
Mean (SD) age	1.01 (0.99–1.02; $p = 0.306$)	0.99 (0.97–1.01; $p = 0.223$)
Gender		
Male	Reference	
Female	2.56 (1.78–3.69; $p < 0.001^*$)	2.83 (1.31–6.15; $p = 0.008^*$)
Pre-admission mRS score		
0–2	Reference	
3–6	1.48 (0.71–3.32; $p = 0.313$)	1.69 (0.35–12.64; $p = 0.547$)
NIHSS score at admission		
Mean (SD)	0.98 (0.96–1.00; $p = 0.064$)	1.00 (0.96–1.05; $p = 0.836$)
Chronic liver disease		
No	Reference	
Yes	1.06 (0.41–2.93; $p = 0.900$)	1.39 (0.30–10.09; $p = 0.701$)
Diabetes		
No	Reference	
Yes	1.19 (0.81–1.77; $p = 0.370$)	2.78 (1.33–6.31; $p = 0.010^*$)
ASPECTS on admission		
Mean (SD)	1.15 (1.04–1.27; $p = 0.007^*$)	1.14 (0.95–1.38; $p = 0.153$)
Door to groin time (min)		
Mean (SD)	1.00 (0.99–1.00; $p = 0.065$)	1.00 (0.99–1.00; $p = 0.249$)
LKN to groin time (min)		
Mean (SD)	1.00 (1.00–1.00; $p = 0.053$)	1.00 (1.00–1.00; $p = 0.906$)
tPA given		
No	Reference	
Yes	0.92 (0.64–1.31; $p = 0.634$)	0.77 (0.38–1.56; $p = 0.472$)

Abbreviations: ASPECTS, Alberta Stroke Program Early Computed Tomography Score; CI, confidence interval; LKN, last known normal; mRS, modified rankin scale; NIHSS, National Institutes of Health Stroke Scale; OR, odds ratio; tPA, tissue plasminogen activator; SD, standard deviation.

*Statistically significant.

and extent of stroke, location of ischemic tissue, time lag to treatment, concomitant pathologies, and other complications all affect post-stroke recovery [34]. The physiological state secondary to COVID-19 appears to further exacerbate damage caused by ischemia [35]. In particular, the heightened prothrombotic and proinflammatory state of COVID-19 may manifest in several ways, including AIS, vasculopathy, myocarditis, arrhythmias, thrombotic microangiopathy, coagulopathy and thrombocytopenia, tropism to endothelial cells via angiotensin-converting enzyme (ACE)-2 receptor, and inhibition of angiotensin (1–7) production [36–61]. It has been proposed that downregulation of ACE-2, leading to both arteriopathy and thrombosis, may play a central role in the development of stroke during COVID-19 [62–64].

Understanding the characteristics of patients who develop LVO in the setting of COVID-19 is important for prognosis and immediate care. To achieve this, we performed one of the first comparative studies of COVID-19 and control patients with LVO. We found that the

mean age of the COVID-19 cohort was younger than that of controls by approximately 10 years, which corroborates prior reported findings from other non-controlled studies [8,27,28,30,31,65,66,67,68]. This apparent incidence is almost fourfold higher than those in the general population. We also found a significantly higher representation of males in the COVID-19 group, which is consistent with previously reported non-controlled studies [28,30].

Elucidating the impact of COVID-19 on the onset and severity of AIS is also an important consideration in the clinical care of this population. This is pertinent, given that 73.3% of the patients who developed LVO with COVID-19 had moderate disease severity according to the WHO classification [21]. Their immune dysregulation may result in a cytokine storm, which is of pathophysiological significance in the development of stroke in COVID-19 disease [69–71]. The time from initial COVID symptomatology to stroke onset was, on average, 9 days in our study [72]. The Global COVID-19 Stroke

TABLE 4 Logistic regression for possible predictors of complete revascularization in patients with large vessel occlusion during the COVID-19 pandemic (mTICI grade 3)

TABLE 5 Logistic regression for possible predictors of unfavorable outcomes (modified Rankin Scale score 3–6)

Predictors	Univariable	Multivariable
	OR (95% CI; <i>p</i> value)	OR (95% CI; <i>p</i> value)
COVID status		
Non-COVID	Reference	
COVID	1.80 (0.94–3.36; <i>p</i> = 0.071)	0.82 (0.23–2.59; <i>p</i> = 0.743)
Age (years)		
Mean (SD)	1.04 (1.02–1.06; <i>p</i> < 0.001 [*])	1.04 (1.02–1.06; <i>p</i> < 0.001 [*])
Gender		
Male	Reference	
Female	0.96 (0.56–1.69; <i>p</i> = 0.887)	1.04 (0.50–2.23; <i>p</i> = 0.923)
Pre-admission mRS score		
0–2	Reference	
3–6	2.58 (0.81–7.98; <i>p</i> = 0.096)	3.28 (0.93–11.72; <i>p</i> = 0.061)
NIHSS score at admission		
Mean (SD)	1.10 (1.06–1.13; <i>p</i> < 0.001 [*])	1.10 (1.06–1.15; <i>p</i> < 0.001 [*])
Chronic liver disease		
No	Reference	
Yes	1.69 (0.43–5.73; <i>p</i> = 0.413)	1.80 (0.35–7.84; <i>p</i> = 0.443)
Diabetes		
No	Reference	
Yes	1.14 (0.64–1.98; <i>p</i> = 0.653)	0.95 (0.48–1.85; <i>p</i> = 0.894)
ASPECTS on admission		
Mean (SD)	0.92 (0.81–1.05; <i>p</i> = 0.206)	0.96 (0.82–1.14; <i>p</i> = 0.640)
tPA given		
No	Reference	
Yes	0.87 (0.51–1.45; <i>p</i> = 0.588)	0.82 (0.44–1.49; <i>p</i> = 0.515)

Abbreviations: ASPECTS, Alberta Stroke Program Early Computed Tomography Score; COVID-19, coronavirus disease 2019; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; tPA, tissue plasminogen activator; SD, standard deviation.

*Statistically significant.

Registry reported a similar latency of approximately 7 days between symptom onset and stroke [30]. The interval from symptom onset to hospital presentation was lower in the COVID-19 group, which may be consistent with previous studies that did not show a delay in endovascular thrombectomy time metrics during the COVID-19 era [73,74]. Factors that can increase the risk of stroke include infection, which can also concurrently worsen the severity of the stroke [75].

There was a significant difference between stroke classification according to TOAST criteria, with a higher proportion of patients with cryptogenic stroke in the COVID-19 group [22]. In our international experience, we found that stroke severity was generally worse in the COVID-19 cohort compared to controls, based on ASPECTS, NIHSS score at presentation, and the number of involved vessels. The etiology of strokes for patients with and without COVID-19 was also significantly different. Confirming the preliminary conclusions of smaller non-controlled studies, we found that COVID-19 LVO was associated with higher rates of strokes due to large vessel atherosclerosis or cryptogenic etiology, whereas non-COVID-19 LVO was more likely to be cardioembolic in etiology [18,76,77]. This further supports the suggestion that COVID-19 is a prothrombotic and

proinflammatory systemic state which may induce LVO. Additionally, the cerebral distribution of strokes was also considerably different. Non-COVID-19 strokes were most frequently observed in the MCA-M1 distribution. However, COVID-19 LVO appeared more likely to occur in the anterior cerebral artery, for reasons that are yet to be elucidated. It should be noted that a non-negligible portion of the occlusion location data was not available in the current study.

When comparing our experience with those of other large analyses, we found that the Get With The Guidelines-Stroke (GWTG-Stroke) registry analysis reported comparable results in patients diagnosed with COVID-19. The authors reported a worse NIHSS score at presentation, and considerably greater proportions of patients with LVO stroke [31]. In our study, the rate of tPA administration was higher amongst non-COVID-19 patients. This may be related to several factors, but one contributor could have been barriers to stroke care for COVID-19 patients. Consistent with other clinical cerebrovascular studies, there was a relative global decline in intravenous thrombolysis, mechanical thrombectomy, ruptured aneurysm treatment, and aneurysmal subarachnoid hemorrhage admissions during the initial wave of the COVID-19 pandemic [15,78].

The complexity of intervention and technical approach are other factors to be considered for mechanical thrombectomy for LVO. We assessed relative complexity between COVID-19 and non-COVID-19 groups by comparing indirect measures based on the duration of the procedure, number of vessels involved, number of passes, rate of complete and favorable revascularization, and technical complications. Our analysis suggests that having LVO with COVID-19 was associated with more involved vessels, longer procedure duration, and a lower rate of complete revascularization at the end of the procedure, albeit with a similar number of passes per occlusion. Furthermore, COVID-19 patients had a 70% lower likelihood of achieving mTICI grade 3. Our results are in line with those reported in prior non-controlled COVID-19 series and are consistent with historic mechanical thrombectomy data [8,67,68]. A possible explanation is the hypercoagulable state in COVID-19 patients, which may cause re-occlusion. In addition, among the COVID-19 patients undergoing thrombectomy, the intravascular clots were prone to fragment and migrate into both new vascular territories and into distal downstream vasculature, at higher rates than are otherwise typical. However, neither of these possible causes emerged as a pattern or were of a frequency great enough to be generalizable. A prospective, large-scale trial could help to answer such a question. Although we found the initial admission ASPECTS to be highly associated with successful revascularization, the association was not significant in the multivariate model. In the same context, time to groin puncture and whether or not tPA was given were also non-influential factors for revascularization in multivariable analysis. Of note, patients with diabetes had 2.78 higher odds of achieving mTICI grade 3 compared to controls, even after adjustment. The reason for the association with diabetes is unknown but possible hypotheses surround differences in etiology of stroke. Diabetic patients may have received treatment for more strict glycemic control, whereas, paradoxically, non-diabetic patients can encounter iatrogenic hyperglycemia with common COVID treatments such as dexamethasone. As a known inhibitor of fibrinolysis, hyperglycemia on admission was the only independent predictor of failed recanalization after tPA treatment [79,80].

Finally, we compared follow-up functional outcomes after Endovascular therapy (EVT) in patients with COVID-19 versus those without. We found, in our combined cohort, that good functional outcome at discharge and follow-up were significantly lower in the COVID-19 group. The mortality rate, when compared to prior published data, was significantly higher. Similarly, the GWTG-Stroke consortium and the Global COVID-19 Stroke Registry studies demonstrated that COVID-19 was an independent predictor of poor outcome and death [30,31]. Other factors such as diabetes and chronic liver disease history, admission ASPECTS, and tPA use were not associated with poor functional outcome on follow-up. In our cohort, we found that 24-h sICH and NIHSS scores were not significantly different with or without COVID-19. However, patients with COVID-19 had a prolonged hospital stay, which suggests that other factors prolong the perioperative and subsequent recovery process.

Strengths of this study include the fact that it is the first global multicenter controlled study for LVO in COVID-19. It highlights the

value of institutional collaboration in addressing clinical questions in a timely and robust manner. The study also has several limitations. Its retrospective design means there is an element of selection bias, particularly when patients are chosen for mechanical thrombectomy. Due to the stress on resources created by the COVID-19 pandemic, some patients may not have received optimal advanced imaging and clinical follow-up, and, as such, there were missing data for several outcomes. Nevertheless, all missing data were within subsidiary variables. Occasionally, control for some cofounders was not possible due to insufficient data in patient records. Future prospective studies are needed to obtain a higher level of evidence.

In conclusion, COVID-19 was an independent predictor of incomplete revascularization in patients with stroke due to LVO in this controlled study. Patients are more often younger, more often male, and have higher morbidity/mortality rates.

AUTHOR CONTRIBUTIONS

Sherief Ghozy: Data curation (lead); formal analysis (lead); project administration (equal); software (equal); validation (equal); visualization (lead); writing – original draft (lead); writing – review and editing (equal). **Ahmad Sweid:** Conceptualization (equal); formal analysis (equal); methodology (equal); resources (equal); validation (equal); writing – review and editing (equal). **Michel Piotin:** Formal analysis (equal); investigation (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Kimon Bekelis:** Conceptualization (equal); data curation (equal); formal analysis (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Nader-Antoine Sourour:** Conceptualization (equal); methodology (equal); resources (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Eytan Raz:** Conceptualization (equal); methodology (equal); resources (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Daniel Vela-Duarte:** Investigation (equal); methodology (equal); project administration (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Italo Linfante:** Conceptualization (equal); data curation (equal); resources (equal); validation (equal); writing – review and editing (equal). **Guilherme Dabus:** Methodology (equal); resources (equal); software (equal); validation (equal); writing – review and editing (equal). **Max Kole:** Conceptualization (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Mario Martinez-Galdmez:** Investigation (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Shahid M. Nimjee:** Investigation (equal); methodology (equal); project administration (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Demetrius K. Lopes:** Data curation (equal); formal analysis (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Ameer E. Hassan:** Conceptualization (equal); investigation (equal); methodology (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Peter Kan:** Formal analysis (equal); investigation (equal); methodology (equal);

resources (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Mohammad Ghorbani:** Conceptualization (equal); data curation (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Michael R. Levitt:** Conceptualization (equal); methodology (equal); resources (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Simon Escalard:** Conceptualization (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Symeon Missios:** Conceptualization (equal); formal analysis (equal); validation (equal); writing – review and editing (equal). **Maksim Shapiro:** Conceptualization (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Frederic Clarenion:** Formal analysis (equal); investigation (equal); methodology (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Mahmoud Elhorany:** Data curation (equal); methodology (equal); project administration (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Rizwan A. Tahir:** Conceptualization (equal); data curation (equal); formal analysis (equal); resources (equal); validation (equal); writing – review and editing (equal). **Patrick P. Youssef:** Data curation (equal); investigation (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Aditya S. Pandey:** Conceptualization (equal); data curation (equal); methodology (equal); resources (equal); validation (equal); writing – review and editing (equal). **Robert M. Starke:** Conceptualization (equal); methodology (equal); project administration (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Kareem El Naamani:** Formal analysis (equal); investigation (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Rawad Abbas:** Conceptualization (equal); methodology (equal); project administration (equal); resources (equal); validation (equal); writing – review and editing (equal). **Ossama Yassin:** Conceptualization (equal); data curation (equal); formal analysis (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Jorge Galvan:** Conceptualization (equal); data curation (equal); project administration (equal); resources (equal); software (equal); writing – review and editing (equal). **Joshua T. Billingsley:** Conceptualization (equal); data curation (equal); formal analysis (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Abolghasem Mortazavi:** Data curation (equal); formal analysis (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Melanie Walker:** Formal analysis (equal); methodology (equal); resources (equal); supervision (equal); validation (equal); writing – review and editing (equal). **Mahmoud Dibas:** Data curation (equal); methodology (equal); project administration (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Fabio Settecase:** Data curation (equal); investigation (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Manraj K.S. Heran:** Conceptualization (equal); data

curation (equal); resources (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Anna L. Kuhn:** Conceptualization (equal); data curation (equal); formal analysis (equal); investigation (equal); validation (equal); writing – review and editing (equal). **Ajit S. Puri:** Formal analysis (equal); methodology (equal); resources (equal); software (equal); validation (equal); writing – review and editing (equal). **B. K. Menon:** Conceptualization (equal); data curation (equal); formal analysis (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Sanjeev Sivakumar:** Conceptualization (equal); formal analysis (equal); methodology (equal); resources (equal); validation (equal); writing – review and editing (equal). **Ashkan Mowla:** Data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Salvatore DAmato:** Conceptualization (equal); data curation (equal); formal analysis (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Alicia M. Zha:** Investigation (equal); methodology (equal); project administration (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Daniel Cooke:** Data curation (equal); methodology (equal); project administration (equal); resources (equal); validation (equal); writing – review and editing (equal). **Justin E. Vranic:** Conceptualization (equal); methodology (equal); resources (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Robert W. Regenhardt:** Investigation (equal); methodology (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **James D. Rabinov:** Conceptualization (equal); data curation (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Christopher J. Stapleton:** Formal analysis (equal); methodology (equal); resources (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Mayank Goyal:** Data curation (equal); investigation (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Hannah Wu:** Investigation (equal); methodology (equal); project administration (equal); resources (equal); validation (equal); writing – review and editing (equal). **Jake Cohen:** Methodology (equal); project administration (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **David Turkel-Parella:** Investigation (equal); methodology (equal); project administration (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Andrew Xavier:** Data curation (equal); formal analysis (equal); methodology (equal); resources (equal); validation (equal); writing – review and editing (equal). **Muhammad Wagas:** Data curation (equal); methodology (equal); resources (equal); software (equal); validation (equal); writing – review and editing (equal). **Vincent Tutino:** Conceptualization (equal); methodology (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Adnan H Siddiqui:** Conceptualization (equal); methodology (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Gaurav Gupta:** Conceptualization (equal); data curation (equal); formal analysis (equal); project

administration (equal); validation (equal); writing – review and editing (equal). **Anil Nanda:** Conceptualization (equal); investigation (equal); methodology (equal); project administration (equal); validation (equal); writing – review and editing (equal). **Priyank Khandelwal:** Investigation (equal); methodology (equal); project administration (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Cristina Tiu:** Conceptualization (equal); data curation (equal); formal analysis (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Pere Cardona:** Conceptualization (equal); methodology (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Natilia Perez de la Ossa:** Investigation (equal); methodology (equal); resources (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Xabier Urrea:** Conceptualization (equal); data curation (equal); formal analysis (equal); investigation (equal); validation (equal); writing – review and editing (equal). **Mercedes De Lera Alfonso:** Conceptualization (equal); investigation (equal); methodology (equal); project administration (equal); validation (equal); writing – review and editing (equal). **JF Arenillas:** Data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Marc Ribo:** Conceptualization (equal); formal analysis (equal); investigation (equal); resources (equal); validation (equal); writing – review and editing (equal). **Manuel Requena:** Data curation (equal); investigation (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Mariangela Piano:** Formal analysis (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Guglielmo Pero:** Conceptualization (equal); data curation (equal); formal analysis (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Keith De Sousa:** Data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Fawaz Al-Mufti:** Conceptualization (equal); investigation (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Zafar Hashim:** Data curation (equal); investigation (equal); software (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Sanjeev Nayak:** Conceptualization (equal); data curation (equal); formal analysis (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Leonardo Renieri:** Conceptualization (equal); methodology (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Rose Du:** Conceptualization (equal); formal analysis (equal); methodology (equal); resources (equal); software (equal); validation (equal); writing – review and editing (equal). **Mohamed A. Aziz-Sultan:** Data curation (equal); formal analysis (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **David Liebeskind:** Conceptualization (equal); data curation (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **R.G. Nogueira:** Conceptualization (equal); data curation (equal); formal analysis

(equal); methodology (equal); validation (equal); writing – review and editing (equal). **Mohamad Abdalkader:** Conceptualization (equal); methodology (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Thanh N Nguyen:** Conceptualization (equal); formal analysis (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Nicholas Vigilante:** Conceptualization (equal); data curation (equal); investigation (equal); resources (equal); validation (equal); writing – review and editing (equal). **James E. Siegler:** Formal analysis (equal); methodology (equal); project administration (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Jonathan A Grossberg:** Conceptualization (equal); formal analysis (equal); investigation (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Hassan Saad:** Investigation (equal); methodology (equal); project administration (equal); validation (equal); writing – original draft (equal); writing – review and editing (equal). **Michael R. Gooch:** Conceptualization (equal); data curation (equal); methodology (equal); resources (equal); validation (equal); writing – review and editing (equal). **Nabeel A. Herial:** Conceptualization (equal); investigation (equal); project administration (equal); resources (equal); validation (equal); writing – review and editing (equal). **Robert H. Rosenwasser:** Conceptualization (equal); formal analysis (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Stavropoula Tjoumakaris:** Conceptualization (lead); methodology (lead); supervision (lead); validation (lead); writing – review and editing (lead). **Aman B. Patel:** Conceptualization (lead); formal analysis (equal); investigation (lead); methodology (lead); validation (lead); visualization (lead); writing – original draft (lead); writing – review and editing (lead). **Ambooj Tiwari:** Conceptualization (lead); formal analysis (lead); supervision (lead); validation (lead); writing – original draft (lead); writing – review and editing (lead). **Pascal M. Jabbour:** Conceptualization (lead); formal analysis (lead); supervision (lead); validation (lead); writing – original draft (lead); writing – review and editing (lead).

CONFLICT OF INTEREST

There are no relevant conflicts of interest.

FUNDING INFORMATION

This research received no grant from any funding agency in public, commercial, or not-for-profit sectors.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ETHICS STATEMENT

All procedures in the studies involving human participants were performed according to our institutional review board ethical standards and national research committee and the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. The study protocol was reviewed and approved by our

University Institutional Review Board. Based on our institutional guidelines, all protected health information was removed, and individual patient consent was not required in the analysis of this case series.

ORCID

Adam A. Dmytriw  <https://orcid.org/0000-0003-0131-5699>

Sherief Ghozy  <https://orcid.org/0000-0001-5629-3023>

Simon Escalard  <https://orcid.org/0000-0003-1306-1008>

Mahmoud Elhorany  <https://orcid.org/0000-0002-0257-2040>

Manuel Requena  <https://orcid.org/0000-0002-5671-6484>

REFERENCES

- Worldometer. COVID-19 Coronavirus Pandemic – United States. Accessed 10 August 2021. <https://www.worldometers.info/coronavirus/country/us/>
- WHO. Accessed 8 November, 2021. https://www.who.int/emergencies/diseases/novel-coronavirus-2019?adgroupsurvey={adgroupsurvey}&gclid=CjwKCAjwn8SLBhAyEiwAHNTJbVoUSPz5QopGJSxtwL2Uev_02TsV3icgoc9umdyGMLkJ6u0xoq4t6hoCXZcQAvD_BwE
- Xiong X, Chi J, Gao Q. Prevalence and risk factors of thrombotic events on patients with COVID-19: a systematic review and meta-analysis. *Thromb J*. 2021;19:32.
- Voicu S, Ketfi C, Stépanian A, et al. Pathophysiological processes underlying the high prevalence of deep vein thrombosis in critically ill COVID-19 patients. *Front Physiol*. 2020;11:608788.
- Pons S, Fodil S, Azoulay E, Zafrani L. The vascular endothelium: the cornerstone of organ dysfunction in severe SARS-CoV-2 infection. *Crit Care*. 2020;24:353.
- Smeeth L, Thomas SL, Hall AJ, Hubbard R, Farrington P, Vallance P. Risk of myocardial infarction and stroke after acute infection or vaccination. *N Engl J Med*. 2004;351:2611-2618.
- Warren-Gash C, Blackburn R, Whitaker H, McMenamin J, Hayward AC. Laboratory-confirmed respiratory infections as triggers for acute myocardial infarction and stroke: a self-controlled case series analysis of national linked datasets from Scotland. *Eur Respir J*. 2018;51:1701794.
- Khandelwal P, Al-Mufti F, Tiwari A, et al. Incidence, characteristics and outcomes of large vessel stroke in COVID-19 cohort: an international multicenter study. *Neurosurgery*. 2021;89:E35-E41.
- Qin C, Zhou L, Hu Z, et al. Clinical characteristics and outcomes of COVID-19 patients with a history of stroke in Wuhan, China. *Stroke*. 2020;51:2219-2223.
- Mao L, Jin H, Wang M, et al. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol*. 2020;77:683-690.
- Lodigiani C, Iapichino G, Carenzo L, et al. Venous and arterial thromboembolic complications in COVID-19 patients admitted to an academic hospital in Milan, Italy. *Thromb Res*. 2020;191:9-14.
- Klok FA, Kruip M, van der Meer NJM, et al. Confirmation of the high cumulative incidence of thrombotic complications in critically ill ICU patients with COVID-19: an updated analysis. *Thromb Res*. 2020;191:148-150.
- Helms J, Tacquard C, Severac F, et al. High risk of thrombosis in patients with severe SARS-CoV-2 infection: a multicenter prospective cohort study. *Intensive Care Med*. 2020;46:1089-1098.
- Jain R, Young M, Dogra S, et al. COVID-19 related neuroimaging findings: a signal of thromboembolic complications and a strong prognostic marker of poor patient outcome. *J Neurol Sci*. 2020;414:116923.
- Nogueira RG, Qureshi MM, Abdalkader M, et al. Global impact of COVID-19 on stroke care and IV thrombolysis. *Neurology*. 2021;96:e2824.
- Nogueira RG, Abdalkader M, Qureshi MM, et al. Global impact of COVID-19 on stroke care. *Int J Stroke*. 2021;16:573-584.
- Kihira S, Schefflein J, Mahmoudi K, et al. Association of coronavirus disease (COVID-19) with large vessel occlusion strokes: a case-control study. *AJR Am J Roentgenol*. 2021;216:150-156.
- Siegler JE, Cardona P, Arenillas JF, et al. Cerebrovascular events and outcomes in hospitalized patients with COVID-19: the SVIN COVID-19 multinational registry. *Int J Stroke*. 2021;16:437-447.
- Cagnazzo F, Piotin M, Escalard S, et al. European multicenter study of ET-COVID-19. *Stroke*. 2021;52:31-39.
- Sasanejad P, Afshar Hezarkhani L, Arsang-Jang S, et al. Safety and outcomes of intravenous thrombolytic therapy in ischemic stroke patients with COVID-19: CASCADE initiative. *J Stroke Cerebrovasc Dis*. 2021;30:106121.
- Son K-B, Lee T-J, Hwang S-S. Disease severity classification and COVID-19 outcomes, Republic of Korea. *Bull World Health Organ*. 2021;99:62-66.
- Adams HP Jr, Bendixen BH, Kappelle LJ, et al. Classification of subtype of acute ischemic stroke. Definitions for use in a multicenter clinical trial. TOAST. Trial of org 10,172 in acute stroke treatment. *Stroke*. 1993;24:35-41.
- Ho D, Imai K, King G, Stuart EA. MatchIt: nonparametric preprocessing for parametric causal inference. *J Stat Softw*. 2011;1(8):2011.
- Oxley TJ, Mocco J, Majidi S, et al. Large-vessel stroke as a presenting feature of Covid-19 in the young. *N Engl J Med*. 2020;382:e60.
- Sweid A, Hammoud B, Bekelis K, et al. Cerebral ischemic and hemorrhagic complications of coronavirus disease 2019. *Int J Stroke*. 2020;15(7):733-742.
- Sweid A, Hammoud B, Weinberg JH, et al. Letter: thrombotic neurovascular disease in COVID-19 patients. *Neurosurgery*. 2020;87(3):E400-E406.
- Wang A, Mandigo GK, Yim PD, Meyers PM, Lavine SD. Stroke and mechanical thrombectomy in patients with COVID-19: technical observations and patient characteristics. *J NeuroInterv Surg*. 2020;12(7):648-653.
- Escalard S, Maier B, Redjem H, et al. Treatment of acute ischemic stroke due to large vessel occlusion with COVID-19: experience from Paris. *Stroke*. 2020;51:2540-2543.
- de Havenon A, Ney JP, Callaghan B, et al. Impact of COVID-19 on outcomes in ischemic stroke patients in the United States. *J Stroke Cerebrovasc Dis*. 2021;30:105535.
- Ntaios G, Michel P, Georgiopoulos G, et al. Characteristics and outcomes in patients with COVID-19 and acute ischemic stroke: the global COVID-19 stroke registry. *Stroke*. 2020;51:e254-e258.
- Srivastava PK, Zhang S, Xian Y, et al. Acute ischemic stroke in patients with COVID-19: an analysis from get with the guidelines-stroke. *Stroke*. 2021;52:1826-1829.
- Belani P, Schefflein J, Kihira S, et al. COVID-19 is an independent risk factor for acute ischemic stroke. *Am J Neuroradiol*. 2020;41:1361-1364.
- Kihira S, Schefflein J, Mahmoudi K, et al. Association of coronavirus disease (COVID-19) with large vessel occlusion strokes: a case-control study. *Am J Roentgenol*. 2020;216(1):150-156.
- Alawieh A, Zhao J, Feng W. Factors affecting post-stroke motor recovery: implications on neurotherapy after brain injury. *Behav Brain Res*. 2018;340:94-101.
- Petrovic V, Radenkovic D, Radenkovic G, Djordjevic V, Banach M. Pathophysiology of cardiovascular complications in COVID-19. *Front Physiol*. 2020;11:575600.
- Mecca AP, Regenhardt RW, O'Connor TE, et al. Cerebroprotection by angiotensin-(1-7) in endothelin-1-induced ischaemic stroke. *Exp Physiol*. 2011;96:1084-1096.

37. Alawieh AM, Spiotta AM. Letter: may cooler heads prevail during a pandemic: stroke in COVID-19 patients or COVID-19 in stroke patients? *Neurosurgery*. 2020;87(4):E522.
38. Mishra AK, Sahu KK, Lal A, Sargent J. Mechanisms of stroke and the role of anticoagulants in COVID-19. *J Formos Med Assoc*. 2020;119(11):1721-1722.
39. Giraudon P, Bernard A. Inflammation in neuroviral diseases. *J Neural Transm*. 2010;117:899-906.
40. Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet*. 2020;395:1417-1418.
41. Barnes BJ, Adrover JM, Baxter-Stoltzfus A, et al. Targeting potential drivers of COVID-19: neutrophil extracellular traps. *J Exp Med*. 2020;217:e20200652.
42. Lorenzo C, Francesca B, Francesco P, Elena C, Luca S, Paolo S. Acute pulmonary embolism in COVID-19 related hypercoagulability. *J Thromb Thrombolysis*. 2020;50(1):223-226.
43. Han H, Yang L, Liu R, et al. Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin Chem Lab Med*. 2020;58(7):1116-1120.
44. Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost*. 2020;18:1094-1099.
45. Campbell CM, Kahwash R. Will complement inhibition be the new target in treating COVID-19-related systemic thrombosis? *Circulation*. 2020;141:1739-1741.
46. Tang N, Li D, Wang X, Sun Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020;18:844-847.
47. Zhang Y, Xiao M, Zhang S, et al. Coagulopathy and antiphospholipid antibodies in patients with Covid-19. *N Engl J Med*. 2020;382:e38.
48. Wang J, Hajizadeh N, Moore EE, et al. Tissue plasminogen activator (tPA) treatment for COVID-19 associated acute respiratory distress syndrome (ARDS): a case series. *J Thromb Haemost*. 2020;18(7):1752-1755.
49. Feldstein LR, Rose EB, Horwitz SM, et al. Multisystem inflammatory syndrome in U.S. children and adolescents. *N Engl J Med*. 2020;383(4):334-346.
50. Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature*. 2005;436:112-116.
51. Kuba K, Imai Y, Rao S, et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat Med*. 2005;11:875-879.
52. Lopez Verrilli MA, Pirola CJ, Pascual MM, Dominici FP, Turyn D, Gironacci MM. Angiotensin-(1-7) through AT2 receptors mediates tyrosine hydroxylase degradation via the ubiquitin-proteasome pathway. *J Neurochem*. 2009;109:326-335.
53. Turner AJ, Hiscox JA, Hooper NM. ACE2: from vasopeptidase to SARS virus receptor. *Trends Pharmacol Sci*. 2004;25:291-294.
54. Sampaio WO, Nascimento AAS, Santos RAS. Systemic and regional hemodynamic effects of angiotensin-(1-7) in rats. *Am J Physiol Heart Circ Physiol*. 2003;284:H1985-H1994.
55. Campagnole-Santos MJ, Diz DI, Santos RA, Khosla MC, Brosnihan KB, Ferrario CM. Cardiovascular effects of angiotensin-(1-7) injected into the dorsal medulla of rats. *Am J Physiol*. 1989;257:H324-H329.
56. Xu P, Sriramula S, Lazartigues E. ACE2/ANG-(1-7)/mas pathway in the brain: the axis of good. *Am J Physiol Regul Integr Comp Physiol*. 2011;300:R804-R817.
57. Chen J, Xiao X, Chen S, et al. Angiotensin-converting enzyme 2 priming enhances the function of endothelial progenitor cells and their therapeutic efficacy. *Hypertension*. 2013;61:681-689.
58. Chen J, Zhao Y, Chen S, et al. Neuronal over-expression of ACE2 protects brain from ischemia-induced damage. *Neuropharmacology*. 2014;79:550-558.
59. Bahouth MN, Venkatesan A. Acute viral illnesses and ischemic stroke: pathophysiological considerations in the era of the COVID-19 pandemic. *Stroke*. 2021;52:1885-1894.
60. Hamming I, Timens W, Bulthuis M, Lely A, Gv N, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol*. 2004;203:631-637.
61. To K, Lo AW. Exploring the pathogenesis of severe acute respiratory syndrome (SARS): the tissue distribution of the coronavirus (SARS-CoV) and its putative receptor, angiotensin-converting enzyme 2 (ACE2). *J Pathol*. 2004;203:740-743.
62. Divani AA, Andalib S, Di Napoli M, et al. Coronavirus disease 2019 and stroke: clinical manifestations and pathophysiological insights. *J Stroke Cerebrovasc Dis*. 2020;29(8):104941.
63. Labò N, Ohnuki H, Tosato G. Vasculopathy and coagulopathy associated with SARS-CoV-2 infection. *Cells*. 2020;9:1583.
64. Kaushik P, Kaushik M, Parveen S, Tabassum H, Parvez S. Cross-talk between key players in patients with COVID-19 and ischemic stroke: a review on neurobiological insight of the pandemic. *Mol Neurobiol*. 2020;57:4921-4928.
65. Majidi S, Fifi JT, Ladner TR, et al. Emergent large vessel occlusion stroke during New York City's COVID-19 outbreak: clinical characteristics and paraclinical findings. *Stroke*. 2020;51:2656-2663.
66. Yaghi S, Ishida K, Torres J, et al. SARS2-CoV-2 and stroke in a New York healthcare system. *Stroke*. 2020;51(7):2002-2011.
67. Altersberger VL, Stolze LJ, Heldner MR, et al. Maintenance of acute stroke care service during the COVID-19 pandemic lockdown. *Stroke*. 2021;52:1693-1701.
68. Rudilosso S, Laredo C, Vera V, et al. Acute stroke care is at risk in the era of COVID-19: experience at a comprehensive stroke Center in Barcelona. *Stroke*. 2020;51:1991-1995.
69. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med*. 2020;46:846-848.
70. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506.
71. Tanaka T, Narazaki M, Kishimoto T. IL-6 in inflammation, immunity, and disease. *Cold Spring Harb Perspect Biol*. 2014;6:a016295.
72. Avula A, Nalleballe K, Narula N, et al. COVID-19 presenting as stroke. *Brain Behav Immun*. 2020;87:115-119.
73. Czap AL, Zha AM, Sebaugh J, et al. Endovascular thrombectomy time metrics in the era of COVID-19: observations from the Society of Vascular and Interventional Neurology Multicenter Collaboration. *J Neurointerv Surg*. 2021. Online ahead of print.
74. Siegler JE, Zha AM, Czap AL, et al. Influence of the COVID-19 pandemic on treatment times for acute ischemic stroke: the Society of Vascular and Interventional Neurology Multicenter Collaboration. *Stroke*. 2021;52:40-47.
75. Elkind MSV, Boehme AK, Smith CJ, Meisel A, Buckwalter MS. Infection as a stroke risk factor and determinant of outcome after stroke. *Stroke*. 2020;51:3156-3168.
76. Ramos-Araque ME, Siegler JE, Ribo M, et al. Stroke etiologies in patients with COVID-19: the SVIN COVID-19 multinational registry. *BMC Neurol*. 2021;21:43.
77. Yaghi S, Ishida K, Torres J, et al. SARS-CoV-2 and stroke in a New York healthcare system. *Stroke*. 2020;51:2002-2011.
78. Nguyen TN, Haussen DC, Qureshi MM, et al. Decline in subarachnoid haemorrhage volumes associated with the first wave of the COVID-19 pandemic. *Stroke Vasc Neurol*. 2021;6(4):542-552.
79. Ribo M, Molina C, Montaner J, et al. Acute hyperglycemia state is associated with lower tPA-induced recanalization rates in stroke patients. *Stroke*. 2005;36:1705-1709.

80. Pandolfi A, Giaccari A, Cilli C, et al. Acute hyperglycemia and acute hyperinsulinemia decrease plasma fibrinolytic activity and increase plasminogen activator inhibitor type 1 in the rat. *Acta Diabetol.* 2001;38:71-76.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Dmytriv AA, Ghozy S, Sweid A, et al. International controlled study of revascularization and outcomes following COVID-positive mechanical thrombectomy. *Eur J Neurol.* 2022;29:3273-3287. doi: [10.1111/ene.15493](https://doi.org/10.1111/ene.15493)