

Supplementary Table 1. Age at and interval between initial questionnaire administration.

SPP version		Age at SPP (yrs)			Age at MGIS <sup>3</sup> (yrs)			Age at BIS <sup>4</sup> (yrs)		
		n	$\bar{x}^{\text{age}}$	SD	n	$\bar{x}^{\text{age}}$	SD	n	$\bar{x}^{\text{age}}$	SD
<b>Girl</b>	Child <sup>1</sup>	22	11.06	1.26	24	10.95	1.33	44	13.63	2.62
	Adolescent <sup>2</sup>	21	15.84	1.37	30	15.48	2.15			
<b>Boy</b>	Child	10	10.56	1.68	12	10.51	1.66	13	13.17	1.83
	Adolescent	5	15.16	1.98	6	14.95	1.93			

Interval (days) between completion of MGIS and SPP				Interval (days) between completion of MGIS and BIS					
	n	$\bar{x}^{\text{days}}$	SD		n	$\bar{x}^{\text{days}}$	SD		
<b>Girl</b>	Child	22	31.41	133.89	<b>Girl</b>	Child	18	76.17	171.36
	Adolescent	21	37.76	316.74		Adolescent	26	17.88	87.18
<b>Boy</b>	Child	10	15.50	246.60	<b>Boy</b>	Child	7	381.57	688.63
	Adolescent	5	42.60	85.55		Adolescent	6	172.83	423.35

SPP, Self-Perception Profile; MGIS, Multidimensional Gender Identity Scale; BIS, Body Image Scale

<sup>1</sup> Child version of SPP is administered to those 8-13 yrs

<sup>2</sup> Adolescent version of SPP is administered to those 14+ yrs

<sup>3</sup> MGIS is administered to those older than 8 years old

<sup>4</sup> BIS is administered to girls 10+ yrs and boys 13+ yrs