






REVIEW

Bariatric Surgery

Prevalence of food addiction among patients undergoing metabolic/bariatric surgery: A systematic review and meta-analysis

Dafiny R. Praxedes¹  | André E. Silva-Júnior²  | Mateus L. Macena¹  |
Ashley N. Gearhardt³  | Nassib B. Bueno^{1,2} 

¹Programa de Pós-Graduação em Nutrição (PPGNUT), Faculdade de Nutrição (FANUT), Universidade Federal de Alagoas (UFAL), Campus AC Simões, Av. Lourival Melo Mota, s/n, Cidade Universitária, Maceió, 57072-900, Brazil

²Programa de Pós-Graduação em Nutrição, Escola Paulista de Medicina, Universidade Federal de São Paulo, Rua Sena Madureira, n.º 1.500, Vila Clementino, São Paulo, 04021-001, Brazil

³Department of Psychology, University of Michigan, 530 Church Street, Ann Arbor, Michigan, 48109-1043, USA

Correspondence

Nassib B. Bueno, Programa de Pós-Graduação em Nutrição (PPGNUT), Faculdade de Nutrição (FANUT), Universidade Federal de Alagoas (UFAL), Campus AC Simões, Av. Lourival Melo Mota, s/n, Cidade Universitária, Maceió AL CEP 57072-900, Alagoas, Brazil. Email: nassib.bueno@fanut.ufal.br

Funding information

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Grant/Award Number: 23065.005919/2021-75

Summary

Candidates for metabolic/bariatric surgery show a high prevalence of food addiction (FA). However, few studies have investigated FA prevalence after bariatric surgery, especially using longitudinal studies. This systematic review with a meta-analysis aimed to determine pre- and postoperative prevalence of FA among patients undergoing metabolic/bariatric surgery. It included both cross-sectional and longitudinal studies that used the Yale Food Addiction Scale (YFAS). The following databases were searched: MEDLINE, ScienceDirect, LILACS, PsycArticles, CENTRAL, greylit.org, and opengrey.eu. Studies that used the YFAS to evaluate FA in pre- or postoperative patients were included. A random-effects meta-analysis was performed with cross-sectional studies to calculate the weighted prevalence of FA at the pre- and postoperative moments. For longitudinal studies, which measured FA at both time points for the same individuals, absolute prevalence reduction (APR) was calculated. Of the 6626 records, 40 studies were included in the meta-analysis. The preoperative weighted prevalence of FA was 32% (95% CI: 27–37%; 33 groups), whereas the postoperative prevalence was 15% (95% CI: 12–18%; 14 groups). Seven longitudinal studies showed a weighted APR of 26 p.p. (95% CI: 10–41 p.p.). Observational data suggest a reduction in the prevalence of FA among patients that undergo bariatric surgery. Interventional studies are needed to confirm these findings.

KEYWORDS

bariatric surgery, obesity, yale food addiction, YFAS

1 | INTRODUCTION

Obesity is a complex and multifactorial disease and is currently considered a global public health problem that affects people of different ages and economic levels.¹ It is closely related to environmental factors, such as the high availability of energy-dense ultra-processed foods rich in sugars and fats,^{2–4} and is associated with an increased risk of morbidity and mortality and reduced life expectancy.¹ In addition, people with obesity may show a higher prevalence of mental

health disorders, such as anxiety and depression^{5,6} and disordered eating behaviors.^{7,8}

Bariatric surgery is the most effective treatment for weight loss in patients with severe obesity and associated comorbidities.⁹ However, some patients who undergo these procedures may not experience satisfactory weight loss and may even present with weight regain,^{10,11} highlighting the need to understand the possible relationships that lead to these conditions in the postoperative period. With the growing interest in food addiction (FA), investigations of the relationships

between bariatric surgery and addiction-like eating behaviors have also increased.¹²⁻¹⁴

Although it is not formally recognized as a clinical condition, FA is defined as the excessive intake of highly processed, hyperpalatable, energy-dense foods with characteristics and effects similar to those of addictive substances.¹⁵ FA is primarily assessed using the Yale Food Addiction Scale (YFAS), which applies the diagnostic criteria for substance use disorders to food intake.¹⁶ Specifically, the YFAS identifies individuals with susceptibility to excessive consumption of ultra-processed and palatable foods that reflects addiction-like behaviors,¹⁷ such as loss of control over intake, intense cravings, continued use despite negative consequences, withdrawal, and tolerance. FA is higher in populations with obesity¹⁸⁻²⁰ and among patients undergoing metabolic/bariatric surgery.^{13,21,22}

The validity of the concept of FA has been the target of intense discussions in the last decade.²³ Many argue that this and other concepts related to eating behavior and overeating investigated through questionnaires examine the same construct that could be defined as “uncontrolled eating.”²⁴ Furthermore, there is an overlap between FA and other eating disorders, such as overeating and binge eating.²⁵ However, some investigations suggest that half of the individuals demonstrating YFAS diagnostic criteria for FA do not present binge eating or bulimia nervosa.²⁶ Another important controversy refers to the diagnostic criteria for FA being intimately related to substance use disorders, as there is no known single agent/compound that acts through specific mechanisms of action to induce the disorder in susceptible individuals.²⁷

Despite these controversies, FA is a rapidly growing area of research, and the last systematic review that examined the prevalence of FA among patients undergoing metabolic/bariatric surgery was published in 2017.²⁸ In addition, Koball et al.¹³ discussed the possible effects of bariatric surgery on FA through a narrative review of the literature. They concluded that FA rates significantly decreased during the first postoperative year, and presurgical FA was not related to postsurgical weight outcomes during the first postoperative year.¹³ However, neither review presented the overall prevalence of FA in the pre- and postoperative bariatric population. Considering the accelerated pace of publications on FA in the scientific literature,²⁹ an update on the estimates is warranted. Hence, it is appropriate to systematically analyze how the YFAS has been applied to the bariatric population and what the possible effects of surgery are on the prevalence of FA. Thus, this systematic review aims to determine the pre- and postoperative prevalence of FA in studies that used the YFAS among patients undergoing metabolic/bariatric surgery.

2 | METHODS

This systematic review with a meta-analysis is part of a larger review that aims to assess the prevalence of FA in different contexts.²⁹ It is reported according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guideline.³⁰ The protocol used in this study was previously published in the PROSPERO database

(<https://www.crd.york.ac.uk/prospero/>) under the registration identification of CRD42020193902.

2.1 | Eligibility criteria

Cross-sectional studies, cohorts, and clinical trials performed with pre-operative and postoperative bariatric surgery populations were included, regardless of age group, clinical condition, or other related variables. Studies that used any of the versions of the YFAS (i.e., YFAS, YFAS 2.0, dYFAS-C 2.0, mYFAS, mYFAS 2.0, YFAS-C, and mYFAS-C) and reported the prevalence of FA were included. All versions of the YFAS apply diagnostic criteria for substance use disorders to the intake of ultra-processed and palatable foods, but the different iterations of the YFAS have been developed to provide updates on diagnostic criteria (YFAS vs. YFAS 2.0), briefer versions (mYFAS and mYFAS 2.0), and developmentally appropriate measures for children/adolescents (YFAS-C, mYFAS-C, and dYFAS-C 2.0). Duplicate publications of studies were excluded.

2.2 | Search strategy

The following electronic bibliographic databases were searched for appropriate studies: Medical Literature Analysis and Retrieval System Online (MEDLINE), ScienceDirect, Latin American and Caribbean Health Sciences Literature (LILACS), American Psychological Association (PsycArticles), and Cochrane Central Register of Controlled Trials (CENTRAL). Likewise, the following gray literature databases were included: Grey Literature Report (GreyLit.org) and the System for Information on Grey Literature in Europe (opengrey.eu). In addition, the reference lists of articles included in the full-text reading were also analyzed to select items that were not retrieved by the search strategy.

The search strategy included terms related to the outcome (FA) and was adapted for each electronic database. The following keywords were used for searching all databases: “Food addiction,” “Eating Addiction,” “Yale Food Addiction Scale,” and “YFAS,” separated by the Boolean operator “OR.” A date restriction (2008–2022), from the year of validation of the first version of YFAS to the current year, but no language restriction, was applied. Before the final analysis, a final search was performed to identify new studies for inclusion in this review. The last database search was conducted on March 27, 2022.

2.3 | Identification and selection of studies

Mendeley v1.19.5[®] software (Elsevier, Netherlands) was used to help manage the references. Three authors (DRP, AESJ, and MLM) who had access to authors and journal titles independently assessed the titles and abstracts of the retrieved articles. Disagreements were resolved in consultation with a senior researcher (NBB). This schematization was repeated in the assessment of the

risk of bias. Full-text versions of potentially eligible articles were retrieved for further evaluation.

2.4 | Data extraction

The following variables were collected as exploratory variables: type of study, the country in which it was carried out, whether it was a validation study, presence or absence of clinical conditions/comorbidities, age group of the sample, gender, the time of surgery evaluated (preoperative, postoperatively, or both), type of surgery, the YFAS version used, the scoring method for YFAS, and the prevalence, if any, of bulimia, anorexia nervosa, binge eating, loss of eating control, and depression. Data were independently extracted by the review authors, and differences were resolved by consensus with a senior researcher (NBB). The necessary information was extracted from published articles, protocols, and comments related to each study.

2.5 | Risk of bias assessment

A risk of bias assessment was performed for all the studies included in the meta-analysis. Three authors independently assessed the potential risk of bias in each study using the Research Triangle Institute Item Bank (RTI-IB)³¹ as most of the studies in this review were observational. The RTI-IB tool was developed to identify sources of distortion and confounding in observational studies; it provides a comprehensive list of 29 questions covering a variety of categories of bias that assesses trends in selection, performance, detection, attrition, confounding, selective reporting of results, and overall study quality. This procedure was used to classify how confident the study was (low, medium, or high) about the proximity of observed effects to actual effects. The score for each study was calculated by dividing the number of completed items by the number of applicable items and later classified using the following cut-off points: 0–0.40 indicating high risk of bias; 0.41–0.70 indicating medium risk of bias, and 0.71–1.00 indicating low risk of bias.

2.6 | Certainty of evidence

The quality of the evidence was analyzed using adaptations of the method proposed by the Grading of Recommendations Assessment, Development and Evaluation (GRADE).³² In the present analysis, this method was adapted for cross-sectional studies. The quality of evidence was classified into three categories—high, moderate, and low—based on criteria such as type of study, methodological limitations, inconsistent results, indirect evidence, imprecision, and publication bias. The quality of the evidence determined by the GRADE system allows the analysis of aggregated results by considering the design and results of the studies included and the pooled effect estimate obtained by the meta-analysis.

2.7 | Data analysis

The “metaprop” command in Stata v.12 software (StataCorp, College Station, Texas) was used for the meta-analysis.³³ According to the YFAS, the primary outcome sought in the studies was the prevalence of FA. The weighted prevalence of FA was calculated separately for preoperative and postoperative studies. A DerSimonian and Laird random-effects model was used with the Freeman–Tukey transformation to stabilize variances. The data analyzed were prevalence evaluated in the other studies using different versions of the YFAS. Longitudinal studies that evaluated the pre- and postsurgery prevalence of FA in the same sample were analyzed in two ways: First, the pre- and postoperative FA prevalence found in these studies was included separately in each meta-analysis of pre- and postoperative studies; second, we analyzed the absolute prevalence reduction (APR) of FA in these longitudinal studies, considering the within-study variation in the FA prevalence data (i.e., the absolute difference in FA prevalence between each study's postoperative and preoperative moment, associated with its confidence interval). For the meta-analyses of the APR, we used the “metan” command with a DerSimonian and Laird random-effects model. Finally, meta-regression analysis was also performed to explore differences in the prevalence of FA between the pre- and postoperative groups, using the “metareg” command in Stata.

3 | RESULTS

3.1 | Search results

A total of 6626 records were identified through the search strategy. With the removal of duplicates and subsequent application of inclusion criteria, 43 studies were selected for full-text screening. After a complete reading, three articles were excluded for the following reasons: Prevalence was not exclusive to the bariatric sample ($n = 1$), and repeated results from a study were already included ($n = 2$). Figure 1 shows a flowchart that illustrates the search and selection of studies.

3.2 | Characteristics of the studies included in the analysis

Of the selected studies, 28 (68.3%) used the original YFAS to assess FA, 12 (29.3%) used YFAS 2.0, and 1 (2.4%) used mYFAS 2.0. No study used the iterations of the YFAS developed specifically for children. Tables 1–3 show the general characteristics of the 40 articles, separated according to the time of data collection (preoperative, postoperative, or both). Twenty-six studies presented preoperative data (Table 1), seven studies presented postoperative data (Table 2), and seven studies performed a longitudinal analysis of both preoperative and postoperative data for the same sample




FIGURE 1 Flowchart of studies included in the review

(Table 3). One study evaluated a sample of women only, whereas the others evaluated both sexes, although the samples were primarily female.

The prevalence of FA among the 26 preoperative studies ranged from 6.3% to 69% (Table 1), whereas it ranged from 0% to 32% in the postoperative studies (Table 2). The APR of the seven prospective studies with preoperative and postoperative measurements of the same sample ranged from 1 p.p. to 59 p.p. (Table 3). Of the seven prospective studies, two evaluated the presence of FA at 24 months after surgery; the others evaluated it at 6 months ($n = 1$), 9 months ($n = 1$), or 12 months ($n = 3$).

3.3 | Risk of bias

The assessment of the risk of bias classified 17 studies as having a low risk of bias, 22 studies as having a moderate risk of bias, and only one study as having a high risk of bias. The most frequent biases in studies with high and moderate risk were selection bias and confounding. The individual classification of the studies is shown in Tables 1–3.

3.4 | Results of meta-analysis

Figure 2 shows the weighted prevalence of FA in the 33 groups evaluated preoperatively, involving the 26 studies that were evaluated exclusively at the preoperative moment and the seven longitudinal studies. The data set included 8304 participants, of whom 2074 presented a positive diagnosis for FA. A pooled FA prevalence of 32% (95% CI: 27–37%, $I^2 = 95.22\%$) was found for the preoperative period.

In the case of the 14 groups of individuals with an evaluation at the postoperative moment (seven studies that exclusively measured FA at the postoperative moment and the seven longitudinal studies with measurements at both moments), the pooled FA prevalence was 15% (95% CI: 12–18%, $I^2 = 71.29\%$) (Figure 3). A total of 1754 individuals were evaluated, with 272 positives for FA.

Finally, in the case of the only seven longitudinal studies with measures of FA at both moments for the same set of individuals, the prevalence of FA at the preoperative moment was 34% (95% CI: 20–50%), whereas it was 9% (95% CI: 6–13%) at the postoperative moment, leading to a weighted APR of 26 p.p. (95% CI: 10–41 p.p., $I^2 = 95.8\%$) (Figure 4).

TABLE 1 Characteristics of included studies that assessed the prevalence of FA using the Yale Food Addiction Scale (YFAS) only in patients who were candidates for bariatric surgery (preoperative) ($n = 26$)

Author, year	YFAS version	Country	Sample (% F)	Prevalence of FA (%)	BMI mean (SD)	Risk of bias
Archi et al., 2021 ³⁴	YFAS 2.0	France	282 (76.6)	26.6	N/R	0.81 (low)
Benzerouk et al., 2018 ³⁵	YFAS 2.0	France	128 (71.1)	25.0	46.1 (6.6)	0.63 (moderate)
Bianciardi et al., 2019 ³⁶	YFAS	Italy	403 (N/R)	31.1	31.72 (6.59)	0.72 (low)
Brunault et al., 2016 ³⁷	YFAS	France	188 (84)	16.5	46.9 (7.8)	0.63 (moderate)
Brunault et al., 2020 ³⁸	YFAS 2.0	France	345 (75.7)	26.1	44.5 (7)	1.00 (low)
Cambiali et al., 2021 ³⁹	YFAS	Italy	141 (63.8)	18.2	43.9 (6.4)	0.54 (moderate)
Çelebi et al., 2021 ⁴⁰	YFAS	Turkey	140 (75)	47.1	47.39 (N/R)	0.81 (low)
Clark et al., 2013 ⁴¹	YFAS	USA	67 (62.7)	53.7	N/R	0.81 (low)
Clark et al., 2019 ⁴²	YFAS 2.0	USA	314 (82.8)	27.4	46.95 (8.16)	0.54 (moderate)
Cullen et al., 2017 ⁴³	YFAS	Australia	23 (74)	52.0	33 (N/R)	0.81 (low)
Dickhut et al., 2021 ²²	YFAS 2.0	Germany	125 (78.9)	39.2	N/R	0.72 (low)
Fuentes et al., 2017 ⁴⁴	YFAS	Chile	112 (54.46)	50.0	35.05 (N/R)	0.45 (moderate)
Gabler et al., 2015 ⁴⁵	YFAS	Mexico	65 (56.9)	58.46	35.35 (N/R)	0.53 (moderate)
Hernandez et al., 2017 ⁴⁶	YFAS	USA	28 (N/R)	13.8	N/R	0.38 (high)
Holgerson et al., 2021 ⁴⁷	YFAS	UK	1034 (71.3)	14.41	47.05 (12.56)	0.76 (low)
Koball et al., 2021 ⁴⁸	YFAS 2.0	USA	1061 (74.3)	18.0	46.9 (13.4)	0.72 (low)
Lawson et al., 2019 ⁴⁹	YFAS	USA	444 (83.50)	35.8	37.23 (15.93)	0.92 (low)
Lescher et al., 2020 ⁵⁰	YFAS 2.0	Germany	107 (77.72)	48.6	N/R	1.00 (low)
Meule et al., 2012 ⁵¹	YFAS	Germany	96 (65.6)	41.7	50.64 (8.99)	0.63 (moderate)
Meule et al., 2014 ⁵²	YFAS	Germany	94 (65.6)	40.42	50.64 (8.99)	0.61 (moderate)
Meule et al., 2017 ⁵³	YFAS 2.0	Germany	133 (78.3)	47.36	49.5 (7.51)	0.92 (low)
MillerMatero et al., 2014 ⁵⁴	YFAS	USA	142 (81)	16.9	49.05 (9.56)	0.72 (low)
Müller et al., 2018 ⁵⁵	YFAS 2.0	Germany	216 (80.10)	27.3	48.3 (7.22)	1.00 (low)
Murray et al., 2019 ⁵⁶	YFAS	USA	16 (93)	6.3	44.3 (4.4)	0.52 (moderate)
Guerrero Pérez et al., 2018 ²¹	YFAS 2.0	Spain	110 (76.36)	26.4	47.3 (8.8)	1.00 (low)
Reslan et al., 2014 ⁵⁷	YFAS	USA	141 (79)	36.0	N/R	0.66 (moderate)

Abbreviations: BMI, body mass index; FA, food addiction; LAGB, laparoscopic adjustable gastric band; N/R, not reported; RYGB, Roux-en-Y gastric bypass; SD, standard deviation; SG, sleeve gastrectomy; UK, United Kingdom; USA, United States; YFAS, Yale Food Addiction Scale; YFAS 2.0, Yale Food Addiction Scale 2.0.

3.5 | Meta-regression

Meta-regression analysis performed for the prevalence between preoperative and postoperative groups showed a significant difference in FA ($\beta = -16\%$; 95% CI: -25% to -7% ; $p = 0.001$).

3.6 | Certainty of evidence

Considering the limitations of the studies included in this review and the inconsistencies in the results, the quality of the evidence was deemed to be very low (Table 4). This classification was based on two criteria: limitations of the studies (quality assessment) and inconsistency in results (heterogeneity). Only these criteria were used due to the inadequacy of the analysis of the traditional criteria “indirect evidence,” “inaccuracy,” and

“publication bias,” given the nature of the studies included in the meta-analysis.

4 | DISCUSSION

4.1 | Summary of the results

The present study provides the results of a systematic review and meta-analysis of the prevalence of FA among individuals undergoing metabolic/bariatric surgery. The analysis included 40 studies, of which 26 presented the cross-sectional prevalence of FA at the preoperative moment and seven at the postoperative moment, and seven longitudinal studies assessed prevalence at both the preoperative and postoperative moments for the same individuals. Most studies evaluated FA using the original YFAS (67.5%). The

TABLE 2 Characteristics of included studies that assessed the prevalence of FA using the Yale Food Addiction Scale (YFAS) only in postoperative bariatric surgery patients ($n = 7$)

Author, year	YFAS version	Country	Type of surgery	Duration (months)	Sample (%Fem)	Prevalence of FA (%)	BMI mean (SD)	Risk of bias
Ames et al., 2017 ⁵⁸	YFAS	USA	SG RYGB	N/R	422 (76.3)	13.74	N/R	0.63 (moderate)
Cassin et al., 2020 ⁵⁹	mYFAS 2.0	Canada	RYGB	N/R	100 (82)	13.0	N/R	0.63 (moderate)
Ivezaj et al., 2019 ⁶⁰	YFAS	USA	SG	N/R	131 (82.8)	17.6	37.7 (7.3)	1.00 (low)
Mousavi et al., 2021 ¹⁴	YFAS	Iran	SG	24	450 (78)	19.8	31.36 (3.87)	0.85 (low)
Nicolau et al., 2021 ⁶¹	YFAS 2.0	Spain	SG	18	134 (71.6)	23.9	33.6 (6.6)	0.54 (moderate)
Steward et al., 2018 ⁶²	YFAS 2.0	Spain	N/R	N/R	69 (100)	24.2	42.4 (6.9)	0.76 (low)
Vidot et al., 2016 ⁶³	YFAS	USA	SG/RYGB/ LAGB	12	50 (76)	9.0	35.2 (10.4)	0.82 (low)

Abbreviations: BMI, body mass index; FA, food addiction; LAGB, laparoscopic adjustable gastric band; N/R, not reported; RYGB, Roux-en-Y gastric bypass; SD, standard deviation; SG, sleeve gastrectomy; UK, United Kingdom; USA, United States; YFAS, Yale Food Addiction Scale; YFAS 2.0, Yale Food Addiction Scale 2.0.

meta-analysis identified a high weighted prevalence of FA in preoperative samples (32%), whereas in postoperative samples, it was reduced to less than half (15%) that of the preoperative samples. Of the seven studies that prospectively measured preoperative and preoperative FA in the same sample, all showed a reduction in FA after surgery, which ranged from 1% to 59%, with a weighted APR of 26 p.p.

4.2 | Comparison with the literature and interpretation of data

The prevalence of FA at the preoperative moment (32%) of individuals was slightly higher than that observed in individuals with obesity but who are not necessarily candidates for bariatric surgery (28%).²⁹ This is consistent with findings in the literature, as there is evidence that individuals with obesity are more likely to meet the YFAS criteria for FA.⁷¹ Nevertheless, it is worth noting that not all individuals with obesity show FA and that other pathways involving physical inactivity, genetics, and the dietary environment may be involved in the genesis and prolongation of obesity.⁷² The weighted prevalence of FA after bariatric surgery (15%) was similar to that in individuals without body weight disorders (16%) and was slightly higher than that in non-clinical samples (13%).²⁹

The meta-regression showed that the difference in FA prevalence between the pre- and postoperative groups was statistically significant. This leads us to assume that bariatric surgery may result in a reduction in FA in this population. Nevertheless, the comparison of prevalence at preoperative and postoperative moments of cross-sectional studies using different samples strongly

decreases our ability to infer any causality. Longitudinal studies are better at determining the possible causal effects of bariatric surgery on FA prevalence. In this sense, the present review also evaluated seven longitudinal prospective studies, which indicated a reasonable weighted APR of 26 p.p. for FA. Such effects were observed as early as after 6 months of postoperative follow-up. However, there was marked variability in the APR; for example, a study with 12 months of follow-up showed an APR of only 1 p.p.⁶⁸ This probably occurred due to the discrepant sample size evaluated preoperatively and postoperatively. Of the 923 individuals included in the preoperative evaluation, which resulted in a low prevalence of FA (15%), only 170 individuals were evaluated in the postoperative period, indicating a significant sample loss and possible generation of selection bias in the sample, given that only the individuals most affected by FA may have returned for the second evaluation. On the other hand, the 24-month study showed an APR of 59 p.p.,⁶⁵ illustrating the potential beneficial effects of bariatric surgery on FA.

The presence of FA is associated with an extra burden in individuals with obesity, such as a lower quality of life,⁷³ impulsivity,⁷³ dysregulation of emotions,⁷³ reward dysfunction,⁷² and a higher likelihood of co-occurring mental health conditions.⁷³ After bariatric surgery, several biological, physiological, and metabolic changes occur, for example in neural transmission,^{74,75} in the regulation of hunger/satiety,^{76,77} on the palate,^{78,79} in food preferences or intolerances,^{80,81} and in the desire to eat.⁸² The observed reduction in the prevalence of FA in the postoperative period may be related to such metabolic changes promoted by bariatric surgery, as they are also associated with FA. Evidence shows that bariatric surgery also decreases other eating disorders, such as loss of eating control and binge eating disorders.^{83,84}

TABLE 3 Characteristics of the included longitudinal prospective studies that evaluated the prevalence of FA using the Yale Food Addiction Scale (YFAS) in the same subjects before and after bariatric surgery (n = 7)

Author, year	Country	Duration (months)	Type of surgery	Preoperative sample size (%Fem)	FA prevalence preoperative period, n (%)	Postoperative sample size, n	FA prevalence postoperative period, n (%)	Absolute prevalence reduction (APR) (p.p.)	95 CI% (p.p.)	Preoperative BMI (SD)	Postoperative BMI (SD)	Dropouts (%)	Risk of bias
Ben-Porat et al., 2022 ⁶⁴	Israel	24	SG	45 (100)	18 (40)	45	4 (14.2)	26.5	8.4; 42.8	44.9 (4.9)	31.5 (4.7)	27.40	0.7 (moderate)
Chiappetta et al., 2020 ⁶⁵	Germany	24	SG/GB	113 (68)	78 (69.0)	50	5 (10.0)	59.0	44.5; 68.3	51.7 (9.8)	N/R	55.75	0.63 (moderate)
Dong et al., 2020 ⁶⁶	USA	6	SG	18 (100)	3 (16.6)	18	0 (0.0)	16.6	-3.9; 39.2	45.5 (4.9)	37.4 (9.7)	No losses	0.63 (moderate)
Holgerson et al., 2018 ⁶⁷	UK	12	SG/RYGB	1586 (73)	238 (15.0)	424	40 (9.4)	5.6	2.0; 8.6	47.22 (N/R)	47.44 (N/R)	73.26	0.92 (low)
Koball et al., 2016 ⁶⁸	USA	12	RYGB	923 (71)	129 (14.0)	169	22 (13.0)	1.0	-5.3; 5.8	45.8 (9)	N/R	42.00	0.76 (low)
Pepino et al., 2014 ⁶⁹	USA	9	RYGB/LAGB/SG	44 (88.63)	14 (31.8)	44	1 (2.2)	29.5	14.3; 44.4	47.85 (8.1)	38.1 (6.75)	No losses	0.86 (low)
Sevinçer et al., 2016 ⁷⁰	Turkey	12	SG/OmegaGB	166 (77.1)	96 (57.8)	134	18 (13.4)	44.4	34.1; 53.0	47.03 (7.1)	30.26 (5.9)	43.38	0.45 (moderate)

Note: All studies used the original YFAS version of the scale.

Abbreviations: APR, absolute prevalence reduction; BMI, body mass index; CI, confidence interval; FA, food addiction; Fem, female; GB, gastric bypass; N/R, no reported; RYGB, Roux-en-Y gastric bypass; SG, standard deviation; SG, sleeve gastrectomy; UK, United Kingdom; USA, United States; YFAS, Yale Food Addiction Scale.




FIGURE 2 Forest plot for the prevalence of food addiction in preoperative samples

Thus, the data from the present review indicate that individuals with FA undergoing bariatric surgery may experience a reduction in the symptoms of this condition, which, in turn, could contribute to the control of obesity in this population. However, more longitudinal studies, preferably with an experimental design with control groups, are needed to assess the negative causal effects of bariatric surgery on FA, especially after 12 months, as weight regain after this period is common.^{85,86} We also highlight the need for longer longitudinal studies to assess whether the effect of bariatric surgery on FA is enduring or transient. This is of great importance from the perspective of treatment of obesity and the recurrence of this condition after surgical intervention.

4.3 | Limitations

A possible limitation of this study is that it did not conduct a specific search for YFAS in bariatric surgery. However, the search strategy

found all studies that used any version of the YFAS (regardless of sample type), suggesting that all studies that evaluated FA in the bariatric population were also included. Another limitation is that the YFAS is a self-report measure, which can be affected by participants' subjective interpretation. Some patients undergoing bariatric surgery may be prone to under- or overreporting of FA due to concerns about how it may influence their access to bariatric surgery. However, the YFAS has been found to have strong psychometric properties in several contexts,⁸⁷⁻⁹⁰ which does increase confidence in the findings.

In addition, the YFAS was the only instrument used to assess FA, which better enables comparison between studies (rather than collate various measures). However, caution should be exercised in the interpretation and applicability of the data because the prevalence of FA reported in the studies showed wide variation, and more than half the studies showed a moderate risk of bias. It is noteworthy that the overall quality of the evidence, as assessed by GRADE,³² was low, mainly due to limitations of the studies and inconsistency in the results.




FIGURE 3 Forest plot for the prevalence of food addiction in postoperative samples




FIGURE 4 Forest plot for the weighted absolute prevalence reduction (APR) of longitudinal studies

Therefore, future research should seek to increase the quality of performance and reporting of the studies.

In addition, most of the studies included in the present review are cross-sectional, making it impossible to determine a causal effect of bariatric surgery on the reduction of FA prevalence. We included a

few longitudinal studies, which, although not experiments, could contribute to the inference of a possible causal effect of bariatric surgery. Nevertheless, such longitudinal studies showed a critical sample loss in the postoperative period, which also introduced bias to our analysis, thereby reinforcing the need for cautious interpretation of our data.

TABLE 4 Evidence quality assessment

Question: What is the prevalence of AA through YFAS in the pre-and postoperative period of bariatric surgery patients?		Results								
Quality assessment		Weighted prevalence	Quality	Importance	Other considerations	Imprecision	Inconsistency	Indirect evidence	Risk of bias	Study design
Number of studies (number of participants)										
Prevalence in the preoperative period	33 (8304)	32% (95% CI [27-37])	⊕○○○ Very low	High	None	N/A	Very severe	N/A	Severe	Observational
Prevalence in the postoperative period	14 (1754)	15% (95% CI [12-18])	⊕○○○ Very low	High	None	N/A	Very severe	N/A	Severe	Observational

Abbreviation: N/A, no applicable.

5 | CONCLUSION

The prevalence of FA was high in the preoperative period among individuals undergoing bariatric surgery. The meta-analysis identified a reduction in the prevalence of FA among patients after metabolic/bariatric surgery, and the difference in prevalence was significant, as shown by the meta-regression analysis. Although this review comprised 40 studies, only seven were longitudinal studies that evaluated the prevalence of FA before and after bariatric surgery. Therefore, further longitudinal studies are needed, especially with an experimental design with control groups, to determine the efficacy of surgical interventions for improving FA outcomes and their effects on long-term obesity control.

ACKNOWLEDGMENT

This work was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) with a research grant for Dafiny Praxedes (under the number of process 23065.005919/2021-75). CAPES had no role in the design, analysis, or writing of this article.

CONFLICT OF INTEREST

No conflict of interest statement.

ORCID

Dafiny R. Praxedes  <https://orcid.org/0000-0002-5345-3869>

André E. Silva-Júnior  <https://orcid.org/0000-0002-1501-171X>

Mateus L. Macena  <https://orcid.org/0000-0002-7168-9605>

Ashley N. Gearhardt  <https://orcid.org/0000-0003-3843-5731>

Nassib B. Bueno  <https://orcid.org/0000-0002-3286-0297>

REFERENCES

- World Health Organization. Obesity and overweight. Fact Sheets. Geneva (CHE); 2018 [cited 2022 Jan 31]. Available from: <http://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- Monteiro CA, Moura EC, Conde WL, Popkin BM. Socioeconomic status and obesity in adult populations of developing countries: a review. *Bull World Health Organ.* 2004;82(12):940-946.
- de Melo Barros D, da Silva AP, de Moura DF, et al. A influência da transição alimentar e nutricional sobre o aumento da prevalência de doenças crônicas não transmissíveis. *Braz J Dev.* 2021;7(7):74647-74664.
- Pagliai G, Dinu M, Madarena MP, Bonaccio M, Iacoviello L, Sofi F. Consumption of ultra-processed foods and health status: a systematic review and meta-analysis. *Br J Nutr.* 2021;125(3):308-318. doi:10.1017/S0007114520002688
- Gariepy G, Nitka D, Schmitz N. The association between obesity and anxiety disorders in the population: a systematic review and meta-analysis. *Int J Obes (Lond).* 2010;34(3):407-419. doi:10.1038/ijo.2009.252
- Tronieri JS, Wurst CM, Pearl RL, Allison KC. Sex differences in obesity and mental health. *Curr Psychiatry Rep.* 2017;19(6):1-11.
- Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Nutrition.* 2007;23(11-12):887-894. doi:10.1016/j.nut.2007.08.008
- Bertoli S, Leone A, Ponissi V, et al. Prevalence of and risk factors for binge eating behaviour in 6930 adults starting a weight loss or maintenance programme. *Public Health Nutr.* 2016;19(1):71-77. doi:10.1017/S1368980015001068

9. Arterburn DE, Telem DA, Kushner RF, Courcoulas AP. Benefits and risks of bariatric surgery in adults: a review. *Jama*. 2020;324(9):879-887. doi:10.1001/jama.2020.12567
10. Lent MR, Wood GC, Cook A, et al. Five-year weight change trajectories in Roux-en-Y gastric bypass patients. *J Patient Cent Res Rev*. 2016;3(3):181.
11. de Souza AC, Gomes DL, de Sá NN, de Carvalho KM. Presença de comorbidades. uso de medicamentos e suplementos nutricionais por mulheres com reganho de peso após 24 meses de bypass gástrico. *RBone-Rev Bras Obes Nutri Emagreciment*. 2018;12(74):738-744.
12. Taba JV, Suzuki MO, Nascimento FSD, et al. The development of feeding and eating disorders after bariatric surgery: a systematic review and meta-analysis. *Nutrients*. 2021;13(7):2396. doi:10.3390/nu13072396
13. Koball AM, Ames G, Goetze RE, Grothe K. Bariatric surgery as a treatment for food addiction? A review of the literature. *Curr Addict Rep*. 2020;7(1):1-8. doi:10.1007/s40429-020-00297-w
14. Mousavi M, Tabesh MR, Khalaj A, Eini-Zinab H, Jahromi SR, Abolhasani M. Food addiction disorder 2 years after sleeve gastrectomy; association with physical activity, body composition, and weight loss outcomes. *Obes Surg*. 2021;31(8):3444-3452. doi:10.1007/s11695-021-05420-y
15. Gearhardt AN, Yokum S, Orr PT, Stice E, Corbin WR, Brownell KD. Neural correlates of food addiction. *Arch Gen Psychiatry*. 2011;68(8):808-816. doi:10.1001/archgenpsychiatry.2011.32
16. Gearhardt AN, Corbin WR, Brownell KD. Preliminary validation of the Yale food addiction scale. *Appetite*. 2009;52(2):430-436. doi:10.1016/j.appet.2008.12.003
17. Gearhardt AN, Corbin WR, Brownell KD. Food addiction: an examination of the diagnostic criteria for dependence. *J Addict Med*. 2009;3(1):1-7. doi:10.1097/ADM.0b013e318193c993
18. Fernandez-Aranda F, Karwautz A, Treasure J. Food addiction: a transdiagnostic construct of increasing interest. *Eur Eat Disord Rev*. 2018;26(6):536-540. doi:10.1002/erv.2645
19. Jiménez-Murcia S, Agüera Z, Paslakis G, et al. Food addiction in eating disorders and obesity: analysis of clusters and implications for treatment. *Nutrients*. 2019;11(11):2633. doi:10.3390/nu11112633
20. Camacho-Barcia L, Munguía L, Lucas I, et al. Metabolic, affective and neurocognitive characterization of metabolic syndrome patients with and without food addiction. Implications for weight progression. *Nutrients*. 2021;13(8):2779. doi:10.3390/nu13082779
21. Guerrero Pérez F, Sánchez-González J, Sánchez I, et al. Food addiction and preoperative weight loss achievement in patients seeking bariatric surgery. *Eur Eat Disord Rev*. 2018;26(6):645-656. doi:10.1002/erv.2649
22. Dickhut C, Hase C, Gruner-Labitzke K, et al. No addiction transfer from preoperative food addiction to other addictive behaviors during the first year after bariatric surgery. *Eur Eat Disord Rev*. 2021;29(6):924-936. doi:10.1002/erv.2857
23. Gearhardt AN, Hebebrand J. The concept of "food addiction" helps inform the understanding of overeating and obesity: debate consensus. *Am J Clin Nutr*. 2021;113(2):274-276. doi:10.1093/ajcn/nqaa345
24. Vainik U, Neseliler S, Konstabel K, Fellows LK, Dagher A. Eating traits questionnaires as a continuum of a single concept. Uncontrolled eating. *Appetite*. 2015;90:229-239. doi:10.1016/j.appet.2015.03.004
25. Fletcher PC, Kenny PJ. Food addiction: a valid concept? *Neuropsychopharmacology*. 2018;43(13):2506-2513. doi:10.1038/s41386-018-0203-9
26. Gearhardt AN, Boswell RG, White MA. The association of "food addiction" with disordered eating and body mass index. *Eat Behav*. 2014;15(3):427-433. doi:10.1016/j.eatbeh.2014.05.001
27. Hebebrand J, Gearhardt AN. The concept of "food addiction" helps inform the understanding of overeating and obesity: NO. *Am J Clin Nutr*. 2021;113(2):268-273. doi:10.1093/ajcn/nqaa344
28. Ivezaj V, Wiedemann AA, Grilo CM. Food addiction and bariatric surgery: a systematic review of the literature. *Obes Rev*. 2017;18(12):1386-1397. doi:10.1111/obr.12600
29. Praxedes DR, Silva-Júnior AE, Macena ML, et al. Prevalence of food addiction determined by the Yale Food Addiction Scale and associated factors: a systematic review with meta-analysis. *Eur Eat Disord Rev*. 2021;30(2):85-95. doi:10.1002/erv.2878
30. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Int J Surg*. 2021;88:105906. doi:10.1016/j.jisu.2021.105906
31. Viswanathan M, Berkman ND. Development of the RTI item bank on risk of bias and precision of observational studies. *J Clin Epidemiol*. 2012;65(2):163-178. doi:10.1016/j.jclinepi.2011.05.008
32. Grade Working Group. The grading of recommendations assessment, development and evaluation. 2017 (accessed April 5, 2022).
33. Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Archives of Public Health*. 2014;72(1):1-10.
34. El Archi S, Brunault P, De Luca A, et al. Do emotion dysregulation, alexithymia and personality dimensions explain the association between attention-deficit/hyperactivity disorder and binge eating among bariatric surgery candidates? *Front Psychol*. 2021;5400.
35. Benzerouk F, Gierski F, Ducluzeau PH, et al. Food addiction, in obese patients seeking bariatric surgery, is associated with higher prevalence of current mood and anxiety disorders and past mood disorders. *Psychiatry Res*. 2018;267:473-479. doi:10.1016/j.psychres.2018.05.087
36. Bianciardi E, Fabbriatore M, Di Lorenzo G, et al. Prevalence of food addiction and binge eating in an Italian sample of bariatric surgery candidates and overweight/obese patients seeking low-energy-diet therapy. *Riv Psichiatr*. 2019;54(3):127-130. doi:10.1708/3181.31602
37. Brunault P, Ducluzeau PH, Bourbao-Tournois C, et al. Food addiction in bariatric surgery candidates: prevalence and risk factors. *Obes Surg*. 2016;26(2):1650-1653. doi:10.1007/s11695-016-2189-x
38. Brunault P, Berthoz S, Gearhardt AN, et al. The Modified Yale Food Addiction Scale 2.0: validation among non-clinical and clinical French-speaking samples and comparison with the Full Yale Food Addiction Scale 2.0. *Front Psych*. 2020;11:892.
39. Cambiali E, Avella MT, Arone A, et al. P. 224 eating behaviours in a sample of 141 candidates for bariatric surgery: clinical and psychopathological correlates. *Eur Neuropsychopharmacol*. 2021;44:S32. doi:10.1016/j.euroneuro.2021.01.053
40. Çelebi C, Sönmez Güngör E, Akvardar Y. Personality dimensions associated with food addiction in a sample of preoperative bariatric surgery patients from Turkey. *Int J Ment Health Addict*. 2021;1-12.
41. Clark SM, Saules KK. Validation of the Yale food addiction scale among a weight-loss surgery population. *Eat Behav*. 2013;14(2):216-219. doi:10.1016/j.eatbeh.2013.01.002
42. Clark SM, Martens K, Smith-Mason CE, Hamann A, Miller-Matero LR. Validation of the Yale Food Addiction Scale 2.0 among a bariatric surgery population. *Obes Surg*. 2019;29(9):2923-2928. doi:10.1007/s11695-019-03927-z
43. Cullen AJ, Barnett A, Komesaroff PA, et al. A qualitative study of overweight and obese Australians' views of food addiction. *Appetite*. 2017;115:62-70. doi:10.1016/j.appet.2017.02.013
44. Fuentes M, Gabler G, Silva J, Olguín P, Rodríguez A. Relation between food addiction and nutritional status in patients candidates for bariatric surgery. 2017.
45. Gabler G, Fuentes M, Rodríguez A, Barbosa AC, Zarazua CJ. P. 6. f. 010 assessing for food addiction in an obese Chilean population seeking for bariatric surgery. *Eur Neuropsychopharmacol*. 2015;25:S633.
46. Hernandez J, Cervoni C, Hymowitz G. The effects of impulsivity on food addiction and eating behaviors in a bariatric sample. *Surg Obes Relat Dis*. 2017;13(10):S93. doi:10.1016/j.soard.2017.09.204

47. Holgerson AA, Clark MM, Frye MA, et al. Symptoms of bipolar disorder are associated with lower bariatric surgery completion rates and higher food addiction. *Eat Behav.* 2021;40:101462.
48. Koball AM, Borgert AJ, Kallies KJ, Grothe K, Ames G, Gearhardt AN. Validation of the Yale Food Addiction Scale 2.0 in patients seeking bariatric surgery. *Obes Surg.* 2021;31(4):1533-1540. doi:10.1007/s11695-020-05148-1
49. Lawson JL, Goldman RL, Swencionis C, Wien R, Persaud A, Parikh M. Examining food addiction and acculturation among a Hispanic bariatric surgery-seeking participant group. *Obes Surg.* 2019;29(7):2151-2157. doi:10.1007/s11695-019-03799-3
50. Lescher M, Wegmann E, Müller SM, et al. A randomized study of food pictures-influenced decision-making under ambiguity in individuals with morbid obesity. *Front Psych.* 2020;11:822.
51. Meule A, Heckel D, Kübler A. Factor structure and item analysis of the Yale Food Addiction Scale in obese candidates for bariatric surgery. *Eur Eat Disord Rev.* 2012;20(5):419-422. doi:10.1002/erv.2189
52. Meule A, Heckel D, Jurowich CF, Vögele C, Kübler A. Correlates of food addiction in obese individuals seeking bariatric surgery. *Clinical Obesity.* 2014;4(4):228-236. doi:10.1111/cob.12065
53. Meule A, de Zwaan M, Müller A. Attentional and motor impulsivity interactively predict 'food addiction' in obese individuals. *Compr Psychiatry.* 2017;72:83-87.
54. Miller-Matero LR, Armstrong R, McCulloch K, Hyde-Nolan M, Eshelman A, Genaw J. To eat or not to eat; is that really the question? An evaluation of problematic eating behaviors and mental health among bariatric surgery candidates. *Eat Weight Disord.* 2014;19(3):377-382. doi:10.1007/s40519-014-0118-3
55. Müller A, Leukefeld C, Hase C, et al. Food addiction and other addictive behaviours in bariatric surgery candidates. *Eur Eat Disord Rev.* 2018;26(6):585-596. doi:10.1002/erv.2629
56. Murray SM, Tweardy S, Geliebter A, Avena NM. A longitudinal preliminary study of addiction-like responses to food and alcohol consumption among individuals undergoing weight loss surgery. *Obes Surg.* 2019;29(8):2700-2703. doi:10.1007/s11695-019-03915-3
57. Reslan S, Saules KK, Greenwald MK, Schuh LM. Substance misuse following Roux-en-Y gastric bypass surgery. *Subst Use Misuse.* 2014;49(4):405-417. doi:10.3109/10826084.2013.841249
58. Ames GE, Heckman MG, Diehl NN, et al. Guiding patients toward the appropriate surgical treatment for obesity: should presurgery psychological correlates influence choice between Roux-en-Y gastric bypass and vertical sleeve gastrectomy? *Obes Surg.* 2017;27(10):2759-2767. doi:10.1007/s11695-017-2876-2
59. Cassin S, Leung S, Hawa R, Wnuk S, Jackson T, Sockalingam S. Food addiction is associated with binge eating and psychiatric distress among postoperative bariatric surgery patients and may improve in response to cognitive behavioural therapy. *Nutrients.* 2020;12(10):2905.
60. Ivezaj V, Wiedemann AA, Lawson JL, Grilo CM. Food addiction in sleeve gastrectomy patients with loss-of-control eating. *Obes Surg.* 2019;29(7):2071-2077. doi:10.1007/s11695-019-03805-8
61. Nicolau J, Dotres K, Ayala L, et al. Long-term prevalence of food addiction among bariatric surgery patients: influence on metabolic and psychological outcomes. *Metab Syndr Relat Disord.* 2021;19(3):152-158. doi:10.1089/met.2020.0079
62. Steward T, Mestre-Bach G, Vintró-Alcaraz C, et al. Food addiction and impaired executive functions in women with obesity. *Eur Eat Disord Rev.* 2018;26(6):574-584. doi:10.1002/erv.2636
63. Vidot DC, Prado G, de la Cruz-Munoz N, Spadola C, Cuesta M, Messiah SE. Postoperative marijuana use and disordered eating among bariatric surgery patients. *Surg Obes Relat Dis.* 2016;12(1):171-178. doi:10.1016/j.soard.2015.06.007
64. Ben-Porat T, Košir U, Peretz S, Sherf-Dagan S, Stojanovic J, Sakran N. Food addiction and binge eating impact on weight loss outcomes two years following sleeve gastrectomy surgery. *Obes Surg.* 2022;1-8.
65. Chiappetta S, Stier C, Hadid MA, et al. Remission of food addiction does not induce cross-addiction after sleeve gastrectomy and gastric bypass: a prospective cohort study. *Obes Facts.* 2020;13(3):307-320. doi:10.1159/000506838
66. Dong TS, Gupta A, Jacobs JP, et al. Improvement in uncontrolled eating behavior after laparoscopic sleeve gastrectomy is associated with alterations in the brain-gut-microbiome axis in obese women. *Nutrients.* 2020;12(10):2924. doi:10.3390/nu12102924
67. Holgerson AA, Clark MM, Ames GE, et al. Association of adverse childhood experiences and food addiction to bariatric surgery completion and weight loss outcome. *Obes Surg.* 2018;28(11):3386-3392. doi:10.1007/s11695-018-3370-1
68. Koball AM, Clark MM, Collazo-Clavell M, et al. The relationship among food addiction, negative mood, and eating-disordered behaviors in patients seeking to have bariatric surgery. *Surg Obes Relat Dis.* 2016;12(1):165-170. doi:10.1016/j.soard.2015.04.009
69. Pepino MY, Stein RI, Eagon JC, Klein S. Bariatric surgery-induced weight loss causes remission of food addiction in extreme obesity. *Obesity.* 2014;22(8):1792-1798. doi:10.1002/oby.20797
70. Sevinçer GM, Konuk N, Bozkurt S, Coşkun H. Food addiction and the outcome of bariatric surgery at 1-year: prospective observational study. *Psychiatry Res.* 2016;244:159-164. doi:10.1016/j.psychres.2016.07.022
71. Gearhardt AN, Corbin WR, Brownell KD. Development of the Yale Food Addiction Scale Version 2.0. *Psychol Addict Behav.* 2016;30:113-112.
72. Gearhardt AN, Schulte EM. Is food addictive? A review of the science. *Annu Rev Nutr.* 2021;41(1):387-410. doi:10.1146/annurev-nutr-110420-111710
73. Minhas M, Murphy CM, Balodis IM, Samokhvalov AV, MacKillop J. Food addiction in a large community sample of Canadian adults: prevalence and relationship with obesity, body composition, quality of life and impulsivity. *Addiction.* 2021;116(10):2870-2879. doi:10.1111/add.15446
74. Orellana ER, Covasa M, Hajnal A. Neuro-hormonal mechanisms underlying changes in reward related behaviors following weight loss surgery: potential pharmacological targets. *Biochem Pharmacol.* 2019;164:106-114. doi:10.1016/j.bcp.2019.04.004
75. Karlsson HK, Tuominen L, Helin S, Salminen P, Nuutila P, Nummenmaa L. Mesolimbic opioid-dopamine interaction is disrupted in obesity but recovered by weight loss following bariatric surgery. *Transl Psychiatry.* 2021;11(1):1, 259-7. doi:10.1038/s41398-021-01370-2
76. Makaronidis JM, Neilson S, Cheung WH, et al. Reported appetite, taste and smell changes following Roux-en-Y gastric bypass and sleeve gastrectomy: effect of gender, type 2 diabetes and relationship to postoperative weight loss. *Appetite.* 2016;107:93-105. doi:10.1016/j.appet.2016.07.029
77. Cazzo E, Pareja JC, Chaim EA, Geloneze B, Barreto MRL, Magro DO. GLP-1 and GLP-2 levels are correlated with satiety regulation after Roux-en-Y gastric bypass: results of an exploratory prospective study. *Obes Surg.* 2017;27(3):703-708. doi:10.1007/s11695-016-2345-3
78. Al-Alsheikh AS, Alabdulkader S, Johnson B, Goldstone AP, Miras AD. Effect of obesity surgery on taste. *Nutrients.* 2022;14(4):866. doi:10.3390/nu14040866
79. Shoar S, Naderan M, Shoar N, Modukuru VR, Mahmoodzadeh H. Alteration pattern of taste perception after bariatric surgery: a systematic review of four taste domains. *Obes Surg.* 2019;29(5):1542-1550. doi:10.1007/s11695-019-03730-w
80. Nielsen MS, Schmidt JB, le Roux CW, Sjödin A. Effects of Roux-en-Y gastric bypass and sleeve gastrectomy on food preferences and potential mechanisms involved. *Curr Obes Rep.* 2019;8(3):292-300. doi:10.1007/s13679-019-00354-0
81. de Almeida Godoy CM, de Araújo Quadros Cunha B, Furtado MC, de Godoy EP, de Souza LBR, Oliveira AG. Relationship of food

- intolerance 2 years after roux-en-y gastric bypass surgery for obesity with masticatory efficiency and protein consumption. *Obes Surg*. 2020;30(8):3093-3098. doi:[10.1007/s11695-020-04669-z](https://doi.org/10.1007/s11695-020-04669-z)
82. Jabbour J, Awada D, Naim N, et al. Impact of bariatric surgery on the healthy eating index, binge eating behavior and food craving in a middle eastern population: a Lebanese experience. *Healthcare*. 2021; 9(11):1416. doi:[10.3390/healthcare9111416](https://doi.org/10.3390/healthcare9111416)
83. White MA, Kalarchian MA, Masheb RM, Marcus MD, Grilo CM. Loss of control over eating predicts outcomes in bariatric surgery patients: a prospective. 24-month follow-up study. *J Clin Psychiatry*. 2009; 70(2):10844.
84. Kalarchian MA, King WC, Devlin MJ, et al. Psychiatric disorders and weight change in a prospective study of bariatric surgery patients: a 3-year follow-up. *Psychosom Med*. 2016;78(3):373-381. doi:[10.1097/PSY.0000000000000277](https://doi.org/10.1097/PSY.0000000000000277)
85. McGrice M, Paul KD. Interventions to improve long-term weight loss in patients following bariatric surgery: challenges and solutions. *Diabetes Metab Syndr Obes: Targets Ther*. 2015;8:263.
86. Velapati SR, Shah M, Kuchkuntla AR, et al. Weight regain after bariatric surgery: prevalence, etiology, and treatment. *Curr Nutr Rep*. 2018; 7(4):329-334. doi:[10.1007/s13668-018-0243-0](https://doi.org/10.1007/s13668-018-0243-0)
87. Torres S, Camacho M, Costa P, et al. Psychometric properties of the Portuguese version of the Yale Food Addiction Scale. *Eat Weight Disord*. 2017;22(2):259-267. doi:[10.1007/s40519-016-0349-6](https://doi.org/10.1007/s40519-016-0349-6)
88. Innamorati M, Imperatori C, Manzoni GM, et al. Psychometric properties of the Italian Yale Food Addiction Scale in overweight and obese patients. *Eat Weight Disord*. 2015;20(1):119-127. doi:[10.1007/s40519-014-0142-3](https://doi.org/10.1007/s40519-014-0142-3)
89. Sevinçer GM, Konuk N, Bozkurt S, Saraçlı Ö, Coşkun H. Psychometric properties of the Turkish version of the Yale Food Addiction Scale among bariatric surgery patients. *Anadolu Psikiyatri Derg*. 2014;16; S44-S53. doi:[10.5455/apd.174345](https://doi.org/10.5455/apd.174345)
90. Panahi A, Haghayegh SA. Developing the Persian version of Yale Food Addiction Scale and assessing its psychometric properties. *Iran J Psychiatry Clin Psychol*. 2020;25(4):454-471. doi:[10.32598/ijpcp.25.4.8](https://doi.org/10.32598/ijpcp.25.4.8)

How to cite this article: Praxedes DR, Silva-Júnior AE, Macena ML, Gearhardt AN, Bueno NB. Prevalence of food addiction among patients undergoing metabolic/bariatric surgery: A systematic review and meta-analysis. *Obesity Reviews*. 2023;24(2):e13529. doi:[10.1111/obr.13529](https://doi.org/10.1111/obr.13529)