

Hahn Samantha L (Orcid ID: 0000-0002-9553-0383)  
Burnette C Blair (Orcid ID: 0000-0002-5003-5856)  
Sonnevile Kendrin R (Orcid ID: 0000-0002-0359-3919)

**Title:** Eating disorder risk in rural US adolescents: What do we know and where do we go?

**Authors:** Samantha L. Hahn, PhD, MPH, RD,<sup>1</sup> C. Blair Burnette, PhD,<sup>2</sup> Kelley A. Borton, PhD, MPH, RDN,<sup>3,4</sup> Lisa Mitchell Carpenter, MA, LPC<sup>4</sup> Kendrin R. Sonnevile, ScD, RD,<sup>5</sup> Beth Bailey, PhD<sup>1</sup>

<sup>1</sup>Central Michigan University College of Medicine, Mount Pleasant, MI, 48859, USA;  
[sam.hahn@cmich.edu](mailto:sam.hahn@cmich.edu); [beth.bailey@cmich.edu](mailto:beth.bailey@cmich.edu)

<sup>2</sup>University of Minnesota School of Public Health, Minneapolis, MN 55455, USA;  
[burne339@umn.edu](mailto:burne339@umn.edu)

<sup>3</sup>Oakland University School of Health Sciences, Rochester, MI 48309, USA;  
[kborton@oakland.edu](mailto:kborton@oakland.edu)

<sup>4</sup>Center of Hope Counseling, Mount Pleasant, MI 48858, USA; [centerofhope2016@gmail.com](mailto:centerofhope2016@gmail.com)

<sup>5</sup>University of Michigan School of Public Health, Ann Arbor, MI 48109, USA;  
[kendrins@umich.edu](mailto:kendrins@umich.edu)

**Corresponding Author:**

Samantha L. Hahn, PhD, MPH, RD  
Assistant Professor and Health Services Researcher  
Central Michigan University College of Medicine  
1280 East Campus Drive, Foust 130  
Mount Pleasant, MI 48859, USA  
[Sam.Hahn@cmich.edu](mailto:Sam.Hahn@cmich.edu)  
(989) 774-7656  
ORCID: 0000-0002-9553-0383

**Author Contributions Statement:** The paper was conceptualized and originally drafted by SLH. All authors reviewed and edited subsequent drafts.

**Conflict of Interest:** The authors declare they have no conflicts of interest.

**Data Availability:** Because there was no primary data used in the present manuscript, no data is available.

**Keywords:** Eating Disorder, Disordered Eating, Rural Health, Epidemiology, Adolescents, Prevention, Health Disparities

**Abstract (unstructured for Spotlight):**

Adolescence is a vulnerable period for the development of eating disorders, but there are disparities in eating disorder risk among adolescents. One population that may be at increased risk but is vastly understudied is adolescents residing in rural regions within the US. Rural communities face many mental and physical health disparities; however, the literature on rural adolescent eating disorder risk is nearly non-existent. In this paper we summarize the scant literature on disordered eating and eating disorder risk and prevalence among rural US adolescents. We also detail eating disorder risk factors that may have unique influence in this population, including socioeconomic status, food insecurity, healthcare access, body image, and weight stigma. Given the presence of numerous eating disorder risk factors, we speculate that rural adolescents may be a particularly vulnerable population for eating disorders and we

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: [10.1002/eat.23843](https://doi.org/10.1002/eat.23843)

This article is protected by copyright. All rights reserved.

propose critical next steps in research for understanding eating disorder risk in among the understudied population of rural adolescents.

## Introduction

Adolescence is a pivotal developmental period for disordered eating and eating disorders, given the physical and social changes that occur during this time.<sup>1,2</sup> However, there are known disparities in eating disorders, putting some adolescents at increased risk,<sup>3</sup> including those who experience food insecurity<sup>4,5</sup> and adverse childhood experiences<sup>4,5</sup> and those who identify as sexual and gender minorities.<sup>6</sup> One population that may have elevated eating disorder risk is rural adolescents. While the definition of what constitutes a rural community varies, rural communities are generally characterized as areas with low population density located outside of metropolitan areas or cities.<sup>7,8</sup> Irrespective of the exact definition used, rural communities are shown to face mental<sup>9-11</sup> and physical health disparities.<sup>12-14</sup> However, adolescents from rural populations are an underserved and understudied population relative to disordered eating and eating disorder risk, and little is known about eating disorder risk among rural adolescents. Therefore, the objective of this paper is to summarize the evidence on the prevalence and unique risk factors of eating disorders among rural adolescents and outline critical future directions for research.

## Prevalence of Disordered Eating and Eating Disorders in Rural Adolescents

Little data exists on the prevalence of disordered eating (e.g., fasting, diet pill use, purging, binge eating, disordered muscle control behaviors, etc.) and eating disorders among rural US adolescents. To our knowledge, there are only two studies that assessed prevalence of disordered eating behaviors and eating disorder risk among rural adolescents in the US. One study of 1,302 students from the Appalachian East Tennessee area found that 24.8% of female and 6.5% of male rural high school students screened as high risk for an eating disorder (as determined by the EAT-40), compared to 14.6% of females, and no males among the least rural communities surveyed in the study.<sup>15</sup> These results therefore suggest that adolescents from rural communities may have elevated eating disorder risk relative to their urban-dwelling peers. The other known study found that in a population of 2,629 high school girls from rural communities within the midwestern state of North Dakota, 11.8% of high school students used diet pills and 9.9% endorsed self-induced vomiting.<sup>16</sup> Comparatively, a study of a nationally representative sample of US high school students found that only 5.1% took diet pills, and 3.8% reported self-induced vomiting, further indicating that rural adolescents may be at increased risk compared to the general population of adolescents.<sup>17</sup> However, the two studies examining eating disorder risk in rural adolescents were limited to high school students from specific communities, and both were conducted nearly 30 years ago. Thus, results may no longer be valid, nor generalizable to other rural communities in the US. Further, these studies did not assess other disordered eating behaviors such as restriction, binge eating, or muscle building behaviors, nor did they assess specific eating disorder diagnoses. They also did not assess factors such as sexual and gender identity, or race/ethnicity. While outside the US context, two studies of rural Canadian adolescents have also reported high prevalence of disordered eating behaviors; specifically, over a quarter reported skipping meals for weight management,<sup>18</sup> 14.2% reported binge eating,<sup>19</sup> 5.2% self-induced vomiting, and 5.5% using laxatives, diet pills, or diuretics. Further, 17% of females and 8.3% of males screened as high risk for an eating disorder (as determined by the EAT-26). However, these studies are nearly 20 years old, and the prevalence of eating disorders has increased in the general population during this period,<sup>20,21</sup> and results may not generalize to the US. Taken together, there is a critical lack of current or sufficient data on the prevalence of either disordered eating or full-threshold eating

disorders among US rural adolescents. Because the little available data suggest rural adolescents may be at increased risk for an eating disorder, high-quality and representative epidemiological data are urgently needed.

## **Unique Risk Factors**

### *Socioeconomic Status and Food Insecurity*

Despite the longstanding assumption that eating disorders are illnesses of affluence, eating disorders may actually be more common among those who are less socioeconomically advantaged.<sup>21,22</sup> Several studies have also shown that lower socioeconomic status (SES) is associated with higher prevalence of disordered eating behaviors, including unhealthy weight control behaviors (e.g. fasting, skipping meals),<sup>23</sup> binge eating, and purging<sup>24</sup> among US adolescents. Rural children are 30% more likely to live in poverty compared to their urban counterparts,<sup>25</sup> and, therefore, may be at heightened eating disorder risk in part due to the increased likelihood for lower SES. Relatedly, food insecurity is also more common among rural populations compared to metropolitan areas,<sup>26</sup> due in part to lower SES, but also driven by decreased food access. Food access may be a particularly salient factor in the potential relationship between food insecurity and eating disorder risk among rural adolescents, as rural communities are less likely to have access to full-service grocery stores that offer fresh fruits and vegetables and are therefore reliant on corner stores and other convenience stores that offer mostly shelf-stable and highly processed foods.<sup>27</sup> Reduced food access due to geographical distance and healthful selection in stores may further contribute to or exacerbate food insecurity in this population, which to disordered eating for a myriad of reasons.<sup>28</sup> Indeed, food insecurity is consistently related to mental and physical health disparities<sup>29,30</sup> including elevated eating disorder risk.<sup>28,31,32</sup> Accordingly, the increased likelihood of poverty, reduced food access, and overall higher levels of food insecurity among rural adolescents, may make rural adolescents particularly susceptible to eating disorders.

### *Health Care Access*

Rural populations are more likely to be medically underserved for numerous reasons, including affordability, distance to providers, lack of providers and services, and cultural perceptions that decrease healthcare utilization.<sup>33</sup> Primary care providers are typically the first, and sometimes only, providers to see patients suffering with eating disorders,<sup>34</sup> and this may be particularly true in rural communities where subspecialty services (i.e. adolescent medicine, gastroenterology, psychiatry) are lacking. However, primary care providers often are not trained in eating disorders and may fail to screen for eating disorders at all or may limit their screening to those who meet antiquated stereotypes about who is most likely to have an eating disorder (e.g. thin white girls).<sup>35</sup> Further, data suggest some primary care providers would choose not to screen for eating disorders when treatment pathways are unavailable.<sup>36</sup> In rural populations, eating disorder treatment options are particularly scarce given the cost and the overall lack of available specialty care.<sup>37</sup> Although telehealth options have risen in recent years and may be a viable option in overcoming the scarcity of treatment options,<sup>38</sup> there may be additional barriers for telehealth in rural communities that must be addressed first, including lack of internet providers servicing rural areas, unreliable internet or cellphone service at their home, or stigma-related concerns. Issues related to health care access in rural populations, may lead to delayed diagnosis, which itself is associated with poorer long-term prognosis.<sup>39</sup> Indeed, a study of

Australian adolescents found that rural teens have more severe physical eating disorder symptoms, likely because they receive treatment later than their more urban peers.<sup>40</sup>

Although healthcare-related stigma is common in the US, particularly behavioral or mental health, rural populations face unique pressures.<sup>33</sup> For example, rural patients often perceive their providers as neighbors and friends and are reluctant to share vulnerable information with providers with whom they are likely to see regularly in their communities.<sup>41</sup> For adolescents, there could be the additional fear that the provider will share sensitive information with their parents given close community connections. Evidence also suggests that individuals from rural communities may be less likely to seek or accept treatment for mental health concerns like eating disorders, believing they can “fix” their own mental health issues and seeing mental health struggles as personal shortcomings.<sup>42</sup> Indeed, an epidemiological study of eating disorder diagnoses and treatment among college students found those from less affluent backgrounds were less likely to perceive a need for treatment.<sup>35</sup> Therefore, rural adolescents may not only be less likely to perceive they need assistance, but may be less likely to be diagnosed and have available treatment options when they do receive a diagnosis or perceive need for treatment.

### *Body Ideals, Body Image, and Weight Stigma*

Rural adolescents have unique experiences of body image and weight stigma given the profound influence of culture on an individual’s construction of body and appearance ideals.<sup>2</sup> Like urban youth, rural adolescents’ body and appearance ideals are heavily influenced by the media.<sup>43</sup> However, rural communities may exert unique influences on adolescents as well. For example, rural populations place an emphasis on the ability to contribute to their community and many individuals do so by working labor-intensive jobs (e.g., farming, manufacturing).<sup>44</sup> These jobs may lead to body ideals emphasizing more muscular physiques, though muscular ideals and disordered muscle building behaviors have not been explored in this population. Alternatively, an emphasis on appreciation of the body’s ability to perform, regardless of physical appearance, could be protective against disordered eating.<sup>45</sup> Research that describes the characteristics and influences of body ideals within this community is needed.

Although research on body image in rural populations is lacking, the presence of weight stigma has been documented. Rural adolescents report experiences of weight-based teasing from peers,<sup>43</sup> and rural adults report experiences of weight stigma in health care settings.<sup>46</sup> Weight stigma is common throughout the US, but may be particularly harmful in rural communities for several reasons. First, the prevalence of higher body weight is more common in rural than urban areas, meaning individuals within rural communities likely experience disproportionate weight bias and discrimination.<sup>47</sup> Next, evidence suggests that rural adolescents believe that having a larger body will lead to loss of belonging among peers and family, and a loss of functionality and productivity.<sup>43</sup> Because of the importance of community and emphasis on productivity as a means of contributing to the community, the perceived consequences of higher weight may be particularly salient and severe in rural populations.<sup>43</sup> Finally, rural adolescents report believing weight is highly malleable, that individuals hold blame for their weight, and that people should be forced to engage in activities to lose weight, suggesting they may see weight loss as easily achievable.<sup>43</sup> Thus, rural adolescents may be particularly likely to experience and internalize weight stigma, thereby feeling compelled to attempt weight loss, and may be especially likely to use disordered eating behaviors to attempt weight loss or to cope with the distress of experiencing weight stigma.<sup>43,48</sup> Future research is

needed to explore influences of body ideals and experiences of weight stigma among this population, including examining differences by source of information (e.g., peers, family, media, and healthcare).

### *Other Risk Factors*

Many other factors not discussed likely affect eating disorder risk among rural adolescents and warrant further investigation. For example, there is evidence that childhood sexual abuse is more strongly associated with self-induced vomiting for weight loss among rural adolescents than urban adolescents.<sup>16</sup> Further, rates of substance use and substance use-associated mortality are higher among rural than urban populations, which some theorize may be secondary to untreated mental health concerns and maladaptive coping mechanisms to difficult experiences.<sup>49</sup> Thus, substance use presents another potential risk factor for rural adolescents given it is strongly linked with disordered eating.<sup>50,51</sup> Importantly, health disparities among rural communities differentially impact people of color, LGBTQ+ individuals, and those experiencing poverty.<sup>52-54</sup> Certain populations, such as American Indian/Native American people, may be both more likely to engage in disordered eating behaviors<sup>55</sup> and to live in rural communities.<sup>56</sup> Thus, the effects of systemic inequities may be compounded for those carrying multiple socially marginalized identities, thereby likely increasing eating disorder risk.<sup>57,58</sup> This is likely true for indigenous populations both within and outside the US.<sup>59</sup>

### **Conclusions and Future Directions**

The research on eating disorder risk and prevalence among rural US adolescents is scant and outdated. Although this paper focused on rural U.S. adolescents, many of the research gaps and factors discussed are likely applicable to rural communities outside of the US. While some risk factors (e.g., healthcare access) likely apply to most rural settings, risk factors seen in rural communities in the U.S. may be rooted more in demographics than in geography, thereby limiting their applicability to non-U.S. contexts. Our review of the literature suggests rural populations may be particularly vulnerable to developing eating disorders and may be at risk for under-diagnosis and under-treatment. Research examining the extent and severity of disordered eating among rural adolescents, as well as unique risk and protective factors within rural communities, is critically needed to inform prevention, identification, and treatment efforts. Specifically, large epidemiologic studies are needed to: assess and monitor prevalence of disordered eating in rural communities; examine if eating disorder risk is higher in rural communities compared to urban and suburban populations, as well as if there are differences in prevalences depending on the definition of rurality used; and examine if certain populations within rural communities are at especially increased risk (e.g., people of color, LGBTQ+, etc.), including examining risk within an intersectional framework. Although existing nationally representative studies of adolescents do not include disordered eating questions and/or do not assess rurality, it may be a feasible option to alter existing studies such as the Youth Risk Behavior Survey to reintroduce disordered eating questions and include direct information on participant rurality within the publicly available dataset. Additionally, because early intervention and specialty care are important prognostic indicators for recovery, research should also focus on establishing best practices for screening and referral in rural communities and identifying innovative approaches to increasing treatment accessibility within this population while addressing financial and geographical restrictions. Moreover, because of the numerous barriers for treatment, prevention is particularly important in rural adolescents. More research specifically examining eating disorder risk among rural adolescents is needed to clarify the

unique risk and protective factors and mechanisms contributing to eating disorder risk in this population. Such research can then be used to inform the development of interventions, policies, and practices that effectively and efficiently prevent eating disorders in this population. Therefore, prevention-focused research in rural communities is needed to reduce the overall disease burden in this population and to prevent the progression of risk behaviors (i.e., dieting, disordered weight control behaviors) to full-threshold eating disorders. Given the well-established disparities in eating disorder symptoms, diagnosis, and treatment, research that uses an intersectional lens to examine unique eating disorder risk and protective factors among rural populations should be a public health priority.

**Public Health Significance:**

Rural adolescents may be at increased risk for eating disorders due to disproportionate burden of known risk factors, though this relationship remains understudied. We present a summary of the literature on prevalence and unique risk factors, proposing that this may be a high-risk population. We detail next steps for research to understand eating disorder risk in this population to inform future prevention, identification, and treatment efforts needed in this community.

## References

1. O'Dea JA, Abraham S. Onset of disordered eating attitudes and behaviors in early adolescence: interplay of pubertal status, gender, weight, and age. *Adolescence*. 1999;34(136):671-679.
2. Markey CN. Invited commentary: Why body image is important to adolescent development. *J Youth Adolesc*. 2010;39(12):1387-1391.
3. Hooper L, Mason SM, Telke S, Larson N, Neumark-Sztainer D. Experiencing Household Food Insecurity During Adolescence Predicts Disordered Eating and Elevated Body Mass Index 8 Years Later. *J Adolesc Health*. 2022;70(5):788-795.
4. Nagata JM, Garber AK, Tabler JL, Murray SB, Bibbins-Domingo K. Differential Risk Factors for Unhealthy Weight Control Behaviors by Sex and Weight Status Among US Adolescents. *J Adolesc Health*. 2018;63(3):335-341.
5. Pauls A, Dimitropoulos G, Marcoux-Louie G, Singh M, Patten SB. Psychological characteristics and childhood adversity of adolescents with atypical anorexia nervosa versus anorexia nervosa. *Eating Disorders*. 2022;30(2):210-222.
6. Roberts SR, Salk RH, Thoma BC, Romito M, Levine MD, Choukas-Bradley S. Disparities in disordered eating between gender minority and cisgender adolescents. *Int J Eat Disord*. 2021;54(7):1135-1146.
7. Zahnd WE, Del Vecchio N, Askelson N, et al. Definition and categorization of rural and assessment of realized access to care. *Health Serv Res*. 2022;57(3):693-702.
8. Long JC, Delamater PL, Holmes GM. Which Definition of Rurality Should I Use?: The Relative Performance of 8 Federal Rural Definitions in Identifying Rural-Urban Disparities. *Med Care*. 2021;59(Suppl 5):S413-S419.
9. Fontanella CA, Hiance-Steelesmith DL, Phillips GS, et al. Widening rural-urban disparities in youth suicides, United States, 1996-2010. *JAMA Pediatr*. 2015;169(5):466-473.
10. Monnat SM, Rigg KK. Examining Rural/Urban Differences in Prescription Opioid Misuse Among US Adolescents. *J Rural Health*. 2016;32(2):204-218.
11. Reiss F. Socioeconomic inequalities and mental health problems in children and adolescents: a systematic review. *Soc Sci Med*. 2013;90:24-31.
12. Ng AS, Kaye K. *Sex in the (Non)City: Teen Childbearing in Rural America*. 2015.
13. Ziller EC, Lenardson JD, Paluso NC, Talbot JA, Daley A. Rural-Urban Differences in the Decline of Adolescent Cigarette Smoking. *Am J Public Health*. 2019;109(5):771-773.
14. Valentine JA, Delgado LF, Haderxhanaj LT, Hogben M. Improving Sexual Health in US Rural Communities: Reducing the Impact of Stigma. *AIDS Behav*. 2022;26(Suppl 1):90-99.
15. Miller MN, Verhegge R, Miller BE, Pumariega AJ. Assessment of risk of eating disorders among adolescents in Appalachia. *J Am Acad Child Adolesc Psychiatry*. 1999;38(4):437-443.
16. Thompson KM, Wonderlich SA, Crosby RD, Mitchell JE. Sexual victimization and adolescent weight regulation practices: a test across three community based samples. *Child Abuse Negl*. 2001;25(2):291-305.
17. Eichen DM, Conner BT, Daly BP, Fauber RL. Weight perception, substance use, and disordered eating behaviors: comparing normal weight and overweight high-school students. *J Youth Adolesc*. 2012;41(1):1-13.
18. Groft JN, Hagen B, Miller NK, Cooper N, Brown S. Adolescent health: a rural community's approach. *Rural Remote Health*. 2005;5(2):366.
19. Jonat LM, Birmingham CL. Disordered eating attitudes and behaviours in the high-school students of a rural Canadian community. *Eat Weight Disord*. 2004;9(4):285-289.



20. Galmiche M, Dechelotte P, Lambert G, Tavolacci MP. Prevalence of eating disorders over the 2000-2018 period: a systematic literature review. *Am J Clin Nutr*. 2019;109(5):1402-1413.
21. Romano KA, Lipson SK, Beccia AL, Quatromoni PA, Gordon AR, Murgueitio J. Changes in the prevalence and sociodemographic correlates of eating disorder symptoms from 2013 to 2020 among a large national sample of US young adults: A repeated cross-sectional study. *Int J Eat Disord*. 2022;55(6):776-789.
22. Mitchison D, Hay PJ. The epidemiology of eating disorders: genetic, environmental, and societal factors. *Clin Epidemiol*. 2014;6:89-97.
23. Larson N, Loth KA, Eisenberg ME, Hazzard VM, Neumark-Sztainer D. Body dissatisfaction and disordered eating are prevalent problems among US young people from diverse socioeconomic backgrounds: Findings from the EAT 2010–2018 study. *Eating Behaviors*. 2021;42:101535.
24. Mitchison D, Hay P, Slewa-Younan S, Mond J. The changing demographic profile of eating disorder behaviors in the community. *BMC Public Health*. 2014;14:943.
25. US Department of Agriculture Economic Research Service. Rural Poverty & Well-Being. 2022; <https://www.ers.usda.gov/topics/rural-economy-population/rural-poverty-well-being/#demographics>. Accessed July 29, 2022.
26. Coleman-Jensen A, Rabbitt MP, Gregory CA, Singh A. *Household Food Security in the United States in 2020*. US Department of Agriculture Economic Research Service,; September 2021 2021. Number 298.
27. Liese AD, Weis KE, Pluto D, Smith E, Lawson A. Food store types, availability, and cost of foods in a rural environment. *J Am Diet Assoc*. 2007;107(11):1916-1923.
28. Hazzard VM, Loth KA, Hooper L, Becker CB. Food Insecurity and Eating Disorders: a Review of Emerging Evidence. *Curr Psychiatry Rep*. 2020;22(12):74.
29. Dush JL. Adolescent food insecurity: A review of contextual and behavioral factors. *Public Health Nurs*. 2020;37(3):327-338.
30. Shanafelt A, Hearst MO, Wang Q, Nanne MS. Food Insecurity and Rural Adolescent Personal Health, Home, and Academic Environments. *J Sch Health*. 2016;86(6):472-480.
31. Hooper L, Telke S, Larson N, Mason SM, Neumark-Sztainer D. Household food insecurity: associations with disordered eating behaviours and overweight in a population-based sample of adolescents. *Public Health Nutr*. 2020;23(17):3126-3135.
32. West CE, Goldschmidt AB, Mason SM, Neumark-Sztainer D. Differences in risk factors for binge eating by socioeconomic status in a community-based sample of adolescents: Findings from Project EAT. *Int J Eat Disord*. 2019;52(6):659-668.
33. Douthit N, Kiv S, Dwolatzky T, Biswas S. Exposing some important barriers to health care access in the rural USA. *Public Health*. 2015;129(6):611-620.
34. Sangvai D. Eating Disorders in the Primary Care Setting. *Primary Care: Clinics in Office Practice*. 2016;43(2):301-312.
35. Sonnevile KR, Lipson SK. Disparities in eating disorder diagnosis and treatment according to weight status, race/ethnicity, socioeconomic background, and sex among college students. *International Journal of Eating Disorders*. 2018;51(6):518-526.
36. Wade TD, Johnson C, Cadman K, Cook L. Turning eating disorders screening in primary practice into treatment: A clinical practice approach. *International Journal of Eating Disorders*. n/a(n/a).
37. Streatfeild J, Hickson J, Austin SB, et al. Social and economic cost of eating disorders in the United States: Evidence to inform policy action. *Int J Eat Disord*. 2021;54(5):851-868.

38. Gorrell S, Reilly EE, Brosof L, Le Grange D. Use of Telehealth in the Management of Adolescent Eating Disorders: Patient Perspectives and Future Directions Suggested from the COVID-19 Pandemic. *Adolesc Health Med Ther.* 2022;13:45-53.
39. Austin A, Flynn M, Richards K, et al. Duration of untreated eating disorder and relationship to outcomes: A systematic review of the literature. *Eur Eat Disord Rev.* 2021;29(3):329-345.
40. Alman J, Hoiles KJ, Watson HJ, et al. A decade of data from a specialist statewide child and adolescent eating disorder service: does local service access correspond with the severity of medical and eating disorder symptoms at presentation? *Journal of Eating Disorders.* 2014;2(1):32.
41. Brems C, Johnson ME, Warner TD, Roberts LW. Barriers to healthcare as reported by rural and urban interprofessional providers. *J Interprof Care.* 2006;20(2):105-118.
42. Willging CE, Salvador M, Kano M. Pragmatic help seeking: How sexual and gender minority groups access mental health care in a rural state. *Psychiatr Serv.* 2006;57(6):871-874.
43. Williams KJ, Taylor CA, Wolf KN, Lawson RF, Crespo R. Cultural perceptions of healthy weight in rural Appalachian youth. *Rural Remote Health.* 2008;8(2):932.
44. McLaughlin DK, Coleman-Jensen AJ. Nonstandard Employment in the Nonmetropolitan United States. *Rural sociology.* 2008;73(4):631-659.
45. Linardon J, McClure Z, Tylka TL, Fuller-Tyszkiewicz M. Body appreciation and its psychological correlates: A systematic review and meta-analysis. *Body Image.* 2022;42:287-296.
46. Hughes K, Bombak AE, Ankomah S. Experiences of weight-related stigma among low-income rural women of higher weights from the midwestern United States. *Qualitative research in medicine & healthcare.* 2019;3(1).
47. Johnson JA, 3rd, Johnson AM. Urban-rural differences in childhood and adolescent obesity in the United States: a systematic review and meta-analysis. *Child Obes.* 2015;11(3):233-241.
48. Ahorsu DK, Lin CY, Imani V, et al. A prospective study on the link between weight-related self-stigma and binge eating: Role of food addiction and psychological distress. *Int J Eat Disord.* 2020;53(3):442-450.
49. Rehder K, Lusk J, Chen JI. Deaths of Despair: Conceptual and Clinical Implications. *Cogn Behav Pract.* 2021;28(1):40-52.
50. Pompili S, Laghi F. Drunkorexia: Disordered eating behaviors and risky alcohol consumption among adolescents. *J Health Psychol.* 2020;25(13-14):2222-2232.
51. Laghi F, Pompili S, Bianchi D, Lonigro A, Baiocco R. Exploring the association between psychological distress and drunkorexia behaviors in non-clinical adolescents: the moderating role of emotional dysregulation. *Eat Weight Disord.* 2021;26(3):797-806.
52. Long AS, Hanlon AL, Pellegrin KL. Socioeconomic variables explain rural disparities in US mortality rates: Implications for rural health research and policy. *SSM - Population Health.* 2018;6:72-74.
53. James CV, Moonesinghe R, Wilson-Frederick SM, Hall JE, Penman-Aguilar A, Bouye K. Racial/Ethnic Health Disparities Among Rural Adults - United States, 2012-2015. *MMWR Surveill Summ.* 2017;66(23):1-9.
54. Rosenkrantz DE, Black WW, Abreu RL, Aleshire ME, Fallin-Bennett K. Health and health care of rural sexual and gender minorities: A systematic review. *Stigma and Health.* 2017;2(3):229-243.
55. Striegel-Moore RH, Rosselli F, Holtzman N, Dierker L, Becker AE, Swaney G. Behavioral symptoms of eating disorders in Native Americans: results from the ADD Health Survey Wave III. *Int J Eat Disord.* 2011;44(6):561-566.

56. US Department of Agriculture Economic Research Service. Rural American Indian and total US population by age group, 2010. 2014.
57. Burke NL, Schaefer LM, Hazzard VM, Rodgers RF. Where identities converge: The importance of intersectionality in eating disorders research. *Int J Eat Disord*. 2020;53(10):1605-1609.
58. Beccia AL, Baek J, Jesdale WM, et al. Risk of disordered eating at the intersection of gender and racial/ethnic identity among US high school students. *Eat Behav*. 2019;34:101299.
59. Hay PJ, Carriage C. Eating disorder features in indigenous Aboriginal and Torres Strait Islander Australian peoples. *BMC Public Health*. 2012;12:233.