

# **Single-Rooted Extraction Socket Classification: A Systematic Review and Proposal of a New Classification System based on Morphologic and Patient-Related Factors**

Hamoun Sabri<sup>1</sup> DMD, PgC, Shayan Barootchi<sup>1</sup> DMD, MS, Teresa Heck<sup>1</sup> DDS, Hom-Lay Wang<sup>1†</sup> DDS, MSD, PhD

<sup>1</sup> Department of Periodontics and Oral Medicine, School of Dentistry, University of Michigan, Ann Arbor, MI, USA

## **† Corresponding Author**

Hom-Lay Wang,

Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, 1011 N. University Ave., Ann Arbor, MI 48109.  
Email: [homlay@umich.edu](mailto:homlay@umich.edu)

**Running title:** Single-rooted extraction socket classifications based on morphologic and patient-related factors.

**Word count:** 4076

**Number of tables and figures:** 12 (5 tables and 7 figures)

**Appendices:** 2

**Number of references:** 55

**Conflict of interest:** None.

## **Abstract**

Taxonomy and classification of a disease contributes to facilitating the diagnosis and treatment planning process and simplifies communication between clinicians. The aim of this study was to provide a critical appraisal based on a systematic review of the single-rooted extraction socket (ES) classifications and subsequently, introduce a new classification system combining the cornerstones of the previously proposed systems and based on the latest consensus in implant dentistry. Following the systematic search process in PubMed, EMBASE and SCOPUS

**This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/jerd.12967**

databases 13 ES classifications were detected. The most repeated hard and soft tissue factors in the previous classifications were buccal bone dehiscence, interproximal bone, gingival recession and soft tissue phenotype. However, there was minimal attention to patient-related factors such as systemic conditions and smoking. Therefore, a new classification system based on the combination of patient-related factors, clinical and radiographical parameters was proposed. This divides an ES into 3 types. Class I and II sockets are candidates for receiving immediate implant placement and conversely, a class III socket includes a compromised condition that requires multiple-stage reconstruction mostly suitable for standard delayed implant placement with alveolar ridge preservation. Within the limitations of this study, the new classification system not only provides comprehensive inclusion of various crucial parameters in implant placement (such as prediction of future implant position and osteotomy difficulty, etc.) but also, in contrast to the previously introduced systems, is able to classify the ES prior to extraction and also, takes into the account the patient-related factors as the class modifiers following the extraction.

**Keywords:** Extraction Socket, Dental Socket, immediate implants, classification

## **1.Introduction**

Tooth extraction is indicated when a tooth has a hopeless prognosis.<sup>1-3</sup> Following the extraction, alveolar ridge resorption is often unavoidable, which may lead to compromised implant placement.<sup>4</sup> Depending on the hard and soft tissue conditions of the extraction socket (ES), various treatment approaches such as alveolar ridge preservation (ARP) or immediate implant placement (IIP) have been attempted.<sup>5</sup> The buccal plate thickness, buccal bone

morphology, overlying soft tissue, and the pathologic condition of the socket are among the most important factors affecting the treatment decision-making and prognosis.<sup>6,7</sup>

Classification of a disease is crucial as it helps clinicians to identify the pathophysiology, symptomatology, diagnosis and treatment approach. Likewise, it could be beneficial for the patients if an ES decision tree can be developed based upon the above available information. Generally, classification serves as a valuable tool for better communication between clinicians and patients and among researchers.<sup>8</sup> Ideally, a classification system should be user-friendly, precise and comprehensive without any overlaps between the disease entities, and are based on the latest knowledge of pathophysiology and biology.<sup>9</sup>

Several single-rooted ES classification systems are available today, most of these classifications aim to predict the IIP according to the remaining buccal bone and/or overlying soft tissue components. However, the presence of many ES classification systems may create unnecessary confusion among involved parties. Moreover, there is lack of consensus with regards to which ES classification should be used. Each of the proposed systems possesses strengths as well as limitations. For instance, one may include a thorough evaluation of the hard tissue without considering the soft tissue elements whereas another one may only focus on soft tissue.<sup>10</sup> Therefore, the aim of this paper was to provide a critical appraisal of current existing ES classifications within the framework of a systematic review and propose a new single-rooted ES classification that taken into consideration of all important factors based on the latest evidence and consensus in implant dentistry.

## **2. Materials and Methods**

### **2.1 Protocol and registration**

The analysis and interpretation methodology of this study were defined within the framework of a protocol and registered prior to initiation in PROSPERO portal (CRD42022345141).

Moreover, the protocol and the search strategy were created based on the Preferred Reporting Items for Systematic Review and Meta-analysis (PRISMA) statement (Appendix 1).

## **2.2 Problem, Intervention, Comparison and Outcome (PICO) statement**

Problem (P): Lack of consensus regarding the single-rooted ES classification

Intervention (I): Evaluation of available ES classification systems

Comparison (C): Comparison of the included variables and factors into each ES classification

Outcome (O): Proposal of a new single-rooted ES classification system to ease the decision-making process

## **2.3 Focused question**

Based on the stated PICO design, the focused question for this study was proposed as follows:

What are the currently available ES classification systems for single-rooted sockets, the factors concerning ES that are considered and the suggested treatment approaches?

## **2.3 Systematic Search Strategy**

A systematic search approach was performed by two authors (HS, SB) in the electronic databases of: PubMed (MEDLINE), Embase and Scopus, aiming to identify all proposed ES classification systems until January 1, 2022. The main keywords were: “extraction socket” OR

“tooth socket” AND “Classification”. The complete performed searching process and keywords are available as the appendix 2.

The inclusion criteria were reserved to the following articles:

- 1- Presenting a new single-rooted ES classification system compared to the previously introduced ones.

On the contrary, the exclusion criteria were as follows:

- 1- Articles in which the authors implemented one of the previously published systems.
- 2- Studies with focus on a different topic besides ES classification.
- 3- Molar (multi-rooted) ES classifications.
- 4- Theses, abstracts, letters to the editors and editorials.

Moreover, no limitations were applied in terms of the language and date of the publication.

The search results were imported into EndNote (version X9) and de-duplicated based on title, and additionally, the automatically identified duplicates were double-checked manually. Two reviewers (HS and SB) screened the results independently against the eligibility criteria using Review manage software (version 5.3.5). The full-text reading of the selected articles was performed searching for the other classification systems (if had not been included) and those detected from the screening of the reference list of the included articles were also added. In case of any discrepancies between the two reviewers, this resolved by referring to the senior reviewer (HLW). The inter-reviewer reliability in the screening and inclusion process were assessed with Cohen's k test. The included articles were thoroughly reviewed and analyzed.

## **2.4 Types of included studies**

This systematic review contained prospective, retrospective, cohort, case-control, review studies without any language and date limitation.

## **2.5 Data Extraction**

Based on the aim of the study, the following data were extracted independently from the included ES classifications: Study design, date of publication, proposed ES types and description in each classification, parameters based on which the ES classification was performed and suggested treatment approach and considerations for each type of socket.

## **2.6 Quality assessment of the included studies**

The full texts of the included ES classifications were determined with regards to their methodological quality and validity. This was performed based on the COnsensus-based Standards for the selection of health Measurements (COSMIN) checklist<sup>11, 12</sup>. Fundamentally, this checklist was applied to thoroughly investigate the methodological quality of each classification<sup>13, 14</sup>. This checklist evaluates three measurement property of reliability, validity and responsiveness. Based on these three components, 10 Boxes have been defined on the COSMIN platform (patient reported outcomes, internal consistency, reliability, measurement error, content validity, structural validity, hypothesis testing, cross-cultural validity, criterion validity and responsiveness) 8 of which were eligible for this study (patient reported outcomes and cross-cultural validity were excluded). Two reviewers performed the quality assessment (HS and SB) and in case of disagreement the third investigator (HLW) confirmed the decision.

## **3. Results**

### **3.1 Search results and study selection**

The literature search process, based on the PRISMA guidelines is shown in Figure 1. This consisted of two stages. Firstly, following the primary search, 740 articles were identified. Following removal of duplicates, 492 records remained for screening by titles and abstracts. After thorough evaluation of the titles and abstracts 17 articles were selected. As the second stage, after the manual screening of the reference list of the articles, 2 additional ES classifications were also detected and included to the study. Following the full-text assessment of these studies and based on predetermined inclusion criteria, 13 articles were included in the qualitative analysis. The reasons for exclusion of the 6 records are provided in table 1. The inter-reviewer reliability in the screening and inclusion process, as assessed with Cohen's  $k$ , corresponded to 0.91 and 0.88 for assessment of titles and abstracts and full-text evaluation respectively.

### **3.1 Findings from the COSMIN quality assessment of the classifications**

Using the COSMIN checklist, the quality of the ES classification systems included in this study was evaluated (table 2). Out of 13 classifications, none of them met the criteria for adequate internal consistency and responsiveness. 10 of the included classifications lacked “adequate” or “very good” properties in any of the 8 evaluated entity<sup>10, 15-23</sup>. Overall, the classification system by Juodzbaly et al.,<sup>6</sup> had “adequate” reliability and testing “measurement error”. Moreover, although the classifications by Chang and Cheng<sup>4</sup> and Kim et al.,<sup>24</sup> yielded “adequate” hypothesis testing and structural validity respectively, all the other tested parameters were either “inadequate” or “doubtful”. Overall, the results of this quality assessment revealed a strong deficiency in terms of the validity and reliability of the existing classification systems.

### **3.2 Description of the included studies**

A summary of all ES classifications including the factors considered, treatment protocols for each subtype is provided in table 3.

### **3.3 Brief history of ES classifications**

The very first attempt to introduce a classification system for single-rooted ESs was proposed by Salama and Salama<sup>15</sup> in 1993. This was within the framework of the regenerative potential based on the guidelines of infrabony periodontal defects, local topography and specifically, the remaining buccal plate. Later on, in 2003, another classification was introduced by Tinti and Parma-Benfenatti<sup>19</sup>. This was based on the remaining bony housing around the future implants and its regenerative potential. Caplanis et al.,<sup>23</sup> however, were the first group to add the soft tissue parameters to the classification system in addition to the hard tissue components.

Later on, Elia et al.,<sup>16</sup> introduced a simplified classification as well as a non-invasive approach for the management of ESs where the soft tissue is present, but the buccal plate is compromised. A sub-classification for this system was introduced in 2015 by Chu et al.,<sup>22</sup> in which they aimed to provide a more detailed description for the type 2 defects. Similar to Elia's classification, Juodzbalys et al.,<sup>6</sup> aimed to classify the ESs based on the quantitative and qualitative evaluations of both soft and hard tissue adjacent to the socket<sup>6</sup>. The only animal study that was included to this review was conducted by Al-Hezaimi et al., in 2011<sup>10</sup>. They concluded that a compromised interdental blood supply, and consequently, the interdental remaining bone contributes to the bone resorption in ES and proposed their classification based on the presence of adjacent teeth and the situation of interdental bone. Another ES classification was introduced by El Chaar et al.,<sup>17</sup> group, mainly based on the bone topography of the socket.



More recently proposed classifications consist of Iyer et al.,<sup>20</sup> in 2019, which was solely based on the hard tissue components, and Chang and Cheng<sup>4</sup>, Kim et al.,<sup>24</sup> and Cardaropoli et al.,<sup>18</sup> classifications all published in 2021. The Chang and Cheng's<sup>4</sup> system is a modification for Elian's classification, which is based on the amount of tissue destruction in all four walls of ESs. The classification by Kim et al.,<sup>24</sup> refers to the pathologically affected, single-rooted ESs. Fundamentally, this was done based on the hard and soft tissue condition of ESs following tooth loss due to periodontal and/or endodontic infection. Similarly, identical variables were also taken into consideration by Cardaropoli's classification<sup>18</sup>.

### **3.3 Included Factors in Existing Classification Systems**

After a thorough evaluation of the detected ES classification systems, the proposed parameters that are taken into account to classify sockets for all the classifications were evaluated. Generally, the parameters which have been used to evaluate the socket prior to the classification can be divided into three groups: hard tissue parameters, soft tissue parameters and patient related factors. Figure 2 shows the pie chart of the included factors to all the selected ES classifications.

#### **3.3.1 Hard tissue parameters**

##### **3.3.1.1 Remaining buccal bone dimensions**

The buccal bone dimensions, including thickness, buccal bone loss such as dehiscence, are taken into account in almost all ES classifications. Only two studies introduced the hard tissue dehiscence as a main factor to consider<sup>15, 21</sup>. However, this consisted of solely qualitative evaluation (presence or absence).

The extent of buccal bone loss was considered as a parameter to classify the socket in 9 studies<sup>4, 6, 16-18, 21-24</sup>. According to the reviewed studies, the amount of acceptable buccal bone loss allowing for IIP is up to 2 mm or 20-25% of resorption<sup>6, 17, 18, 23</sup>. Moreover, one study also added the amount of bone loss on the palatal aspect in addition to buccal<sup>24</sup>. The thickness of the buccal plate is also included in 4 systems<sup>6, 20, 21, 23</sup>. All classification systems considered at least 2 mm of buccal bone thickness as an acceptable parameter for IIP.

### **3.3.1.2 Defect walls**

This parameter has taken into account in four systems<sup>15, 23, 20, 24</sup>. Overall, it can be stated that based on the included classifications, the regenerative potential as well as the vascularity of the socket decreases in 3- or less- wall defects compared to a 4-wall intact bony structure and the prognosis of an IIP in 4-wall defects, provided that the other parameters are also in optimum levels, can be considered as “good”<sup>17, 20, 21</sup>.

### **3.3.1.3 Apical topography**

Three of the selected ES classification systems considered the apical topography as a main factor to classify defects<sup>6, 15, 19</sup>. Moreover, the rationale to bear in mind is the amount of remaining bone in the apical region to be engaged with the implant. For instance, the minimum amount of remaining bone is around 3 to 4 mm, to be in contact with the implant<sup>6, 17</sup>.

### **3.3.1.4 Future peri-implant hard tissue**

The foreseeable amount of bone housing around the future implants were considered by Tinti and Parma-Benfenati<sup>19</sup> for the single-rooted socket classification. This refers to the importance of an intact envelope of bone for clot stability.

Finally, El Chaar et al.,<sup>17</sup> and Al-Yafi et al.,<sup>21</sup> added the interproximal bone parameter to the previous criteria, as the level of interproximal bone dictates the presence or absence of the soft tissue and interproximal papillae.

### **3.3.2 Soft tissue**

#### **3.3.2.1 Soft tissue phenotype (previously named biotype)**

Four ES classifications pointed out to the important role of the tissue phenotype<sup>6, 17, 21, 23</sup>. This is because tissue phenotype plays an important role in the implant esthetics. In general, thick tissue phenotype often achieve better esthetic outcomes as well as to be more incline for IIP<sup>6</sup>. However, for a thin tissue phenotype, the more conservative approaches are often suggested to minimize the potential esthetic challenges.

#### **3.3.2.2 Buccal soft tissue level/loss**

The destruction and amount of the remaining soft tissue was included in 6 classifications<sup>4, 6, 16, 18, 21, 24</sup>. This variable was assessed qualitatively in all studies except, in the classification by Juodzbalys et al.,<sup>6</sup> it was stated that a soft tissue loss of more than 2 mm contributes to a poor prognosis for ESs.

#### **3.3.2.3 Soft tissue quality**

This consisted of soft tissue predictability which was proposed by Caplanis et al.,<sup>23</sup> and the soft tissue quality by Juoudzbalys et al.<sup>6</sup>. The former comprises evaluation of various factors affecting the outcomes of future soft tissue and the latter refers to qualitative features of the soft tissue such as consistency, color and contour.

#### **3.3.2.4 Blood supply**

One of the included systems took the blood supply to the ES into account in the classification<sup>10</sup>. This concept was investigated by Al-Hezaimi et al.,<sup>10</sup> and they suggested that the blood supply to the ES is derived from interdental bone (the internal walls of the socket) and this is an important factor in terms of the soft tissue contours and prevention of bone resorption. Thus, the presence of adjacent (proximal) teeth serves an important consideration in maintaining the blood supply to the area.

### **3.3.3 Patient- and tooth- related factors**

#### **3.3.3.1 Etiology, pathology and systemic factors**

The presence of socket pathology prior to extraction and the etiologic factors were only considered in 3 classifications<sup>4, 23, 24</sup>. This mainly consisted of pre-extraction evaluation of the systemic health and risk factors and the cause of extraction (e.g. infection, fracture, etc.) which can affect the prognosis of the treatment. Generally, none of the classification systems clearly mentioned the exact factors to consider. Finally, in one classification system the authors considered the esthetic concern of the patient as one of the main factors<sup>21</sup>.

### **3.4 Evaluation tools**

All classification systems performed the socket evaluation using clinical and radiographical findings. Moreover, some classification systems specifically mentioned the CBCT images should be taken and evaluated<sup>18</sup> whereas in others the necessity of CBCT image acquisition was not stated. In one classification, the authors utilized a prefabricated prosthetic guide to evaluate the hard and soft tissue around the socket.<sup>23</sup>

## **4. Proposal of a new classification system**

Based on the proposed quality assessment and critical appraisal, and also, taking the latest consensus reports<sup>7, 25-28</sup> into consideration, a new single-rooted ES classification was proposed. The new classification system is presented in table 4 and figure 3. This consists of three main steps to apply as follows:

The first two steps determine the sockets' class based on the morphologic and anatomical features. The first step is determining clinical factors with regards to ES (figure 4):

- 1- **Determining the etiology of extraction:** extractions with the etiology of excessive caries, endodontic failure, root fractures yield superior prognosis compared to tooth loss due to severe periodontitis or severe endo-perio lesions (figure 4a).<sup>25</sup>
- 2- **The amount of gingival recession at the extraction site.** A gingival recession of more than 3 mm considered to be associated with risk of soft tissue deficiency following IIP<sup>6, 29</sup> (figure 4b).
- 3- **Determination of soft tissue phenotype:** this parameter can be either thin or thick and plays a key role in determination of future peri-implant soft tissue<sup>25, 27</sup>.

Following clinical examination, the radiographic examination can be performed based on available radiographs and CBCT as the second step (figure 5).

- 1- **Buccal Bone:** the thickness and amount of dehiscence should be considered. Up to 50% of buccal plate loss could be manageable if IIP is considered (figures 5a and b).<sup>25</sup>

- 2- **Interproximal bone loss:** is especially important in the esthetic zone as it contributes to future papilla fill and prevents future interproximal soft tissue defects.<sup>30</sup>
- 3- **Apical Lesions:** current evidence indicates favorable success rates for IIP in sockets with periodontal lesion and/or periapical pathology<sup>31, 32</sup>. However, before placing the implant careful and thorough decontamination and removal of the infected tissue is required<sup>25, 27</sup> (figure 5c).
- 4- **Root position:** this parameter predicts the future three-dimensional position of the implant. In IIP more palatal/lingual positioning of the implant is desired to avoid excessive contact with the residual buccal plate<sup>25, 27</sup> to fulfill “prosthodontically driven” concept<sup>33, 34</sup>(figures 3 and 5).

Following the second step the initial classification of the socket can be achieved. This allows the dentist to preliminarily diagnose and perform treatment planning. Nevertheless, if the tooth is extracted, the third step (figure 6) can be initiated, which considers possible class modifiers that are only examinable following removal of the tooth and by inspection of the residual socket, based on possible events during extraction surgery and future implant osteotomy factors, patient-related factors, and also any adverse events during the extraction following criteria:

- 1- **Presence of poorly controlled systemic disease:** factors such as diabetes mellitus, smoking and advanced autoimmune conditions may affect the socket healing process as well as pregnancy or adolescence. Therefore, in such scenarios caution required.<sup>25</sup>

- 2- **Smoking:** smoking more than 10 cigarettes per day would be considered as a risk indicator in the IIP.<sup>32</sup>
- 3- **Medication history:** if the patient takes any medications which can affect favorable healing of the socket, despite scarcity of the literature supporting this point of view, in certain cases (such as bisphosphonate, chemotherapy agent etc.), caution is necessary.<sup>25, 35</sup>
- 4- **Presence of active periodontitis in the same sextant.** Although data regarding hazardous impact of previous periodontitis on IIP is controversial<sup>27, 36, 37</sup> presence of active periodontitis within the same sextant could serve an additional risk for IIP.<sup>38, 39</sup>
- 5- **Evaluation of oral hygiene:** poor oral hygiene may increase the failure and complications in IIP.<sup>5</sup>
- 6- **Any major trauma during the procedure;** which causes failure in preservation of hard and soft tissue quality.<sup>40</sup>
- 7- **Occurrence of iatrogenic complications:** such as sinus membrane perforation, buccal plate fracture etc.
- 8- **Re-evaluation of buccal bone thickness and bone quality.** This step is advised to be followed in order to re-evaluate and confirm the pre-extraction diagnosis and apply any changes if needed.
- 9- **Osteotomy related factors:** including the presence of possible limitations in osteotomy sequence of implant (e.g. risk of damage to the adjacent root or nerves<sup>41</sup>, location of nasopalatine canal etc.).<sup>28, 42</sup>

Lastly, based on the proposed classification system, a decision-making flowchart is presented in figure 7 demonstrating the suggested approach in the management of ESs.

## **5. Discussion**

This article systematically reviewed all single-rooted ES classifications and proposed a new classification based on a critical appraisal and quality assessment of the previously available systems and with the aim of providing a system with combination of all crucial parameters to consider while performing dental extractions in the esthetic zone.

It is important to be able to discuss the treatment planning prior to the extraction of the tooth. This emphasizes the need for a classification system allowing to classify before the extraction. However, most of the existing ES classification systems lack this feature. Thereby, the introduced new classification system in this paper, allows clinicians to perform the initial classification prior to extraction (as demonstrated in figures 4 and 5) and next, if proceeded to extraction, one can modify the class by the mentioned class modifiers accordingly (figure 6). Needless to mention that a critical step (step 3) consists of clinically confirming the pre-extraction determined class by inspection.

Unfortunately, the importance of possible multiple ESs is underappreciated.<sup>43</sup> Although Al-Hezaimi et al,<sup>10</sup> intended to consider this, the nature of the animal study leaves room for the further research. As a possible approach, condition of the interproximal walls between the adjacent extraction socket was entered into the new ES classification.

All the previous systems, to some extent, covered the anatomical factors affecting the treatment approach. However, it should be noted that there was little attention to the patient



related factors such as systemic diseases or smoking or even diabetes, which can easily transform a favorable ES to a questionable despite its undamaged structure. Therefore, one of the main goals of the newly proposed system was to include systemic conditions, as possible class modifiers, following the examination of the anatomical and topographic factors. In a study by Urban et al.,<sup>44</sup> it is indicated that smoking can be a risk factor for molar area IIP. Similarly, it is reported in the literature that despite its acceptable success rate, infectious ES needs adjunctive therapy and additional considerations if IIP is planned.<sup>39, 45</sup> Also, it is elaborated throughout the studies that health-related systemic conditions such as diabetes and hypertension could possibly affect the extraction socket healing process and therefore, alter the expected outcomes.<sup>46, 47</sup> Therefore, all the aforementioned parameters included as the class modifiers the novel ES classification system.

An important aim of using classifications is to facilitate the communication among all involved parties. This needs the implementation of well-described and precise variables comprising the classification system.<sup>8, 48</sup> However, it can be noted that most of the proposed ES classifications assess the defects qualitatively or at best a combination of qualitative and quantitative. This causes discrepancies in terms of diagnosis as it leads to intra- and inter-observer bias as well as several grey zones defining a defect and considering it adequate or compromised<sup>49</sup>. Further investigation on the repeatability of the classifications is suggested. In our classification system, we attempted to provide quantitative and/or dichotomous values for the parameters which increases the reproducibility and quality of assessment.

All included studies implemented radiographic images, either peri-apical, panoramic and CBCT or combination of these, for the examination of the hard tissue situation and periapical diagnosis. Nevertheless, in terms of the soft tissue evaluation there is a lack of standardized

techniques to diagnose. Overall, CBCT images seem to be one of the cornerstones of the ES classification, thus, it is strongly suggested to perform classification by the means of this image modality. And for soft tissue parameters, currently the conventional clinical measurements are suggested.<sup>50</sup>

Lastly, it is suggested that a thorough soft and hard tissue evaluation in combination with patient related factors should be followed. Subsequently, the decision for either IIP or DIP with the suitable modification can be made and applied. Figure 7 illustrates the factors and the flowchart that is suggested to be take into account when performing tooth extractions using the new classification system. That being mentioned, within the limitation of systematic reviews, in this paper we aimed to merely provide an overview concerning the available classifications and possible gaps within them and describe the characteristics of each one as well as the parameters that are included into each system. Moreover, the readers should bear in mind that the newly introduced classification system is solely based on the most recent consensus reports in implant dentistry<sup>25, 28</sup> and despite its benefits in terms of updating the previous systems, it requires further studies to evaluate its validity, responsiveness and reliability.<sup>11, 13, 14</sup>

## **6. Conclusions**

The present study provided a systematic review and a critical appraisal on the previous single-rooted extraction socket classifications and proposed a new classification system. This classification revises and updates the definitions and criteria from the former systems. An important feature is including the factors affecting future implant treatment, especially in the esthetic zone. Likewise, the most recent consensus-based criteria for immediate implantation such as soft tissue

Author Manuscript

esthetic considerations as well as the sagittal root position was taken into consideration. Lastly, this classification considered the patient-related and extraction-related factors for the first time, as the class modifiers in case they have an impact on the prognosis and treatment planning following the extraction.

## 7. Conflict of Interest

The authors declare no conflict of interest.

## 8. Funding

None.

## References

1. Avila G, Galindo-Moreno P, Soehren S, Misch CE, Morelli T, Wang HL. A novel decision-making process for tooth retention or extraction. *Journal of periodontology* 2009;80:476-491.
2. Tarazona B, Tarazona-Alvarez P, Peñarrocha-Oltra D, Penarrocha-Diago M. Relationship between indication for tooth extraction and outcome of immediate implants: a retrospective study with 5 years of follow-up. *Journal of clinical and experimental dentistry* 2014;6:e384.
3. Council NR. Toward precision medicine: building a knowledge network for biomedical research and a new taxonomy of disease. 2011.
4. Chang L-C, Cheng Y-M. The Effect of Different Socket Types on Implant Therapy While Using Flapless Ridge Preservation. *Applied Sciences* 2021;11:970.
5. Noelken R, Kunkel M, Wagner W. Immediate implant placement and provisionalization after long-axis root fracture and complete loss of the facial bony lamella. *Int J Periodontics Restorative Dent* 2011;31:175-183.
6. Juodzbaly G, Sakavicius D, Wang HL. Classification of extraction sockets based upon soft and hard tissue components. *J Periodontol* 2008;79:413-424.
7. Barootchi S, Wang HL, Ravida A, et al. Ridge preservation techniques to avoid invasive bone reconstruction: A systematic review and meta-analysis: Naples Consensus Report Working Group C. *Int J Oral Implantol (Berl)* 2019;12:399-416.
8. Klasser GD, Manfredini D, Goulet JP, De Laat A. Oro-facial pain and temporomandibular disorders classification systems: A critical appraisal and future directions. *J Oral Rehabil* 2018;45:258-268.
9. Jakob R. Disease Classification. In: Quah SR, ed. *International Encyclopedia of Public Health (Second Edition)*. Oxford: Academic Press, 2017:332-337.


10. Al-Hezaimi K, Levi P, Rudy R, Al-Jandan B, Al-Rasheed A. An extraction socket classification developed using analysis of bone type and blood supply to the buccal bone in monkeys. *Int J Periodontics Restorative Dent* 2011;31:421-427.
11. Mokkink LB, Terwee CB, Patrick DL, et al. The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res* 2010;19:539-549.
12. Mokkink LB, de Vet HCW, Prinsen CAC, et al. COSMIN Risk of Bias checklist for systematic reviews of Patient-Reported Outcome Measures. *Qual Life Res* 2018;27:1171-1179.
13. Prinsen CAC, Mokkink LB, Bouter LM, et al. COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018;27:1147-1157.
14. Terwee CB, Prinsen CAC, Chiarotto A, et al. COSMIN methodology for evaluating the content validity of patient-reported outcome measures: a Delphi study. *Qual Life Res* 2018;27:1159-1170.
15. Salama H, Salama M. The role of orthodontic extrusive remodeling in the enhancement of soft and hard tissue profiles prior to implant placement: a systematic approach to the management of extraction site defects. *Int J Periodontics Restorative Dent* 1993;13:312-333.
16. Elian N, Cho S, Froum S, Smith RB, Tarnow DP. A simplified socket classification and repair technique. *Practical Procedures and Aesthetic Dentistry* 2007;19:99.
17. El Chaar E, Oshman S. Single-Rooted Extraction Sockets: Classification and Treatment Protocol. *Compendium of continuing education in dentistry (Jamesburg, NJ: 1995)* 2016;37:537-541; quiz542.
18. Cardaropoli D, Nevins M, Casentini P. A Clinical Classification System for the Treatment of Postextraction Sites. *Int J Periodontics Restorative Dent* 2021;41:227-232.
19. Tinti C, Parma-Benfenati S. Clinical classification of bone defects concerning the placement of dental implants. *Int J Periodontics Restorative Dent* 2003;23:147-155.
20. Iyer S, Haribabu PK, Xing Y. Part II. Minimizing alveolar bone loss during and after extractions. Protocol and techniques for alveolar bone preservation. *Alpha Omegan* 2014;107:14-25.
21. Al Yafi F, Alchawaf B, Nelson K. What is the Optimum for Alveolar Ridge Preservation? *Dent Clin North Am* 2019;63:399-418.
22. Chu S, Sarnachiaro G, Hochman M, Tarnow D. Subclassification and clinical management of extraction sockets with labial dentoalveolar dehiscence defects. *Compendium* 2015;36:516-525.
23. Caplanis N, Lozada J, Kan J. Extraction Defect: Assessment, Classification and Management. *Journal of the California Dental Association* 2005;33:853-863.
24. Kim JJ, Ben Amara H, Chung I, Koo KT. Compromised extraction sockets: a new classification and prevalence involving both soft and hard tissue loss. *J Periodontal Implant Sci* 2021;51:100-113.
25. Tonetti MS, Jung RE, Avila-Ortiz G, et al. Management of the extraction socket and timing of implant placement: Consensus report and clinical recommendations of group 3 of the XV European Workshop in Periodontology. *Journal of Clinical Periodontology* 2019;46:183-194.

26. Kan JYK, Rungcharassaeng K, Deflorian M, Weinstein T, Wang HL, Testori T. Immediate implant placement and provisionalization of maxillary anterior single implants. *Periodontol 2000* 2018;77:197-212.
27. Blanco J, Carral C, Argibay O, Liñares A. Implant placement in fresh extraction sockets. *Periodontology 2000* 2019;79:151-167.
28. Buser D, Chappuis V, Belser UC, Chen S. Implant placement post extraction in esthetic single tooth sites: when immediate, when early, when late? *Periodontology 2000* 2017;73:84-102.
29. Groenendijk E, Bronkhorst EM, Meijer GJ. Does the pre-operative buccal soft tissue level at teeth or gingival phenotype dictate the aesthetic outcome after flapless immediate implant placement and provisionalization? Analysis of a prospective clinical case series. *Int J Implant Dent* 2021;7:84-84.
30. Teughels W, Merheb J, Quirynen M. Critical horizontal dimensions of interproximal and buccal bone around implants for optimal aesthetic outcomes: a systematic review. *Clinical Oral Implants Research* 2009;20:134-145.
31. Crespi R, Capparé P, Crespi G, Lo Giudice G, Gastaldi G, Gherlone E. Immediate implant placement in sockets with asymptomatic apical periodontitis. *Clinical implant dentistry and related research* 2017;19:20-27.
32. Lee C-T, Chuang S-K, Stoupel J. Survival analysis and other clinical outcomes of immediate implant placement in sites with periapical lesions: systematic review. *International Journal of Oral & Maxillofacial Implants* 2015;30.
33. Demircan S. Prosthetically driven immediate implant placement at lower molar area; an anatomical study. *European Oral Research* 2020;54:25-30.
34. Katsoulis J, Pazera P, Mericske-Stern R. Prosthetically driven, computer-guided implant planning for the edentulous maxilla: a model study. *Clinical Implant Dentistry and Related Research* 2009;11:238-245.
35. Mohammadi A, Dehkordi NR, Mahmoudi S, et al. Effects of Drugs and Chemotherapeutic Agents on Dental Implant Osseointegration: Narrative Review. *Current Reviews in Clinical and Experimental Pharmacology* 2022.
36. Renvert S, Persson GR. Periodontitis as a potential risk factor for peri-implantitis. *J Clin Periodontol* 2009;36 Suppl 10:9-14.
37. Wagenberg B, Froum SJ. A retrospective study of 1925 consecutively placed immediate implants from 1988 to 2004. *Int J Oral Maxillofac Implants* 2006;21:71-80.
38. Lee J, Park D, Koo K-T, Seol Y-J, Lee Y-M. Comparison of immediate implant placement in infected and non-infected extraction sockets: a systematic review and meta-analysis. *Acta Odontologica Scandinavica* 2018;76:338-345.
39. Saijeva A, Juodzbaly G. Immediate Implant Placement in Non-Infected Sockets versus Infected Sockets: a Systematic Review and Meta-Analysis. *J Oral Maxillofac Res* 2020;11:e1-e1.
40. Lee W. Immediate implant placement in fresh extraction sockets. *J Korean Assoc Oral Maxillofac Surg* 2021;47:57-61.
41. Greenstein G, Tarnow D. The mental foramen and nerve: clinical and anatomical factors related to dental implant placement: a literature review. *Journal of periodontology* 2006;77:1933-1943.

42. Lin M-H, Mau L-P, Cochran DL, Shieh Y-S, Huang P-H, Huang R-Y. Risk assessment of inferior alveolar nerve injury for immediate implant placement in the posterior mandible: a virtual implant placement study. *Journal of dentistry* 2014;42:263-270.
43. Testori T, Weinstein T, Scutellà F, Wang H-L, Zucchelli G. Implant placement in the esthetic area: criteria for positioning single and multiple implants. *Periodontology 2000* 2018;77:176-196.
44. Urban T, Kostopoulos L, Wenzel A. Immediate implant placement in molar regions: risk factors for early failure. *Clin Oral Implants Res* 2012;23:220-227.
45. Chrcanovic BR, Martins MD, Wennerberg A. Immediate placement of implants into infected sites: a systematic review. *Clin Implant Dent Relat Res* 2015;17 Suppl 1:e1-e16.
46. Fernandes KS, Glick M, de Souza MS, Kokron CM, Gallottini M. Association between immunologic parameters, glycemic control, and postextraction complications in patients with type 2 diabetes. *The Journal of the American Dental Association* 2015;146:592-599.
47. Ra'ed M. The effect of cigarette smoking on the severity of pain, swelling and trismus after the surgical extraction of impacted mandibular third molar. *Journal of Clinical and Experimental Dentistry* 2013;5:e117.
48. Webster AJ, Gaitskell K, Turnbull I, Cairns BJ, Clarke R. Characterisation, identification, clustering, and classification of disease. *Sci Rep* 2021;11:5405.
49. Fillingim RB, Bruehl S, Dworkin RH, et al. The ACTION-American Pain Society Pain Taxonomy (AAPT): an evidence-based and multidimensional approach to classifying chronic pain conditions. *The Journal of Pain* 2014;15:241-249.
50. Juodzbalys G, Wang HL. Soft and hard tissue assessment of immediate implant placement: a case series. *Clinical Oral Implants Research* 2007;18:237-243.
51. Kumar PR, Kher U. Shield the socket: Procedure, case report and classification. *J Indian Soc Periodontol* 2018;22:266-272.
52. Al-Shabeeb MS, Al-Askar M, Al-Rasheed A, et al. Alveolar Bone Remodeling Around Immediate Implants Placed in Accordance With the Extraction Socket Classification: A Three-Dimensional Microcomputed Tomography Analysis. *Journal of Periodontology* 2012;83:981-987.
53. Juodzbalys G, Wang H-L. Socket Morphology--Based Treatment for Implant Esthetics: A Pilot Study. *International Journal of Oral & Maxillofacial Implants* 2010;25.
54. Smith RB, Tarnow DP. Classification of molar extraction sites for immediate dental implant placement. *International Journal of Oral & Maxillofacial Implants* 2013;28.
55. Bleyan S, Gaspar J, Huwais S, et al. Molar Septum Expansion with Osseodensification for Immediate Implant Placement, Retrospective Multicenter Study with Up-to-5-Year Follow-Up, Introducing a New Molar Socket Classification. *J Funct Biomater* 2021;12.


## Figure and Table Legends

**Figure 1.** The PRISMA chart of the identification, screening and selection process of the present systematic review.



**Figure 2.** The pie chart of all parameters that are taken into account by the previous classification systems included to this study.


■ Apical Topography ■ Soft Tissue Destruction ■ Phenotype ■ Blood Supply ■ Defect Walls ■ Soft tissue Quality  
■ Patient-related Factors ■ Hard Tissue Dehiscence ■ Peri-implant Bone ■ Buccal Plate Loss ■ Pathology  
■ Buccal Plate Thickness ■ Local Anatomy ■ Palatal Bone Loss




**Figure 3.** the new single-rooted extraction socket classification system. (note that the presence of even one criterion from each class will put a socket into that group. For instance, more than 50% of buccal bone deficiency, even without presence of




gingival recession of >3mm would still be considered as a class III socket.)




**Figure 4.** Clinical evaluation of the ES prior to the extraction. (a) a chronic fistula and presence of interproximal attachment loss. Based on clinical findings this would be put into class II. (b) more than 3 mm of gingival recession and interproximal attachment loss in a class III socket tooth.



**Figure 5.** Examples of radiographic evaluation of the future ES prior to extraction. (a) a future class II extraction socket with thin (<1mm) buccal bone thickness and moderate buccal plate dehiscence (<50%). (b) a class II socket with buccal bone thickness between 1 and 2 mm and sagittal root position adjacent to the vestibular plate. (c) a class III socket with more than 50% of buccal bone dehiscence and an apical lesion.




**Figure 6.** the third step of the extraction socket classification: evaluation of the post-extraction socket to confirm the classification. This consists of confirming the amount of remaining buccal bone as well as interproximal plates and the apical topography.



**Figure 7.** The Suggested Flow-chart to follow for the management of extraction sockets.

**Abbreviations.** IIP: Immediate Implant Placement, DIP: Delayed Implant Placement



Author Manuscript

**Table 1.** Excluded studies and the reasons for exclusion.

<b>Authors</b>	<b>Type of Study</b>	<b>Reason for Exclusion</b>
Kumar and Kher <sup>51</sup>	Case report and review	Socket shield classification
Juodzbaly et al., <sup>39</sup>	Systematic Review	Review of socket augmentation and ARP
Al-Shabeeb et al., <sup>52</sup>	Pilot Animal Study	An existing ES classification was implied
Juodzbaly and Wang <sup>53</sup>	Pilot clinical study	An existing ES classification was implied
Smith and Tarnow <sup>54</sup>	Technical note	Molar extraction socket classification
Bleyan and Gaspar <sup>55</sup>	Retrospective	Molar extraction socket classification



	Apical residual bone ST phenotype								
Al-Yafi et al., 2019 <sup>21</sup>	Dehiscence and fenestration Buccal plate thickness Interproximal bone ST phenotype Recession Esthetic concern	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate
Chang and Cheng, 2021 <sup>4</sup>	HT Destruction ST Destruction Infection Systemic disease	Inadequate	Inadequate	Doubtful	Doubtful	Inadequate	Adequate	Inadequate	Inadequate
Kim et al., 2021 <sup>24</sup>	Buccal bone loss Palatal bone loss Remaining HT walls ST Destruction Etiology of extraction	Inadequate	Inadequate	Inadequate	Doubtful	Adequate	Inadequate	Doubtful	Doubtful
Cardaropoli et al., 2021 <sup>18</sup>	ST level Buccal Plate resorption Local bone anatomy	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate	Inadequate

\*Patient reported outcome results and cross-cultural validity were not applicable to this study.

**Abbreviations.** N/A: Not available, HT: Hard tissue, ST: Soft tissue

**Table 3.** The characteristics of the included single-rooted extraction socket classification systems.

Classification	Study Design	Soft tissue Parameters	Hard Tissue Parameters	Patient Related Factors	Subtypes	Treatment Approach	Considerations
<b>Salama and Salama, 1993<sup>15</sup></b>	Case series	None	Residual bone around apex Defect Walls Dehiscence	None	T1	IIP/ Adjunctive treatments if compromised or severe case	-
					T2	DIP + ARP or Forced eruption	-
					T3	DIP + regeneration with DFDB allografts + tetracycline covered by a membrane	-
<b>Tinti and Parma-Benfenatti, 2003<sup>19</sup></b>	Case series	None	Bone Housing around the placed implant	None	C1	N/A	-
					C2	N/A	-
<b>Caplanis et al., 2005<sup>23</sup></b>	Review	Tissue phenotype Soft tissue predictability	Number of affected walls HT Loss	Past medical and dental history Detection of systemic risk factors	EDS-1	IIP	-
					EDS-2	ARP + IIP or ARP+DIP	-
					EDS-3	ARP+DIP	-
					EDS-4	ARP + Site development+ DIP (three stage)	-
<b>Elian et al., 2007<sup>16</sup></b>	Case series	Remaining buccal ST	Remaining Buccal HT	None	TI	IIP/ DIP	The easiest and most predictable



					TII	Staged approach with HT and with or without ST augmentation	Risk of misdiagnosing as type I/ Difficult to diagnose
					TIII	Staged Approach with HT with or without ST augmentation	Require experience, dexterity and time
<b>Juodzbaly et al., 2008<sup>6</sup></b>	Case series	Tissue Phenotype ST Quality (Color, consistency, contour)	Alveolar Process height Residual bone around apex Labial Plate thickness and position Pathology	None	TI	IIP	Optimal esthetic outcomes expected
					TII	IIP or DIP with ST or HT augmentation	-
					TIII	DIP after ST or HT augmentation or Orthodontic forced eruption	-
<b>Al-Hezaimi et al., 2011<sup>10</sup></b>	Animal Study	Blood supply to the area	Presence of adjacent teeth	None	CI	N/A	-
					CII	N/A	-
					CIII	N/A	-
<b>Iyer et al., 2014<sup>20</sup></b>	Review and case series	None	Buccal plate thickness Number of affected walls	None	CI	IIP without ARP/ ARP + DIP	-
					CII	IIP + graft/ DIP + graft or ARP	
					CIII	Particulate graft + IIP if primary stability achievable	
					CIV	Socket walls regeneration +DIP	
					CV	Augmentation with autogenic or allogenic grafts + DIP	
					CVI	Major interventions (e.g. distraction osteogenesis and etc.) + DIP	
<b>Chu et al., 2015<sup>22</sup></b>	Case series	Intact Labial ST in all types	Residual Buccal Bone	None	T2A	IIP + GBR (caution on T2C sockets)	-
					T2B		-

					T2C		-
<b>El Chaar et al., 2016<sup>17</sup></b>	Case series	ST phenotype	Apical topography Buccal plate loss% Interproximal bone	None	GI	IIP with without Provisionalization and bone graft	-
					GII	Thin phenotype: DIP+ARP Thick Phenotype: IIP without temporization	-
					GIII	ARP or Forced eruption	In case of compromised apical topography→ DIP+GBR
<b>Al-Yafi et al., 2019<sup>21</sup></b>	Review	Tissue phenotype Recession	Dehiscence and fenestration Buccal plate thickness Interproximal bone height	Esthetic concern	CIA	IIP/ 2 Stage	3 Stage approach: ARP + Site Development + implant Placement
					CIB	IIP/ 2 Stage	
					CIIA	IIP with grafting/ 2 Stage	
					CIIB	2 Stage/ 3 Stage	
					CIIAA	IIP with grafting/ 2 stage	
					CIIBB	2 Stage/ 3 Stage	
					CIVA	IIP with grafting/ 2 stage	
					CIVB	3 Stage	
<b>Chang and Cheng, 2021<sup>4</sup></b>	retrospective	ST destruction	HT destruction	Systemic disease Infection	CI	Varies based on the patient related factors: Natural healing to ridge augmentation	IIP can be performed if primary stability is achievable
					CII		
					CIII		
					CIV		
<b>Kim et al., 2021<sup>24</sup></b>	Cohort	Buccal ST level	Number of remaining walls Buccal and palatal bone loss	Etiology of the tooth loss	TI	IIP	if no infection or thin buccal plate is present
					TII	IIP + GBR or DIP	Risk of compromised esthetics in IIP DIP preferred in extensive destruction
					TIII	ARP + CTG or FGG	-

					TIV	Ridge augmentation	Augmentation beyond the present envelope of bone
					TV	Extensive ridge augmentation (complementary bone augmentation after healing)	-
<b>Cardaropoli et al., 2021<sup>18</sup></b>	Case series	ST level	Buccal Plate resorption Local bone anatomy	None	CI	IIP/ ARP + DIP	-
					CII	Ridge augmentation + DIP/ DIP + simultaneous bone augmentation	
					CIII	Ridge augmentation W/WO ST augmentation + DIP/ DIP with simultaneous bone augmentation	
					CIV	Ridge augmentation W/WO ST augmentation + DIP/ Delayed bone augmentation + DIP/	

**Abbreviations.** T: Type, C: Class, IIP: Immediate Implant Placement, ARP: Alveolar Ridge Preservation, DIP: Delayed Implant Placement, DFDB: Demineralized Freeze-Derived Bone, HT: Hard Tissue, ES: Extraction Socket, ST: Soft Tissue, EDS: Extraction Defect Sounding, G: Grade, GBR: Guided Bone Regeneration, CTG: Connective Tissue Graft, FGG: Free Gingival Graft

**Table 4.** The new extraction socket classification. **Class I:** refers to a socket with ideal condition and able to receive IIP. The etiology for extraction is not periodontitis-related and mostly includes endodontic-related origins and excessive caries or fracture. The amount of gingival recession does not exceed 3 mm. the soft tissue phenotype is thick. In the radiographic images, at least 2mm of buccal bone thickness without dehiscence, interproximal bone loss and apical pathology can be seen. The root position is ideal for IIP planning. **Class II:** Whenever the ES includes at least one of the proposed criteria in this class, it will be considered as a Class II socket. This consists of a mildly affected socket. A thin phenotype can be detected. Radiographic parameters include less than 2mm of buccal bone thickness and less than 50% dehiscence with or without interproximal bone loss and/or apical lesion. The root position is adjacent to the palatal plate. Mild periodontal or endo-perio origin can be a feature of class II sockets. **Class III:** The etiology of a class III socket can be severe perio or endo-perio lesions. The gingiva has more than 3mm of recession also severe loss of buccal plate in terms of dehiscence puts a socket into class III. The root position is unfavorable with only 2/3 of the root engaging the buccal and palatal plates. In order to facilitate classification process, even one criteria that meets the features of each class will put the ES into the respective socket type. For instance, a socket with ideal clinical and radiographic parameters but gingival recession of more than 4 mm would be considered as a class III.

### Single-Rooted Extraction Socket Classification

	Class I	Class II	Class III
<b>Etiology</b>	<b>Non Periodontal</b>	<b>Mild periodontal or endo-perio lesion</b>	<b>Severe periodontal or endo-perio lesion</b>
<b>Gingival recession (mm)</b>	<b>≤3</b>	<b>-</b>	<b>&gt;3</b>
<b>Soft tissue phenotype</b>	<b>Thick</b>	<b>Thin</b>	<b>-</b>
<b>Buccal Bone Width (mm)</b>	<b>≥2</b>	<b>&lt;2</b>	<b>-</b>
<b>Buccal bone loss</b>	<b>Intact</b>	<b>&lt;50%</b>	<b>&gt;50%</b>
<b>Interproximal bone loss</b>	<b>No</b>	<b>Yes</b>	<b>-</b>
<b>Apical pathology</b>	<b>No</b>	<b>Yes</b>	<b>-</b>
<b>Root position</b>	<b>Adjacent to vestibular plate/ at the center</b>	<b>Adjacent to palatal plate</b>	<b>At least 2/3 of the root engaging both buccal and palatal plates</b>

**Table 5. Extraction Socket Class Modifiers.** The modification proceeds the classification step. This aims to include factors that are not properly examinable prior to the extraction and designed to adjust the initial classification if required. These can be divided into extraction- and osteotomy-related factors. If the extraction process occurs invasively and cause any damage to the adjacent structure this will transform class I and II to class III. Similar scenario is applicable for iatrogenic complications such as nerve damage or sinus floor perforation. Finally, post extraction evaluation of the socket is required to determine whether it is possible to place implant in the correct position in correspondence to adjacent structure (nerve proximity etc.) and if not possible, classes I and II will be considered as class III.

<b>Post-extraction Class Modifiers</b>			
<b>Patient-related factors</b>	Active periodontitis in the same sextant		Class I and II to III
	Poor oral hygiene		Class I and II to III
	Medications affecting healing		Class II to III
	Poorly controlled systemic disease		Class II to III
	Smoking	More than 10/day	Class I and II to III
<b>Extraction related factors</b>	Invasively traumatic extraction (extensive bone removal)		Class I and II to III
	Iatrogenic complications (sinus floor damage, nerve damage, Buccal plate fracture)		Class I and II to III
	Post-extraction evaluation of buccal bone thickness and bone quality		If compromised, Class I and II to III
<b>Osteotomy related Factors</b>	Possible limitations in implant osteotomy (nerve proximity, adjacent roots etc.)		If IIP not possible, Class I and II to III

