

Koebnick Corinna (Orcid ID: 0000-0001-8274-0309)
Resnicow Kenneth (Orcid ID: 0000-0003-1416-9627)
Young Deborah Rohm (Orcid ID: 0000-0002-8335-7851)
Woolford Susan J (Orcid ID: 0000-0003-4431-0652)

Disparities in weight changes during the COVID-19 pandemic-related lockdown among youths

Corinna Koebnick*¹, PhD; Margo A. Sidell,*¹ ScD; ¹ Xia Li¹, MSc; Ken Resnicow,² PhD; Poornima Kunani,³ MD; Deborah R Young¹, PhD; Susan J. Woolford,⁴ MD
*Shared equal first authors

¹ Department of Research & Evaluation, Kaiser Permanente Southern California, Pasadena, CA,

² School of Public Health, University of Michigan Health Behavior and Health Education, Ann Arbor, MI

³ Department of Pediatrics, Kaiser Permanente Manhattan Beach Medical Office, Manhattan Beach, CA

⁴ Child Health Evaluation and Research Center, Department of Pediatrics, University of Michigan, Ann Arbor, MI

Keywords: childhood obesity, Body mass index, weight gain, SARS-COV-2, disparities

Running title: Weight gain in youth during the COVID-19 pandemic

Contact information: Corinna Koebnick, PhD
Department of Research & Evaluation, Kaiser Permanente Southern California,
100 S. Los Robles, 2nd Floor, Pasadena, CA 91101
Corinna.Koebnick@kp.org, Phone: (626) 564-3693.

Word count: 3253

Funding/Support: All phases of this study were supported by Kaiser Permanente Community Funds.

Disclosures: The authors declared no conflict of interest.

Author contributions: All authors conceptualized and designed the study and supervised the data collection and the statistical analyses. Dr. Sidell and Mrs. Li conducted the statistical analyses. Dr. Koebnick and Dr. Woolford drafted the initial manuscript. All authors contributed to the data interpretation and reviewed and revised the manuscript. All authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Abbreviations: BMI – body mass index; KPSC – Kaiser Permanente Southern California; RR – relative risk; COVID-19 – coronavirus disease 2019, EHR - electronic health record.

This is the author manuscript accepted for publication and has undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/oby.23645

This article is protected by copyright. All rights reserved.

What is already known about this subject

- The pandemic contributed to excess weight gain among Southern California children
- Overall childhood obesity during this period increased by almost 5.1%

What are the new findings in your manuscript?

- We observed significant racial and social disparities in weight gain during the COVID-19 pandemic lockdown
- Black and Hispanic youth gained more body weight than their white counterparts
- Youth from low-income families and those with limited access to neighborhood parks also gained more weight.

How might your results change the direction of research or the focus of clinical practice?

- Programs that promote healthy eating and physical activity should be developed that account for the unique stressors and lifestyle changes during the COVID-19 pandemic lockdown and will likely persist in ways we currently do not understand.

ABSTRACT

Objective: This study evaluates whether changes in weight among school-aged youth in California due to the COVID-19 lockdown vary by social constructs of race/ethnicity and associated social factors.

Methods: Including 160,472 youth ages 5-17 years enrolled at Kaiser Permanente Southern California, mixed effects models stratified by age group were fitted to estimate changes in distance from the median body mass index (BMI)-for-age from 3/2020 to 1/2021 (lockdown) compared to the same period prepandemic.

Results: Excess pandemic weight gain was higher among Black and Hispanic youth 5-17 years than among white and Asian youth; this difference was most pronounced in those ages 5-11 years. In youth 5-11 years, the distance from the median BMI-for-age increased by 1.72 kg/m² (95% CI 1.61, 1.84) in Hispanic and 1.70 kg/m² (95% CI 1.47, 1.94) in Black youth during the lockdown compared to 1.16 kg/m² (95% CI 1.02, 1.29) in non-Hispanic white youth. The excess weight gain was also higher in youth with fewer neighborhood parks and those with state-subsidized health insurance.

Conclusions: The COVID-19 pandemic lockdown led to a gain of excess body weight, particularly for Black and Hispanic youth; this weight gain varied by social factors associated with race and ethnicity.

INTRODUCTION

The COVID-19 pandemic has dramatically altered many aspects of life across all population sectors, including children and adolescents.¹ Due to the pandemic-related lockdown, schools were intermittently closed, learning was moved online, recreational activities were curtailed, social activities were restricted, and the food supply was impacted.^{2,3} These changes, combined with the emotional and financial stress caused by the pandemic, may affect energy balance and, consequently, body weight.⁴ We recently reported that the pandemic contributed to accelerated weight gain among Southern California children and led to an almost 5.1% increase in the prevalence of obesity, the same increase observed nationally over the last nearly two decades combined.^{5,6}

In the US, childhood obesity (\geq 95th percentile of body mass index, BMI-for-age) disproportionately affects Hispanic and Black youth compared to non-Hispanic white youth (25.8%, 22.0% vs. 14.1%, respectively) and children from families with low income.^{7,8} While the COVID-19 pandemic has affected the country as a whole, Hispanic, non-Hispanic Black and populations with low income and socioeconomic status have shouldered the burden of higher case rates, hospitalizations, and deaths.^{9,10} The same populations may also experience a higher burden of weight gain during the pandemic.

The present study investigates disparities in weight changes among youths 5 to 17 years of age of different racial/ethnic and social groups among members of a Kaiser Permanente health plan in Southern California. Early in the pandemic, California had one of the world's highest per capita rates of COVID-19 infection, followed by one of the most prolonged and strict lockdown orders in the US, lasting from March 19, 2020 to January 25, 2021.

METHODS

Study design and setting

To compare BMI in youth before and during the COVID-19 pandemic-related lockdown

(including the stay-home period) in California, we conducted a retrospective cohort study among pediatric health plan members of Kaiser Permanente Southern California (KPSC), an integrated managed healthcare system that serves approximately 4.7 million members in Southern California (~19.5% of the Southern California population). We compared electronic health record (EHR) data during the lockdown (March 01, 2020 through January 31, 2021) with data from the same youth before the lockdown (March 01, 2019 through January 31, 2020) as the comparison period. KPSC members receive care in medical offices and hospitals owned by KPSC throughout the nine-county region. The membership is diverse and similar in socioeconomic characteristics to the region's census demographics.¹¹ The KPSC Institutional Review Board approved the study and granted a waiver for informed consent.

Study population

We identified school-aged youth between 5 and 17 years enrolled at KPSC on March 16, 2020 (n = 701,082, **Figure 1**).¹² To be eligible for the present study, youth were required to have a continuous health plan membership between 1/1/2020 and 12/31/2020 and in the 24 months prior to March 16, 2020 (allowing a 90-day gap in coverage) and to be free of any pediatric complex chronic conditions such as cerebral palsy, congenital and chromosomal abnormalities.^{13,14} Eligible youth (n=569,350) were required to have one or more valid BMI measurements between June 16, 2020 (i.e., about three months after the start of the lockdown in California) and the end of the study period, January 31, 2021 (n=197,707). We excluded youth without a BMI measurement during the comparison period (March 2019 to Jan 2020 = 37,235, 3.2%). The final analytical cohort comprised 160,472 youth (667,554 BMI measurements).

Body weight and height

Weight and height measurements, routinely obtained at clinic visits by trained medical staff, were extracted from the EMR; BMI was calculated as weight (kilograms)

divided by the square of the height (meters). Well-child visits during the pandemic were conducted as in-person or phone/video visits with follow-up in a nurse clinic where weight and height were measured (mainly to update vaccination). In July 2020, in-person visits with weight and height measurements were at 76.4 % of the prepandemic in-person visit numbers (**Table S1**). BMI percentiles are based on sex-specific BMI-for-age according to growth charts developed by the Centers for Disease Control and Prevention (**Table S2**).¹⁵⁻

18

Our primary measure of increased adiposity was the absolute difference in the distance from the median BMI-for-age and sex.¹⁶ This metric is a more reliable measure of change in adiposity, particularly for individuals in the upper end of the BMI distribution, compared to other methods such as BMI z-score. It does not have an upper limit (as has BMI-for-age percentile) and can be used to assess adiposity across the entire BMI spectrum. In addition, we also provide data on change in body weight adjusted for height and the prevalence of overweight and obesity because these measures of weight status are more commonly used by clinicians.^{15,19}

Main exposures and covariates

All covariates are defined in **Table S2**.²⁰ We categorized self-reported race/ethnicity from various sources such as registration records, clinical visit records, and birth certificates as non-Hispanic White (hereafter, White), Hispanic, or Latino (regardless of race, hereafter, Hispanic), Black or African American (hereafter, Black), Asian or Pacific Islander, and other or unknown race/ethnicity. Non-Hispanic White, Black, Hispanic, Asian/Pacific Islander, other or unknown. Data for this study included US adults who self-reported as non-Hispanic Black (hereafter, Black), Hispanic or Latino, and non-Hispanic White (hereafter, White) individuals. Health insurance coverage through government health care assistance programs such as Medicaid (Yes/No) was used as a proxy for socioeconomic status. Neighborhood education and income were derived from geocoding patients' address information and based on the 2010 US census block group and annual

American Community Survey information. Neighborhood park information was derived using ArcGIS software (Version 10.1, ESRI, Redlands, CA), Claritas LLC (Cincinnati, OH), and the ESRI USA 2019 parks layer package.

Statistical analysis

Standardized mean differences (SMD) were used to compare the KPSC source cohort to the analytical cohort; SMD values ≤ 0.2 were considered small.²¹ We fit mixed-effect linear models for distance from the median BMI-for-age and body weight; Poisson regression was used for overweight and obesity. All models accounted for repeated measures within each child, using an autoregressive correlation structure and maximum likelihood estimation of covariance parameters to assess each outcome during the 11-month COVID-19 pandemic lockdown (March 2020 to January 2021) compared to the same children in the same calendar months before COVID-19 (March 2019 to January 2020). Similar to an interrupted time series design, we included a binary indicator representing the periods before or during the pandemic plus a calendar month by period interaction term. Additional details on the model development are described in the

Supplemental Methods. All final models included three 3-way interactions for a pandemic by time by race/ethnicity, parks, and state-subsidized health insurance, as well as their main effects and 2-way interactions and adjust for sex, neighborhood education, and income. BMI and body weight models were also adjusted for BMI-for-age group at baseline, and the model for body weight was additionally adjusted for height. All analyses were performed with $\alpha = 0.05$ for two-sided tests using SAS version 9.4 (SAS Institute, Inc., Cary, NC, USA).

RESULTS

The analytical cohort was racially/ethnically diverse (50.7% Hispanic, 25.2% non-Hispanic White, 6.9% non-Hispanic Black, 10.4% Asian/Pacific Islander) and comparable to the overall KPSC pediatric population regarding to sex, age, race/ethnicity, and

socioeconomic factors (**Table S3, Table 1**). Before the pandemic, 39.2% of children in the analytic cohort lived with overweight, obesity, or severe obesity compared to 39.4% of children in the KPSC source population.

The COVID-19 pandemic lockdown was associated with a higher distance from the median BMI-for-age compared to the pre-pandemic period which varied by race/ethnicity, socioeconomic status, and number of neighborhood parks (**Table 2**). Excess weight gain compared to the same period before the pandemic lockdown was most pronounced among Hispanic and Black youth with an excess increase in the distance from the median BMI-for-age by 1.72 kg/m² (95% CI 1.61, 1.84) and 1.70 kg/m² (95% CI 1.47, 1.94), respectively, during the pandemic compared to 1.16 kg/m² (95% CI 1.02, 1.29) in non-Hispanic white youth (P for lockdown x time x race/ethnicity < .001 for 5-11 years, P < .001 for 12-15 years, and P = .004 for 16-17 years, respectively). While the excess weight gain overall was lower in youth 12-15 years and 16 – 17 years, Hispanic and Black youth consistently gained more than non-Hispanic white youth (**Table 2**).

Youth with state-subsidized health insurance gained more body weight than those with other insurance types (P for lockdown x time x state-subsidized insurance < .001 for 5-11 years, P = .02 for 12-15 years, and P = .39 for 16-17 years, respectively). Among 5-11 years olds, youth with state-subsidized health insurance increased their distance from the median BMI-for-age by 1.39 kg/m² (95% CI 1.21, 1.56) compared to 1.16 kg/m² (95% CI 1.02, 1.29) in those with commercial or privately funded health plans.

Youth across all ages who lived in neighborhoods with 2 or more parks gained less excess body weight during the lockdown than during the pre-pandemic period compared to those who lived in neighborhoods with 1 park or no parks (P for lockdown x time x parks P <.001 for 5-11 years, P =.03 and 12-15 years, P = .02 for 16-17, respectively).

In models using body weight (rather than distance from the median), the excess weight gain during the pandemic lockdown compared to prepandemic weight gain ranged between 1.90 kg (95% CI 1.74, 2.06) in White, 2.63 kg (95% CI 2.50, 2.76) in Hispanic, and 2.79 kg (95% CI 2.51, 3.06) in Black youth 5-11 years old, respectively (**Table S3**).

Among 5-11 year-olds, youth with state-subsidized health insurance gained 2.20 kg (95% CI 1.99, 2.41) compared to 1.90 kg (95% CI 1.74, 2.06) in those on commercial and private health plans. Among 5-11 year-olds, youth with two or more neighborhood parks gained 1.55 kg (1.39, 1.71) compared to 1.90 kg (95% CI 1.74, 2.06) and 1.81 kg (95% CI 1.65, 1.98) for those with one park or no parks, respectively. The adjustment for height ensures that changes in weight reflect changes independent of changes in height.

The prevalence of obesity increased among white youths 5-11 years old at the time of the lockdown from 15.2% before the lockdown to 22.0% during the pandemic lockdown, an absolute excess increase of 5.3% (95% CI 3.8, 6.7%) compared to the reference period (**Table 3**). This compared to an absolute excess increase of 7.0% (95% CI 4.5, 9.6) and 8.2% (95% CI 7.3, 9.2) among Black and Hispanic youth 5-11 years old, respectively. Among youth 12-17 years, Black and Hispanic youth had a higher prepandemic prevalence of obesity than whites; and although we found higher excess gains for Black and Hispanic youth aged 12 to 15, they did not differ in the excess increase in obesity rates among older age groups (P for lockdown x time x race and ethnicity = .49 for 5-11 years, P = .86 for 12-15 years, and P = .58 for 16-17 years, respectively). Among 5-11 year olds, the increase in obesity was marginally lower in youth from neighborhoods with 2 or more parks (5.0% vs 6.9% and 6.5%, respectively, for 2+ vs. 0 and 1 park; P for lockdown x time x parks: P = .06 for 5-11 years); this association was not observed for the older age groups (P for lockdown x time x parks: P = .89 for 12-15 years, and P = .30 for 16-17 years, respectively). No significant difference in the increase in obesity during the pandemic between those with and without state-subsidized health insurance was observed (P for lockdown x time x state-subsidized health insurance: P = .63 for 5-11 years, P = .94 for 12-15 years, and P = .12 for 16-17 years, respectively).

To visualize the additive effects of factors associated with excess weight gain among 5-11 year old youth, we compared a reference youth who live with overweight and have social advantages (non-Hispanic White, without state-subsidized insurance, from

neighborhoods with 2 or more parks) to youth who live with overweight without these social advantages (Hispanic and Black youth who are overweight, had state-subsidized insurance, and lived in neighborhoods without parks, **Figure 2**). For example, non-Hispanic white youth with social advantages increased their distance from the median BMI by 0.84 kg/m² (95% CI 0.71, 0.97) compared to non-Hispanic white youth without these advantages who gained 1.34 kg/m² (95% CI 1.17, 1.51). In contrast, Hispanic and Black youth without those social advantages gained 1.91 kg/m² (95% CI 1.78, 2.04) and 1.89 (95% CI 1.65, 2.12), respectively.

DISCUSSION

Communities of color and low income populations have been disproportionately affected by the COVID-19 pandemic in the U.S.^{9,22-24} including California.²⁵⁻²⁷ These populations carried a higher burden of infection, hospitalizations, and deaths from COVID-19. Since the beginning of the pandemic, public health experts were concerned that obesity rates might increase as a result of COVID-19 mitigation strategies such as schools closings, limitations on team sports, social, and recreational activities, alterations in eating habits, and food supply, as well as stress and social isolation.²⁸

We and others showed a dramatic excess pandemic weight gain amongst US youth during the COVID-19 lockdown period when compared to the same period before the pandemic lockdown.^{5,29-32} The present study examines racial, ethnic, and social, and neighborhood differences in weight changes during the during the COVID-19 pandemic lockdown. Among ~160,000 youth in southern California, the lockdown was associated with significantly higher excess weight gain among Hispanic and Black youth and those living in families with low socioeconomic status and youth from neighborhoods without parks. While not a national sample, our findings together with findings of others^{5,29-32} raise concerns of a national trend that existing gaps in youth health based on race or ethnicity, income level, and environmental inequalities have widened during the lockdown.

Socioeconomic, educational, and environmental disadvantages have historically affected vulnerable communities with strong links to disparities in nutrition and obesity.³³ Over the last almost two decades, data from the National Health and Nutrition Examination Surveys showed an annual increase in obesity of 0.20 % in 6-11 year olds and 0.35% in 12-19 year olds.³⁴ In contrast, we found that the increase in obesity during the 11-month pandemic lockdown in Southern California ranged from 5.1% in non-Hispanic white to 8.1% in Hispanic youth among 5-11 year olds among; this difference was slightly lower in older children. This means that the increase in obesity rates during the lockdown was almost equivalent to the increase observed over the last two decades. The higher excess weight gain in youth of color will, if they persist, contribute to widening of the existing disparities in childhood obesity.³⁵

Access to parks and playgrounds is inversely associated with childhood obesity³⁶⁻³⁸ and cardiorespiratory fitness in children,³⁹ especially in youth living in families with low income.^{36,38} In the present study, living in communities with fewer parks was associated with higher weight gain during the pandemic lockdown, a time when schools were closed, and most youth sports and other recreational activities were suspended. A higher number of neighborhood parks may have alleviated the excess weight increase in youth, especially among 5-11 year olds. Additionally, youth between 5-11 from families with low socioeconomic background gained more than their peers. However, due to the nature of the study this association may also reflect associations with other neighborhood factors and reflect economic, financial, and health inequalities.

Structural racism has patterned wealth and shaped environments such that youth of color are more likely to live in families with low income and with limited access to parks.⁴⁰ Racial and ethnic disparities in obesity are driven by socioeconomic and environmental disadvantages and well-established in adults and children.^{33,34,41} In 2016, a study conducted in Los Angeles County, the largest counties in southern California, revealed

that more than half of the county had very little greenspace; about 82% of these “park-poor” areas were located in communities of color.⁴² The present study demonstrates how the combination of these risk factors such as low income and limited parks access add to the observed racial disparities in excess weight gain. Our data show that youth who live with social advantages gained significantly less than those without these social advantages. Because youth from communities of color will be less likely to have social advantages,^{33,43,44} this may contribute to a further widening of the health gap associated with racial disparities.

Other aspects of structural racism may explain the greater pandemic lockdown-related weight gain in Hispanic and Black youths than in their white counterparts. The emotional and financial stress faced by Black and Hispanic families caused by the COVID-19 pandemic, job losses, over-representation in frontline jobs that cannot be done remotely, closed schools with limited or no access to school lunch, lack of childcare, housing and food insecurity may impact energy balance and, consequently, body weight.^{4,45} Food insecurity increased during the pandemic and was most pronounced among Black and Hispanic families than any other group.⁴⁶ Food insecurity can lead to lower dietary quality and greater consumption of calorie-dense foods, which can predispose to weight gain.⁴⁷⁻⁵¹ A full discussion of the complex ways communities of color were affected in comparison to primarily white communities is beyond the scope of this study.

These findings should be considered within the context of certain limitations. In the absence of an unexposed control group in 2020, our analyses depended on comparisons to weight patterns among the same youth in the year before the pandemic lockdown. Our analysis also only included those children with in-person appointments during the time period of interest. However, in-person well-child visits resumed within 2 months (in part to ensure standard child vaccinations remained up to date), suggesting that the BMI data

collected during this time reflect the BMI of the source population. While the possibility of selection bias exists, our analytical cohort was strikingly similar to the overall population of KPSC pediatric patients in all relevant characteristics including BMI. We used state-subsidized health insurance as proxy for socioeconomic status and number of parks in the neighborhood of the primary address of the patient a proxy for access to parks and playgrounds. Finally, in this observational study conducted in Southern California, we cannot establish causation or determine which factors other than the COVID-19 pandemic accounted for the excess weight gain noted during 2020. Our results also may not reflect populations outside of the study area.

The pandemic had a devastating effect on the physical, psychologic, and economic health of millions in the US. While the direct impact of the COVID-19 pandemic on morbidity and mortality was less severe for the pediatric population, the indirect impact on the weight of children may have long-term detrimental health consequences and exacerbate already existing racial and social disparities. It may place children of color, at risk for comorbidities associated with excess weight such as diabetes and hypertension and with higher incidence rates and at an earlier age, especially when they are from families with low socioeconomic background. In turn, this could make children more susceptible to poor outcomes after infection with SARS-CoV-2 or other similar pathogens in the future. It will be important to track this cohort to see if the excess weight gain observed during COVID-19 is maintained. Future studies will have to show if similar trends are observed in other states and areas outside of Southern California. Analysis of diet and activity patterns (including the increase in screen time associated with at-home education) during COVID-19 will be helpful in understanding the drivers of the weight gain. Diet and activity interventions should be developed that account for the unique stressors and lifestyle changes that occurred during the COVID-19 pandemic lockdown and will likely persist in ways we currently do not understand.

Acknowledgments

Thank you to the patients of Kaiser Permanente for helping to improve care with information collected through our electronic health record systems.

TABLE 1: Characteristics of the analytical cohort by age group.

	Analytical cohort	Age at baseline** (years)		
		5-11	12-15	16-17
N	160,472	84,762	51,635	24,075
Age at baseline				
Mean (SD)	11.5 (3.8)	8.4 (2.2)	14.1 (1.2)	16.9 (0.6)
Sex				
Girls	79496 (49.5)	40895 (48.2)	25620 (49.6)	12981 (53.9)
Boys	80976 (50.5)	43867 (51.8)	26015 (50.4)	11094 (46.1)
Race/ethnicity				
Non-Hispanic White	40487 (25.2)	21324 (25.2)	13029 (25.2)	6134 (25.5)
Non-Hispanic Black	11052 (6.9)	5546 (6.5)	3699 (7.2)	1807 (7.5)
Hispanic	81421 (50.7)	42524 (50.2)	26504 (51.3)	12393 (51.5)
Asian/Pacific Islanders	16740 (10.4)	9454 (11.2)	5131 (9.9)	2155 (9.0)
Other/Unknown	10772 (6.7)	5914 (7.0)	3272 (6.3)	1586 (6.6)
State-subsidized health insurance				
No	126118 (78.6)	65883 (77.7)	40998 (79.4)	19237 (79.9)
Yes	34354 (21.4)	18879 (22.3)	10637 (20.6)	4838 (20.1)
Neighborhood Education				
Less than high school	31364 (19.5)	16853 (19.9)	9898 (19.2)	4613 (19.2)
High school	35961 (22.4)	19085 (22.5)	11499 (22.3)	5376 (22.3)
Some college or higher	93147 (58.0)	48824 (57.6)	30237 (58.6)	14086 (58.5)
Neighborhood Income (\$)				
<\$35,000	36034 (22.5)	19303 (22.8)	11409 (22.1)	5322 (22.1)
\$35,000 to \$49,999	17973 (11.2)	9599 (11.3)	5707 (11.1)	2667 (11.1)
\$50,000 to \$74,999	25971 (16.2)	13810 (16.3)	8292 (16.1)	3869 (16.1)
\$75,000 to \$99,999	21225 (13.2)	11215 (13.2)	6828 (13.2)	3182 (13.2)
\$100,000 to \$149,999	29194 (18.2)	15320 (18.1)	9465 (18.3)	4408 (18.3)
\$150,000 or more	30075 (18.7)	15514 (18.3)	9933 (19.2)	4628 (19.2)
Parks				
0	55545 (34.6)	29866 (35.2)	17540 (34.0)	8139 (33.8)
1 (includes missing)	57825 (36.0)	30472 (36.0)	18643 (36.1)	8710 (36.2)

2+	47102 (29.4)	24424 (28.8)	15452 (29.9)	7226 (30.0)
BMI before March, 2020				
N	160472	84762	51635	24075
Mean (SD)	20.8 (5.4)	18.4 (4.0)	22.9 (5.3)	24.9 (5.7)
% BMI distance to median				
Mean (SD)	19.3 (28.2)	18.2 (28.4)	20.8 (27.8)	20.2 (27.9)
BMI distance to median				
Mean (SD)	4.3 (6.3)	4.1 (6.4)	4.6 (6.2)	4.5 (6.2)
BMI-for-age #				
Underweight	3228 (2.0)	2013 (2.4)	852 (1.7)	363 (1.5)
Normal	94419 (58.8)	50804 (59.9)	29552 (57.2)	14063 (58.4)
Overweight	29702 (18.5)	14772 (17.4)	10138 (19.6)	4792 (19.9)
Obesity	22386 (14.0)	12132 (14.3)	7203 (13.9)	3051 (12.7)
Severe Obesity	10737 (6.7)	5041 (5.9)	3890 (7.5)	1806 (7.5)
Missing	0	0	0	0

* Baseline: March 01, 2020

Youth were categorized at baseline as living with underweight (BMI-for-age <5th percentile), normal weight (BMI-for-age ≥5th to <85th percentile), overweight (BMI-for-age ≥85th to <95th percentile), moderate obesity (BMI-for age ≥95th to <1.2x95th percentile), and severe obesity (BMI-for age ≥1.2x95th percentile).

† Standardized mean differences (SMD) were tested for each variable between two KPSC source populations. All SMD values were smaller than 0.2, suggesting no difference between the two groups.

TABLE 2: Predicted means and change in the distance from the median BMI-for-age in youth before and during the pandemic by race/ethnicity and socioeconomic factors

Characteristic	Prepandemic (March 2019 to January 2020)			Pandemic (March 2020 to January 2021)			$\Delta 2 - \Delta 1$ (95% CI)
	Start Mean (SD.)	End Mean (SD.)	Change 1 ($\Delta 1$, 95% CI)	Start Mean (SD.)	End Mean (SD.)	Change 2 ($\Delta 2$, 95% CI)	
5-11 years							
Race/Ethnicity							
NH White	0.35 (0.03)	0.39 (0.03)	0.04 (-0.03, 0.12)	0.43 (0.04)	1.64 (0.04)	1.20 (1.09, 1.32)	1.16 (1.02, 1.29)
Hispanic	0.52 (0.03)	0.76 (0.03)	0.24 (0.18, 0.29)	0.78 (0.04)	2.74 (0.03)	1.96 (1.86, 2.06)	1.72 (1.61, 1.84)
NH Black	0.44 (0.05)	0.78 (0.05)	0.34 (0.22, 0.47)	0.68 (0.07)	2.73 (0.06)	2.05 (1.85, 2.25)	1.70 (1.47, 1.94)
API	0.00 (0.04)	0.25 (0.04)	0.25 (0.15, 0.35)	0.07 (0.06)	1.67 (0.05)	1.60 (1.44, 1.77)	1.36 (1.16, 1.55)
Other/Unknown	0.22 (0.05)	0.47 (0.05)	0.25 (0.13, 0.38)	0.53 (0.07)	1.86 (0.06)	1.32 (1.12, 1.52)	1.07 (0.84, 1.30)
State-subsidized health care							
No	0.42 (0.03)	0.47 (0.03)	0.04 (-0.03, 0.12)	0.51 (0.04)	1.71 (0.04)	1.20 (1.09, 1.32)	1.16 (1.02, 1.29)
Yes	0.56 (0.04)	0.62 (0.04)	0.07 (-0.02, 0.16)	0.74 (0.06)	2.20 (0.05)	1.46 (1.31, 1.61)	1.39 (1.21, 1.56)
Neighborhood parks							
No park	0.39 (0.03)	0.50 (0.03)	0.10 (0.03, 0.18)	0.59 (0.05)	1.80 (0.04)	1.22 (1.10, 1.34)	1.11 (0.97, 1.25)
1 park	0.46 (0.03)	0.50 (0.03)	0.04 (-0.03, 0.12)	0.54 (0.05)	1.74 (0.04)	1.20 (1.09, 1.32)	1.16 (1.02, 1.29)

2 or more parks	0.35 (0.03)	0.42 (0.03)	0.07 (-0.00, 0.14)	0.55 (0.04)	1.46 (0.04)	0.91 (0.80, 1.02)	0.84 (0.71, 0.97)
12-15 years							
Race/Ethnicity							
NH White	0.36 (0.04)	0.52 (0.04)	0.16 (0.09, 0.23)	0.47 (0.05)	1.17 (0.04)	0.70 (0.58, 0.81)	0.54 (0.40, 0.67)
Hispanic	0.78 (0.03)	0.76 (0.03)	-0.02 (-0.08, 0.04)	0.74 (0.04)	1.68 (0.03)	0.93 (0.84, 1.03)	0.95 (0.84, 1.07)
NH Black	0.85 (0.06)	0.85 (0.06)	-0.00 (-0.13, 0.13)	0.89 (0.08)	1.96 (0.06)	1.08 (0.88, 1.27)	1.08 (0.85, 1.30)
API	0.28 (0.05)	0.29 (0.05)	0.01 (-0.10, 0.13)	0.17 (0.07)	0.77 (0.06)	0.60 (0.42, 0.78)	0.58 (0.37, 0.80)
Other/Unknown	0.60 (0.06)	0.54 (0.06)	-0.06 (-0.20, 0.07)	0.51 (0.08)	1.31 (0.07)	0.80 (0.60, 1.01)	0.87 (0.62, 1.11)
State-subsidized health care							
No	0.42 (0.04)	0.58 (0.04)	0.16 (0.09, 0.23)	0.54 (0.05)	1.24 (0.04)	0.70 (0.58, 0.81)	0.54 (0.40, 0.67)
Yes	0.51 (0.05)	0.71 (0.05)	0.20 (0.11, 0.30)	0.68 (0.06)	1.57 (0.05)	0.90 (0.75, 1.05)	0.69 (0.52, 0.87)
Neighborhood parks							
No park	0.46 (0.04)	0.62 (0.04)	0.16 (0.08, 0.24)	0.58 (0.05)	1.26 (0.04)	0.68 (0.57, 0.80)	0.52 (0.38, 0.66)
1 park	0.45 (0.04)	0.61 (0.04)	0.16 (0.09, 0.23)	0.56 (0.05)	1.26 (0.04)	0.70 (0.58, 0.81)	0.54 (0.40, 0.67)
2 or more parks	0.33 (0.04)	0.50 (0.04)	0.17 (0.09, 0.24)	0.49 (0.05)	1.02 (0.04)	0.53 (0.42, 0.65)	0.37 (0.24, 0.50)
16-17 years							
Race/Ethnicity							
NH White	0.42 (0.05)	0.28 (0.05)	-0.14 (-0.24, -0.03)	0.34 (0.06)	0.40 (0.06)	0.06 (-0.09, 0.22)	0.20 (0.02, 0.39)
Hispanic	0.64 (0.05)	0.47 (0.05)	-0.18 (-0.26, -0.09)	0.46 (0.06)	0.82 (0.05)	0.36 (0.24, 0.49)	0.54 (0.39, 0.69)

NH Black	0.86 (0.08)	0.66 (0.08)	-0.21 (-0.38, -0.04)	0.72 (0.10)	1.13 (0.09)	0.41 (0.17, 0.66)	0.62 (0.32, 0.92)
API	0.28 (0.08)	0.06 (0.08)	-0.23 (-0.40, -0.05)	-0.06 (0.10)	-0.02 (0.08)	0.04 (-0.21, 0.30)	0.27 (-0.04, 0.58)
Other/Unknown	0.55 (0.09)	0.24 (0.09)	-0.31 (-0.50, -0.13)	0.44 (0.11)	0.52 (0.09)	0.08 (-0.18, 0.35)	0.40 (0.08, 0.72)
State-subsidized health care							
No	0.47 (0.05)	0.33 (0.05)	-0.14 (-0.24, -0.03)	0.39 (0.07)	0.45 (0.06)	0.06 (-0.09, 0.22)	0.20 (0.02, 0.39)
Yes	0.65 (0.07)	0.42 (0.07)	-0.23 (-0.37, -0.09)	0.55 (0.09)	0.60 (0.07)	0.05 (-0.15, 0.25)	0.28 (0.04, 0.53)
Neighborhood parks							
No park	0.53 (0.06)	0.32 (0.06)	-0.21 (-0.32, -0.10)	0.39 (0.07)	0.55 (0.06)	0.16 (0.00, 0.32)	0.37 (0.18, 0.56)
1 park	0.49 (0.05)	0.35 (0.05)	-0.14 (-0.24, -0.03)	0.41 (0.07)	0.47 (0.06)	0.06 (-0.09, 0.22)	0.20 (0.02, 0.39)
2 or more parks	0.38 (0.05)	0.32 (0.05)	-0.06 (-0.16, 0.04)	0.32 (0.06)	0.36 (0.06)	0.04 (-0.11, 0.18)	0.10 (-0.08, 0.28)

Total BMI measures included in the models were 425,855 from March 2019-January 2020 (2.65 BMI measures/youth) compared to 241,699 from March 2020-January 2021 (1.51 BMI measures/youth). After initial decrease, in-person well-child visits were back to 84% of prepandemic visits by June 2020.

All models are adjusted for race/ethnicity (Hispanic, non-Hispanic black, Asian or Pacific Islander, other, reference: white), state-subsidized health care (reference: none), parks (no parks, 2 or more parks, reference: 1 park), neighborhood education, neighborhood income, and baseline weight class (<5th, 5th to 84th, 85 to 94th; 95 to 97th, ≥97th; reference: 5th to ≤ 85th). Estimates are shown for the reference group.

Interaction terms: Pandemic x time x race/ethnicity P = 0.001 for 5-11 years, P < 0.001 for 12-15 years, P = 0.004 for 16-17 years, Pandemic x time x State-subsidized insurance P < 0.001 for 5-11 years, P = 0.02 for 12-15 years, P = 0.39 for 16-17 years, Pandemic x time x parks P < 0.001 for 5-11 years, P = 0.03 for 12-15 years, P = 0.02 for 16-17 years.

TABLE 3: YOUTH LIVING WITH OBESITY IN BEFORE AND DURING THE PANDEMIC

Race/Ethnicity	Prepandemic			Pandemic			
	Start Mean (SD.)	End Mean (SD.)	Change 1 ($\Delta 1$, 95% CI)	Start Mean (SD.)	End Mean (SD.)	Change 2 ($\Delta 2$, 95% CI)	$\Delta 2 - \Delta 1$ (95% CI)
Obesity ($\geq 95^{\text{th}}$ percentile), Mean (SD.)							
5-11 years							
NH White	14.17 (0.32)	15.15 (0.33)	0.99 (0.27,1.70)	15.71 (0.42)	21.95 (0.44)	6.24 (5.04,7.44)	5.25 (3.84,6.66)
Hispanic	23.53 (0.25)	24.91 (0.26)	1.38 (0.92,1.84)	25.37 (0.31)	34.97 (0.34)	9.59 (8.77,10.42)	8.21 (7.27,9.16)
NH Black	19.66 (0.60)	21.77 (0.64)	2.11 (0.81,3.40)	22.32 (0.80)	31.46 (0.84)	9.14 (6.88,11.40)	7.03 (4.50,9.55)
API	13.38 (0.46)	14.43 (0.46)	1.06 (0.03,2.09)	14.02 (0.61)	21.14 (0.64)	7.11 (5.28,8.94)	6.06 (3.98,8.14)
Other/Unknown	16.43 (0.61)	18.32 (0.65)	1.88 (0.44,3.32)	19.04 (0.84)	25.19 (0.80)	6.16 (3.82,8.49)	4.28 (1.59,6.96)
12-15 years							
NH White	16.27 (0.42)	17.01 (0.43)	0.74 (-0.09,1.57)	16.46 (0.51)	21.46 (0.54)	5.00 (3.63,6.37)	4.26 (2.64,5.88)
Hispanic	24.68 (0.33)	24.39 (0.33)	-0.29 (-0.86,0.29)	24.08 (0.40)	29.39 (0.39)	5.31 (4.34,6.28)	5.60 (4.45,6.74)
NH Black	24.90 (0.79)	24.60 (0.79)	-0.30 (-1.75,1.16)	24.53 (1.01)	29.89 (0.99)	5.37 (2.67,8.07)	5.67 (2.56,8.78)
API	12.77 (0.59)	13.24 (0.64)	0.47 (-0.93,1.88)	11.65 (0.75)	16.21 (0.76)	4.56 (2.42,6.70)	4.09 (1.55,6.63)
Other/Unknown	19.37 (0.83)	18.66 (0.82)	-0.71 (-2.31,0.89)	18.75 (1.10)	23.88 (1.06)	5.14 (2.09,8.18)	5.85 (2.32,9.37)
16-17 years							
NH White	16.72 (0.64)	15.63 (0.60)	-1.09 (-2.26,0.08)	16.12 (0.72)	18.43 (0.72)	2.31 (0.54,4.09)	3.40 (1.26,5.54)

Hispanic	23.38 (0.49)	23.34 (0.48)	-0.04 (-0.89,0.82)	23.43 (0.57)	25.87 (0.54)	2.45 (1.14,3.76)	2.49 (0.93,4.04)
NH Black	22.15 (1.11)	22.67 (1.08)	0.52 (-1.28,2.32)	23.16 (1.35)	25.52 (1.22)	2.36 (-0.65,5.37)	1.84 (-1.69,5.37)
API	12.34 (0.88)	12.01 (0.85)	-0.33 (-2.01,1.35)	11.40 (1.24)	12.82 (1.13)	1.42 (-2.16,5.00)	1.75 (-2.07,5.56)
Other/Unknown	18.34 (1.26)	17.84 (1.18)	-0.50 (-3.08,2.08)	19.59 (1.60)	20.14 (1.34)	0.55 (-3.38,4.48)	1.05 (-3.47,5.57)
State-subsidized health care							
5-11 years							
No	15.96 (0.21)	17.37 (0.22)	1.41 (0.94,1.89)	17.81 (0.29)	24.86 (0.29)	7.05 (6.23,7.87)	5.64 (4.70,6.57)
Yes	18.21 (0.34)	19.72 (0.35)	1.50 (0.81,2.20)	19.93 (0.43)	28.07 (0.45)	8.13 (6.92,9.34)	6.63 (5.23,8.02)
12-15 years							
No	18.29 (0.28)	18.19 (0.29)	-0.10 (-0.69,0.50)	17.50 (0.36)	22.22 (0.35)	4.72 (3.72,5.73)	4.82 (3.64,6.00)
Yes	19.73 (0.45)	19.99 (0.46)	0.26 (-0.60,1.13)	19.41 (0.57)	25.04 (0.57)	5.64 (4.13,7.14)	5.37 (3.62,7.13)
16-17 years							
No	17.07 (0.41)	16.87 (0.40)	-0.20 (-1.00,0.60)	17.56 (0.56)	18.70 (0.49)	1.14 (-0.32,2.61)	1.34 (-0.30,2.98)
Yes	19.26 (0.69)	18.72 (0.66)	-0.54 (-1.77,0.69)	18.70 (0.84)	21.24 (0.78)	2.54 (0.45,4.62)	3.08 (0.73,5.42)
Neighborhood parks							
5-11 years							
No park	17.61 (0.30)	18.86 (0.31)	1.26 (0.62,1.89)	19.32 (0.39)	27.03 (0.40)	7.70 (6.61,8.79)	6.45 (5.19,7.70)
1 park	17.42 (0.29)	18.71 (0.30)	1.29 (0.67,1.92)	18.69 (0.38)	26.93 (0.40)	8.24 (7.14,9.33)	6.94 (5.69,8.19)
2 or more parks	16.17 (0.32)	17.97 (0.33)	1.80 (1.11,2.50)	18.53 (0.43)	25.33 (0.43)	6.80 (5.61,7.99)	5.00 (3.63,6.36)

12-15 years

No park	20.02 (0.41)	19.90 (0.41)	-0.12 (-0.92,0.68)	19.19 (0.51)	24.59 (0.51)	5.40 (4.02,6.78)	5.52 (3.91,7.13)
1 park	19.63 (0.40)	19.67 (0.41)	0.04 (-0.76,0.83)	19.13 (0.51)	24.41 (0.50)	5.29 (3.91,6.66)	5.25 (3.65,6.85)
2 or more parks	17.43 (0.41)	17.71 (0.42)	0.28 (-0.55,1.11)	17.04 (0.51)	21.86 (0.52)	4.82 (3.43,6.21)	4.54 (2.90,6.18)

16-17 years

No park	18.40 (0.59)	17.86 (0.57)	-0.54 (-1.61,0.52)	18.06 (0.73)	20.43 (0.68)	2.37 (0.55,4.19)	2.92 (0.86,4.97)
1 park	18.61 (0.60)	18.25 (0.56)	-0.36 (-1.49,0.77)	18.24 (0.73)	20.49 (0.67)	2.25 (0.41,4.10)	2.62 (0.47,4.76)
2 or more parks	17.40 (0.62)	17.22 (0.60)	-0.18 (-1.36,0.99)	18.07 (0.83)	18.91 (0.72)	0.84 (-1.27,2.96)	1.02 (-1.33,3.38)

Total BMI measures included in the models were 425,855 from March 2019-January 2020 (2.65 BMI measures/youth) compared to 241699 from March 2020-January 2021 (1.51 BMI measures/youth). After initial decrease, in-person well-child visits were back to 84% of prepandemic visits by June 2020.

All models are adjusted for race/ethnicity (Hispanic, non-Hispanic black, Asian or Pacific Islander, other, reference: White), state-subsidized health care (reference: none), parks (no parks, 2 or more parks, reference: 1 park), neighborhood education, neighborhood income, and 3-way interactions pandemic x time x race, pandemic x time x state-subsidized health care, pandemic x time x parks. Estimates are shown for the reference group.

Interaction terms: Pandemic x time x race/ethnicity P =0.49 for 5-11 years, P =0.86 for 12-15 years, P = 0.58 for 16-17 years, Pandemic x time x State-subsidized insurance P =0.63 for 5-11 years, P = 0.94 for 12-15 years, P = 0.12 for 16-17 years, Pandemic x time x parks P =0.06 for 5-11 years, P = 0.89 for 12-15 years, P = 0.58 for 16-17 years.

Figure Legends:

Figure 1: Study flow chart


Figure 2: Changes in the distance from median BMI-for-age before and during the COVID-19 lockdown in white, Hispanic or Black overweight boys ages 5-11 years who live in neighborhoods with 2 or more parks and without state subsidized health insurance (Figure 2A) compared to those who live in neighborhoods without parks and have state subsidized health insurance (Figure 2B)

REFERENCES

1. Ghosh R, Dubey MJ, Chatterjee S, Dubey S. Impact of COVID -19 on children: special focus on the psychosocial aspect. *Minerva pediatrica*. 2020;72(3):226-235.
2. Education Commission of the States. COVID Response. Available from: <https://www.fina.org/rules-guidance/key-topics/covid-19/shelter-in-place> Accessed 4/6/2021. 2020.
3. Dunn CG, Kenney E, Fleischhacker SE, Bleich SN. Feeding Low-Income Children during the Covid-19 Pandemic. *N Engl J Med*. 2020;382(18):e40.
4. Gundersen C, Mahatmya D, Garasky S, Lohman B. Linking psychosocial stressors and childhood obesity. *Obes Rev*. 2011;12(5):e54-63.
5. Woolford SJ, Sidell M, Li X, et al. Changes in Body Mass Index Among Children and Adolescents During the COVID-19 Pandemic. *JAMA*. 2021;326(14):1434-1436.
6. Stierman B, Ogden CL, Yanovski JA, Martin CB, Sarafrazi N, Hales CM. Changes in adiposity among children and adolescents in the United States, 1999-2006 to 2011-2018. *Am J Clin Nutr*. 2021.
7. Hales CM, Carroll MD, Fryar CD, Ogden CL. Prevalence of Obesity and Severe Obesity Among Adults: United States, 2017-2018. *NCHS Data Brief*. 2020(360):1-8.
8. Division of Nutrition PA, and Obesity, National Center for Chronic Disease Prevention and Health Promotion,. Childhood Obesity Facts: Prevalence of Childhood Obesity in the United States. <https://www.cdc.gov/obesity/data/childhood.html>, accessed 04/06/2021. 2021.
9. Centers for Disease Control and Prevention. Risk for COVID-19 Infection, Hospitalization, and Death By Race/Ethnicity, accessed 04/02/2021. <https://www.cdc.gov/coronavirus/2019-ncov/covid-data/investigations-discovery/hospitalization-death-by-race-ethnicity.html>. 2021.
10. Tung EL, Peek ME, Rivas MA, Yang JP, Volerman A. Association Of Neighborhood Disadvantage With Racial Disparities In COVID-19 Positivity In Chicago. *Health Aff (Millwood)*. 2021;40(11):1784-1791.
11. Koebnick C, Langer-Gould AM, Gould MK, et al. Sociodemographic characteristics of members of a large, integrated health care system: comparison with US Census Bureau data. *Perm J*. 2012;16(3):37-41.
12. Koebnick C, Coleman KJ, Black MH, et al. Cohort profile: the KPSC Children's Health Study, a population-based study of 920 000 children and adolescents in southern California. *Int J Epidemiol*. 2012;41(3):627-633.
13. Feinstein JA, Russell S, DeWitt PE, Feudtner C, Dai D, Bennett TD. R Package for Pediatric Complex Chronic Condition Classification. *JAMA pediatrics*. 2018;172(6):596-598.
14. Feudtner C, Feinstein JA, Zhong W, Hall M, Dai D. Pediatric complex chronic conditions classification system version 2: updated for ICD-10 and complex medical technology dependence and transplantation. *BMC Pediatr*. 2014;14:199.
15. Kuczmarski RJ, Ogden CL, Guo SS, et al. 2000 CDC Growth Charts for the United States: methods and development. *Vital Health Stat 11*. 2002;11(246):1-190.
16. Ryder JR, Kelly AS, Freedman DS. Metrics matter: Toward consensus reporting of BMI and weight-related outcomes in pediatric obesity clinical trials. *Obesity (Silver Spring)*. 2022;30(3):571-572.
17. Freedman DS, Butte NF, Taveras EM, et al. BMI z-Scores are a poor indicator of adiposity among 2- to 19-year-olds with very high BMIs, NHANES 1999-2000 to 2013-2014. *Obesity (Silver Spring)*. 2017;25(4):739-746.
18. Freedman DS, Woo JG, Ogden CL, Xu JH, Cole TJ. Distance and percentage distance from median BMI as alternatives to BMI z score. *Br J Nutr*. 2020;124(5):493-500.
19. Freedman DS, Berenson GS. Tracking of BMI z Scores for Severe Obesity. *Pediatrics*. 2017;140(3).

20. Claritas LLC. Claritas Demographic Update 2021 Methodology. Accessed at <https://environicanalytics.com/docs/default-source/us---data-product-support-documents/claritas-demographic-update-methodology-eapdf> on 04/06/2021. 2021.
21. Andrade C. Mean Difference, Standardized Mean Difference (SMD), and Their Use in Meta-Analysis: As Simple as It Gets. *The Journal of clinical psychiatry*. 2020;81(5).
22. Beech BM, Ford C, Thorpe RJ, Jr., Bruce MA, Norris KC. Poverty, Racism, and the Public Health Crisis in America. *Frontiers in public health*. 2021;9:699049.
23. Acosta AM, Garg S, Pham H, et al. Racial and Ethnic Disparities in Rates of COVID-19-Associated Hospitalization, Intensive Care Unit Admission, and In-Hospital Death in the United States From March 2020 to February 2021. *JAMA Netw Open*. 2021;4(10):e2130479.
24. Magesh S, John D, Li WT, et al. Disparities in COVID-19 Outcomes by Race, Ethnicity, and Socioeconomic Status: A Systematic-Review and Meta-analysis. *JAMA Netw Open*. 2021;4(11):e2134147.
25. Azar KMJ, Shen Z, Romanelli RJ, et al. Disparities In Outcomes Among COVID-19 Patients In A Large Health Care System In California. *Health Aff (Millwood)*. 2020;39(7):1253-1262.
26. Escobar GJ, Adams AS, Liu VX, et al. Racial Disparities in COVID-19 Testing and Outcomes : Retrospective Cohort Study in an Integrated Health System. *Ann Intern Med*. 2021;174(6):786-793.
27. Nau C, Bruxvoort K, Navarro RA, et al. COVID-19 Inequities Across Multiple Racial and Ethnic Groups: Results From an Integrated Health Care Organization. *Ann Intern Med*. 2021;174(8):1183-1186.
28. Browne NT, Snethen JA, Greenberg CS, et al. When Pandemics Collide: The Impact of COVID-19 on Childhood Obesity. *Journal of pediatric nursing*. 2021;56:90-98.
29. Lange SJ, Kompaniyets L, Freedman DS, et al. Longitudinal Trends in Body Mass Index Before and During the COVID-19 Pandemic Among Persons Aged 2-19 Years - United States, 2018-2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(37):1278-1283.
30. Vogel M, Geserick M, Gausche R, et al. Age- and weight group-specific weight gain patterns in children and adolescents during the 15 years before and during the COVID-19 pandemic. *Int J Obes (Lond)*. 2021:1-9.
31. Brooks CG, Spencer JR, Sprafka JM, et al. Pediatric BMI changes during COVID-19 pandemic: An electronic health record-based retrospective cohort study. *EClinicalMedicine*. 2021;38:101026.
32. Wu AJ, Aris IM, Hivert MF, et al. Association of Changes in Obesity Prevalence With the COVID-19 Pandemic in Youth in Massachusetts. *JAMA pediatrics*. 2021.
33. Belanger MJ, Hill MA, Angelidi AM, Dalamaga M, Sowers JR, Mantzoros CS. Covid-19 and Disparities in Nutrition and Obesity. *N Engl J Med*. 2020;383(11):e69.
34. Ogden CL, Fryar CD, Martin CB, et al. Trends in Obesity Prevalence by Race and Hispanic Origin-1999-2000 to 2017-2018. *JAMA*. 2020;324(12):1208-1210.
35. Child I, Loza A, Doolittle B. Collateral Damage from the COVID-19 Pandemic: Pediatric Obesity. *Population health management*. 2022.
36. Morgan Hughey S, Kaczynski AT, Child S, Moore JB, Porter D, Hibbert J. Green and lean: Is neighborhood park and playground availability associated with youth obesity? Variations by gender, socioeconomic status, and race/ethnicity. *Preventive medicine*. 2017;95 Suppl:S101-s108.
37. McCarthy SM, Hughey SM, Kaczynski AT. Examining Sociodemographic Differences in Playground Availability and Quality and Associations with Childhood Obesity. *Childhood obesity*. 2017;13(4):324-331.
38. White MJ, McClure E, Killeen J, et al. Changes in the Recreational Built Environment and Youth Body Mass Index. *Acad Pediatr*. 2021;21(1):76-83.
39. D'Agostino EM, Patel HH, Ahmed Z, et al. Impact of change in neighborhood racial/ethnic segregation on cardiovascular health in minority youth attending a park-based afterschool program. *Soc Sci Med*. 2018;205:116-129.

40. Nardone A, Rudolph KE, Morello-Frosch R, Casey JA. Redlines and Greenspace: The Relationship between Historical Redlining and 2010 Greenspace across the United States. *Environ Health Perspect*. 2021;129(1):17006.
41. Nobari TZ, Whaley SE, Prelip ML, Crespi CM, Wang MC. Trends in Socioeconomic Disparities in Obesity Prevalence among Low-Income Children Aged 2-4 Years in Los Angeles County, 2003-2014. *Childhood obesity*. 2018.
42. Hingorani A. LA County park equity groups fight to turn park-poor community “red to green.” (Advancement Project 1.22.19.). <https://www.advancementprojectca.org/blog/la-county-park-equity-groups-fight-to-turn-park-poor-communities-red-to-green>. Accessed 11/29/2021.
43. Bailey ZD, Feldman JM, Bassett MT. How Structural Racism Works — Racist Policies as a Root Cause of U.S. Racial Health Inequities. *N Engl J Med*. 2020;384(8):768-773.
44. Bleich SN, Ard JD. COVID-19, Obesity, and Structural Racism: Understanding the Past and Identifying Solutions for the Future. *Cell Metab*. 2021;33(2):234-241.
45. Ames JL, Ferrara A, Avalos LA, et al. COVID-19 prevalence, symptoms, and sociodemographic disparities in infection among insured pregnant women in Northern California. *PLoS One*. 2021;16(9):e0256891.
46. Center on Budget and Policy Priorities. Tracking the COVID-19 economy’s effects on food, housing, and employment hardships. <https://www.cbpp.org/research/poverty-and-inequality/tracking-the-covid-19-economys-effects-on-food-housing-and>. Accessed on 11/10/2021.
47. Jansen E, Thapaliya G, Aghababian A, Sadler J, Smith K, Carnell S. Parental stress, food parenting practices and child snack intake during the COVID-19 pandemic. *Appetite*. 2021;161:105119.
48. Hardcastle SJ, Caraher M. The role of Foodbanks in the context of food insecurity: Experiences and eating behaviours amongst users. *Appetite*. 2021:105208.
49. Stinson EJ, Votruba SB, Venti C, Perez M, Krakoff J, Gluck ME. Food Insecurity is Associated with Maladaptive Eating Behaviors and Objectively Measured Overeating. *Obesity (Silver Spring)*. 2018;26(12):1841-1848.
50. von Hippel PT, Workman J. From Kindergarten Through Second Grade, U.S. Children's Obesity Prevalence Grows Only During Summer Vacations. *Obesity (Silver Spring)*. 2016;24(11):2296-2300.
51. Wang YC, Vine S, Hsiao A, Rundle A, Goldsmith J. Weight-related behaviors when children are in school versus on summer breaks: does income matter? *J Sch Health*. 2015;85(7):458-466.



OBY_23645_Koebnick-CovidDisparitiesObesity-Figure 1 Revision.tif

Figure 2A: “Advantaged” youth (Male, overweight, 2+ parks, no Medicaid*)





Figure 2B: “Disadvantaged” youth (Male, overweight, no parks, Medicaid*)



Change in distance from the median BMI-for-age (kg/m²) (95% CI)

	prepandemic ($\Delta 1$, 95% CI)	pandemic lockdown ($\Delta 2$, 95% CI)	$\Delta 2 - \Delta 1$ (95% CI)
Figure 2A			
Race/ethnicity			
Non-Hispanic White	0.07 (0.00, 0.14)	0.91 (0.80, 1.02)	0.84 (0.71, 0.97)
Non-Hispanic Black	0.37 (0.24, 0.50)	1.76 (1.55, 1.96)	1.39 (1.15, 1.63)
Hispanic	0.26 (0.19, 0.33)	1.67 (1.56, 1.78)	1.41 (1.28, 1.53)
Asian/Pacific Islanders	0.28 (0.17, 0.38)	1.31 (1.15, 1.48)	1.04 (0.84, 1.23)
Other/Unknown	0.28 (0.15, 0.40)	1.03 (0.83, 1.23)	0.75 (0.52, 0.98)
Figure 2B			
Race/ethnicity			
Non-Hispanic White	0.13 (0.04, 0.22)	1.47 (1.32, 1.62)	1.34 (1.17, 1.51)
Non-Hispanic Black	0.43 (0.30, 0.56)	2.32 (2.11, 2.52)	1.89 (1.65, 2.12)
Hispanic	0.32 (0.25, 0.39)	2.23 (2.11, 2.34)	1.91 (1.78, 2.04)
Asian/Pacific Islanders	0.33 (0.22, 0.45)	1.87 (1.68, 2.06)	1.54 (1.31, 1.76)
Other/Unknown	0.34 (0.20, 0.47)	1.59 (1.37, 1.81)	1.25 (1.00, 1.51)