

MRA use in renal dialysis : will evidence from prospective randomized trials confirm the results from “ real world “ evidence?

1. Université de Lorraine, Centre d'Investigations Cliniques- Plurithématique 14-33, Inserm U1116, CHRU Nancy, and F-CRIN INI-CRCT (Cardiovascular and Renal Clinical Trialists), Nancy, France
2. Medicine and Nephrology-dialysis departments, Princess Grace Hospital, and Monaco Private Hemodialysis Centre, Monaco, Monaco
3. University of Michigan School of Medicine, Ann Arbor, Michigan, USA.

Keywords: mineralocorticoid receptor antagonists, hyperkalemia, heart failure, endstage renal disease

In patients with end-stage renal disease (ESRD) on chronic renal dialysis the cardiovascular (CV) mortality rate is 10 to 20 times higher than in the general population and among adults younger than 45 years of age the mortality rate is approximately 100 times higher than that in the general population. Hence, treatments to improve survival, more specifically CV outcomes, in this population are urgently needed (1,2).

Activation of mineralocorticoid receptors (MR) in patients with chronic renal disease (CKD) leads to adverse outcomes, especially hospitalization for heart failure (HF), ventricular arrhythmias, sudden cardiac death as well as progression of CKD to ESRD (2,3) through several mechanisms including: inflammation; oxidative stress; endothelial dysfunction; vascular fibrosis, stiffening and calcification (3-5) all of which are relevant in patients undergoing renal dialysis (2). MR antagonists (MRAs) are currently approved and or are recommended in European and international guidelines in patients with resistant hypertension, diabetic nephropathy and in HF with a reduced ejection fraction (HFrEF), where they are one of the “four pillars” along with angiotensin converting enzyme inhibitors (ACEis) /angiotensin receptor blockers (ARBs)/sacubitril valsartan; beta blockers; and SGLT2 inhibitors. MRAs are therefore prime candidates to be assessed in prospective randomized clinical trials in patients undergoing renal dialysis. However, in view of the risk of developing hyperkalaemia, especially in those on an ACEI or ARB, there has been a reluctance to initiate therapy with a MRA in patients with ESRD undergoing renal dialysis. The concomitant use of one of the new potassium binders, such as patiromer, as demonstrated in clinical trials (6,7) along with a MRA may help to reduce the risk of hyperkalaemia but does not eliminate it. Despite the lack of evidence regarding the effectiveness and safety of MRAs in patients undergoing renal dialysis and the fact that they are currently contraindicated some clinicians have initiated their off-label use. In this issue of the journal Shu-Han Lin et al evaluated the effectiveness and safety of MRAs on CV outcomes in patients with HF undergoing maintenance renal dialysis in a real-world setting followed for slightly over 2 years (8). They performed a retrospective analysis within the Taiwan National Health Insurance Research Database (NHIRD), using propensity score matching during the years 2001 to 2013. They concluded that in patients undergoing renal dialysis who are diagnosed with HF the use of MRAs was associated with a lower risk of all-cause mortality (Hazard ratio (HR) 0.88, 95% CI 0.83–0.94) and CV death (HR 0.88, 95% CI 0.80–0.95) but not hospitalizations for HF, myocardial infarction or stroke in comparison to those not on a MRA. The benefits of MRA administration appeared to be greater in those patients undergoing hemodialysis as compared to those undergoing peritoneal dialysis. However the number of patients undergoing peritoneal dialysis was rather small and therefore it is difficult to reach any definitive conclusion as to the relative

efficacy of MRAs in patients undergoing hemo vs peritoneal dialysis. The benefits of MRAs in these patients with ESRD undergoing renal dialysis was attributed to their effect on reducing ventricular arrhythmias and sudden cardiac death. However, while the use of MRAs was shown to be beneficial in these high-risk patients the risk of new-onset hyperkalemia, defined as a serum potassium >6 meq/l, was significantly higher in the MRA group (HR 1.34, 95% CI 1.17–1.54) compared to those not on a MRA, which could limit the clinical application of MRAS in patients with ESRD.

While this provocative study is supported by robust statistical methodology there are however inherent limitations to such retrospective analyses, even with propensity matching. As pointed out by the authors unmeasured confounders could have a profound effect on the physicians' choice to administer a MRA. Therefore, as acknowledged by the authors, further prospective randomized evidence is warranted in patients with ESRD, both with and without concomitant HF, before we can confidently recommend the use of MRAs in patients undergoing renal dialysis. Interestingly, a prior meta-analysis (9) gathering data from 14 randomized trials with steroidal MRAs in 1309 patients with ESRD also revealed a significant reduction in both CV mortality (RR, 0.41; 95% CI, 0.24 to 0.70; $P=0.001$; $N=975$) and all-cause mortality (RR, 0.44; 95% CI, 0.30 to 0.66; $P<0.001$, $N=1246$) in patients receiving MRAs compared to the control group. Similar to the analysis by Lin et al there was no evidence of a reduction in stroke or non-fatal cardiovascular events. However, in contrast to the analysis by Lin et al (8), the use of a MRA was not associated with an increased risk of hyperkalemia (RR, 1.12; 95% CI, 0.91 to 1.36; $P=0.29$). Moreover, no obvious subgroup difference was observed ($I^2=0\%$; $P=0.48$).” In the discussion, the authors point out that their results were heavily influenced by just 2 studies of modest size and therefore they point out the need for further evidence regarding the effectiveness of MRAs in patients with ESRD. The authors of this meta analysis also acknowledge that their result on hyperkalemia “contradicts that of a previous meta-analysis(10) that described a significantly higher risk of hyperkalaemia in MRA-treated dialysis patients. However, after incorporating more recent trials, the risk of hyperkalaemia was negated”. These two trials (Spin-D and MiREnDa trials) were actually phase 2 trials (2). Importantly, the largest trial considered in the meta analysis (Matsumoto et al(11)) was an open label Japanese only trial of 309 patients (corresponding to half of the initial recruitment target)(12)). Thus, despite the retrospective evidence presented by Lin et al (8) and the meta analysis of randomized trials by Chen et al (9) more evidence based on large prospective international double-blind placebo controlled CV outcome trials is warranted. Thus, we must await the results of prospective randomized trials of the steroidal MRA spironolactone in patients with ESRD undergoing renal dialysis. The ALCHEMIST trial (NCT01848639) which randomized patients

to spironolactone up to 25 mg/day or placebo was completed at the end of 2022 and enrolled its scheduled 825 high risk CV haemodialysis patients. In the ALCHEMIST trial hyperkalaemia, defined by a serum potassium > 5.5 mmol/l within 2 weeks is an exclusion criteria. The ACHIEVE trial (NCT03020303) plans to randomize 2750 patients, likely low CV risk patients on hemo -or peritoneal dialysis to 25mg/day of spironolactone or placebo and excludes patients with hyperkalaemia, defined as a serum potassium >5.8 mmol/l within 6 weeks, is scheduled for completion of recruitment in December 2024 (according to clinicaltrial.gov, accessed January 14th, 2023). It should however be emphasized that these 2 trials are randomising patients with a demonstrated tolerance to spironolactone (25 mg qod in ALCHEMIST: drop out if HK ≥ 5.5 mmol/L twice) or 25 mg/d in ACHIEVE: randomisation if no HK > 6 mmol/L and 80 % adherence) in terms of hyperkalemia during a run-in phase of 4 and 8 weeks, respectively. Therefore, a number of patients intolerant to a steroidal MRA due to hyperkalaemia will be excluded from these trials. Furthermore, should the incidence of hyperkalemia be increased in patients randomized to spironolactone in these trials this could limit its long-term use despite a significant reduction in CV mortality or morbidity, since hyperkalemia is highly prevalent in patients undergoing chronic haemodialysis(13) and clinicians may be reluctant to initiate spironolactone serially monitor serum potassium and adjust its dose accordingly . Therefore, the combined use of potassium mitigating strategies, with either potassium binders (used by 9.5 % of the patients included in the analysis by Lin et al in Taiwan(8)) or low dialysis potassium concentrations might need to be considered(13). Alternatively, a selective non-steroidal MRA, such as Finerenone (14), or an aldosterone synthase inhibitor (ASI) such as Baxdrostat (13), both of which appear to be less prone to induce hyperkalaemia than spironolactone, might provide greater safety with similar or possibly even greater efficacy than spironolactone. Further data will however be required before considering larger scale phase 3 registration trials to assess the safety, tolerability, and to provide evidence of efficacy of these new non-steroidal MRAs and ASIs in patients with ESRD undergoing renal dialysis. For example, the FIDELIO and FIGARO phase III trials of finerenone (15) in patients with diabetic nephropathy and an eGFR above 25 ml/min/1.73m² demonstrated nephroprotection along with CV prevention, with a relatively low incidence and easily manageable burden of hyperkalemia. The efficacy and safety of Finerenone however has not as yet been evaluated in patients with ESRD undergoing renal dialysis.

In conclusion, real-world evidence from the off-label use of MRAs in patients with HF undergoing renal dialysis in Taiwan (8) and a meta analysis of relatively small randomized trials (9) provide reassuring but still preliminary evidence regarding the benefits and safety

of MRAs in patients undergoing renal dialysis. Before widespread clinical application of MRAs to patients with ESRD undergoing renal dialysis we should await the results of the recently completed ALCHEMIST and still ongoing ACHIEVE trials. Should the results of these trials show that spironolactone is effective and safe in patients undergoing renal dialysis, they could result in a major change in clinical practice. Spironolactone is universally available, generic and therefore relatively inexpensive. Therefore, proof of its efficacy and safety in high-risk patients with ESRD undergoing renal dialysis could have a profound effect on CV morbidity, mortality, quality of life and health care costs. Let's stay tuned !

Disclosures:

PR: reports consulting for Idorsia, G3P, honoraria from AstraZeneca, Bayer, Boehringer-Ingelheim, Cincor, CVRx, Fresenius, KBP biosciences, Novartis, NovoNordisk, Relypsa, Servier, and Vifor Fresenius Medical Care Renal Pharma; and travel grants from AstraZeneca, Bayer, CVRx, Novartis, and Vifor Fresenius Medical Care Renal Pharma; Cofounder: CardioRenal

BP: consultant : Bayer , Astra- Zeneca, Boehringer Ingelheim, Merck, Lexicon,, KBP biosciences*, Vifor*, sarfez*, scpharmaceuticals*, sqinnovations*, G3pharmaceuticls^, proton intel*, cereno scientific*, brainstorm medical*

*=stock options

Us patent 9931422 - site specific delivery of eplerenone to the myocardium . us patent pending63/045,783 Histone acetylation modulating agents for the protection and treatment of organ damage .

1. Rossignol P, Agarwal R, Canaud B et al. Cardiovascular outcome trials in patients with chronic kidney disease: challenges associated with selection of patients and endpoints. *Eur Heart J* 2019;40:880-886.
2. Rossignol P, Frimat L, Zannad F. The safety of mineralocorticoid antagonists in maintenance hemodialysis patients: two steps forward. *Kidney Int* 2019;95:747-749.
3. Zannad F, Rossignol P. Cardiorenal Syndrome Revisited. *Circulation* 2018;138:929-944.

4. Ortiz A, Ferro CJ, Balafa O et al. Mineralocorticoid receptor antagonists for nephroprotection and cardioprotection in patients with diabetes mellitus and chronic kidney disease. *Nephrol Dial Transplant* 2021.
5. Rossignol P, Hernandez AF, Solomon SD, Zannad F. Heart failure drug treatment. *Lancet* 2019;393:1034-1044.
6. Agarwal R, Rossignol P, Romero A et al. Patiromer versus placebo to enable spironolactone use in patients with resistant hypertension and chronic kidney disease (AMBER): a phase 2, randomised, double-blind, placebo-controlled trial. *Lancet* 2019;394:1540-1550.
7. Butler J, Anker SD, Siddiqi TJ et al. Patiromer for the management of hyperkalaemia in patients receiving renin-angiotensin-aldosterone system inhibitors for heart failure: design and rationale of the DIAMOND trial. *Eur J Heart Fail* 2022;24:230-238.
8. Lin DS, Lin FJ, Lin YS, Lee JK, Lin YH. The effects of mineralocorticoid receptor antagonists on cardiovascular outcomes in patients with end-stage renal disease and heart failure. *Eur J Heart Fail* 2022.
9. Chen KT, Kang YN, Lin YC et al. Efficacy and Safety of Mineralocorticoid Receptor Antagonists in Kidney Failure Patients Treated with Dialysis: A Systematic Review and Meta-Analysis. *Clin J Am Soc Nephrol* 2021;16:916-925.
10. Quach K, Ltvyn L, Baigent C et al. The Safety and Efficacy of Mineralocorticoid Receptor Antagonists in Patients Who Require Dialysis: A Systematic Review and Meta-analysis. *Am J Kidney Dis* 2016;68:591-598.
11. Matsumoto Y, Mori Y, Kageyama S et al. Spironolactone reduces cardiovascular and cerebrovascular morbidity and mortality in hemodialysis patients. *J Am Coll Cardiol* 2014;63:528-36.
12. Pitt B, Rossignol P. Mineralocorticoid receptor antagonists in patients with end-stage renal disease on chronic hemodialysis. *J Am Coll Cardiol* 2014;63:537-8.
13. Rossignol P, Lamiral Z, Frimat L et al. Hyperkalaemia prevalence, recurrence and management in chronic haemodialysis: a prospective multicentre French regional registry 2-year survey. *Nephrol Dial Transplant* 2017;32:2112-2118.
14. Bakris G, Yang YF, Pitt B. Mineralocorticoid Receptor Antagonists for Hypertension Management in Advanced Chronic Kidney Disease: BLOCK-CKD Trial. *Hypertension* 2020;76:144-149.
15. Agarwal R, Filippatos G, Pitt B et al. Cardiovascular and kidney outcomes with finerenone in patients with type 2 diabetes and chronic kidney disease: the FIDELITY pooled analysis. *Eur Heart J* 2021.