







REVIEW

The peri-implant mucosa color: A systematic appraisal of methods for its assessment and clinical significance

Leonardo Mancini DDS^{1,2,3,4}  | Shayan Barootchi DMD, MS^{3,5}  |
Daniel S. Thoma DMD^{2,6}  | Ronald E. Jung DMD, PhD² |
German O. Gallucci DMD, PhD⁷  | Hom-Lay Wang DDS, MS, PhD⁵  |
Lorenzo Tavelli DDS, MS^{3,4,5} 

¹Department of Life, Health and Environmental Sciences, University of L'Aquila, L'Aquila, Italy

²Clinic of Reconstructive Dentistry, University of Zurich, Zurich, Switzerland

³Center for Clinical Research and Evidence Synthesis in Oral Tissue Regeneration (CRITERION), Boston, Massachusetts, USA

⁴Department of Oral Medicine, Infection, and Immunity, Division of Periodontology, Harvard School of Dental Medicine, Boston, Massachusetts, USA

⁵Department of Periodontics & Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, Michigan, USA

⁶Department of Periodontology, Research Institute for Periodontal Regeneration, College of Dentistry, Yonsei University, Seoul, South Korea

⁷Department of Restorative Dentistry and Biomaterial Science, Harvard School of Dental Medicine, Boston, Massachusetts, USA

Correspondence

Lorenzo Tavelli, Department of Oral Medicine, Infection, and Immunity, Division of Periodontology, Harvard School of Dental Medicine, 188 Longwood Avenue, Boston, MA 02115 USA.
Email: lorenzo_tavelli@hsdm.harvard.edu

Abstract

Background: Peri-implant mucosa color (PMC) seems to be one of the main parameters affecting the esthetic outcome of implant therapy. However, more emphasis should be given to its assessment and reporting.

Purpose: To describe the available evidence on methods to assess and report the color of the peri-implant mucosa (PMC) and the respective clinical relevance.

Material and Methods: A comprehensive electronic and manual search was performed to identify clinical studies reporting on PMC.

Results: A total of 121 studies were included. PMC was evaluated at the time of the follow-up visit (chairside) in 45.5% studies. PMC assessment was performed qualitatively, by comparing PMC with adjacent and/or contralateral gingiva (78.6%) or quantitatively, using spectrophotometry (20.7%) or a software on clinical photographs (0.8%). The most performed method to assess PMC was through esthetic indices (76.9%), either at the time of the follow-up visit (chairside) or at later time point using photographs. Quantitative reporting of PMC included averages of points from esthetic indices or color differences to natural gingiva expressed with the CIELAB color system. PMC assessment allowed describing color discrepancies compared to natural gingiva, evaluating color changes over time, and comparing the outcomes of different treatment modalities. PMC assessment through spectrophotometry was additionally utilized to assess the role of mucosal thickness (MT) on PMC.

Conclusions: Various methods for PMC assessment and reporting were described, including visual assessment, mainly through esthetic indices, and spectrophotometry. PMC evaluation has allowed to demonstrate the factors affecting the color of the peri-implant soft tissue, such as the type of abutment/restoration, MT, and soft tissue augmentation.

KEYWORDS

color, dental implants, esthetics, soft tissue, spectrophotometry, systematic review

SUMMARY BOX**What is known**

- Peri-implant mucosa color (PMC) is one of the main parameters affecting the esthetic outcome of implant therapy.
- Several methods for assessing and reporting PMC have been described in the literature.

What this study adds

- The present review describes the available evidence on methods to assess and report the color of the peri-implant mucosa and the respective clinical relevance.
- Objective methods for assessing PMC mainly involve the use of spectrophotometry or professional esthetic indices.
- The type and color of the abutment, the type of restoration, soft tissue augmentation and peri-implant mucosal thickness, affect PMC.
- The use of professional indices allows to assess the color match of the peri-implant mucosa with the adjacent soft tissue and/or gingiva of the contralateral tooth and to evaluate the stability of PMC over time.

1 | INTRODUCTION

Implant therapy has had a significant evolution over the years.¹ Advances in research, implant systems, techniques, and biomaterials have contributed to increase the predictability and outcomes of dental implants.¹⁻³ From a patient's perspective, one of the most crucial parameters in implant therapy is the esthetics, with the appearance of the implant-supported crown and the peri-implant soft tissue that needs to mimic the ones of the natural dentition.^{4,5} The esthetic outcomes of implant therapy depend on several factors. Severe bony and soft tissue dehiscences prior to implant placement are considered challenging clinical conditions for obtaining an ideal esthetic result.⁶ It has been demonstrated that the esthetics of dental implants is often suboptimal, with an average incidence of peri-implant soft tissue dehiscence (PSTD)—defined as the apical shift of the peri-implant soft tissue margin compared to the cemento-enamel junction (CEJ) of the homologous contralateral tooth—of 54.2% and 56.8% on a patient and implant level, respectively.⁷ Limited mucosal thickness (MT), reduced/lack keratinized mucosa and buccal bone dehiscence were among the factors associated with this condition.⁷ It has also been suggested that immediate implant therapy may increase the risk of esthetic complications.^{8,9}

On the other hand, the esthetics of dental implants depends also on the characteristics of the peri-implant mucosa in terms of keratinization, thickness, texture, and color.¹⁰⁻¹² A study by Bonino and coworkers demonstrated a statistically significant association between patient-reported esthetic satisfaction following implant therapy and the presence of a band of keratinized mucosa.¹³ The thickness of peri-implant mucosa has also been found to play a key role on the implant esthetic outcomes, with thin tissue often displaying grayish discoloration due to the implant components underneath.^{10,14,15}

Although peri-implant mucosa color (PMC) seems to be one of the main parameters affecting the esthetic outcome of implant therapy, little emphasis has been given on its assessment. PMC has often

been reported as “similar” or “different” in comparison with the adjacent or contralateral natural gingiva and incorporated in esthetic indices with different scores.¹⁶⁻¹⁸ Moreover, several authors have advocated the use of spectrophotometry to quantify color differences between peri-implant mucosa and the gingival of the adjacent or contralateral tooth,^{15,19-21} showing high accuracy and reproducibility for this technology.²¹⁻²⁴

In this scenario, guidelines for assessing and reporting PMC in clinical research are missing.

Therefore, the aim of the present manuscript was to describe the available evidence on methods utilized for assessing and reporting PMC and the respective clinical relevance.

2 | MATERIAL AND METHODS

2.1 | Protocol registration and reporting format

The protocol for the present review was designed according to the Cochrane guidelines²⁵ and reported with the Preferred Reporting Items for Systematic reviews and Meta-Analysis Extension (PRISMA)²⁶ statement for systematic reviews incorporating network meta-analyses for health care interventions.^{27,28} The study protocol was registered and allocated the identification number CRD42021264941 in the PROSPERO database, hosted by the National Institute for Health Research, University of York, Center for Reviews and Dissemination (www.crd.york.ac.uk/PROSPERO).

2.2 | Objectives

The goal of this review was to address the following focused question: What are the methods described for assessing and reporting PMC-related parameters and what is their clinical significance?

2.3 | PICOT question

The following Population, Intervention, Comparison, Outcomes, and Time (PICOT) framework²⁹ was used to guide the inclusion and exclusion of studies for the above-mentioned focused questions:

- **Population (P):** Patients with dental implants or in need of dental implant therapy.
- **Intervention (I):** Implant-related intervention (e.g., immediate or delayed implant placement, hard or soft tissue augmentation at implant sites, prosthetic restoration of dental implants, etc.) or visits in which PMC was assessed.
- **Comparison (C):** Any comparison among the included studies in terms of the methods for evaluating PMC and their reliability assessment was evaluated.
- **Outcome (O):** The current methods for assessing PMC, including direct comparison with adjacent or contralateral sites, professional indices/score systems, and spectrophotometry were evaluated.
- **Time (T):** No time restrictions were applied.

2.4 | Inclusion criteria

- Human studies.
- Randomized clinical trial (RCT), prospective non-randomized trials and observational studies with at least five subjects.
- Quantitative or qualitative assessment of PMC.

2.5 | Exclusion criteria

- Reviews, case reports, retrospective, in-vitro, or animal studies.
- Studies with less than five subjects.
- Studies providing the final score of an esthetic index involving the assessment of PMC, without reporting actual data on PMC.
- Studies assessing the color of the implant-supported restoration only.

2.6 | Outcome measures

Any quantitative outcomes describing PMC was assessed.

Details on the search strategy, study selection, data extraction, and risk of bias assessment are reported in the Supplementary Appendix S1.

3 | RESULTS

3.1 | Search results and study selection

The literature search process is shown in Figure 1. Following removal of duplicates, 865 records were screened on the basis of titles and

abstracts. Full-text assessment was performed for 327 articles. Based on the predetermined inclusion criteria, 121 articles were included.^{14-20,22,23,30-141} The reason for exclusion of the other 196 articles is available in the Appendix (Table S1). The inter-reviewer reliability in the screening and inclusion process, assessed with Cohen's κ , corresponded to 0.87 and 0.94 for assessment of titles and abstracts and full-text evaluation, respectively.

3.2 | Characteristics of the included studies

A total of 53 studies were RCTs, 9 non RCTs, 32 case series, and 27 observational studies (Table S2 of the Appendix). Among the RCTs, 44 were conducted with a parallel arm design, 8 were split-mouth and 1 trial was a combination of a parallel group and a split-mouth study design. Overall, 95 studies were performed in a single center, while the remaining 26 were multicenter studies. A total of 100 studies were conducted at university, 15 in private practice and 6 in both university and private practice settings. A total of 85 studies were performed in Europe, 13 in Asia, 9 in North America, 6 in Europe and North America, 3 in Asia and North America, 2 in Africa, 2 in Oceania, and 1 in South America. Smokers were excluded in 14 studies. Fifty articles included smokers based on the number of cigarettes consumed per day and 18 studies included smokers without any restrictions. The remaining 39 studies did not report this information. No funding from companies were reported in 96 studies, while the remaining 25 studies were funded by companies. Detailed study characteristics are described in the Table S2 of the Appendix.

A total of 93 studies assessed the outcomes of bone level implants, 6 study the outcomes of tissue level implants, 1 article investigated both bone level and tissue level implants, and the remaining 21 studies did not report information on the implant type (Table S3 of the Appendix).

Among the interventional studies, 37 articles evaluated the outcomes of immediate implant placement, 17 conventional implant placement, 14 implant placement followed by different abutments, 8 implant placement with guided bone regeneration, 9 peri-implant soft tissue augmentation with different graft materials, 3 flapless implant placement, 9 implant placement followed by different restoration material or protocol, 2 treatment of PSTD and one study assessed the outcomes following second stage performed either with scalpel or laser (Table S3 of the Appendix).

3.3 | Methods for assessing PMC

Peri-implant mucosa color was evaluated either in person, at the time of the visit (direct assessment, chairside) (45.5% of the included studies) or after the appointment on the collected photographs (indirect assessment) (54.5% of the included studies). The direct evaluation of PMC was conducted either with spectrophotometry (20.7% of the included studies) or visual comparison with adjacent and/or contralateral natural gingiva (24.8% of the included studies).




FIGURE 1 PRISMA flowchart

Peri-implant mucosa color assessment was performed qualitatively or quantitatively. The qualitative evaluation involves the definition of PMC as equal or different from the color of the adjacent and/or contralateral natural gingiva. This method was employed in two articles (1.7%).^{101,130} Another qualitative evaluation is based on the visual observation of PMC that is then graded using esthetic score systems (qualitative PMC evaluation and quantitative PMC reporting). A total of 93 studies (76.9%) described PMC using esthetic indices with predetermined grading scores.^{16-18,31-36,38,39,42-48,50-67,70-77,80-82,84-93,95-101,103,104,106-113,115-123,125,128,129,134-137,140-142}

Peri-implant mucosa color was assessed quantitatively using spectrophotometry in 25 studies (20.7%).^{14,15,19,20,22,23,36,39,40,48,67,77,78,82,86,93,104,113,123,125,126,129,131,132,138} The outcomes of this evaluation utilized the CIELAB parameters (Commission Internationale de l'Eclairage; *L* = lightness, *a* = chroma along red-green axis, *b* = chroma along yellow-blue axis), in terms of ΔE , ΔL , Δa , and Δb , for describing color differences in comparison with the adjacent and/or contralateral gingiva. One study (0.8%) utilized a software to import clinical photographs

and to measure the lightness, green-red, and blue-yellow shades of PMC, that were then converted to CIELAB color system.¹³⁷ Figure 2 summarizes the modalities for PMC assessment and reporting described in the included studies.

3.3.1 | PMC assessment using esthetic indices

The pink esthetic score (PES)¹⁶ was utilized in 58 studies (62.4% of the studies assessing PMC with an index),^{16,17,33-35,46,49-54,56-66,69-71,74,76,79,81,83,84,94,96-100,105,107-112,114-116,118-122,124,127,128,133,134} while the combined pink and white esthetic score (PES/WES)¹⁴² was employed in 26 articles (28.0%).^{17,31,37,38,41-45,47,55,72,73,80,90-92,102,103,106,117,135,136,139-141} Other esthetic indices utilized include the Copenhagen index score (CIS),⁸⁵⁻⁸⁸ the implant crown esthetic index (ICAI),^{17,75} the complex esthetic index (CEI),^{17,32,95} the implant esthetic score (IAS),¹⁷ the implant restoration esthetic index (IREI),¹⁰⁰ the mucosal scarring index (MSI)^{54,143} and the implant soft




FIGURE 2 Summary of the modalities for assessing and reporting peri-implant mucosa color. The percentage of the included articles utilizing the different modalities are reported in gray.

tissue dehiscence coverage esthetic score (IDES).¹⁸ One study utilized a modified gingival index,³⁰ one article followed a 3-point scoring system (0 for obvious color differences, 1 for moderate difference and 2 for a natural color/no differences)⁶⁸ and one study mentioned a 0–2 point grading for PMC, without specifying the criteria for this evaluation⁸⁹ (Tables S10 of the Appendix and 1).

The assessment of PMC in these above-mentioned esthetic indices was described in detail in the Supplementary Appendix, and illustrated in Figure 3.

3.3.2 | Significance of PMC assessment with esthetic indices

In 14 studies, the evaluation of PMC—as a parameter of the PES—provided information related to the color match with the gingiva of the reference tooth, without additional comparisons.^{16,17,50,52,70,76,83,84,94,98–100,107,108} PMC assessment as a part of the PES evaluation was also utilized for assessing changes/stability of the color of peri-implant soft tissue over time in 15 studies,^{34,35,50,70,72,80,97,98,110–113,115,116,119} while the majority ($N = 29$) of the articles reporting PMC from the PES evaluation utilized this parameter for comparing different treatment groups or interventions (e.g. immediate vs. delayed implants, different implant systems, implants restored with different type of abutment, etc.).^{35,46,51,53,54,56–66,74,81,105,116,119–122,124,127,128,133,134} Similarly, 10 studies reporting PMC as a parameter of the PES/WES, provided information on color discrepancy with contralateral reference tooth only, without additional comparisons.^{17,31,37,38,41,43,47,55,90,117} PMC as a part of the PES/WES assessment

was utilized to evaluate color changes over time in 7 studies^{42,45,80,102,103,106,139} and to compare color differences among different groups in 9 studies.^{44,72,73,91,92,136,140,141,143} The other esthetic indices were mainly used for assessing PMC compared to adjacent gingiva/gingiva of the contralateral tooth^{18,75,85,95,100} or for evaluating the outcomes of different treatment protocols on the color of the peri-implant mucosa.^{32,54,86–88,143}

It can be summarized that the assessment of PMC as a component of esthetic indices has allowed: (i) to describe the color match/discrepancy in relation to adjacent and/or contralateral natural gingiva, (ii) to evaluate color changes of the same implant site at different time points, and (iii) to compare the effects of different treatments on the esthetic outcomes of dental implants.

3.3.3 | PMC assessment using spectrophotometry

The characteristics of the 25 studies evaluating PMC with spectrophotometry are depicted in Table 2 and in the Supplementary Appendix S1. Some heterogeneity in the definition of the region of interest (ROI) for the use of the spectrophotometer was observed. The outcomes of the spectrophotometric analysis were expressed as ΔE , ΔL , Δa , and Δb , using the CIELAB parameters (Table S4 of the Appendix).

Five studies were designed to compare peri-implant and peri-odontal soft tissue color as primary outcomes, while the other studies focused on PMC with different abutments, restorations, or interventions, incorporating the assessment of the gingival color at adjacent or contralateral sites as a secondary endpoint. In particular, 15 studies

TABLE 1 Characteristics of PMC within the esthetic indices utilized in the included studies.

Index	Is PMC assessed as an individual parameter?	Reference for PMC	Categories for PMC (points)	Maximum score for the index (points)
PES (Furhauser et al., 2005)	Yes	Gingiva of the contralateral tooth	Obvious difference (0) Moderate difference (1) No difference (2)	14
ICAI (Meijer et al., 2005)	No (together with the surface of the mucosa)	Gingiva of the contralateral tooth	Gross mismatch (1) Slight mismatch (2) No mismatch (3)	35
IAS (Testori et al., 2005)	Yes	Healthy gingiva	Completely different (0) Not similar but acceptable (1) Similar (2)	9
PES/WES (Belser et al., 2009)	No (together with root convexity and soft tissue texture)	Gingiva of the contralateral tooth	Major discrepancy (0) Minor discrepancy (1) No discrepancy (2)	20
CIS (Dueded et al., 2009)	Yes	Adjacent soft tissue	No visible discoloration (1) Light grayish discoloration (2) Distinct grayish discoloration (3) Discoloration with visible metal (4)	24
CEI (Juodzbaly & Wang, 2010)	No (together with soft tissue texture)	Adjacent soft tissue	Deficient (0 ^a) Compromised (10 ^a) Adequate (20 ^a)	100 ^a
IREI (Li et al., 2019)	No (together with soft tissue texture)	Gingiva of the adjacent and contralateral tooth	No categories, PMC assessed using a VAS	100
MSI (Wessels et al., 2019)	Yes	Gingiva of the adjacent and contralateral tooth	Perfect color match (0) Slight mismatch (1) Obvious mismatch (2)	10
IDES (Zucchelli et al., 2021)	Yes	Adjacent soft tissue	Distinguishable from adjacent soft tissue (0) Not distinguishable from adjacent soft tissue (1)	10

Abbreviations: CEI, complex esthetic index; CIS, Copenhagen index score; IAS, implant esthetic index; ICAI, implant crown esthetic index; IDES, implant soft tissue dehiscence coverage esthetic score; IREI, implant restoration esthetic index; MSI, mucosal scarring index; PES, pink esthetic score; PES/WES, combined pink and white esthetic score.

^aRefers to a percentage and not to points.

[Correction added on 07 February 2023, after first online publication: 'Giannobile et al.' was changed to (2) in six places, (0) was changed to (1) beside 'Not distinguishable from the adjacent soft tissue' under 'Categories for PMC (points)' column in this version.]

utilized spectrophotometry for evaluating the effects of different abutments on PMC, four articles reported the PMC of implants with different necks, one study investigated the correlations between PMC and peri-implant mucosa characteristics, one trial assessed differences in PMC at sites that received connective tissue graft versus non-augmented sites and one RCT evaluated PMC at implant sites augmented with connective tissue graft or acellular dermal matrix (Tables 2 and S4 of the Appendix).

The use of spectrophotometer allowed demonstrating that the color of the peri-implant mucosa significantly differ from the color of the natural gingiva in 17 studies.^{14,20,22,23,36,39,40,48,67,78,104,113,123,129,131,132,138} Through spectrophotometric analysis, 4 studies were able to conclude that zirconia abutments induce a different, and less marked, color change in peri-implant mucosa than titanium abutments as compared to natural gingiva,^{14,39,48,132} while two studies did not confirm this finding.^{40,67} Anodized pink abutment had a significant different PMC than gray abutment according to Gil et al.⁷⁷ Another study from the same group showed that implants with pink neck and abutment had similar PMC than natural gingiva.⁷⁸ This finding was confirmed also by Bittner and coworker using

spectrophotometry.³⁶ Two studies showed that different restorations can result in different PMC.^{15,113} Similarly, the use of spectrophotometer allowed to conclude that other factors, including veneering or not zirconia abutments or augmenting peri-implant soft tissue with connective tissue graft, can also play a role on PMC.^{86,125} According to Hosseini, a better color match between peri-implant mucosa and natural gingiva was observed for implant sites augmented with connective tissue graft that also tended to maintain this color match over time, while non-augmented implant sites showed an increase color mismatch over the years compared to natural gingiva.⁸⁶ (Table S5 of the Appendix).

3.3.4 | Influence of mucosal thickness on PMC assessed with spectrophotometry

Twelve studies investigated possible correlations between MT and PMC using spectrophotometry.^{14,22,39,40,48,67,86,93,104,123,125,126} While three trials did not find a correlation between MT and PMC,^{39,40,86} nine studies demonstrated that MT has a significant impact on




FIGURE 3 Illustration of the professional esthetic indices involving the assessment of peri-implant mucosa color.

PMC.^{14,22,48,67,93,104,123,125,126} Benic and coworkers observed an inverse correlation between MT and PMC, in terms of ΔE (total color difference).²² A similar conclusion was also reached by Jun et al.⁹³ Two studies observed that the type of abutment significantly affected PMC only when MT was ≤ 2 mm.^{14,67} According to Lops, gold and zirconium abutments showed better PMC outcomes than titanium abutment in presence of MT ≤ 2 mm.¹⁴ Similarly, Martinz-Rus and coworkers reported a correlation between MT and PMC for titanium and pink-anodized titanium abutments only,¹⁰⁴ while Sailer and coworkers observed that MT had a significant impact on PMC at sites restored with zirconia and not titanium abutments.¹²³ Two studies from Thoma and coworkers further showed that MT has an impact on discoloration of the peri-implant soft tissue, although this association was not explored statistically, but it was obtained from sub-analyses dividing the sites in MT < 2 and MT ≥ 2 .^{125,126} (Table S5 of the Appendix).

3.3.5 | Significance of PMC assessment with spectrophotometry

Quantitative assessment of PMC using spectrophotometry has allowed: (i) to compare the esthetic outcomes of different abutments,

restorations, or interventions, (ii) to compare the color of natural gingiva of adjacent or contralateral dentition to PMC following different treatment protocols and (iii) to investigate the effect of MT on PMC.

3.4 | Risk of bias assessment

The risk of bias is reported in detail in the Tables S6–S9 of the Appendix.

4 | DISCUSSION

The esthetic outcome of dental implants remains one of the most widely discussed topics in implant therapy, with patients and clinicians having the common goal of rehabilitating the edentulous site(s) with an implant-supported restoration unnoticeable from natural dentition.^{44–146}

It has been largely discussed that the esthetics of dental implants depend on several parameters, including but not limited to the position, thickness and appearance of the soft tissue phenotypes,^{16,147,148} the presence/height of the papillae^{18,95,149} and the color, shape and texture of the implant-supported crown.^{42,50} Although a great deal of

TABLE 2 Characteristics of intervention among the studies quantitatively evaluating PMC using spectrophotometry.

Publication, reference	Study comparison	Treatment group	Specifics of the spectrophotometer	Landmarks for the ROI	PMC outcomes	Multiple follow-up visits
(Benic et al., 2017)	PMC versus color of gingiva in natural dentition	All ceramic Metal-ceramic Metal	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	Circular area of 1 mm Ø (1 mm apical to the mid-buccal mucosal/gingival margin)	ΔE ΔL Δa Δb	No
(Bittner et al., 2020)	PMC of implants with different necks	Pink neck Gray abutments	Crystaleye®, Olympus	Three incremental areas (1 × 1 mm ²) from the gingival margin in the apical direction (cervical, middle, apical)	ΔE ΔL Δa Δb	3 weeks and 6 months
(Bressan et al., 2011)	PMC with different abutments and comparison with color of gingiva in natural dentition	Abutments: Titanium Gold-alloy Zirconia	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	5 mm area around the gingival margin of the implant and contralateral tooth	ΔE ΔL Δa Δb	No
(Büchi et al., 2014)	PMC with different abutments	Abutments: Pink zirconia Zirconia	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	1 mm below the gingival margin of the implant and contralateral tooth	ΔE ΔL Δa Δb	BL, 5 min, 10 min and 1 week
(Cosgarea et al., 2015)	PMC with different abutments	Abutments: Titanium Zirconia	Shade vision system, X-Rite Inc.	1 mm ² located on the buccal gingival aspect of the abutments/crowns/tooth at 1 mm, 2 mm, 3 mm from the gingival margin	ΔE ΔL Δa Δb	20 min and 1 week
(Ferrari et al., 2017)	PMC with different abutments	Abutments: Titanium Titanium nitride Zirconia	VITA easy shade	1 mm below the gingival margin of the implant and contralateral tooth	ΔE ΔL Δa Δb	No
(Gil et al., 2017)	PMC with different abutments and implants with different necks	Abutments: Gray implant and gray abutment, Pink implant and gray abutment, Pink implant and Pink abutment	Crystaleye®, (Olympus Co., Tokyo, Japan)	Three incremental areas (1 × 1 mm ²) from the gingival margin in the apical direction (cervical, middle, apical) of implant and contralateral tooth	ΔL Δa Δb	No
(Gil et al., 2019)	PMC with different abutments and implants with different necks	Pink neck Gray abutments Pink abutments	Crystaleye®, (Olympus Co., Tokyo, Japan)	Three incremental areas (1 × 1 mm) from the gingival margin in the apical direction (cervical, middle, apical) of implant and contralateral tooth	ΔL Δa Δb	2 and 3 weeks
(Happe et al., 2022)	PMC of immediate implant augmented with different soft tissue grafts	Acellular dermal matrix versus Connective tissue graft	SpectroShade, Type 71.3000, MHT Optic Research AG, Switzerland	Not reported	ΔE	12 months

(Continues)

TABLE 2 (Continued)

Publication, reference	Study comparison	Treatment group	Specifics of the spectrophotometer	Landmarks for the ROI	PMC outcomes	Multiple follow-up visits
(Hosseini et al., 2020)	PMC of augmented versus non-augmented implants	Connective tissue graft versus non-augmented sites	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	Circular area of 3 mm Ø on the marginal implant mucosa and of the adjacent tooth	ΔE ΔL Δa Δb	BL, 1, 3, and 5 years
(Ishikawa-Nagai et al., 2007)	PMC of implants with different necks	Color strips White Black Light pink Pink Light orange Orange Gold violet	Handy-MSC, (Olympus Co., Tokyo, Japan)	Circular area of 20 mm Ø on the marginal implant mucosa and of adjacent or contralateral tooth	ΔL Δa Δb	BL and 5 min
(Jun et al., 2013)	PMC with different abutments	Abutments: Black White	Shadepilot, Degudent	5 mm ² area around the gingival margin	ΔL Δa Δb	No
(Jung et al., 2008)	PMC of implants with different restorations	Ceramic on Al ₂ O ₃ abutments Fused to metal (PFM) with titanium or gold abutment	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	Circular area of 3 mm Ø on the marginal implant mucosa and of the adjacent tooth	ΔE ΔL Δa Δb	BL, 1 and 2 weeks
(Lops et al., 2017)	PMC with different abutments	Abutments: Titanium Gold-alloy Zirconia	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	1 mm below the gingival margin 4 mm ² area around the implant mucosa and adjacent tooth	ΔE ΔL Δa Δb	No
(Martinez-Rus et al., 2017)	PMC with different abutments	Abutments: Pink anodized Titanium Gold anodized titanium Zirconia dioxide	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	1 mm below the gingival margin Circular area of 3 mm Ø around the implant mucosa and adjacent tooth	ΔE	No
(Paniz et al., 2014)	PMC versus color of gingiva in natural dentition	Esthetic evaluation	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	5 mm ² area around the gingival margin of implant mucosa and adjacent tooth	ΔE ΔL Δa Δb	No
(Park et al., 2007)	PMC versus color of gingiva in natural dentition	Implant tooth	Handy-MSC, (Olympus Co., Tokyo, Japan)	Circular area of 20 mm Ø (five incremental areas in 1 mm increments) on the marginal implant mucosa and of adjacent or contralateral tooth	ΔE ΔL Δa Δb	No
(Peng et al., 2017)	PMC with different abutments	Abutments: Titanium, Gold-gold with zirconia coping,	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	Circular area of 3 mm Ø on the marginal implant mucosa and of adjacent or contralateral tooth	ΔE ΔL Δa Δb	No

TABLE 2 (Continued)

Publication, reference	Study comparison	Treatment group	Specifics of the spectrophotometer	Landmarks for the ROI	PMC outcomes	Multiple follow-up visits
(Sailer et al., 2009)	PMC with different abutments	Titanium with metal coping, Titanium with gold-gold hue and metal coping, Zirconia Abutments: Titanium, Customized zirconia	Spectroshade™ Micro Device; Medical High Technologies, Verona, Italy	1 mm below the gingival margin Circular area of 3 mm Ø around the implant mucosa and contralateral tooth	ΔE ΔL Δa Δb	BL, 6 and 12 months
(Thoma et al., 2016)	PMC with different abutments	Abutments: White zirconia, Pink-veneered zirconia	Spectroshade, MHT Optic Research AG, Niederhasli, Switzerland	1 mm below the gingival margin around the implant mucosa and contralateral tooth	ΔE ΔL Δa Δb	BL, 5 and 10 min
(Thoma et al., 2017)	PMC with different abutments	Abutments: Fluorescent Zirconia, Zirconia	Spectroshade, MHT Optic Research AG, Niederhasli, Switzerland	1 mm below the gingival margin around the implant mucosa and adjacent tooth	ΔE ΔL Δa Δb	BL, abutment try-in, final crown
(Varoni et al., 2017)	PMC versus color of gingiva in natural dentition	Implant tooth	Portable UV-Vis-NIR spectrophotometer (HR4000; Ocean Optics, Dunedin, FL) Fiber optics reflectance spectroscopy with a 45°x45° geometry	1 mm below the gingival margin Circular area of 2.25 mm Ø around the implant mucosa and contralateral tooth (three incremental areas in 2 mm increments)	ΔE ΔL Δa Δb	No
(Waller et al., 2020)	PMC versus color of gingiva in natural dentition	Esthetic evaluation	Spectroshade, MHT Optic Research AG, Niederhasli, Switzerland	1 mm below the gingival margin Circular area of 5 mm Ø around the implant mucosa and contralateral tooth	ΔE	No
(Wang et al., 2020)	PMC with different abutments	Abutments: Gold titanium Pink titanium Titanium zirconia	Crystaleye®, (Olympus Co., Tokyo, Japan)	Are of 4 mm ² around the implant mucosa and contralateral tooth	ΔE ΔL Δa Δb	No
(Zembic et al., 2009)	PMC with different abutments	Abutments: Titanium zirconia	Spectroshade, MHT Optic Research AG, Niederhasli, Switzerland	1 mm below the gingival margin Circular area of 3 mm Ø around the implant mucosa and contralateral tooth	ΔE ΔL Δa Δb	No

Abbreviations: BL, baseline; PMC, peri-implant; ROI, region of interest; Δa, color difference on the red/green axis according to the CIELAB (Commission Internationale de l'Eclair) system; Δb, color difference on the yellow/blue axis according to the CIELAB system; ΔE, total color difference value according to the CIELAB system; ΔL, color difference in lightness/darkness according to the CIELAB system.

interest has been centered on PMC, there are no currently guidelines on the methods recommended for its assessment and reporting in clinical research.

In the present manuscript, we conducted a systematic appraisal of method for PMC assessment. Overall, PMC is frequently investigated in clinical studies, however its outcome is often not reported.

Among the 121 included articles, PMC assessment was more often performed after the research visit utilizing clinical photographs (54.5% of the studies). The evaluation of PMC in the remaining studies occurred during the research appointment (chairside) (45.5%), either with a visual examination (24.8%), or with a spectrophotometer (20.7%). While there are no studies comparing direct versus indirect PMC evaluation within the same patient population, one may speculate that the timing of the examination can affect the esthetic outcomes. The use of clinical photographs for esthetic assessment has several advantages compared to direct examination, including high resolution of anatomical structures, the possibility of obtaining evaluations from several operators with different background/expertise and also comparing the esthetic outcomes at different time points. Therefore, it's not surprising that several esthetic indices that are nowadays routinely used have been proposed and validated with clinical photographs.^{16,18,143}

Nevertheless, standardization of clinical photographs presents several challenges. While certain aspects of dental photography, mainly related to equipment, setting, patient position and operator, can be standardized and reproduced, there are several non-standardizable factors related to the hardware of the camera, such as resolution, color space, bit depth, absolute color rendition, etc. that are device dependent.¹⁵⁰ The illumination is also a factor that can affect the color of photographs even if taken with the same camera. Future studies are needed to compare the outcomes of direct versus indirect PMC assessment.

In terms of outcome measure, PMC was mostly reported as a number (98.4%), either as an average score from the points assigned to each case according to the utilized esthetic index (76.9%) or as mean color difference (ΔE , ΔL , Δa , and Δb according to the CIELAB system) with the adjacent and/or contralateral gingiva (21.5%). There are no doubts that grading PMC within predetermined esthetic indices is an easier, inexpensive, and less time-consuming method compared to spectrophotometry. Many esthetic indices—such as the PES, the PES/WES, the ICAI and the MSI^{16,142,143,151}—utilize a three-point scale to rate PMC, based on obvious, moderate or lack of color mismatch between peri-implant mucosa and adjacent and/or contralateral gingiva. This simplicity in defining PMC may also represent a limit for the detection of less marked color differences among two investigated sites. In addition, the use of several indices with heterogeneous criteria and scores for PMC often prevent the comparison of the esthetic outcomes among different studies in the literature.

On the other hand, spectrophotometry is the only currently available tool able to assess PMC in a quantitative manner. Intraoral scanners may represent an alternative—and more feasible—option to spectrophotometry to quantify PMC, however, more studies are needed to establish their accuracy in color reproduction.^{152,153}

Intraoral digital scanning appears to be a relatively easy and fast method for PMC assessment, but it has also to be considered that the calculation of PMC from the obtained digital models would require a certain level of experience with specific imaging software. Future studies in this direction are therefore advocated. On the other hand, spectrophotometry has been shown to be able to capture color differences that are not perceived by the human eye.^{21,23,154} In the studies included in the present review, the use of spectrophotometry allowed to evaluate PMC following different abutments, restorations, and interventions, and also to investigate the effect of MT on the appearance of the peri-implant mucosa. Overall, there is evidence that PMC differs from the color of natural gingiva, which is probably due to the structural differences within the soft tissue in terms of cellularity, fiber orientation, and vascularity, together with other factors related to the site (peri-implant soft tissue phenotype) and implant position.^{10,155–158} However, by using spectrophotometry to assess PMC can reduce these color differences between peri-implant mucosa and the adjacent/contralateral gingiva. Hence, spectrophotometry may be a good tool for assessing PMC especially when it was used to compare the color of natural gingiva.

Dental implants with zirconia abutments tend to have less discoloration than titanium abutments^{14,39,48,132} and that veneering zirconia abutments can further reduce color mismatch with contralateral gingiva.¹²⁵ Soft tissue augmentation at implant sites seems also to positively affect color match with adjacent and contralateral sites and the maintenance of this outcome over time,⁸⁶ which is probably due to the increased MT that can prevent discoloration of the soft tissue from underlying implant components.^{11,14} Most of the included studies investigating a correlation between PMC and MT using spectrophotometry concluded that MT has a significant impact on the color of peri-implant mucosa.^{14,22,48,67,93,105,124,126,127} Current recommendations for improving esthetic outcomes of implant therapy advocate a MT of at least 2 mm for avoiding discoloration of the peri-implant mucosa.^{10–12}

Finally, it should be mentioned that the literature has shown an inconsistent correlation between professional and patient esthetic evaluation.^{5,23,145,157} The impact of PMC on patient-reported esthetic evaluation should be further evaluated.

5 | RECOMMENDATION FOR FUTURE STUDIES AND PROPOSAL OF NEW GUIDELINES FOR EVALUATING PMC

Based on the present systematic appraisal, the following guidelines are suggested for evaluating and reporting PMC in clinical trials in dental implantology:

1. PMC evaluation should include the use of a spectrophotometer performed at different time points by a pre-calibrated operator: the color of the mucosa should be compared both with adjacent and contralateral gingiva and the result from these comparisons could be expressed using the CIELAB color system (ΔE , ΔL , Δa ,

and Δb). Methods for standardizing PMC assessment with spectrophotometry, including determination of the ROI and use of stents for reproducing the same position of the device at different time points, required further investigation.

2. PMC evaluation should also include the use of two or more professional esthetic indices (PES, PES/WES, ICAI, IAS, CIS, CEI, IREI, and MSI) performed at different time points by a pre-calibrated operator: if the study involves treatment of PSTDs, using the IDES is advocated. Direct qualitative evaluation of PMC at the time of the appointment should be preferred over indirect assessment of PMC from photographs.
3. Patient-reported subjective assessment of PMC using a visual analogue scale should also be reported: this outcome should be compared with professional evaluation of PMC (spectrophotometry and esthetic indices).
4. Intraoral digital scanners can be used to assess of PMC and soft tissue texture and shape. Future studies should take into consideration these recommendations for assessing PMC to improve reproducibility of PMC assessment and comparison of PMC outcomes among different studies and treatment modalities. The accuracy of intraoral digital scanners for PMC and esthetic assessment needs to be explored in future research.

6 | CONCLUSIONS

Based on the currently available evidence, and the limitations within this research, the following conclusions can be drawn:

1. The color of peri-implant mucosa is often investigated in clinical studies as one of the main esthetic outcomes.
2. Objective methods for assessing PMC mainly involve the use of spectrophotometry (quantitative assessment using the CIELAB color system) or professional esthetic indices (qualitative assessment).
3. Spectrophotometric assessment of PMC demonstrated that the color of the soft tissue around dental implants and natural teeth often differs. Several factors, including the type and color of the abutment, the type of restoration, soft tissue augmentation, and peri-implant MT, were found to affect PMC and to have the potential of promoting a better color match between peri-implant mucosa and natural gingiva.
4. There is evidence that MT is correlated to PMC, with thin mucosa (<2 mm) showing greater chance of discoloration compared to the natural gingiva of adjacent or contralateral teeth.
5. Several professional indices introduced for assessing the esthetic outcomes of dental implants (PES, PES/WES, ICAI, IAS, CIS, CEI, IREI, MSI, and IDES) involves the evaluation of PMC, which is rated based on the color match with the adjacent soft tissue and/or gingiva of the contralateral homologous tooth. The use of these indices also allows to evaluate the stability of PMC over time and to compare different intervention/treatment protocols in terms of final PMC.

6. The high level of heterogeneity observed between the included studies assessing PMC, in terms of study design, methods for PMC assessment, and reported outcomes, render comparisons among different studies and protocols challenging at the present time.

AUTHOR CONTRIBUTIONS

Lorenzo Tavelli, Leonardo Mancini, and Shayan Barootchi involved in conception and design of the study; Lorenzo Tavelli, Leonardo Mancini, and Shayan Barootchi involved in literature search; Lorenzo Tavelli and Leonardo Mancini involved in data extraction; Daniel S. Thoma, Ronald E. Jung, German O. Gallucci and Hom-Lay Wang involved in data interpretation, manuscript drafting and critical manuscript reviewal; Lorenzo Tavelli, Leonardo Mancini and Shayan Barootchi involved in the initial and final drafting of the work; all the authors were involved in the critical manuscript reviewal, and are accountable for all aspects of the work.

ACKNOWLEDGMENTS

The study was self-supported.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

ORCID

Leonardo Mancini  <https://orcid.org/0000-0002-7030-155X>

Shayan Barootchi  <https://orcid.org/0000-0002-5347-6577>

Daniel S. Thoma  <https://orcid.org/0000-0002-1764-7447>

German O. Gallucci  <https://orcid.org/0000-0001-6386-594X>

Hom-Lay Wang  <https://orcid.org/0000-0003-4238-1799>

Lorenzo Tavelli  <https://orcid.org/0000-0003-4864-3964>

REFERENCES

1. Elani HW, Starr JR, Da Silva JD, Gallucci GO. Trends in dental implant use in the U.S., 1999-2016, and projections to 2026. *J Dent Res.* 2018;97(13):1424-1430.
2. Buser D, Sennerby L, De Bruyn H. Modern implant dentistry based on osseointegration: 50 years of progress, current trends and open questions. *Periodontol 2000.* 2017;73(1):7-21.
3. Duong HY, Rocuzzo A, Stahli A, Salvi GE, Lang NP, Sculean A. Oral health-related quality of life of patients rehabilitated with fixed and removable implant-supported dental prostheses. *Periodontol 2000.* 2022;88(1):201-237.
4. Zucchelli G, Sharma P, Mounssif I. Esthetics in periodontics and implantology. *Periodontol 2000.* 2018;77(1):7-18.
5. Cosyn J, Thoma DS, Hammerle CH, De Bruyn H. Esthetic assessments in implant dentistry: objective and subjective criteria for clinicians and patients. *Periodontol 2000.* 2017;73(1):193-202.
6. Hammerle CHF, Tarnow D. The etiology of hard- and soft-tissue deficiencies at dental implants: a narrative review. *J Clin Periodontol.* 2018;45(20):S267-S277.
7. Tavelli L, Barootchi S, Majzoub J, et al. Prevalence and risk indicators of midfacial peri-implant soft tissue dehiscence at single site in the

- esthetic zone: a cross-sectional clinical and ultrasonographic study. *J Periodontol.* 2022;93(6):857-866.
8. Tonetti MS, Cortellini P, Graziani F, et al. Immediate versus delayed implant placement after anterior single tooth extraction: the timing randomized controlled clinical trial. *J Clin Periodontol.* 2017;44(2):215-224.
 9. Chen ST, Buser D. Clinical and esthetic outcomes of implants placed in postextraction sites. *Int J Oral Maxillofac Implants.* 2009;24:186-217.
 10. Wang II, Barootchi S, Tavelli L, Wang HL. The peri-implant phenotype and implant esthetic complications. Contemporary overview. *J Esthet Restor Dent.* 2021;33(1):212-223.
 11. Thoma DS, Muhlemann S, Jung RE. Critical soft-tissue dimensions with dental implants and treatment concepts. *Periodontol 2000.* 2014;66(1):106-118.
 12. Avila-Ortiz G, Gonzalez-Martin O, Couso-Queiruga E, Wang HL. The peri-implant phenotype. *J Periodontol.* 2020;91(3):283-288.
 13. Bonino F, Steffensen B, Natto Z, Hur Y, Holtzman LP, Weber HP. Prospective study of the impact of peri-implant soft tissue properties on patient-reported and clinically assessed outcomes. *J Periodontol.* 2018;89(9):1025-1032.
 14. Lops D, Stellini E, Sbricoli L, Cea N, Romeo E, Bressan E. Influence of abutment material on peri-implant soft tissues in anterior areas with thin gingival biotype: a multicentric prospective study. *Clin Oral Implants Res.* 2017;28(10):1263-1268.
 15. Jung RE, Holderegger C, Sailer I, Khraisat A, Suter A, Hammerle CH. The effect of all-ceramic and porcelain-fused-to-metal restorations on marginal peri-implant soft tissue color: a randomized controlled clinical trial. *Int J Periodontics Restorative Dent.* 2008;28(4):357-365.
 16. Furhauser R, Florescu D, Benesch T, Haas R, Mailath G, Watzek G. Evaluation of soft tissue around single-tooth implant crowns: the pink esthetic score. *Clin Oral Implants Res.* 2005;16(6):639-644.
 17. Hof M, Umar N, Budas N, Seemann R, Pommer B, Zechner W. Evaluation of implant esthetics using eight objective indices-comparative analysis of reliability and validity. *Clin Oral Implants Res.* 2018;29(7):697-706.
 18. Zucchelli G, Barootchi S, Tavelli L, Stefanini M, Rasperini G, Wang HL. Implant soft tissue dehiscence coverage esthetic score (IDES): a pilot within- and between-rater analysis of consistency in objective and subjective scores. *Clin Oral Implants Res.* 2021;32:349-358.
 19. Ishikawa-Nagai S, Da Silva J, Weber H, Park S. Optical phenomenon of peri-implant soft tissue. Part II. Preferred implant neck color to improve soft tissue esthetics. *Clin Oral Implants Res.* 2007;18(5):575-580.
 20. Park SE, Da Silva JD, Weber HP, Ishikawa-Nagai S. Optical phenomenon of peri-implant soft tissue. Part I. spectrophotometric assessment of natural tooth gingiva and peri-implant mucosa. *Clin Oral Implants Res.* 2007;18(5):569-574.
 21. Benic GI, Elmasry M, Hammerle CH. Novel digital imaging techniques to assess the outcome in oral rehabilitation with dental implants: a narrative review. *Clin Oral Implants Res.* 2015;26(11):86-96.
 22. Benic GI, Scherrer D, Sancho-Puchades M, Thoma DS, Hämmerle CH. Spectrophotometric and visual evaluation of peri-implant soft tissue color. *Clin Oral Implants Res.* 2017;28(2):192-200.
 23. Paniz G, Bressan E, Stellini E, Romeo E, Lops D. Correlation between subjective and objective evaluation of peri-implant soft tissue color. *Clin Oral Implants Res.* 2014;25(8):992-996.
 24. Gehrke P, Riekeberg U, Fackler O, Dhom G. Comparison of in vivo visual, spectrophotometric and colorimetric shade determination of teeth and implant-supported crowns. *Int J Comput Dent.* 2009;12(3):247-263.
 25. Higgins JPT, Thomas J, Chandler J, et al. *Cochrane Handbook for Systematic Reviews of Interventions version 6.2 (updated February 2021).* Cochrane; 2021 www.training.cochrane.org/handbook
 26. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *J Clin Epidemiol.* 2021;134:178-189.
 27. Hutton B, Salanti G, Caldwell DM, et al. The PRISMA extension statement for reporting of systematic reviews incorporating network meta-analyses of health care interventions: checklist and explanations. *Ann Intern Med.* 2015;162(11):777-784.
 28. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ.* 2015;350:g7647.
 29. Stillwell SB, Fineout-Overholt E, Melnyk BM, Williamson KM. Evidence-based practice, step by step: asking the clinical question: a key step in evidence-based practice. *Am J Nurs.* 2010;110(3):58-61.
 30. Al-Delayme RMA. Preservation of keratinized gingiva around dental implants using a diode laser when uncovering implants for second stage surgery. *Eur Oral Res.* 2019;53(3):106-112.
 31. Al-Dosari A, Al-Rowis R, Moslem F, Alshehri F, Ballo AM. Esthetic outcome for maxillary anterior single implants assessed by different dental specialists. *J Adv Prosthodont.* 2016;8(5):345-353.
 32. Anderson L, Dds MS, Inglehart M, et al. Implant associated soft tissue defects in the anterior maxilla: a randomized control trial comparing subepithelial connective tissue graft and acellular dermal matrix allograft. *Implant Dent.* 2014;23(4):416-425.
 33. Arora H, Ivanovski S. Correlation between pre-operative buccal bone thickness and soft tissue changes around immediately placed and restored implants in the maxillary anterior region: a 2-year prospective study. *Clin Oral Implants Res.* 2017;28(10):1188-1194.
 34. Arora H, Khzam N, Roberts D, Bruce WL, Ivanovski S. Immediate implant placement and restoration in the anterior maxilla: tissue dimensional changes after 2-5 year follow up. *Clin Implant Dent Relat Res.* 2017;19(4):694-702.
 35. Atef M, El Barbary A, Dahrous MSE, Zahran AF. Comparison of the soft and hard peri-implant tissue dimensional changes around single immediate implants in the esthetic zone with socket shield technique versus using xenograft: a randomized controlled clinical trial. *Clin Implant Dent Relat Res.* 2021;23(3):456-465.
 36. Bittner N, Schulze-Spate U, Silva C, et al. Comparison of peri-implant soft tissue color with the use of pink-neck vs gray implants and abutments based on soft tissue thickness: a 6-month follow-up study. *Int J Prosthodont.* 2020;33(1):29-38.
 37. Boon L, De Mars G, Favril C, Duyck J, Quirynen M, Vandamme K. Esthetic evaluation of single implant restorations, adjacent single implant restorations, and implant-supported fixed partial dentures: a 1-year prospective study. *Clin Implant Dent Relat Res.* 2020;22(1):128-137.
 38. Borgonovo AE, Corrocher G, Dolci M, Censi R, Vavassori V, Maiorana C. Clinical evaluation of zirconium dental implants placed in esthetic areas: a case series study. *Eur J Esthet Dent.* 2013;8(4):532-545.
 39. Bressan E, Paniz G, Lops D, Corazza B, Romeo E, Favero G. Influence of abutment material on the gingival color of implant-supported all-ceramic restorations: a prospective multicenter study. *Clin Oral Implants Res.* 2011;22(6):631-637.
 40. Büchi DL, Sailer I, Fehmer V, Hämmerle CH, Thoma DS. All-ceramic single-tooth implant reconstructions using modified zirconia abutments: a prospective randomized controlled clinical trial of the effect of pink veneering ceramic on the esthetic outcomes. *Int J Periodontics Restorative Dent.* 2014;34(1):29-37.
 41. Buser D, Halbritter S, Hart C, et al. Early implant placement with simultaneous guided bone regeneration following single-tooth extraction in the esthetic zone: 12-month results of a prospective study with 20 consecutive patients. *J Periodontol.* 2009;80(1):152-162.
 42. Buser D, Wittneben J, Bornstein MM, Grütter L, Chappuis V, Belser UC. Stability of contour augmentation and esthetic outcomes

- of implant-supported single crowns in the esthetic zone: 3-year results of a prospective study with early implant placement postextraction. *J Periodontol.* 2011;82(3):342-349.
43. Canullo L, Menini M, Covani U, Pesce P. Clinical outcomes of using a prosthetic protocol to rehabilitate tissue-level implants with a convergent collar in the esthetic zone: a 3-year prospective study. *J Prosthet Dent.* 2020;123(2):246-251.
 44. Cappare P, Ferrini F, Ruscica C, Pantaleo G, Tete G, Gherlone EF. Digital versus traditional workflow for immediate loading in single-implant restoration: a randomized clinical trial. *Biology (Basel).* 2021; 10(12):1281.
 45. Chappuis V, Rahman L, Buser R, Janner SFM, Belser UC, Buser D. Effectiveness of contour augmentation with guided bone regeneration: 10-year results. *J Dent Res.* 2018;97(3):266-274.
 46. Checchi V, Felice P, Zucchelli G, et al. Wide diameter immediate post-extractive implants vs delayed placement of normal-diameter implants in preserved sockets in the molar region: 1-year post-loading outcome of a randomised controlled trial. *Eur J Oral Implantol.* 2017;10(3):263-278.
 47. Cho HL, Lee JK, Um HS, Chang BS. Esthetic evaluation of maxillary single-tooth implants in the esthetic zone. *J Periodontal Implant Sci.* 2010;40(4):188-193.
 48. Cosgarea R, Gasparik C, Dudea D, Culic B, Dannewitz B, Sculean A. Peri-implant soft tissue colour around titanium and zirconia abutments: a prospective randomized controlled clinical study. *Clin Oral Implants Res.* 2015;26(5):537-544.
 49. Cosyn J, De Bruyn H, Cleymaet R. Soft tissue preservation and pink aesthetics around single immediate implant restorations: a 1-year prospective study. *Clin Implant Dent Relat Res.* 2013;15(6):847-857.
 50. Cosyn J, Eghbali A, De Bruyn H, Colllys K, Cleymaet R, De Rouck T. Immediate single-tooth implants in the anterior maxilla: 3-year results of a case series on hard and soft tissue response and aesthetics. *J Clin Periodontol.* 2011;38(8):746-753.
 51. Cosyn J, Eghbali A, De Bruyn H, Dierens M, De Rouck T. Single implant treatment in healing versus healed sites of the anterior maxilla: an aesthetic evaluation. *Clin Implant Dent Relat Res.* 2012;14(4): 517-526.
 52. Cutrim ES, Peruzzo DC, Benatti B. Evaluation of soft tissues around single tooth implants in the anterior maxilla restored with cemented and screw-retained crowns. *J Oral Implantol.* 2012;38(6):700-705.
 53. De Angelis N, Felice P, Pellegrino G, Camurati A, Gambino P, Esposito M. Guided bone regeneration with and without a bone substitute at single post-extractive implants: 1-year post-loading results from a pragmatic multicentre randomised controlled trial. *Eur J Oral Implantol.* 2011;4(4):313-325.
 54. De Bruyckere T, Cosyn J, Younes F, et al. A randomized controlled study comparing guided bone regeneration with connective tissue graft to re-establish buccal convexity: one-year aesthetic and patient-reported outcomes. *Clin Oral Implants Res.* 2020;31(6): 507-516.
 55. den Hartog L, Raghoobar GM, Slater JJ, Stellingsma K, Vissink A, Meijer HJ. Single-tooth implants with different neck designs: a randomized clinical trial evaluating the aesthetic outcome. *Clin Implant Dent Relat Res.* 2013;15(3):311-321.
 56. Donos N, Horvath A, Calciolari E, Mardas N. Immediate provisionalization of bone level implants with a hydrophilic surface. A five-year follow-up of a randomized controlled clinical trial. *Clin Oral Implants Res.* 2019;30(2):139-149.
 57. Elaskary AT, Maebed MA, Cho SC, El Tantawi M. A novel method for immediate implant placement in defective fresh extraction sites. *Int J Oral Maxillofac Implants.* 2020;35(4):799-807.
 58. Esposito M, Barausse C, Pistilli R, et al. Immediate loading of post-extractive versus delayed placed single implants in the anterior maxilla: outcome of a pragmatic multicenter randomised controlled trial 1-year after loading. *Eur J Oral Implantol.* 2015;8(4):347-358.
 59. Esposito M, Bressan E, Grusovin MG, et al. Do repeated changes of abutments have any influence on the stability of peri-implant tissues? One-year post-loading results from a multicentre randomised controlled trial. *Eur J Oral Implantol.* 2017;10(1):57-72.
 60. Esposito M, Cardaropoli D, Gobbato L, et al. The role of dental implant abutment design on the aesthetic outcome: preliminary 3-month post-loading results from a multicentre split-mouth randomised controlled trial comparing two different abutment designs. *Eur J Oral Implantol.* 2018;11(1):77-87.
 61. Esposito M, González-García A, Peñarrocha Diago M, et al. Natural or palatal positioning of immediate post-extractive implants in the aesthetic zone? 1-year results of a multicentre randomised controlled trial. *Eur J Oral Implantol.* 2018;11(2):189-200.
 62. Esposito M, Tallarico M, Trullenque-Eriksson A, Gianserra R. Endodontic retreatment vs dental implants of teeth with an uncertain endodontic prognosis: 1-year results from a randomised controlled trial. *Eur J Oral Implantol.* 2017;10(3):293-308.
 63. Esposito M, Trullenque-Eriksson A, Blasone R, et al. Clinical evaluation of a novel dental implant system as single implants under immediate loading conditions - 4-month post-loading results from a multicentre randomised controlled trial. *Eur J Oral Implantol.* 2016; 9(4):367-379.
 64. Esposito M, Zucchelli G, Cannizzaro G, et al. Immediate, immediate-delayed (6 weeks) and delayed (4 months) post-extractive single implants: 1-year post-loading data from a randomised controlled trial. *Eur J Oral Implantol.* 2017;10(1):11-26.
 65. Felice P, Pistilli R, Barausse C, Trullenque-Eriksson A, Esposito M. Immediate non-occlusal loading of immediate post-extractive versus delayed placement of single implants in preserved sockets of the anterior maxilla: 1-year post-loading outcome of a randomised controlled trial. *Eur J Oral Implantol.* 2015;8(4):361-372.
 66. Felice P, Zucchelli G, Cannizzaro G, et al. Immediate, immediate-delayed (6 weeks) and delayed (4 months) post-extractive single implants: 4-month post-loading data from a randomised controlled trial. *Eur J Oral Implantol.* 2016;9(3):233-247.
 67. Ferrari M, Carrabba M, Vichi A, Goracci C, Cagidiaco MC. Influence of abutment color and mucosal thickness on soft tissue color. *Int J Oral Maxillofac Implants.* 2017;32(2):393-399.
 68. Froum SJ, Khoully I, Tarnow DP, et al. The use of a xenogeneic collagen matrix at the time of implant placement to increase the volume of buccal soft tissue. *Int J Periodontics Restorative Dent.* 2015;35(2): 179-189.
 69. Fögl A, Zechner W, Pozzi A, et al. An open prospective single cohort multicenter study evaluating the novel, tapered, conical connection implants supporting single crowns in the anterior and premolar maxilla: interim 1-year results. *Clin Oral Investig.* 2017;21(6):2133-2142.
 70. Fürhauser R, Mailath-Pokorny G, Haas R, Busenlechner D, Watzek G, Pommer B. Esthetics of flapless single-tooth implants in the anterior maxilla using guided surgery: association of three-dimensional accuracy and pink esthetic score. *Clin Implant Dent Relat Res.* 2015;17(2):e427-e433.
 71. Fürhauser R, Mailath-Pokorny G, Haas R, Busenlechner D, Watzek G, Pommer B. Immediate restoration of immediate implants in the esthetic zone of the maxilla via the copy-abutment technique: 5-year follow-up of pink esthetic scores. *Clin Implant Dent Relat Res.* 2017;19(1):28-37.
 72. Furze D, Byrne A, Alam S, Brägger U, Wismeijer D, Wittneben JG. Influence of the fixed implant-supported provisional phase on the esthetic final outcome of implant-supported crowns: 3-year results of a randomized controlled clinical trial. *Clin Implant Dent Relat Res.* 2019;21(4):649-655.
 73. Gallucci GO, Grütter L, Nedir R, Bischof M, Belser UC. Esthetic outcomes with porcelain-fused-to-ceramic and all-ceramic single-implant crowns: a randomized clinical trial. *Clin Oral Implants Res.* 2011;22(1):62-69.

74. Garcia-Sanchez R, Mardas N, Buti J, Ortiz Ruiz AJ, Pardo ZG. Immediate implant placement in fresh alveolar sockets with a minimal split-thickness envelope flap: a randomised controlled clinical trial. *Clin Oral Implants Res.* 2021;32(9):1115-1126.
75. Gehrke P, Degidi M, Lulay-Saad Z, Dhom G. Reproducibility of the implant crown aesthetic index-rating aesthetics of single-implant crowns and adjacent soft tissues with regard to observer dental specialization. *Clin Implant Dent Relat Res.* 2009;11(3):201-213.
76. Gehrke P, Lobert M, Dhom G. Reproducibility of the pink esthetic score-rating soft tissue esthetics around single-implant restorations with regard to dental observer specialization. *J Esthet Restor Dent.* 2008;20(6):375-384. Discussion 385.
77. Gil MS, Ishikawa-Nagai S, Elani HW, et al. A prospective clinical trial to assess the optical efficacy of pink neck implants and pink abutments on soft tissue esthetics. *J Esthet Restor Dent.* 2017;29(6):409-415.
78. Gil MS, Ishikawa-Nagai S, Elani HW, et al. Comparison of the color appearance of peri-implant soft tissue with natural gingiva using anodized pink-neck implants and pink abutments: a prospective clinical trial. *Int J Oral Maxillofac Implants.* 2019;34(3):752-758.
79. Groenendijk E, Staas TA, Bronkhorst E, Raghoobar GM, Meijer GJ. Immediate implant placement and provisionalization: aesthetic outcome 1 year after implant placement. A prospective clinical multicenter study. *Clin Implant Dent Relat Res.* 2020;22(2):193-200.
80. Gu YX, Shi JY, Zhuang LF, Qiao SC, Xu YY, Lai HC. Esthetic outcome and alterations of soft tissue around single implant crowns: a 2-year prospective study. *Clin Oral Implants Res.* 2015;26(8):909-914.
81. Gualini F, Salina S, Rigotti F, et al. Subcrestal placement of dental implants with an internal conical connection of 0.5 mm versus 1.5 mm: outcome of a multicentre randomised controlled trial 1 year after loading. *Eur J Oral Implantol.* 2017;10(1):73-82.
82. Happe A, Schmidt A, Neugebauer J. Peri-implant soft-tissue esthetic outcome after immediate implant placement in conjunction with xenogeneic acellular dermal matrix or connective tissue graft: a randomized controlled clinical study. *J Esthet Restor Dent.* 2022;34(1): 215-225.
83. Hartlev J, Kohberg P, Ahlmann S, Andersen NT, Schou S, Isidor F. Patient satisfaction and esthetic outcome after immediate placement and provisionalization of single-tooth implants involving a definitive individual abutment. *Clin Oral Implants Res.* 2014;25(11): 1245-1250.
84. Hof M, Pommer B, Strbac GD, Sütö D, Watzek G, Zechner W. Esthetic evaluation of single-tooth implants in the anterior maxilla following autologous bone augmentation. *Clin Oral Implants Res.* 2013;24:88-93.
85. Hosseini M, Gotfredsen K. A feasible, aesthetic quality evaluation of implant-supported single crowns: an analysis of validity and reliability. *Clin Oral Implants Res.* 2012;23(4):453-458.
86. Hosseini M, Worsaae N, Gotfredsen K. Tissue changes at implant sites in the anterior maxilla with and without connective tissue grafting: a five-year prospective study. *Clin Oral Implants Res.* 2020;31(1):18-28.
87. Hosseini M, Worsaae N, Schiodt M, Gotfredsen K. A 1-year randomised controlled trial comparing zirconia versus metal-ceramic implant supported single-tooth restorations. *Eur J Oral Implantol.* 2011;4(4): 347-361.
88. Hosseini M, Worsaae N, Schiodt M, Gotfredsen K. A 3-year prospective study of implant-supported, single-tooth restorations of all-ceramic and metal-ceramic materials in patients with tooth agenesis. *Clin Oral Implants Res.* 2013;24(10):1078-1087.
89. Huang JP, Liu JM, Wu YM, et al. Clinical evaluation of xenogeneic collagen matrix versus free gingival grafts for keratinized mucosa augmentation around dental implants: a randomized controlled clinical trial. *J Clin Periodontol.* 2021;48:1293-1301.
90. Jones AR, Martin W. Comparing pink and white esthetic scores to layperson perception in the single-tooth implant patient. *Int J Oral Maxillofac Implants.* 2014;29(6):1348-1353.
91. Jonker BP, Wolvius EB, van der Tas J, Pijpe J. The effect of resorbable membranes on 1-stage ridge augmentation in anterior single-tooth replacement: a randomized, controlled clinical trial. *J Oral Maxillofac Surg.* 2019;77(9):e27.
92. Jonker BP, Wolvius EB, van der Tas JT, Tahmaseb A, Pijpe J. Esthetics and patient-reported outcomes of implants placed with guided bone regeneration and complete native bone: a prospective controlled clinical trial. *Int J Oral Maxillofac Implants.* 2020;35(2):406-414.
93. Jun SH, Ahn JS, Chang BM, Lee JD, Ryu JJ, Kwon JJ. In vivo measurements of human gingival translucency parameters. *Int J Periodontics Restorative Dent.* 2013;33(4):427-434.
94. Juodzbalys G, Wang HL. Soft and hard tissue assessment of immediate implant placement: a case series. *Clin Oral Implants Res.* 2007; 18(2):237-243.
95. Juodzbalys G, Wang HL. Esthetic index for anterior maxillary implant-supported restorations. *J Periodontol.* 2010;81(1):34-42.
96. Kolinski M, Hess P, Leziy S, et al. Immediate provisionalization in the esthetic zone: 1-year interim results from a prospective single-cohort multicenter study evaluating 3.0-mm-diameter tapered implants. *Clin Oral Investig.* 2018;22(6):2299-2308.
97. Konstantinidis IK, Siormpas KD, Kontsiotou-Siormpa E, Mitsias ME, Kotsakis GA. Long-term esthetic evaluation of the roll flap technique in the implant rehabilitation of patients with agenesis of maxillary lateral incisors: 10-year follow-up. *Int J Oral Maxillofac Implants.* 2016;31(4):820-826.
98. Kunavisarut C, Buranajanyakul L, Kitisubkanchana J, Pumpaluk P. A pilot study of small-diameter one-piece ceramic implants placed in anterior regions: clinical and esthetic outcomes at 1-year follow-up. *Int J Oral Maxillofac Implants.* 2020;35(5):965-973.
99. Lai HC, Zhang ZY, Wang F, Zhuang LF, Liu X, Pu YP. Evaluation of soft-tissue alteration around implant-supported single-tooth restoration in the anterior maxilla: the pink esthetic score. *Clin Oral Implants Res.* 2008;19(6):560-564.
100. Li X, Wu B, Cheng X, Li Y, Xie X, Deng F. Esthetic evaluation of implant-supported single crowns: the implant restoration esthetic index and patient perception. *J Prosthodont.* 2019;28(1):e51-e58.
101. Lorenzo R, Garcia V, Orsini M, Martin C, Sanz M. Clinical efficacy of a xenogeneic collagen matrix in augmenting keratinized mucosa around implants: a randomized controlled prospective clinical trial. *Clin Oral Implants Res.* 2012;23(3):316-324.
102. Mangano C, Levrini L, Mangano A, Mangano F, Macchi A, Caprioglio A. Esthetic evaluation of implants placed after orthodontic treatment in patients with congenitally missing lateral incisors. *J Esthet Restor Dent.* 2014;26(1):61-71.
103. Marconcini S, Giammarinaro E, Derchi G, Alfonsi F, Covani U, Barone A. Clinical outcomes of implants placed in ridge-preserved versus nonpreserved sites: a 4-year randomized clinical trial. *Clin Implant Dent Relat Res.* 2018;20(6):906-914.
104. Martinez-Rus F, Prieto M, Salido MP, Madrigal C, Ozcan M, Pradies G. A clinical study assessing the influence of anodized titanium and zirconium dioxide abutments and peri-implant soft tissue thickness on the optical outcome of implant-supported lithium disilicate single crowns. *Int J Oral Maxillofac Implants.* 2017;32(1):156-163.
105. Mau JL, Grodin E, Lin JJ, Chen MC, Ho CH, Cochran D. A comparative, randomized, prospective, two-center clinical study to evaluate the clinical and esthetic outcomes of two different bone grafting techniques in early implant placement. *J Periodontol.* 2019;90(3): 247-255.
106. Meijndert CM, Raghoobar GM, Santing HJ, Vissink A, Meijer HJA. Performance of bone-level implants with conical connections in the anterior maxilla: a 5-year prospective cohort study. *Clin Oral Implants Res.* 2020;31(2):173-180.
107. Nissen KJ, Starch-Jensen T. Maxillary sinus floor augmentation with autogenous bone graft from the ascending mandibular ramus. *Implant Dent.* 2019;28(1):46-53.

108. Noelken R, Kunkel M, Wagner W. Immediate implant placement and provisionalization after long-axis root fracture and complete loss of the facial bony lamella. *Int J Periodontics Restorative Dent.* 2011; 31(2):175-183.
109. Noelken R, Moergel M, Kunkel M, Wagner W. Immediate and flap-less implant insertion and provisionalization using autogenous bone grafts in the esthetic zone: 5-year results. *Clin Oral Implants Res.* 2018;29(3):320-327.
110. Noelken R, Neffe BA, Kunkel M, Wagner W. Maintenance of marginal bone support and soft tissue esthetics at immediately provisionalized OsseoSpeed implants placed into extraction sites: 2-year results. *Clin Oral Implants Res.* 2014;25(2):214-220.
111. Noelken R, Oberhansl F, Kunkel M, Wagner W. Immediately provisionalized OsseoSpeed™ profile implants inserted into extraction sockets: 3-year results. *Clin Oral Implants Res.* 2016;27(6):744-749.
112. Östman PO, Chu SJ, Drago C, Saito H, Nevins M. Clinical outcomes of maxillary anterior postextraction socket implants with immediate provisional restorations using a novel macro-hybrid implant design: an 18- to 24-month single-cohort prospective study. *Int J Periodontics Restorative Dent.* 2020;40(3):355-363.
113. Peng M, Zhao WJ, Hosseini M, Zhou WJ, Xiao T, Chuan JL. Influence of restorative materials on color of implant-supported single crowns in esthetic zone: a spectrophotometric evaluation. *Biomed Res Int.* 2017;2017:5034358.
114. Pieri F, Aldini NN, Marchetti C, Corinaldesi G. Esthetic outcome and tissue stability of maxillary anterior single-tooth implants following reconstruction with mandibular block grafts: a 5-year prospective study. *Int J Oral Maxillofac Implants.* 2013;28(1):270-280.
115. Pieri F, Siroli L, Forlivesi C, Corinaldesi G. Clinical, esthetic, and radiographic evaluation of small-diameter (3.0-mm) implants supporting single crowns in the anterior region: a 3-year prospective study. *Int J Periodontics Restorative Dent.* 2014;34(6):825-832.
116. Pohl V, Cede J, Pokorny GBM, Haas R, Pohl S. Esthetic outcomes for immediate implant placement with immediate provisionalization in the anterior maxilla with buccal dehiscence: results of a comparative pilot study. *Int J Oral Maxillofac Implants.* 2022;37(3):508-514.
117. Pollini A, Morton D, Arunyanak SP, Harris BT, Lin WS. Evaluation of esthetic parameters related to a single implant restoration by laypeople and dentists. *J Prosthet Dent.* 2020;124(1):94-99.
118. Prati C, Zamparini F, Canullo L, Pirani C, Botticelli D, Gandolfi MG. Factors affecting soft and hard tissues around two-piece transmucosal implants: a 3-year prospective cohort study. *Int J Oral Maxillofac Implants.* 2020;35(5):1022-1036.
119. Puisys A, Auzbikaviciute V, Vindasiute-Narbutė E, Pranskunas M, Razukevicius D, Linkevicius T. Immediate implant placement vs. early implant treatment in the esthetic area. A 1-year randomized clinical trial. *Clin Oral Implants Res.* 2022;33(6):634-655.
120. Puisys A, Deikuvienė J, Vindasiute-Narbutė E, Razukevicius D, Zvirblis T, Linkevicius T. Connective tissue graft vs porcine collagen matrix after immediate implant placement in esthetic area: a randomized clinical trial. *Clin Implant Dent Relat Res.* 2022;24(2):141-150.
121. Raes F, Cosyn J, Crommelinck E, Coessens P, De Bruyn H. Immediate and conventional single implant treatment in the anterior maxilla: 1-year results of a case series on hard and soft tissue response and aesthetics. *J Clin Periodontol.* 2011;38(4):385-394.
122. Rieder D, Eggert J, Krafft T, Weber HP, Wichmann MG, Heckmann SM. Impact of placement and restoration timing on single-implant esthetic outcome - a randomized clinical trial. *Clin Oral Implants Res.* 2016;27(2):e80-e86.
123. Sailer I, Zembic A, Jung RE, Siegenthaler D, Holderegger C, Hämmerle CH. Randomized controlled clinical trial of customized zirconia and titanium implant abutments for canine and posterior single-tooth implant reconstructions: preliminary results at 1 year of function. *Clin Oral Implants Res.* 2009;20(3):219-225.
124. Sun C, Zhao J, Liu Z, et al. Comparing conventional flap-less immediate implantation and socket-shield technique for esthetic and clinical outcomes: a randomized clinical study. *Clin Oral Implants Res.* 2020; 31(2):181-191.
125. Thoma DS, Brandenberg F, Fehmer V, Knechtle N, Hämmerle CH, Sailer I. The esthetic effect of veneered zirconia abutments for single-tooth implant reconstructions: a randomized controlled clinical trial. *Clin Implant Dent Relat Res.* 2016;18(6):1210-1217.
126. Thoma DS, Gamper FB, Sapata VM, Voce G, Hämmerle CHF, Sailer I. Spectrophotometric analysis of fluorescent zirconia abutments compared to "conventional" zirconia abutments: a within subject controlled clinical trial. *Clin Implant Dent Relat Res.* 2017;19(4): 760-766.
127. Urban IA, Wessing B, Aláñez N, et al. A multicenter randomized controlled trial using a novel collagen membrane for guided bone regeneration at dehiscence single implant sites: outcome at prosthetic delivery and at 1-year follow-up. *Clin Oral Implants Res.* 2019;30(6): 487-497.
128. van Nimwegen WG, Raghoobar GM, Zuiderveld EG, Jung RE, Meijer HJA, Mühlemann S. Immediate placement and provisionalization of implants in the aesthetic zone with or without a connective tissue graft: a 1-year randomized controlled trial and volumetric study. *Clin Oral Implants Res.* 2018;29(7):671-678.
129. Varoni EM, Moltrasio G, Gargano M, Ludwig N, Lodi G, Scaringi R. Color analysis of periimplant soft tissues focusing on implant system: a case series. *Implant Dent.* 2017;26(2):217-224.
130. Vellis J, Dmd MS, Kutkut A, Dds MS, Al-Sabbagh M, Dds MS. Comparison of xenogeneic collagen matrix vs. free gingival grafts to increase the zone of keratinized mucosa around functioning implants. *Implant Dent.* 2019;28(1):20-27.
131. Waller T, Evcı E, Hämmerle CH, Hüslér J, Jung RE, Thoma DS. Perceptibility and acceptability of color differences of single-tooth implants at the restoration and mucosa levels: an exploratory clinical study. *Int J Prosthodont.* 2020;33(5):487-492.
132. Wang T, Wang L, Lu Q, Fan Z. Influence of anodized titanium abutments on the esthetics of the peri-implant soft tissue: a clinical study. *J Prosthet Dent.* 2021;125(3):445-452.
133. Wanis RW, Hosny MM, ElNahass H. Clinical evaluation of the buccal aspect around immediate implant using dual zone therapeutic concept versus buccal gap fill to bone level: a randomized controlled clinical trial. *Clin Implant Dent Relat Res.* 2022;24(3): 307-319.
134. Weinländer M, Lekovic V, Spadijer-Gostovic S, Milicic B, Wegscheider WA, Piehslinger E. Soft tissue development around abutments with a circular macro-groove in healed sites of partially edentulous posterior maxillae and mandibles: a clinical pilot study. *Clin Oral Implants Res.* 2011;22(7):743-752.
135. Wessels R, Vervaeke S, Seyssens L, Eghbali A, Cosyn J. A 5-year cohort study on early implant placement with guided bone regeneration or alveolar ridge preservation with connective tissue graft. *Clin Implant Dent Relat Res.* 2020;22(6):697-705.
136. Wittneben JG, Gavric J, Belser UC, et al. Esthetic and clinical performance of implant-supported all-ceramic crowns made with prefabricated or CAD/CAM zirconia abutments: a randomized, multicenter clinical trial. *J Dent Res.* 2017;96(2):163-170.
137. Yao JW, Wang HL. Assessment of peri-implant soft tissue adaptive pressure and time after provisional restorations. *Int J Periodontics Restorative Dent.* 2019;39(6):809-815.
138. Zembic A, Sailer I, Jung RE, Hammerle CH. Randomized-controlled clinical trial of customized zirconia and titanium implant abutments for single-tooth implants in canine and posterior regions: 3-year results. *Clin Oral Implants Res.* 2009;20(8):802-808.
139. Zucchelli G, Felice P, Mazzotti C, et al. 5-year outcomes after coverage of soft tissue dehiscence around single implants: a prospective cohort study. *Eur J Oral Implantol.* 2018;11(2):215-224.

140. Zuiderveld E, Meijer H, Vissink A, Raghoobar G. The influence of different soft-tissue grafting procedures at single implant placement on esthetics: a randomized controlled trial. *J Periodontol*. 2018;89(8):903-914.
141. Zuiderveld EG, Meijer HJ, Vissink A, Raghoobar GM. Outcome of treatment with single implants in preserved versus nonpreserved alveolar ridges: a 1-year cohort study. *Int J Oral Maxillofac Implants*. 2019;34(6):1457-1465.
142. Belsler UC, Grütter L, Vailati F, Bornstein MM, Weber HP, Buser D. Outcome evaluation of early placed maxillary anterior single-tooth implants using objective esthetic criteria: a cross-sectional, retrospective study in 45 patients with a 2- to 4-year follow-up using pink and white esthetic scores. *J Periodontol*. 2009;80(1):140-151.
143. Wessels R, De Rooze S, De Bruyckere T, et al. The mucosal scarring index: reliability of a new composite index for assessing scarring following oral surgery. *Clin Oral Investig*. 2019;23(3):1209-1215.
144. Stefanini M, Felice P, Mazzotti C, Mounssif I, Marzadori M, Zucchelli G. Esthetic evaluation and patient-centered outcomes in single-tooth implant rehabilitation in the esthetic area. *Periodontol 2000*. 2018;77(1):150-164.
145. Rocuzzo A, Imber JC, Marruganti C, Salvi GE, Ramieri G, Rocuzzo M. Clinical outcomes of dental implants in patients with and without history of periodontitis: a 20-year prospective study. *J Clin Periodontol*. 2022;49(12):1346-1356.
146. Rocuzzo A, Imber JC, Lempert J, Hosseini M, Jensen SS. Narrow diameter implants to replace congenital missing maxillary lateral incisors: a 1-year prospective, controlled, clinical study. *Clin Oral Implants Res*. 2022;33(8):844-857.
147. Zucchelli G, Tavelli L, McGuire MK, et al. Autogenous soft tissue grafting for periodontal and peri-implant plastic surgical reconstruction. *J Periodontol*. 2020;91(1):9-16.
148. Zucchelli G, Tavelli L, Stefanini M, et al. Classification of facial peri-implant soft tissue dehiscence/deficiencies at single implant sites in the esthetic zone. *J Periodontol*. 2019;90(10):1116-1124.
149. Dueled E, Gottfredsen K, Trab Damsgaard M, Hede B. Professional and patient-based evaluation of oral rehabilitation in patients with tooth agenesis. *Clin Oral Implants Res*. 2009;20(7):729-736.
150. Ahmad I. Standardization for dental photography. Part 1. *J Cosmetic Dent*. 2020;36(2):26-35.
151. Meijer HJ, Stellingsma K, Meijndert L, Raghoobar GM. A new index for rating aesthetics of implant-supported single crowns and adjacent soft tissues—the implant crown aesthetic index. *Clin Oral Implants Res*. 2005;16(6):645-649.
152. Liu Y, Zhang R, Ye H, et al. The development of a 3D colour reproduction system of digital impressions with an intraoral scanner and a 3D printer: a preliminary study. *Sci Rep*. 2019;9(1):20052.
153. Deferm JT, Schreurs R, Baan F, et al. Validation of 3D documentation of palatal soft tissue shape, color, and irregularity with intraoral scanning. *Clin Oral Investig*. 2018;22(3):1303-1309.
154. Jung RE, Sailer I, Hammerle CH, Attin T, Schmidlin P. In vitro color changes of soft tissues caused by restorative materials. *Int J Periodontics Restorative Dent*. 2007;27(3):251-257.
155. Sculean A, Gruber R, Bosshardt DD. Soft tissue wound healing around teeth and dental implants. *J Clin Periodontol*. 2014;41(15):S6-S22.
156. Ivanovski S, Lee R. Comparison of peri-implant and periodontal marginal soft tissues in health and disease. *Periodontol 2000*. 2018;76(1):116-130.
157. De Bruyn H, Raes S, Matthys C, Cosyn J. The current use of patient-centered/reported outcomes in implant dentistry: a systematic review. *Clin Oral Implants Res*. 2015;26(11):45-56.
158. Sailer I, Fehmer V, Ioannidis A, Hammerle CH, Thoma DS. Threshold value for the perception of color changes of human gingiva. *Int J Periodontics Restorative Dent*. 2014;34(6):757-762.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Mancini L, Barootchi S, Thoma DS, et al. The peri-implant mucosa color: A systematic appraisal of methods for its assessment and clinical significance. *Clin Implant Dent Relat Res*. 2023;25(2):224-240. doi:[10.1111/cid.13180](https://doi.org/10.1111/cid.13180)