

Infectivity of the parasite *Metschnikowia bicuspidata* is decreased by time spent as a transmission spore, but exposure to phycotoxins in the water column has no effect

Kristel F. Sánchez¹  | Baili Zhong^{1,2}  | Jorge A. Agudelo^{1,3} | Meghan A. Duffy¹ 

¹Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, U.S.A.

²Chemistry Department, Pomona College, Claremont, California, U.S.A.

³Department of Chemistry and Biochemistry, Florida International University, Miami, Florida, U.S.A.

Correspondence

Kristel F. Sánchez, Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, U.S.A.
Email: kfsanche@umich.edu

Funding information

Gordon and Betty Moore Foundation, Grant/Award Number: GBMF9202; National Science Foundation, Grant/Award Number: DEB-1655856; University of Michigan

Abstract

1. Transmission from one host to another is a crucial component of parasite fitness. For some aquatic parasites, transmission occurs via a free-living stage that spends time in the water, awaiting an encounter with a new host. These parasite transmission stages can be impacted by biotic and abiotic factors that influence the parasite's ability to successfully infect or grow in a new host.
2. Here we tested whether time spent in the water column and/or exposure to common cyanobacterial toxins impacted parasite transmission stages. More specifically, we tested whether the infectivity, within host growth, and virulence of the fungal parasite *Metschnikowia bicuspidata* changed as a result of time spent in the water or from exposure to cyanotoxins in the water column. We exposed parasite transmission spores to different levels of one of two ecologically important cyanotoxins, microcystin-LR and anatoxin-a, and factorially manipulated the amount of time spores were incubated in water. We removed the toxins and used those same spores to infect one genotype of the common lake zooplankton *Daphnia dentifera*.
3. We found that cyanotoxins did not impact parasite fitness (infection prevalence and spore yield per infected host) or virulence (host lifetime reproduction and survivorship) at the tested concentrations (10 and 30 µg/L). However, we found that spending longer as a transmission spore decreased a spore's chances for successful infection: spores that were only incubated for 24 hr infected approximately 75% of exposed hosts, whereas spores incubated for 10 days infected less than 50% of exposed hosts.
4. We also found a negative relationship between the final spore yield from infected hosts and the proportion of hosts that became infected. In treatments where spores spent longer in the water column prior to encountering a host, infection prevalence was lower (indicating lower per spore infectivity), but each infected host yielded more spores at the end of infection. We hypothesise that this pattern may result from intraspecific parasite competition within the host.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *Freshwater Biology* published by John Wiley & Sons Ltd.

5. Overall, these results suggest that transmission spores of this parasite are not strongly influenced by cyanotoxins in the water column, but that other aspects of spending time in the water strongly influence parasite fitness.

KEYWORDS

anatoxin, cyanobacteria, *Daphnia*, host-parasite, microcystin

1 | INTRODUCTION

A crucial component of parasite fitness is successfully transmitting from one host to another. For many parasites, transmission occurs via a free-living stage that spends time in the environment until it encounters a suitable host. In aquatic ecosystems, these parasite transmission spores spend time in the water column (before sinking out or moving on to their next life history stage), where they are exposed to abiotic and biotic factors that can ultimately affect the timing, size, and duration of an epidemic. While we know these transmission stages are very common for aquatic parasites and that many biotic and abiotic factors impact their transmission to hosts, we know less about how prolonged exposure to environmental factors such as cyanotoxins influences their ability to infect and propagate in new hosts.

One factor that is likely to influence the fitness of parasite transmission stages is time spent in the environment. In addition to the risks of spore predation (Thieltges et al., 2013) and settling out of the water column (Brookes et al., 2004), spores may lose viability over time, perhaps due to depleting the resources they acquired from their host (Amigó et al., 1996; Duffy & Hunsberger, 2019; Malone et al., 2001). Longer time in the environment also increases spore exposure to stressors such as light, temperature, secondary metabolites produced by other organisms, and anthropogenic pollutants (Amigó et al., 1996; Hallmann & Sikora, 1996; Heagle, 1973; Shaw et al., 2020; Vasemagi et al., 2017). Together, this suggests that increased time in the water column should decrease parasite fitness.

When they are in the water column, parasite spores are exposed to secondary metabolites. These are often released from phytoplankton cells, becoming dissolved in water (Huisman et al., 2018); some are known to have antibacterial and antifungal properties (Borowitzka, 1995; Leflaive & Ten-Hage, 2007; Ostensvik et al., 2002; Volk & Furkert, 2006). While there is no clear consensus on why cyanobacteria produce these phycotoxins, one possibility is that these antimicrobial compounds are released in the water as part of an allelopathic attack by phytoplankton to deter enemies (Leflaive & Ten-Hage, 2007); these same compounds may have a negative impact on the transmission spores of parasites. If so, this could impact parasite epidemics in aquatic ecosystems (Lafferty & Holt, 2003). Thus, we were interested in understanding how cyanotoxins might impact parasite transmission stages in the water.

Understanding the impact of cyanotoxins on parasite transmission stages is particularly important given that cyanobacterial harmful algal blooms (CHABs) are expected to increase in

frequency and intensity in the coming years with climate change (Huisman et al., 2018). CHABs and their toxins can increase infection rates (by making hosts more vulnerable to infection) and increase the negative effects of parasites on their hosts (Harvell et al., 2004; Landsberg, 2002) including important zooplankton in freshwater ecosystems (Tellenbach et al., 2016); however, positive effects on aquatic hosts of consuming chemically defended cyanobacteria have also been observed (Coopman et al., 2014; Manzi et al., 2019; Sánchez et al., 2019). When considering the potential impact of CHABs on parasitism, to date, the focus has been on one main mechanism: that toxins may alter host susceptibility or tolerance (Andersen et al., 2016; Coopman et al., 2014; Duperron et al., 2019; Harvell et al., 2004; Landsberg, 2002; Lassudrie et al., 2020; Penczykowski et al., 2014; Tellenbach et al., 2016). Here, because some cyanotoxins are known to have antimicrobial properties, we consider the possibility that cyanotoxins have direct negative effects on parasite transmission spores in the water column.

Using the *Daphnia*-*Metschnikowia* host-parasite system, we tested whether time spent in the water column and cyanotoxins have negative consequences on the parasite's ability to infect and grow within a host, both of which are key components of parasite fitness. A variety of biotic and abiotic factors are known to impact *Metschnikowia* epidemics in lakes (Cáceres et al., 2006; Hall et al., 2010). In this study, we were concerned with how the duration of exposure to the environment impacts the free-living stage of the parasite with an emphasis on the duration of exposure to certain cyanotoxins. We also looked for impacts on parasite virulence, specifically looking at host reproduction and lifespan, both of which are influenced by infections.

2 | METHODS

In this study, we used the zooplankton *Daphnia dentifera*, which is common in stratified lakes in temperate North America (Tessier & Woodruff, 2002). For our experiments, we used the Midland 37 (MID37) genotype, which was isolated from Midland Lake in Greene County, Indiana and has been used in several prior experiments (e.g., Auld et al., 2012, 2014). We also used the common fungal parasite *Metschnikowia bicuspidata* Standard isolate, originally isolated from Baker Lake in Barry County, Michigan. *Daphnia* become infected by inadvertently consuming transmission spores they encounter in the water column when feeding. By *transmission spore*, we refer to the mature, needle-shaped ascus that

contains the ascospore (Codreanu & Codreanu-Balcescu, 1981; Metschnikoff, 1884). After consumption by the host, infection can begin if the needle-shape spore crosses the gut barrier and is not fought off by a host haemocyte response (Metschnikoff, 1884; Stewart Merrill & Cáceres, 2018). Once infection has taken hold, the fungus replicates within the haemolymph of the host (Stewart Merrill & Cáceres, 2018). The parasite reduces the fecundity and lifespan of infected hosts (Auld et al., 2012). *Metschnikowia* is an obligate killer, meaning it must kill its host in order to transmit to a new host (Ebert, 2005); transmission spores are released into the environment after host death, after which they can be consumed by a new host, completing the parasite's life cycle.

We incubated fungal transmission spores for different lengths of time in filtered lake water (Pall AE filters, 1 µm pore size). The incubation times were 24 hr, 3 days, 5 days, 7 days, and 10 days. We also added two common cyanobacterial toxins, microcystin-LR and anatoxin-a, to the water during the incubations. We chose these cyanotoxins because they are commonly produced during blooms (Huisman et al., 2018) and because prior research suggested that they reduced infection prevalence (in the case of microcystin-LR) or that production of them can increase in the presence of *Metschnikowia* (in the case of anatoxin; Sánchez et al., 2019). Microcystin-LR is produced by some members of the genus *Microcystis*, which has been extensively studied due to concerns over its geographical expansion and capability of producing CHABs in both marine and freshwater ecosystems (Huisman et al., 2018). Anatoxin-a is produced by some members of the genus *Anabaena*. Microcystin-LR is a hepatotoxin while anatoxin-a is considered a potent neurotoxin in vertebrate models (Christensen & Khan, 2020). Both toxins are also produced by other genera of cyanobacteria such as *Planktothrix*, *Oscillatoria*, *Aphanizomenon*, *Cylindrospermum*, and *Dolichospermum*, all capable of producing CHABs (Huisman et al., 2018). We also included two types of negative controls: a solvent control of 0.01% acetic acid (see toxin preparation for explanation) and a negative control with no toxin or solvent added. The incubation times were crossed factorially with the toxin treatments, as described below. After the appropriate incubation time, we carried out infection assays in which we exposed *Daphnia* hosts to these spores and measured infection, spore production, host reproduction, and host mortality over time.

2.1 | Toxin preparation

Pure microcystin-LR standard was purchased from Cayman Chemical and suspended in 1 ml of nano-pure water for a concentration of 1 mg/ml. Anatoxin-a standard was acquired from Abraxis. The anatoxin-a comes in a solution of 3:1 water and methanol and 0.01% acetic acid. We placed 1.5 ml of the anatoxin solution in a 2-ml Eppendorf tube and evaporated the methanol using an Eppendorf Vacufuge (Eppendorf) at 23°C. Once evaporation had occurred, we added nanopure water to restore to the

original volume/concentration. Toxins were frozen in between uses during the exposure period.

2.2 | Spore preparation

Spores for the experiment were grown in vivo by infection of *D. dentifera* (Standard genotype) in the laboratory. Infected animals with well-developed late-stage infections were collected from laboratory cultures and placed in 2-ml Eppendorf tubes with 100–500 µl of milliQ water, then stored in the refrigerator at 4°C. For this experiment, we used spores from animals that had been stored in the refrigerator for 2 weeks. To generate the spore slurry for experimental infections, we crushed infected animals to release spores, and then determined the density of mature ascospores using a haemocytometer (Hausser Scientific 3100) and a compound microscope (Olympus BX53) at 400× magnification. Each incubation treatment was initiated with a new spore slurry (made from infected animals that were harvested live from the laboratory cultures, then stored in the refrigerator for 2 weeks) because we know that *Metschnikowia* spores lose infectivity over time, even in the refrigerator (Duffy & Hunsberger, 2019).

2.3 | Transmission spores exposure to toxins

On the first day of the experiment (day 1), we initiated the longest incubation treatment (10 days) by placing 5,000 mature transmission spores of *M. bicuspidata* in 15-ml Falcon tubes with 10 ml of lake water that had been filtered through a Pall AE filter (Pall Corporation). We then added microcystin-LR, anatoxin-a, acetic acid (negative control), or no toxin/chemical; for both microcystin-LR and anatoxin-a, we had two toxin levels: 10 and 30 µg/L. The concentrations we chose for this study are below and/or well within the range observed during natural CHABs blooms (Ha et al., 2009; Ibelings et al., 2005; Park et al., 1998; Pawlik-Skowrońska et al., 2004), and therefore ecologically relevant. These treatment doses are below the LC50s reported in previous *Daphnia* toxicology experiments for microcystins and anatoxin-a (DeMott et al., 1991; Pawlik-Skowrońska et al., 2019). Even though we did not expose hosts to these toxins, we chose these concentrations because they should not cause high levels of stress and mortality in *Daphnia*, so any impact to pathogens could have a substantial impact on parasite–host interactions. While these concentrations are likely to be on the high side of what spores are likely to encounter in nature, if there is no impact of the cyanotoxins at these levels, it suggests that they are unlikely to significantly impact the free-living stages of this parasite in the water column. There were 10 replicates of each treatment (including the negative control treatment of acetic acid), with the exception of the no toxin controls (0 µg/L), which had 20 replicates; this yielded a total of 70 experimental units per incubation time treatment. The tubes with spores were left uncapped for 10 days inside a large plastic tote covered with a lid at 20°C with a 16:8 light: dark (L:D) photoperiod. On day 4, the same procedure from day 1 was repeated. In this treatment, spores were incubated for 7 days at 20°C with a

16:8L:D photoperiod, yielding the 7-day incubation treatment. On day 6, the same procedure was used to initiate the 5-day incubation treatment. Additionally, on this day, adult *Daphnia* were set-up so that we could use their offspring in the infection assays. Adult *Daphnia* of the MID37 genotype were placed in 150-ml beakers (5 adults per beaker) with 100ml of filtered lake water. Beakers were placed in incubators at 20°C with a 16:8L:D photoperiod for 24 hr. On day 7, we collected neonates (0–24 hr old) produced from mothers that had been set-up the previous day. We placed 10 neonates per beaker in 250-ml beakers with 150ml filtered lake water for a total of 400 animals. Each beaker received 2mg C/L of *Ankistrodesmus falcatus* food and was placed in incubators at 20°C with a 16:8L:D photoperiod. After that, 2mg C/L of *Ankistrodesmus* was added to each beaker each day until the day the experimental animals were exposed to parasites. On days 8 and 10, we used the same procedure as described above to initiate the 3- and 1-day incubation treatments, respectively.

2.4 | Infection assays

On day 11, we carried out infection assays, in which we exposed *Daphnia* to transmission spores that had been incubated for different time lengths and with varying levels of exposure to toxins/chemicals. All Falcon tubes, containing spores with different exposure times and toxin levels, were collected and placed in a centrifuge (Sorvall ST 16; Thermo Scientific) to spin down the spores. The tubes were spun at 1,690 g for 10min. We decanted the tubes using a 10-ml pipette without disrupting the spore pellet at the bottom. In a pilot experiment, we confirmed that the original concentration of spores was recovered from vials after a 24-hr incubation followed by centrifugation. After the water was removed (which also removed the toxin), we resuspended the spores by adding 10ml of filtered lake water and disturbing the pellet by vigorously pipetting the water in the tube. Then, we placed one 6–7-day old *Daphnia* in each Falcon tube and allowed the tubes to incubate again at 20°C with 16:8L:D photoperiod. Because we had originally placed 5,000 mature transmission spores in each tube, this yielded an exposure dose of 500 spores/ml. Hosts were fed 1mg C/ml of *Ankistrodesmus* on this day; using this lower level of food on the day of exposure is common in infection assays because it promotes infection. After 24hr (that is, on day 12), each *Daphnia* individual was removed from the tubes with spores and placed in a 50-ml beaker with 30ml filtered lake water that did not contain spores (one animal per beaker). Animals were fed 2mg/L C of *A. falcatus* ad libitum for the rest of the experiment (20 days post-infection, 32 days from day 1 of entire experiment).

During those 20 days, we tracked mortality in each of the beakers 5 days per week (Mondays to Fridays). Water changes were done twice a week; during these, we counted offspring in each beaker; offspring were removed from beakers and discarded. Any animals that died during the trial were placed in 2-ml Eppendorf tubes with 100µl of nanopure water and stored at 4°C for later spore counts. At the end of the experiment (20 days post-infection, 32 days from day 1 of entire experiment), any remaining live animals (130 total) were placed in a 2-ml Eppendorf

tube with 100µl of nanopure water. Animals were then ground to release spores, and spores in the ascus stage were counted using a haemocytometer under a compound microscope at 400× magnification.

2.5 | Data analysis

For all our models, we analysed data from the microcystin-LR and anatoxin-a treatments separately. This means that for each analysis described below, there was one performed for the microcystin-LR relevant data and another for the anatoxin-a data. The same no-toxin control data (0µg/L; 20 replicates per incubation time) were used for the two sets of data (microcystin-LR and anatoxin-a). When analysing the anatoxin-a data, the acetic acid treatments were included in the analyses (as an additional negative control) and were treated in our analyses as a low concentration (0.01%); statistical analysis did not find differences between the no toxin controls and the acetic acid controls (data not shown). Prior to analyses, the data were checked for normality using the Shapiro–Wilk test. Data that did not meet normality were analysed using appropriate family error distribution link functions and checking for overdispersion. All analyses were carried out in R Studio Version 1.2.1335 using the stats v3.6.1 package.

We assessed environmental effects on the parasite's ability to infect by comparing differences in the number of animals that developed terminal infections (that is, infections that produced asci [Stewart Merrill & Cáceres, 2018]). In this analysis, our response variable had two outcomes (terminal infection or not). We performed a generalised linear model (GLM) with incubation time and toxin concentration as explanatory variables, using a binomial family error distribution.

We also evaluated if toxin concentration and incubation time (that is, the time spent as a transmission spore in the water) affected the number of mature transmission spores produced per infected host individual. For this analysis, we ran a GLM; here the number of spores was our response variable and, similarly to the analysis of infections, toxin concentration and incubation time were used as the explanatory variables. In this model, we used a Gaussian error distribution.

To evaluate the effects of incubation time and toxin exposure on the parasite's virulence, we measured host offspring production, host survival over the 20-day experiment, and proportion of hosts who died within 20 days of parasite exposure. For these analyses, we ran GLMs with toxin concentration and time of exposure as explanatory variables; for the two former response variables—the number of offspring each host individual produced and the number of days each host survived (up to 20 days post infection)—we used a Poisson family error distribution. For the proportion of dead hosts at the end of the experiment, we used a binomial family distribution.

We ran two linear models analysing the relationship between the proportion of infected individuals and the mean number of spores produced at the end of infection. In the first, we averaged across the different toxin exposure treatments, yielding one value for each incubation time ($n=5$). In the second, we averaged within the toxin

treatments, yielding five values per toxin treatment (one per incubation time treatment, total $n=20$).

We also ran a second experiment to evaluate whether plastic absorbed considerable amounts of microcystin in the water and therefore negated the toxic effects on the parasite's transmission spores. Studies have reported adsorption of microcystins by plastics (Hyenstrand et al., 2001; Moura et al., 2022), which raised the possibility that the results of our first experiment may have been due to toxin concentrations that were lower than we intended. Therefore, in the second experiment, we evaluated whether the plastic vessels we used in the first block adsorbed microcystin-LR from the water and whether using glass versus plastic vessels for the incubations impacted infections. Additional methods and results from that experiment can be found in the Appendix S1. Briefly, we did not find any effect of vessel (that is, glass vs. plastic tubes) on toxin concentration, infection prevalence, or spore production.

3 | RESULTS

3.1 | Time spent in the water decreased infectivity, but toxin exposure did not

Transmission spores that spent longer in the water were less infectious, but toxin exposure did not significantly influence infectivity (Figure 1, Table 1 Infection prevalence). The number of hosts that

became infected decreased with increased incubation time: spores that were only incubated for 24hr infected around 75% of exposed hosts, whereas spores incubated for 10days infected less than 50% of exposed hosts. The reductions in infectivity with increasing incubation time were consistent across different toxin concentrations (as indicated by a non-significant concentration \times incubation interaction term in the GLM: microcystins, $Z=1.160$, $p=0.246$; anatoxin, $Z=0.816$, $p=0.415$).

3.2 | Spore yield was not affected by toxin exposure

Neither microcystin nor anatoxin dose significantly influenced the number of spores produced per infected host (Figure 2; Table 1). However, in the anatoxin treatments (but not the microcystin treatments), infected hosts exposed to spores that were incubated for longer periods of time yielded more spores at death (or, for those that did not die within 20days of exposure, at 20days post-exposure; Figure 2; Table 1).

3.3 | Virulence was not affected by toxin concentration or incubation time

Neither toxin concentration nor incubation time impacted host offspring production, lifespan, or survivorship measured up to 20days post-infection. Host lifetime offspring production was consistent




FIGURE 1 Infection prevalence of hosts exposed to transmission spores treated with different concentrations of pure cyanotoxins and incubated for different amounts of time. This figure shows the proportion of individuals that became infected in each treatment group and standard errors. For both toxins and in all concentration treatments, the number of individual hosts that became infected decreased with the amount of time the transmission spores spent in the water before being used for host exposure. Note that the 0 $\mu\text{g/L}$ concentration treatment is the same set of data for both panels.

TABLE 1 Statistical results of generalised linear models on the effects of toxin concentration and incubation time on the parasites fitness and virulence.

Variable	Estimate	SE	Z-value	p-Value
Infection prevalence				
Microcystin-LR				
Toxin concentration	-0.012	0.025	-0.497	0.619
Incubation time	-0.382	0.083	-4.630	<0.0001*
Conc×Incub	0.005	0.005	1.160	0.246
Anatoxin-a				
Toxin concentration	-0.003	0.026	-0.107	0.915
Incubation time	-0.527	0.087	-6.071	<0.0001*
Conc×Incub	0.004	0.005	0.816	0.415
Spore reproduction				
Microcystin-LR				
Toxin concentration	466.76	386.33	1.208	0.231
Incubation time	2,230.73	1,528.03	1.460	0.149
Conc×Incub	-87.23	85.90	-1.016	0.313
Anatoxin-a				
Toxin concentration	326.10	355.22	0.918	0.361
Incubation time	4,337.31	1,508.30	2.876	<0.01*
Conc×Incub	-21.27	83.29	-0.255	0.799
Host lifetime reproduction				
Microcystin-LR				
Toxin concentration	-0.0007	0.005	-0.140	0.889
Incubation time	0.012	0.019	0.643	0.520
Conc×Incub	0.0008	0.001	0.790	0.430
Anatoxin-a				
Toxin concentration	-0.003	0.005	-0.753	0.452
Incubation time	0.007	0.019	0.353	0.724
Conc×Incub	0.0005	0.001	0.506	0.613
Host survivorship				
Microcystin-LR				
Toxin concentration	0.0008	0.004	0.227	0.820
Incubation time	0.013	0.015	0.854	0.393
Conc×Incub	-0.0002	0.0008	-0.327	0.744
Anatoxin-a				
Toxin concentration	7.074e-04	3.485e-03	0.203	0.839
Incubation time	1.451e-02	1.447e-02	1.003	0.316
Conc×Incub	-9.278e-05	7.997e-04	-0.116	0.908

Note: Host reproduction and host survivorship are used here as proxies of virulence. Conc×Incub indicates the toxin concentration×incubation time interaction term. * indicates significant at $p < 0.05$.

across the microcystin-LR treatments (Figure 3, left panel) and across the anatoxin-a treatments (Figure 3, right panel), and across incubation times (Figure 3, Table 1). Similarly, neither toxin concentration nor incubation time significantly influenced the number of days each host survived (up to 20 days post-infection) or the proportion of dead individuals at day 20 (Table 1). Thus, there is no evidence that exposure of transmission spores to toxins in the water, or the duration of time spent in the water column, influenced the virulence of this fungal parasite.

3.4 | Greater parasites infectiousness is associated with lower spore yield

Treatments that had higher infection prevalence yielded fewer spores per infected host (Figure 4). When averaged across the different toxin treatments within incubation times, longer incubation times had fewer infections that yielded more spores per infected host ($R^2 = 0.80$, $p = 0.02$). If we average within toxin treatments by




FIGURE 2 Parasite reproduction within infected hosts exposed to transmission spores treated with different concentrations of pure cyanotoxins that were incubated for different lengths of time. (Left panel) Spores that were treated with microcystin-LR did not appear to be affected by the concentration of toxin nor the amount of time these spores spend in the water. (Right panel) The number of mature transmission spores produced in all concentration treatments increased with longer periods of incubation time for transmission spores. Note that the 0 µg/L concentration treatment is the same set of data for both panels. Small symbols represent individual data points while large symbols represent means for a given treatment group.

incubation times, we observe a similar pattern but with a lower amount of variance explained ($R^2=0.40$, $p=0.001$).

4 | DISCUSSION

In this study, we found that the amount of time transmission spores spend in the water impacted their fitness. If spores encountered a new host within a few days of being released from their dead host, the number of hosts that became infected was high, suggesting high per-spore infectivity. However, the high prevalence did not translate into high spore yield; instead, hosts from treatments with high infectivity produced low numbers of mature transmission spores. By contrast, spores that spent longer in the water column after being released from a dead host infected relatively few hosts, but each infected host yielded more mature transmission spores as compared to a host infected with spores that spent less time in the water column.

A notable finding from our study is the absence of toxic effects of two common cyanotoxins on parasite transmission spores while in the water column. We predicted that long exposures to high concentrations of cyanotoxin would decrease parasite fitness and virulence. However, our analyses did not detect any effect of toxins. Previous studies have found that consuming toxin-producing cyanobacteria can make hosts more susceptible to parasites (Tellenbach et al., 2016), can decrease the number of spores produced within the host (Manzi et al., 2019) and, in certain cases,

protects hosts from infection (Sánchez et al., 2019). However, the conditions of parasite toxin exposure in the previous experiments were different from our current study. In the previous studies, parasites were exposed to these cyanobacteria as part of their diet, so interactions between the toxins and parasites would occur in the host gut. The key difference is that, in our study, the toxin exposure only occurred in the water column, and hosts were not exposed to these compounds. This suggests that the location of exposure of spores to cyanotoxins is important. One possible reason why we did not observe effects of toxins on parasite fitness might be the structure of the transmission spore. *Metschnikowia* transmission spores are composed of an ascus, which is an outer structure surrounding the ascospore; the ascospore is the actual transmission spore that pierces the host gut and therefore causes the infection. One possibility is that the ascus, which protects the spore from chemical and physical damage (Lachance et al., 1976), also shields against secondary metabolites in the water, which would explain the lack of effects from the cyanotoxins in this study. Taken together, the results of our study combined with other studies that have found effects of toxin and non-toxin producing cyanobacteria on parasite fitness (Coopman et al., 2014; Manzi et al., 2019; Sánchez et al., 2019; Tellenbach et al., 2016) suggest that the interaction between parasites and secondary metabolites may be particularly important within the gut of the host. If true, this would suggest that, for parasites with a protective structure that envelops




FIGURE 3 Lifetime reproduction of hosts that were infected with transmission spores treated with different concentrations of pure cyanotoxins and incubated for different lengths of time. Virulence of the parasite did not change based on the transmission spore treatments. Note that the 0 µg/L concentration treatment is the same set of data for both panels. Small symbols represent individual data points while large symbols represent means for a given treatment group.

transmission spores, the interaction of toxins with parasites might primarily be important in parasite life stages that lack the protective coating.

There was an apparent trade-off between per-spore infectivity and the number of spores yielded per infected host. One mechanism that could explain this result is within-host spore competition (Ebert et al., 2000). Spores that spent little time in the water column had a high per-spore infectivity, which probably led to multiple spores infecting an individual host; individual *Daphnia* can be infected by multiple *Metschnikowia* spores (Stewart Merrill & Cáceres, 2018). Infection by multiple spores would result in resource competition among spores inside the host, decreasing parasite spore production. In contrast, spores that spent longer in the water column were less viable; this would mean fewer spores were able to infect an individual host, decreasing resource competition and increasing within host reproduction. Prior work has shown that not all spores that make it to the gut pierce the gut—a critical first step of infection that must occur prior to growing in the host haemolymph (Stewart Merrill & Cáceres, 2018). Our results and this hypothesised mechanism are consistent with an earlier study using the same parasite but a different species of *Daphnia*; that study found parasite reproductive success decreased with increased spore dosage at the time of infection (Ebert et al., 2000). It is also possible that spores that survived longer were, on average, stronger than the spores that died during the incubation, meaning a non-random sample of spores generated the infections in the longer incubation periods.

It remains to be determined why spores lost infectivity so rapidly in the water column. One possibility is that light played a role. Our parasite incubations for the toxin exposures were done in the light (using a 16:8 L:D photoperiod), and other studies have found that light, including UV and photosynthetically active radiation, decreases infectivity of *Metschnikowia* spores (Overholt et al., 2012; Shaw et al., 2020). Therefore, spores may have retained high infectivity for longer if our experiment had been done in the dark. However, in a previous experiment, even spores that were maintained in the refrigerator in the dark rapidly lost infectivity (Duffy & Hunsberger, 2019); thus, it is possible that multiple factors explain the rapid loss in infectivity of *Metschnikowia* spores in the water column.

One limitation of our study is that we only used a single host genotype and a single parasite genotype. There is substantial diversity in the susceptibility of *D. dentifera* to *Metschnikowia* (e.g., Auld et al., 2017; Duffy et al., 2012; Duffy & Sivers-Becker, 2007). While there is much less diversity in *Metschnikowia*, there is some (Shaw, 2019). Future work exploring whether *Metschnikowia* genotypes vary in their sensitivity to phycotoxins and/or time spent in the water column would be valuable, as would studies assessing whether the patterns we found in this study are consistent when other host genotypes (and species) are exposed. Interestingly, a new study that incubated *Metschnikowia* spores with cyanobacterial extracts or a control solution, then exposed them to two genotypes of *Daphnia galeata* × *longispina* (Manzi et al., 2022) suggests that these results might hold broadly. Consistent with our study, they did not




FIGURE 4 Linear regressions of the proportion of individuals that became infected versus the mean number of mature transmission spores produced in each treatment group. (Left panel) The data points in this regression are averaged by incubation time of transmission spores in all toxin treatments. The number in the label for each point represents the number of days those spores were incubated, $R^2=0.80$, $p=0.02$. (Right panel) This correlation was done by averaging the data by toxin treatment and incubation period, $R^2=0.40$, $p=0.001$. For both correlations, we observed a decreasing number of spores produced within a host as the number of individuals that become infected in a treatment increased. Error bars in both panels represent standard error for infection prevalence and spore reproduction for the horizontal and vertical bars, respectively.

find reduced infectivity of spores that had been incubated with the cyanobacterial extract; in fact, for one of the two host genotypes, infectivity of spores that had been incubated with the cyanobacterial extract was actually higher.

Here, we found that parasite spores rapidly lose infectivity in the water, but that two common cyanotoxins had no detectable effect on spores. We propose that *Metschnikowia* spores may be protected from toxins in the environment by their ascus structure. We also found that, while spores that had spent more time in the water were less infectious, they yielded more spores per infected host, results that might be driven by resource competition. This raises the intriguing possibility that there might be an intermediate spore age (or dose) that is most likely to fuel large epidemics. Given that spores spend substantial time in the sediment between outbreaks (Decaestecker et al., 2004), the joint impacts of these mechanisms are likely to be important drivers of *Metschnikowia* epidemics in lakes.

AUTHOR CONTRIBUTIONS

Conceptualisation: K.F.S., M.A.D. Developing methods: K.F.S., M.A.D. Conducting the research: K.F.S., B.Z., J.A.A. Data analysis: K.F.S., B.Z., J.A.A., M.A.D. Data interpretation: K.F.S., B.Z., J.A.A., M.A.D. Preparation of figures and tables: K.F.S., B.Z., J.A.A. Writing: K.F.S., B.Z., J.A.A., M.A.D.

ACKNOWLEDGMENTS

We would like to thank M. Hunter for providing lab equipment and space and for his valuable input, P. Yajnik for his advice on statistical analysis, and G. Kling and four anonymous reviewers for helpful feedback on earlier drafts of this manuscript. K. Sánchez was supported by an NSF Graduate Research Fellowship. B. Zhong and J. Agudelo were supported by the Doris Duke Conservation Scholars Program at the University of Michigan. This work was supported by NSF grant DEB-1655856 to MAD, Block Grant funds from the University of Michigan Department of Ecology & Evolutionary Biology, and by the Moore Foundation (GBMF9202; DOI: [10.37807/GBMF9202](https://doi.org/10.37807/GBMF9202)).

DATA AVAILABILITY STATEMENT

Data is accessible through DRYAD digital repository [10.5061/dryad.612jm6420](https://doi.org/10.5061/dryad.612jm6420).

ORCID

Kristel F. Sánchez [ORCID](https://orcid.org/0000-0003-4273-6887) <https://orcid.org/0000-0003-4273-6887>

Baili Zhong [ORCID](https://orcid.org/0000-0002-1801-385X) <https://orcid.org/0000-0002-1801-385X>

Meghan A. Duffy [ORCID](https://orcid.org/0000-0002-8142-0802) <https://orcid.org/0000-0002-8142-0802>

REFERENCES

Amigó, J. M., Gracia, M. P., Rius, M., Salvadó, H., Maillo, P. A., & Vivarés, C. P. (1996). Longevity and effects of temperature on the viability

- and polar-tube extrusion of spores of *Glugea stephani*, a microsporidian parasite of commercial flatfish. *Parasitology Research*, 82, 211–214. <https://doi.org/10.1007/s004360050097>
- Andersen, N. G., Lorenzen, E., Boutrup, T. S., Hansen, P. J., & Lorenzen, N. (2016). Sublethal concentrations of ichthyotoxic alga *Prymnesium parvum* affect rainbow trout susceptibility to viral haemorrhagic septicaemia virus. *Diseases of Aquatic Organisms*, 117, 187–195. <https://doi.org/10.3354/dao02946>
- Auld, S. K. J. R., Hall, S. R., & Duffy, M. A. (2012). Epidemiology of a *Daphnia*-multiparasite system and its implications for the red queen. *PLoS One*, 7, e39564. <https://doi.org/10.1371/journal.pone.0039564>
- Auld, S. K. J. R., Hall, S. R., Ochs, J. H., Sebastian, M., & Duffy, M. A. (2014). Predators and patterns of within-host growth can mediate both among-host competition and evolution of transmission potential of parasites. *The American Naturalist*, 184, S77–S90. <https://doi.org/10.1086/676927>
- Auld, S. K. J. R., Searle, C. L., & Duffy, M. A. (2017). Parasite transmission in a natural multihost-multiparasite community. *Philosophical Transactions of the Royal Society, B: Biological Sciences*, 372, 20160097. <https://doi.org/10.1098/rstb.2016.0097>
- Borowitzka, M. A. (1995). Microalgae as sources of pharmaceuticals and other biologically active compounds. *Journal of Applied Phycology*, 7, 3–15. <https://doi.org/10.1007/BF00003544>
- Brookes, J. D., Antenucci, J., Hipsey, M., Burch, M. D., Ashbolt, N. J., & Ferguson, C. (2004). Fate and transport of pathogens in lakes and reservoirs. *Environment International*, 30, 741–759. <https://doi.org/10.1016/j.envint.2003.11.006>
- Cáceres, C. E., Hall, S. R., Duffy, M. A., Tessier, A. J., Helmle, C., & Macintyre, S. (2006). Physical structure of lakes constrains epidemics in *daphnia* populations. *Ecology*, 87, 1438–1444.
- Christensen, V. G., & Khan, E. (2020). Freshwater neurotoxins and concerns for human, animal, and ecosystem health: A review of anatoxin-a and saxitoxin. *Science of the Total Environment*, 736, 139515. <https://doi.org/10.1016/j.scitotenv.2020.139515>
- Codreanu, R., & Codreanu-Balcescu, D. (1981). On two *Metschnikowia* yeast species producing infections in *Daphnia magna* and *Artemia salina* (Crustacea, Phyllozoa) from Romania. *Journal of Invertebrate Pathology*, 37, 22–27.
- Coopman, M., Muylaert, K., Lange, B., Reyserhove, L., & Decaestecker, E. (2014). Context dependency of infectious disease: The cyanobacterium *Microcystis aeruginosa* decreases white bacterial disease in *Daphnia magna*. *Freshwater Biology*, 59, 714–723. <https://doi.org/10.1111/fwb.12420>
- Decaestecker, E., Lefever, C., De Meester, L., & Ebert, D. (2004). Haunted by the past: Evidence for dormant stage banks of microparasites and epibiotics of *Daphnia*. *Limnology and Oceanography*, 49, 1355–1364. https://doi.org/10.4319/lo.2004.49.4_part_2.1355
- DeMott, W. R., Zhang, Q., & Carmichael, W. W. (1991). Effects of toxic cyanobacteria and purified toxins on the survival and feeding of a copepod and three species of *Daphnia*. *Limnology and Oceanography*, 36, 1346–1357.
- Duffy, M. A., & Hunsberger, K. K. (2019). Infectivity is influenced by parasite spore age and exposure to freezing: Do shallow waters provide *Daphnia* a refuge from some parasites? *Journal of Plankton Research*, 41, 12–16. <https://doi.org/10.1093/plankt/fby046>
- Duffy, M. A., Ochs, J. H., Penczykowski, R. M., Civitello, D. J., Klausmeier, C. A., & Hall, S. R. (2012). Size and parasite-driven evolution. *Science*, 335, 1636–1638.
- Duffy, M. A., & Sivars-Becker, L. (2007). Rapid evolution and ecological host-parasite dynamics. *Ecology Letters*, 10, 44–53. <https://doi.org/10.1111/j.1461-0248.2006.00995.x>
- Duperron, S., Halary, S., Habiballah, M., Gallet, A., Huet, H., Duval, C., Bernard, C., & Marie, B. (2019). Response of fish gut microbiota to toxin-containing cyanobacterial extracts: A microcosm study on the medaka (*Oryzias latipes*). *Environmental Science & Technology Letters*, 6, 341–347. <https://doi.org/10.1021/acs.estlett.9b00297>
- Ebert, D. (2005). *Ecology, epidemiology, and evolution of parasitism in Daphnia*. National Center for Biotechnology Information.
- Ebert, D., Zschokke-Rohringer, C. D., & Hans, J. C. (2000). Dose effects and density-dependent regulation of two microparasites of *Daphnia magna*. *Ecology*, 122, 200–209.
- Ha, J. H., Hidaka, T., & Tsuno, H. (2009). Quantification of toxic microcystin and evaluation of its dominance ratio in blooms using real-time PCR. *Environmental Science and Technology*, 43, 812–818. <https://doi.org/10.1021/es801265f>
- Hall, S. R., Smyth, R., Becker, C. R., Duffy, M. A., Knight, C. J., MacIntyre, S., Tessier, A. J., & Cáceres, C. E. (2010). Why are *Daphnia* in some lakes sicker? Disease ecology, habitat structure, and the plankton. *Bioscience*, 60, 363–375. <https://doi.org/10.1525/bio.2010.60.5.6>
- Hallmann, J., & Sikora, R. A. (1996). Toxicity of fungal endophyte secondary metabolites to plant parasitic nematodes and soil-borne plant pathogenic fungi. *European Journal of Plant Pathology*, 102, 155–162.
- Harvell, D., Aronson, R., Baron, N., Connell, J., Dobson, A., Ellner, S., Gerber, L., Kim, K., Kuris, A., McCallum, H., Lafferty, K., McKay, B., Porter, J., Pascual, M., Smith, G., Sutherland, K., & Ward, J. (2004). The rising tide of ocean diseases: Unsolved problems and research priorities. *Frontiers in Ecology and the Environment*, 2, 375–382.
- Heagle, A. S. (1973). Interactions between air pollutants and plant parasites. *Annual Review of Phytopathology*, 11, 365–388. <https://doi.org/10.1146/annurev.py.11.090173.002053>
- Huisman, J., Codd, G. A., Paerl, H. W., Ibelings, B. W., Verspagen, J. M. H., & Visser, P. M. (2018). Cyanobacterial blooms. *Nature Reviews Microbiology*, 16, 471–483. <https://doi.org/10.1038/s41579-018-0040-1>
- Hyenstrand, P., Metcalf, J., Beattie, K., & Codd, G. A. (2001). Effects of adsorption to plastics and solvent conditions in the analysis of the cyanobacterial toxin microcystin-LR by high performance liquid chromatography. *Water Research*, 35, 3508–3511. [https://doi.org/10.1016/S0043-1354\(01\)00068-9](https://doi.org/10.1016/S0043-1354(01)00068-9)
- Ibelings, B. W., Bruning, K., de Jonge, J., Wolfstein, K., Pires, L. M. D., Postma, J., & Burger, T. (2005). Distribution of microcystins in a Lake Foodweb: No evidence for biomagnification. *Microbial Ecology*, 49, 487–500. <https://doi.org/10.1007/s00248-004-0014-x>
- Lachance, M.-A., Miranda, M., Miller, M. W., & Phaff, H. J. (1976). Dehiscence and active spore release in pathogenic strains of the yeast *Metschnikowia bicuspidatavar. Australis*: Possible predatory implication. *Canadian Journal of Microbiology*, 22, 1756–1761.
- Lafferty, K. D., & Holt, R. D. (2003). How should environmental stress affect the population dynamics of disease? *Ecology Letters*, 6, 654–664.
- Landsberg, J. H. (2002). The effects of harmful algal blooms on aquatic organisms. *Reviews in Fisheries Science*, 10, 1064–1262. <https://doi.org/10.1080/20026491051695>
- Lassudrie, M., Hégaret, H., Wikfors, G. H., & da Silva, P. M. (2020). Effects of marine harmful algal blooms on bivalve cellular immunity and infectious diseases: A review. *Developmental and Comparative Immunology*, 108, 103660. <https://doi.org/10.1016/j.dci.2020.103660>
- Leflaive, J., & Ten-Hage, L. (2007). Algal and cyanobacterial secondary metabolites in freshwaters: A comparison of allelopathic compounds and toxins. *Freshwater Biology*, 52, 199–214. <https://doi.org/10.1111/j.1365-2427.2006.01689.x>
- Malone, L. A., Gatehouse, H. S., & Tregidga, E. L. (2001). Effects of time, temperature, and honey on *Nosema apis* (Microsporidia: Nosematidae), a parasite of the honeybee, *Apis mellifera* (Hymenoptera: Apidae). *Journal of Invertebrate Pathology*, 77, 258–268. <https://doi.org/10.1006/jipa.2001.5028>
- Manzi, F., Agha, R., Lu, Y., Ben-Ami, F., & Wolinska, J. (2019). Temperature and host diet jointly influence the outcome of infection in a

- Daphnia-fungal parasite system. *Freshwater Biology*, 65, 757–767. <https://doi.org/10.1111/fwb.13464>
- Manzi, F., Agha, R., Mühlenhaupt, M., & Wolinska, J. (2022). Prior exposure of a fungal parasite to cyanobacterial extracts does not impair infection of its Daphnia host. *Hydrobiologia*, 849, 2731–2744. <https://doi.org/10.1007/s10750-022-04889-7>
- Metschnikoff, V. E. (1884). Ueber eine Sprosspilzkrankheit der Daphnien Beitrag zur Lehre über den Kampf der Phagocyten gegen Krankheitserreger. *Archiv für Pathologische Anatomie und Physiologie und für Klinische Medizin*, 96, 177–195. <https://doi.org/10.1007/BF02361555>
- Moura, D. S., Pestana, C. J., Moffat, C. F., Hui, J., Irvine, J. T., Edwards, C., & Lawton, L. A. (2022). Adsorption of cyanotoxins on polypropylene and polyethylene terephthalate: Microplastics as vector of eight microcystin analogues. *Environmental Pollution*, 303, 119135. <https://doi.org/10.1016/j.envpol.2022.119135>
- Ostensvik, O., Skulberg, O. M., Underdal, B., & Hormazabal, V. (2002). Antibacterial properties of extracts from selected planktonic freshwater cyanobacteria – A comparative study of bacterial bioassays. *Journal of Applied Microbiology*, 84, 1117–1124.
- Overholt, E. P., Hall, S. R., Williamson, C. E., Meikle, C. K., Duffy, M. A., & Cáceres, C. E. (2012). Solar radiation decreases parasitism in Daphnia. *Ecology Letters*, 15, 47–54. <https://doi.org/10.1111/j.1461-0248.2011.01707.x>
- Park, H. D., Kim, B., Kim, E., & Okino, T. (1998). Hepatotoxic microcystins and neurotoxic anatoxin-a in cyanobacterial blooms from Korean lakes. *Environmental Toxicology and Water Quality*, 13, 225–234. [https://doi.org/10.1002/\(SICI\)1098-2256\(1998\)13:3<225::AID-TOX4>3.0.CO;2-9](https://doi.org/10.1002/(SICI)1098-2256(1998)13:3<225::AID-TOX4>3.0.CO;2-9)
- Pawlik-Skowrońska, B., Skowroński, T., Pirszel, J., & Adamczyk, A. (2004). Relationship between cyanobacterial bloom composition and anatoxin-A and microcystin occurrence in the eutrophic dam reservoir (se Poland). *Polish Journal of Ecology*, 52, 479–490.
- Pawlik-Skowrońska, B., Toporowska, M., & Mazur-Marzec, H. (2019). Effects of secondary metabolites produced by different cyanobacterial populations on the freshwater zooplankters *Brachionus calyciflorus* and *Daphnia pulex*. *Environmental Science and Pollution Research*, 26, 11793–11804. <https://doi.org/10.1007/s11356-019-04543-1>
- Penczykowski, R. M., Lemanski, B. C. P., Sieg, R. D., Hall, S. R., Housley, O. J., Kubanek, J., & Duffy, M. A. (2014). Poor resource quality lowers transmission potential by changing foraging behaviour. *Functional Ecology*, 28, 1245–1255. <https://doi.org/10.1111/1365-2435.12238>
- Sánchez, K. F., Huntley, N., Duffy, M. A., & Hunter, M. D. (2019). Toxins or medicines? Phytoplankton diets mediate host and parasite fitness in a freshwater system. *Proceedings of the Royal Society B: Biological Sciences*, 286, 20182231. <https://doi.org/10.1098/rspb.2018.2231>
- Shaw, C. L. (2019). *Drivers of epidemic timing and size in a natural aquatic system*. University of Michigan.
- Shaw, C. L., Overholt, E., Williamson, C., Cáceres, C. E., Hall, S. R., & Duffy, M. A. (2020). Shedding light on environmentally transmitted parasites: Lighter conditions within lakes restrict epidemic size. *Ecology*, 101, e03168. <https://doi.org/10.1002/ECY.3168>
- Stewart Merrill, T. E., & Cáceres, C. E. (2018). Within-host complexity of a plankton-parasite interaction. *Ecology*, 99, 2864–2867. <https://doi.org/10.1002/ecy.2483>
- Tellenbach, C., Tardent, N., Pomati, F., Keller, B., Hairston, N. G., Wolinska, J., & Spaak, P. (2016). Cyanobacteria facilitate parasite epidemics in daphnia. *Ecology*, 97, 3422–3432. <https://doi.org/10.1002/ecy.1576>
- Tessier, A. J., & Woodruff, P. (2002). Cryptic trophic cascade along a gradient of lake size. *Ecology*, 83, 1263–1270.
- Thieltges, D. W., Amundsen, P.-A., Hechinger, R. F., Johnson, P. T. J., Lafferty, K. D., Mouritsen, K. N., Preston, D. L., Reise, K., Zander, C. D., & Poulin, R. (2013). Parasites as prey in aquatic food webs: Implications for predator infection and parasite transmission. *Oikos*, 122, 1473–1482. <https://doi.org/10.1111/j.1600-0706.2013.00243.x>
- Vasemagi, A., Visse, M., & Kisand, V. (2017). Effect of environmental factors and an emerging parasitic disease on gut microbiome of wild salmonid fish. *mSphere*, 2, e00418–17. <https://doi.org/10.1128/mSphere.00418-17>
- Volk, R. B., & Furkert, F. H. (2006). Antialgal, antibacterial and antifungal activity of two metabolites produced and excreted by cyanobacteria during growth. *Microbiological Research*, 161, 180–186. <https://doi.org/10.1016/j.micres.2005.08.005>

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Sánchez, K. F., Zhong, B., Agudelo, J. A., & Duffy, M. A. (2023). Infectivity of the parasite *Metschnikowia bicuspidata* is decreased by time spent as a transmission spore, but exposure to phycotoxins in the water column has no effect. *Freshwater Biology*, 68, 1020–1030. <https://doi.org/10.1111/fwb.14082>