



CASE SERIES

Minimally invasive gingival phenotype modification in gingival recession associated with a non-carious cervical lesion using the root plastique technique: A quasi-experimental one-group pretest-posttest study

Luigi Romano¹ | Michele Paolantonio¹ | Paolo De Ninis² |
Muhammad H. A. Saleh³ | Bruna Sinjari¹ | Edit Xhajanka⁴ |
Beatrice Femminella¹ | Hom-lay Wang³ | Andrea Ravidá⁵

¹Department of Medical, Oral, and Biotechnological Sciences, G. D'Annunzio University, Chieti-Pescara, Italy

²Luisa D'Annunzio Institute for High Culture, Pescara, Italy

³Department of Periodontics and Oral Medicine, University of Michigan School of Dentistry, Ann Arbor, Michigan, USA

⁴Department of Dental Medicine, Medical University of Tirana, Rruga e Dibrës, Tirana, Albania

⁵Department of Periodontics & Oral Medicine, University of Pittsburgh, Pittsburgh, Pennsylvania, USA

Correspondence

Andrea Ravidá, Department of Periodontics & Oral Medicine, North University Avenue, University of Pittsburgh, Pittsburgh, PA, USA.
Email: Andrearavida@pitt.edu

Abstract

Background: This study introduces the root plastique technique (RPT), the aim of which is to modify the gingival phenotype of sites with gingival recessions (GRs) associated with non-carious cervical lesions (NCCLs) prior to surgical treatment.

Methods: RPT was performed in 22 subjects with 53 RT1 A/B + GRs. Changes in keratinized tissue thickness (KTT), keratinized tissue width (KTW), relative gingival recession (RGR), relative clinical attachment level (RCAL), and probing pocket depth (PPD) were measured at baseline (T0) and 2 months (T1) after the procedure was performed. All analyses were performed by means of hierarchical models.

Results: The study revealed statistically significant changes ($P < 0.01$) in KTT (0.45 ± 0.04 mm), RGR (0.80 ± 0.13 mm), KTW (0.67 ± 0.07 mm), and RCAL (-0.72 ± 0.16 mm). No changes in PPD ($P > 0.05$) were observed. Regression analyses of KTT increase and RGR reduction at T0 showed statistically significant correlation between the two variables ($P < 0.05$). All the teeth with a KTT of < 0.8 mm at T0 ($N = 14$) reached or surpassed this threshold at T1.

Conclusion: RPT increases KTT and KTW. In most of the sites, a reduction in GR was also achieved.

KEYWORDS

cemento-enamel junction, gingival recession, oral mucosa, tooth abrasion

1 | INTRODUCTION

Gingival recession (GR) is defined as the apical shift of the gingival margin with a consequent exposure of the root.¹ Mid-buccal GRs affect almost the entire population of the United States, with an approximate preva-

lence of 91% across the oral cavity and 70% when only the aesthetic zone is considered.² The attachment loss associated with GR may lead to an increased risk for root caries,³ tooth sensitivity, and compromised patient aesthetics.^{4,5} Multiple morphological and functional factors have been associated with the etiopathogenesis of



GR including toothbrush-related trauma, inflammation induced by plaque accumulation, orthodontic tooth movement, and lack of keratinized tissue thickness (KTT).⁶ GRs are frequently concomitant with wedge-shaped, concave, flattened, and irregular defects in the cervical area of the teeth known as non-carious cervical lesions (NCCLs).^{6,7} The periodontal literature reports that the prevalence of NCCLs ranges from 11% to 62% in the general population.^{8,9} As a result of the combined defects with GRs, the probability of achieving complete root coverage following root coverage therapy is decreased.¹⁰ The presence of an NCCL leads to alterations in both crown and root surface that often develop into root concavities (or steps) of variable extensions and depths, with the possibility of an indistinguishable cemento-enamel junction (CEJ).¹¹ Pini-Prato and coworkers classified the presence/absence of the CEJ as Class A (detectable CEJ) or Class B (undetectable CEJ), and the presence/absence of a cervical step at the apical extent of the NCCL into either the presence of a cervical step that is >0.5 mm (denoted as a +) or the absence of a cervical step (denoted as a -).¹¹ As aforementioned, where an NCCL is present, surgical root coverage procedures possess a lesser chance of success.¹⁰ In two randomized clinical trials investigating the surgical coverage of an NCCL, restorative materials in conjunction with a coronally advanced flap (CAF) with or without a subepithelial connective tissue graft (CTG) were used.^{12,13} The addition of a CTG demonstrated higher combined defect coverage (83%) compared with a CAF alone (43%), most likely due to the increased tissue thickness provided by the CTG.¹⁴ However, multiple studies revealed that the use of a CTG could be avoided at sites with tissue thickness >0.8 mm, where the use of a CAF alone is associated with similar clinical outcomes and superior aesthetics.^{8,14,15} Hence, the aim of the present study is to present in a cohort of consecutively treated patients, a minimally invasive technique (root plastique technique; RPT) which aims to modify the gingival phenotype of sites with GRs associated with NCCLs prior to CAF.

2 | MATERIALS AND METHODS

The one-group pretest-posttest design is a type of quasi-experiment study in which the outcome of interest is measured two times: once before and once after exposing a non-random group of participants to a certain intervention/treatment. This study was conceived as a proof of concept for a new treatment. While aware that a randomized controlled study would be needed to make a confirmatory claim, we realized that blinding the examiner would be impossible in this case. Hence, we performed this exploratory trial to study the new treatment efficacy

in detail by relying on the properties of its within-subject design, in which each patient is the control of himself. We designed this study specifying all the requirements needed for a future pivotal study. This investigation was conducted in agreement with the Helsinki Declaration of 1975 (World Medical Association, 1975) as revised in 2013 (World Medical Association, 2013). The protocol was approved by the Medical University of Tirana, Department of Dental Medicine, Institutional Review Board for Human Studies. Participants received an information sheet and provided their informed consent in accordance with the EU General Data Protection Regulation GDPR (UE) n. 2016/679 before beginning the rehabilitation. This prospective pilot study involved consecutive patients screened and treated in the time between February 2019 and October 2020.

2.1 | Study population

The present study was conducted in combined defects (GR associated with an NCCL). The patients were selected according to the following eligibility criteria: (1) Presence of an ≥ 1 RT1 plus A/B+ NCCL combined defect; (2) Absence of caries and/or restorations in the area to be treated; (3) Absence of mucogingival surgery or surgical treatment for periodontal disease at the same site in the previous 12 months; (4) Full-mouth plaque score (FMPS) $<20\%$ and full-mouth bleeding score (FMBS) $<20\%$ ¹⁶; and (5) Probing Pocket depth (PPD) of <4 mm. Patients were excluded if: (1) they suffered any major systemic disease that may interfere with healing; (2) they were smokers or recreational drug users; (3) the defect was an RT1 GR without the presence of a gingival step (4) the GR was a class RT2 or RT3 defect; (5) there was evidence of trauma for occlusion in the area to be treated.

2.2 | Clinical assessments

Before data acquisition, each patient underwent a professional mechanical plaque removal session and individualized oral hygiene instructions. After this initial therapy, the following parameters were recorded:

1. Probing pocket depth (PPD): Assessed as the distance from the gingival margin to the apical end of the gingival sulcus.
2. Relative gingival recession (RGR): Measured as distance from the gingival margin to the incisal border of the tooth of the involved tooth.¹⁷
3. Relative clinical attachment level (RCAL): Measured as distance from the apical end of the gingival sulcus to the incisal border of the involved tooth.¹⁷

4. Relative cervical lesion apical step (RCLAS): Measured as distance from the lower margin of the NCCL to the incisal border of the involved tooth.
5. Non-carious cervical lesion depth (CLD): Distance between the deepest point on the facial wall of the NCCL and the coronal projection of the most external point of the apical border of the non-carious cervical lesion. This parameter was measured using an endodontic k file, a silicone stop fixed with a cyanoacrylate adhesive, and a digital caliper.¹⁰
6. Keratinized tissue width (KTW): Measured as the distance from the gingival margin to the mucogingival junction with a periodontal probe (University of North Carolina Probe).
7. KTT: Gingival thickness at baseline was measured 1.5 mm apical to the gingival margin using an endodontic k file inserted perpendicular to the tissue surface with a silicon stop over the gingival surface. The silicon disc stop was then placed in tight contact with the soft tissue surface and fixed with a drop of cyanoacrylate adhesive, and a digital caliper used to measure the distance between the tip of the file and the stopper.¹⁴

Figure 1 schematically depicts the measurements performed in the present study. All measurements were performed at the time of the minimally invasive therapy (T0) under 5× magnification loops. All parameters were measured at baseline. PPD, RGR, RCAL, KTW, and KTT were also measured at 2 months after the procedure was performed (T1) by a single examiner (LR). Prior to the beginning of the clinical study, the examiner (LR) measured the RGR and RCAL of all patients twice within 24 h, with at least 60 min between examinations. The Kappa index was calculated for PD and KKT, resulting in 94% and 97% of reproducibility, and intraclass correlation was calculated for RGR, resulting in 92% of agreement.¹⁰

2.3 | Minimally invasive procedure

The technique performed is simple and fast, resonating the canons of minimally invasive therapies. After the administration of local anesthesia, the apical step of the NCCL (Figure 2A) was eliminated using rotary burs mounted on a low-speed handpiece. The first cutter used was a rugby ball-shaped cutter with a diameter of 0.16 white ring, ultra-fine grain with a maximum grain size of 14 microns (Figure 2B). The bur was inserted into the sulcus under abundant irrigation, until it invades the supracrestal tissue attachment (i.e., biological width). This was achieved by calculating the measurement of RBL and PPD. Eliminating the apical step was done in such a way to flatten the NCCL only in the apical portion. Namely, the coronal projection

of the most external point of the apical border of the NCCL will be removed, reaching with the bur the bottom of the NCCL. In this way, the depth of the NCCL was eliminated (Figure 2C). Subsequently, a final preparation drill with a diameter of 0.12 white ring, ultrafine grain with a maximum grain size of 8 microns was used (Figure 2D). The bone crest was reached using the bur gently while checking the working depth with the probe. An ultra-fine finishing bur was used to ensure minimal cutting on the expense of the soft tissue and maximum smoothness of the root surface.

The use of this drill made it possible to refine and extend the preparation of the root from the mesial line angle to the distal line angle of the tooth affected (Figure 2E). After the passage of the diamond burs, a flame Arkansas bur was used to refine and polish the root surface. The whole procedure was performed with the aid of a 5× magnifying lens. Through the invasion of supracrestal tissue attachment, the stimulation of bleeding was obtained and never required the use of sutures. This procedure was performed to eliminate any obstacle to the ascent of the gingival tissue and, at the same time, to create the space needed for the thickening of the soft tissues, a space that was initially filled only by the blood clot (Figure 2F). It was recommended that patients adopt a soft and cold diet following the procedure, avoiding biting any food with the treated teeth. All patients were instructed to stop brushing the site and the two contiguous teeth (mesial and distal) for 30 days to avoid mechanical trauma to the tissues. The area was cleaned with chlorhexidine gel two times a day instead. The patient was reassured that the brownish discoloration of the tooth surface was extrinsic and reversible after a professional oral hygiene session. After 15 and 30 days the patient was re-evaluated (professional plaque control and oral hygiene instruction). After 2 months, all patients were examined, and clinical parameters reassessed (Figure 2G–2H). A professional oral hygiene session was also performed. At that time, all the patients reinstated brushing in the affected area using an ultra-soft toothbrush for 2 weeks, then a soft toothbrush for another month.

2.4 | Sample size

According to our insight of the biological phenomenon, the population SD of the GT Gain in the most well-known treatments lies somewhere in between the 0.09 estimate of the CAF treatment,¹⁸ the one having the smaller probabilities of a substantial change, and the 0.47 of CTG treatment,¹⁹ which shows a greater variance probably because of the biological material addition. Even though an intermediate value seemed the most likely, the upper bound of this range was entered for the sample-size




FIGURE 1 (A-B) Schematic drawings illustrating the clinical anatomic characteristics measured at the buccal aspect of the defect

calculation, to warrant a sufficient power for the worst-case scenario. The minimal clinically important difference to detect was deemed 0.3 mm since it appeared the convenient size to bring cases with baseline thin phenotype around 0.5 mm up to the 0.8 mm threshold individuated by Cairo¹⁴ as guarantee of a better prognosis against recession relapses. A sample-size of $N = 22$ patients was calculated for a single-level analysis, setting $\alpha = 0.05$ and $\beta = 0.2$. Assuming a mean cluster-size = 2.5, 22 patients ensure to the highest level of the mixed model an effective sample-size equivalent to 26.²⁰

2.5 | Statistical analyses

All analyses were performed by means of Mixed Effect Models. A model with maximal random effects structure²¹

entailed a time effect with two repeated measurements within tooth site at the first level, tooth site within patient at the second level and the patient at the third level. After the selection procedure, only an intercept per tooth site and an intercept and a time slope, constrained to be equal, per patient were added as random effects since any other parameter would result in over-specified models. For most regressions, the Time factor was the only included fixed effect. Additional predictors as age, sex and tooth type most often resulted in not significant coefficients, so they were dropped out from the relative models. Only for some response variables the tooth type main effect and time by tooth type interaction proved to be significant, so were included. The time by tooth type interaction, although not significant, was included in models aimed to get more realistic effect predictions having the pre-post change as




FIGURE 2 Step-by-step description of the root plastique technique

response and baseline values as covariate. The R 4.0.1 software package was used.[‡]

A power analysis by simulations (using the R package *simr*) was planned to know the range of possible effect-size with power equal or greater than 80% and the influence of patients' number on power.

[‡] R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>

3 | RESULTS

Twenty-two subjects (12 males and 10 females) aged from 26 to 66 years (mean age of 45 years) were consecutively included in the present study. A total of 53 recessions were treated since each patient contributed with one to four recessions. Overall, five incisors, 17 canines, 23 premolars and eight molars were included in the study. All the subjects healed uneventfully and none of them were excluded from the study.



TABLE 1 Clinical variables change from T0 to T1

Parameter	Time 0	Time 1	Difference T1-T0
	Mean \pm SE (95% CI)	Mean \pm SE (95% CI)	Mean \pm SE (95% CI)
Keratinized tissue thickness	0.90 \pm 0.045 mm (0.812 to 0.988)	1.35 \pm 0.056 mm (1.238 to 1.461)	0.45 \pm 0.043 mm (0.366 to 0.533)
			$P < 0.001^{***}$
Relative gingival recession	11.18 \pm 0.214 mm (10.755 to 11.602)	10.38 \pm 0.242 mm (9.901 to 10.861)	-0.798 \pm 0.132 mm (-1.057 to -0.538)
			$P < 0.001^{***}$
Keratinized tissue width	3.01 \pm 0.228 mm (2.556 to 3.459)	3.68 \pm 0.228 mm (3.225 to 4.129)	0.67 \pm 0.074 mm (0.525 to 0.815)
			$P < 0.001^{***}$
Relative clinical attachment level	12.74 \pm 0.219 mm (12.308 to 13.177)	12.03 \pm 0.255 mm (11.521 to 12.534)	-0.715 \pm 0.16 (-1.028 to -0.401)
			$P < 0.001^{***}$
Bone level	14.34 \pm 0.186 mm (13.970 to 14.709)	14.26 \pm 0.187 mm (13.894 to 14.634)	-0.076 \pm 0.042 mm (-0.157 to 0.006)
			$P > 0.05$
Probing pocket depth	1.58 \pm 0.089 mm (1.400 to 1.755)	1.55 \pm 0.111 mm (1.335 to 1.775)	-0.023 \pm 0.09 mm (-0.199 to 0.153)
			$P > 0.05$

Note: The T1-T0 means differences are conditional to the estimated models, so result of a partial pooling.

The Power Analysis showed the study had power greater than 80% in detecting an effect right as small as 0.13 mm and greater than 99% for effects larger than 0.25 mm (see Figure S1 in online *Journal of Periodontology*).

3.1 | Keratinized tissue thickness

Overall, 98.5% of the sites exhibited KTT gain during the follow-up period. Statistically significant difference ($P < 0.01$) was found from T0 to T1 for the KTT with a mean difference of 0.45 ± 0.04 mm (Table 1). Changes in the KTT by patient and recession are shown in Figure 3A. Tooth type, sex and age were not related to an increase in KTT from T0 to T1 ($P > 0.05$). All the teeth with a KTT of < 0.8 mm at T0 ($N = 14$) reached this threshold at T1 (Figure 3B) and 12 out of 14 (85.7%) passed this threshold. Regression analyses of increased tissue thickness by CLD at T0 showed statistically significant correlation between the two variables ($P < 0.001$) (Figure 4). Furthermore, a statistically significant correlation was found between CLD depth and patient age ($P < 0.01$) (see Figure S2 in online *Journal of Periodontology*).

3.2 | Relative gingival recession

Statistically significant difference ($P < 0.01$) was found from T0 to T1 for the RGR reduction with a mean difference

of 0.80 ± 0.132 (Table 1). Changes in the RGR by patient and recession are shown in Figure 5. Overall, 81% of the sites ($n = 43$) had a RGR reduction, 15% ($n = 8$) had no reduction and 3.8% ($n = 2$) had an increase in the recession. Tooth type was related to a decrease of RGR ($P < 0.001$) while age and sex were not ($P > 0.05$). The mean RGR reduction was 0.51 mm for the incisors, 1 mm for the canines and 0.70 and 0.74 mm for premolars and molars respectively (see Figure S3 in online *Journal of Periodontology*). Statistically significant difference ($P < 0.01$) was found between canines-premolars, canine-molars, and molars-incisors ($P < 0.05$). A strong trend was shown between premolar and incisors ($P = 0.056$). Regression analyses of RGR reduction by CLD at T0 showed statistically significant correlation between the two variables ($P < 0.05$) (Figure 4).

3.3 | Keratinized tissue width

Statistically significant difference ($P < 0.01$) was found from T0 to T1 for the KTW increase with a mean difference of 0.67 ± 0.07 (Table 1). Changes in the KTW by patient and recession are shown in (see Figure S4 in online *Journal of Periodontology*). Tooth type, sex and age were not related to an increase in KTW ($P > 0.05$) from T0 to T1. Linear regression analysis showed a correlation between the RGR reduction and increase of KTW (see Figure S5 in online *Journal of Periodontology*). None of the sites had a KTW of 0 at T0. Post-hoc Subgroup-analyses comparing changes in




FIGURE 3 (A) Changes in the keratinized tissue thickness by patient and recession from T0 to T1. (B) Changes in the keratinized tissue thickness (only recession with initial KTT of <0.8 mm) by patient and recession from T0 to T1




FIGURE 4 Regression analysis of the relative gingival recession reduction and keratinized tissue thickness increase by non-carious cervical lesion depth at baseline




FIGURE 5 Changes in the relative gingival recession by patient and recession from T0 to T1

RGR, KTW, and KTT according to the KTW at T0 (<2 vs. ≥ 2 mm) reported no statistically significant difference in any of the parameters (see Figure S6 in online *Journal of Periodontology*).

3.4 | Relative clinical attachment level

Statistically significant difference ($P < 0.01$) was found from T0 to T1 for the increase of RCAL with a mean difference of -0.72 ± 0.16 (Table 1). Changes in the RCAL by patient and recession are shown in (see Figure S7 in online *Journal of Periodontology*). Tooth type was related to a decrease of RGR ($P < 0.001$) while age and sex were not ($P > 0.05$). The mean RCAL increase was 0.59 mm for the incisors, 1 mm for the canines and 0.45 and 0.71 mm for premolars and molars respectively (see Figure S8 in online *Journal of Periodontology*). Statistically significant difference ($P < 0.01$) was found between canines-premolars and canine-molars ($P < 0.01$).

3.5 | Probing pocket depth

No statistically significant difference ($P > 0.05$) was found from T0 to T1 for the KTW increase with a mean difference of 0.23 ± 0.09 (Table 1). Changes in the PPD by patient and recession are shown in (Figure S9 in online *Journal of Periodontology*). Tooth type, sex, and age were not related to an increase in PPD ($P > 0.05$).

4 | DISCUSSION

The present study evaluates whether the newly presented RPT led to an increased gingival phenotype (with a primary focus on the tissue thickness) after 2 months of follow-up. The result of the study clearly shows how this technique enables gingival thickness augmentation by diminishing root convexity apical to the NCCL. This provides a concave space for the formation and stability of a blood clot that will eventually lead to increased tissue thickness, and in most cases, partial reduction of the GR and increase in KTW. The described technique has a similar biological foundation as the biologically oriented preparation technique (BOPT) previously introduced by Loi and Di Felice.²² The BOPT is a protocol in which vertical (featheredge) tooth preparation in a flapless approach is performed, which eliminates the anatomical emergence profile of the crown corresponding to the CEJ. This allows the contiguous soft tissue to adjust its form and location to host the new prosthetic profile, thereby directing the healing, reattachment, and gingival thickness

augmentation. A similar procedure and clinical concept has also been utilized in implant therapy by employing abutments with a shoulder-less design. The IBOPT abutments do not have a finish line, and it is the buccal gingival margin of the crown that creates the soft tissue form. The reduced buccal width of the abutment provides additional space to the gingival thickness thus promoting stability.²³

The first observation that can be extrapolated from the present study is that 98.5% of the sites exhibited KTT gain during the follow-up period with an average gain of 0.5 mm. Of high clinical relevance is that 85.7% of the recession with an initial KTT of <0.8 mm overcame this threshold during the follow-up period, while the remaining teeth reached exactly 0.8 mm. This possibly means that the current minimally invasive approach may facilitate root coverage procedures with promising results without the need of using a collagen matrix which would increase the cost of the procedure²⁴ or harvesting a CTG. Indeed, studies showed that CAF alone was associated with similar clinical outcomes and better final aesthetics compared with CAF + CTG at sites with tissue thickness ≥ 0.8 mm.¹⁴ On the other hand, it should be said that the patient must undergo two different procedures (RPT first and CAF after). Another important consideration is that none of the sites presented with a KTW of 0 at T0. Hence, we could not examine if in complete absence of KTW, the clot formation after the procedure would have the ability to create keratinized gingiva. However, when we divided the sites with lowest KTW values (<2 mm) and compared their performance (changes in RGR, KTW, and KTT) with those with ≥ 2 mm, no difference was found, although these P -values > 0.05 are not confirmatory results. As expected, REC reduction was observed only in the presence of KTW gain, whereas no reduction occurred when KTW did not show an increase. This finding is in accordance with the well-known concept of muco-gingival junction position stability over time.²⁵

Despite the primary goal of the RPT being to increase tissue thickness, most of the treated teeth displayed partial root coverage (81% of the sites with a mean root coverage of 0.8 mm). Another noteworthy observation is the statistically significant association noted between CLD and RGR. A biological explanation of this event is that in plastic surgery the convexity of the surface of the root may negatively impact the root coverage procedure. In this study, a higher CLD corresponded to a greater portion of root structure removed. Hence, the presence of a deep cervical lesion will lead to the creation of a greater space for the clot to form and, therefore, an increase in KTT and reduction in recession depth. The same concept could explain the higher mean root coverage achieved in the canine area, since the entire labial surface of this tooth is



markedly convex. Furthermore, the reported association between increased CLD and older age is a common finding in that the prevalence and severity of NCCLs appear to increase with age.^{8,9,26}

The present study is not exempt from limitations. Clinically, removal of root structure may lead to increased postoperative sensitivity. We did not collect the patient reported outcomes with a visual analog scale, but none of the patients included in the trial complained of excessive sensitivity. Since the RPT would serve as a preparatory (see Figure S10A–E in online *Journal of Periodontology* S10A–E) procedure to be followed by CAF (see Figure S10F in online *Journal of Periodontology*), sensitivity is expected to diminish after the surgical procedure. Moreover, the technique utilized here is essentially a flapless technique, which means it is technique sensitive and may involve a learning curve to attain the presented results. In this regard, it may be useful to consider future use of single-sided diamond tips mounted on piezo surgical handpieces to counterweight the technique sensitivity. Furthermore, RPT followed by CAF should be compared to CAF plus connective tissue graft (as the current gold standard therapy). This comparison would elaborate whether the two techniques achieve similar clinical results, with the advantage for the RPT of avoiding the need to harvest an autogenous graft when a GR defect is associated with an NCCL.

From the methodological perspective, five validity threats are usually indicated due to their potential to impact the one-sample pretest-posttest design. *History* is an implausible rival hypothesis since the trial was conducted under experimental conditions. *Maturation* would be in the opposite direction (decreased tissue thickness) compared to treatment.²⁷ *Patient selection* not representative of the population might have biased the size of the effect. With a standard deviation of 0.2 and a $\rho = 0.07$ as in our sample, and with an assumed population averages ranging between 0.5 and 1.4 mm,^{27,28} in a hypothetical scenario where all patients have baseline values below the cutoff of 0.9 mm, the bias due to the regression to the mean effect would not exceed 0.15 mm. This would guarantee an average treatment effect not smaller than 0.3 mm^{29,30} (see Figure S11 in online *Journal of Periodontology*). *Placebo effects* (when neither patients nor examiner are blinded) may be a cause of overoptimistic results; if any, this bias is expected to be bounded in the 0.1 mm rounding range.

5 | CONCLUSION

Within the limitations of the present study, it can be concluded that the RPT can increase keratinized tissue

thickness and width. Moreover, in most of the teeth, an improvement in gingival recession was achieved. This procedure is especially recommended in patients with thin phenotype to reduce the need of using a collagen matrix or harvesting a CTG. Increased depth of the NCCL was related to better clinical outcomes after the RPT procedure.

AUTHOR CONTRIBUTIONS

All authors have made substantial contributions to conception and design of the study. Luigi Romano, Bruna Sinjari, Edit Xhajanka, and Beatrice Femminella have been involved in data collection and data analysis. Andrea Ravidá, Luigi Romano, Michele Paolantonio, Paolo De Ninis, and Muhammad H. A. Saleh have been involved in data interpretation, drafting the manuscript, and revising it critically and have given final approval of the version to be published. Paolo De Ninis performed the statistical analysis.

CONFLICT OF INTEREST

The authors have no conflicts of interest to disclose.

ORCID

Muhammad H. A. Saleh  <https://orcid.org/0000-0001-5067-7317>

Hom-lay Wang  <https://orcid.org/0000-0003-4238-1799>

Andrea Ravidá  <https://orcid.org/0000-0002-3029-8130>

REFERENCES

- Jepsen S, Caton JG, Albandar JM, et al. Periodontal manifestations of systemic diseases and developmental and acquired conditions: consensus report of workgroup 3 of the 2017 World Workshop on the classification of periodontal and peri-implant diseases and conditions. *J Periodontol*. 2018;89(Suppl 1):S237–S248.
- Romandini M, Soldini MC, Montero E, Sanz M. Epidemiology of mid-buccal gingival recessions in NHANES according to the 2018 World Workshop Classification System. *J Clin Periodontol*. 2020;47:1180–1190.
- Bignozzi I, Crea A, Capri D, Littarru C, Lajolo C, Tatakis DN. Root caries: a periodontal perspective. *J Periodontol Res*. 2014;49:143–163.
- Nieri M, Pini Prato GP, Giani M, Magnani N, Pagliaro U, Rotundo R. Patient perceptions of buccal gingival recessions and requests for treatment. *J Clin Periodontol*. 2013;40:707–712.
- Vignoletti F, Di Martino M, Clementini M, Di Domenico GL, de Sanctis M. Prevalence and risk indicators of gingival recessions in an Italian school of dentistry and dental hygiene: a cross-sectional study. *Clin Oral Investig*. 2020;24:991–1000.
- Cortellini P, Bissada NF. Mucogingival conditions in the natural dentition: narrative review, case definitions, and diagnostic considerations. *J Periodontol*. 2018;89(Suppl 1):S204–S213.
- Sangnes G, Gjermo P. Prevalence of oral soft and hard tissue lesions related to mechanical tooth cleansing procedures. *Community Dent Oral Epidemiol*. 1976;4:77–83.



8. Bartlett DW, Shah P. A critical review of non-carious cervical (wear) lesions and the role of abfraction, erosion, and abrasion. *J Dent Res.* 2006;85:306-312.
9. Pecie R, Krejci I, Garcia-Godoy F, Bortolotto T. Noncarious cervical lesions (NCCL)—a clinical concept based on the literature review. Part 2: restoration. *Am J Dent.* 2011;24:183-192.
10. Rasperini G, Acunzo R, Pellegrini G, et al. Predictor factors for long-term outcomes stability of coronally advanced flap with or without connective tissue graft in the treatment of single maxillary gingival recessions: 9 years results of a randomized controlled clinical trial. *J Clin Periodontol.* 2018;45:1107-1117.
11. Pini-Prato G, Franceschi D, Cairo F, Nieri M, Rotundo R. Classification of dental surface defects in areas of gingival recession. *J Periodontol.* 2010;81:885-890.
12. Santamaria MP, da Silva Feitosa D, Casati MZ, Nociti FH, Jr., Sallum AW, Sallum EA. Randomized controlled clinical trial evaluating connective tissue graft plus resin-modified glass ionomer restoration for the treatment of gingival recession associated with non-carious cervical lesion: 2-year follow-up. *J Periodontol.* 2013;84:e1-e8.
13. Santamaria MP, da Silva Feitosa D, Nociti FH, Jr., Casati MZ, Sallum AW, Sallum EA. Cervical restoration and the amount of soft tissue coverage achieved by coronally advanced flap: a 2-year follow-up randomized-controlled clinical trial. *J Clin Periodontol.* 2009;36:434-441.
14. Cairo F, Cortellini P, Pilloni A, et al. Clinical efficacy of coronally advanced flap with or without connective tissue graft for the treatment of multiple adjacent gingival recessions in the aesthetic area: a randomized controlled clinical trial. *J Clin Periodontol.* 2016;43:849-856.
15. Berlucchi I, Francetti L, Del Fabbro M, Basso M, Weinstein RL. The influence of anatomical features on the outcome of gingival recessions treated with coronally advanced flap and enamel matrix derivative: a 1-year prospective study. *J Periodontol.* 2005;76:899-907.
16. Ainamo J, Bay I. Problems and proposals for recording gingivitis and plaque. *Int Dent J.* 1975;25:229-235.
17. Santamaria MP, Ambrosano GM, Casati MZ, Nociti FH, Jr., Sallum AW, Sallum EA. The influence of local anatomy on the outcome of treatment of gingival recession associated with non-carious cervical lesions. *J Periodontol.* 2010;81:1027-1034.
18. Bozkurt Dogan S, Ongoz Dede F, Balli U, Atalay EN, Durmuslar MC. Concentrated growth factor in the treatment of adjacent multiple gingival recessions: a split-mouth randomized clinical trial. *J Clin Periodontol.* 2015;42:868-875.
19. Zangrando MSR, Eustachio RR, de Rezende MLR, Sant'ana ACP, Damante CA, Greggi SLA. Clinical and patient-centered outcomes using two types of subepithelial connective tissue grafts: a split-mouth randomized clinical trial. *J Periodontol.* 2021;92:814-822.
20. Snijders TAB, Bosker RJ. *Multilevel Analysis: An Introduction to Basic and Advanced Multilevel Modeling.* 2nd ed. SAGE Publications Ltd; 2011.
21. Barr DJ, Levy R, Scheepers C, Tily HJ. Random effects structure for confirmatory hypothesis testing: keep it maximal. *J Mem Lang.* 2013;68(3):10.
22. Loi I, Di Felice A. Biologically oriented preparation technique (BOPT): a new approach for prosthetic restoration of periodontically healthy teeth. *Eur J Esthet Dent.* 2013;8:10-23.
23. O'Keefe M. Peri-implant tissue remodeling: scientific background and clinical implication. *Br Dental J.* 2012;213:142-142.
24. Mathias-Santamaria IF, Silveira CA, Rossato A, Sampaio de Melo MA, Bresciani E, Santamaria MP. Single gingival recession associated with non-carious cervical lesion treated by partial restoration and coronally advanced flap with or without xenogenous collagen matrix: a randomized clinical trial evaluating the coverage procedures and restorative protocol. *J Periodontol.* 2022;93:504-514.
25. Ainamo J, Talari A. The increase with age of the width of attached gingiva. *J Periodontol Res.* 1976;11:182-188.
26. Heasman PA, Holliday R, Bryant A, Preshaw PM. Evidence for the occurrence of gingival recession and non-carious cervical lesions as a consequence of traumatic toothbrushing. *J Clin Periodontol.* 2015;42(Suppl 16):S237-S255.
27. Kolte R, Kolte A, Mahajan A. Assessment of gingival thickness with regards to age, gender and arch location. *J Indian Soc Periodontol.* 2014;18:478-481.
28. Holtzman LP, Blasi G, Rivera E, Herrero F, Downton K, Oates T. Gingival thickness and outcome of periodontal plastic surgery procedures: a meta-regression analysis. *JDR Clin Trans Res.* 2021;6:295-310.
29. Davis CE. The effect of regression to the mean in epidemiologic and clinical studies. *Am J Epidemiol.* 1976;104:493-498.
30. Barnett AG, van der Pols JC, Dobson AJ. Regression to the mean: what it is and how to deal with it. *Int J Epidemiol.* 2005;34:215-220.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Romano L, Paolantonio M, De Ninis P, et al. Minimally invasive gingival phenotype modification in gingival recession associated with a non-carious cervical lesion using the root plastique technique: A quasi-experimental one-group pretest-posttest study. *J Periodontol.* 2023;94:641–651.

<https://doi.org/10.1002/JPER.22-0414>