

Complexes of Tetracyanobiimidazole. 3. Platinum and Palladium Species

P. G. RASMUSSEN*, O. H. BAILEY

Department of Chemistry, University of Michigan, Ann Arbor, Mich. 48109, U.S.A.

and J. C. BAYÓN*

Department de Química Inorgànica, Facultat de Ciències, Universitat Autònoma de Barcelona, Bellaterra, Barcelona, Spain

Received October 25, 1983


In this paper we describe monomeric complexes of Pd(II) and Pt(II) with 4,4',5,5'-tetracyano-2,2'-biimidazole ($H_2Tcbiim$). These compounds always contain the dianion ($Tcbiim^{2-}$) in accord with the greater acidity expected for the cyanosubstituted ring system. The structure of $[N(Bu)_4]_2[Pt(CN)_2Tcbiim]$ was determined by x-ray crystallography. The composition $PtC_{44}H_{72}N_{12}$ crystallizes in the $P\bar{1}$ triclinic space group with $a = 16.664(3)$ Å, $b = 12.422(4)$ Å, $c = 13.842(4)$ Å, $\alpha = 112.47(2)$, $\beta = 111.01(2)$, $\gamma = 89.40(2)$ and $Z = 2$ molecules/cell. A final agreement of $R = 0.038$ was obtained for 5359 observed reflections. The structure confirms planar bidentate chelation by $Tcbiim^{2-}$ and an irregular square planar coordination for the platinum atom. The preparation of a series of salts $M[Pt(CN)_2(Tcbiim)]$, $M = NBu_4^+$, NH_4^+ , Ba^{2+} , and for K^+ both the platinum and palladium analogs, are described. The neutral molecule $Pt(COD)Tcbiim$ was also prepared. Activity of these species under oxidation conditions are discussed in relation to the π accepting properties of the title ligand.

Introduction

We have previously reported [1] the synthesis and some of the properties of a novel new compound 4,4',5,5'-tetracyano-2,2'-biimidazole ($H_2Tcbiim$), 1. We have also reported its use in forming complexes with Rh(I), Ir(I) and Mo(0) [2–4]. In this communication we report on compounds formed with divalent platinum and palladium.

These compounds were prepared as part of an effort to study deliberately synthesized analogues to partially oxidized, anisotropically conducting materials such as the Krogmann salts [5]. This field was reviewed recently by Williams [6], and it is evident that the range of examples is relatively narrow. We

were particularly interested in exploring the use of the planar tetracyanobiimidazole dianion as a substitute for two cyanide ligands since its π accepting properties should be similar. We were also interested in elaborating the use of tetracyanobiimidazole a π accepting anionic ligand in concert with other ligands of low donor strength for synthetic purposes in organoplatinum chemistry.



Experimental

The compounds $cis-Pt(CN)_2(NH_3)_2$, $Pt(COD)Cl_2$, and $H_2Tcbiim$ were prepared following published procedures [1, 7, 8]. Elemental analyses were performed by Spang Microanalytical Laboratory, Eagle Harbor, Michigan. Infrared spectra were recorded as Nujol mulls on a Perkin Elmer 1330 spectrophotometer. Fourier transform NMR were obtained on a Bruker WM-360 spectrometer calibrated with ^{13}C and ^{31}P shifts as positive downfield relative to TMS and H_3PO_4 .

Preparation of $K_2[Pd(CN)_2Tcbiim]$ 2

500 mg of $K_2[Pd(CN)_4] \cdot 3H_2O$ (1.46 mmol) and 335 mg of $H_2Tcbiim$ (1.43 mmol) were refluxed in 25 ml of water. After 20 minutes the solution turned clear. Further refluxing for 3 hours, followed by cooling on ice produced a white precipitate. The ppt was filtered and washed with 3 ml of cold water and 10 ml of acetone. Yield 600 mg (90%). *Anal.* Calcd for $PdK_2C_{12}N_{10}$: C, 30.74; N, 29.88. Found: C, 30.66; N, 29.77%.

Preparation of $(NH_4)_2[Pt(CN)_2Tcbiim]$ 3

The method described for the palladium complex above gives yields of less than 5% for the platinum

*Authors to whom all correspondence should be addressed.

complex. Addition of varying quantities of HCl did not improve the yields. Instead, 600 mg of *cis*-Pt(NH₃)₂(CN)₂ (2.13 mmol) were added in small portions to a solution of 500 mg of H₂Tcbiim (2.13 mmol) in 200 ml of boiling water over a period of 15 minutes with strong stirring. The volume was reduced to 30 ml, filtered to remove traces of Pt(CN)₂ and furthered reduced to 10 ml. Addition of 15 ml methanol–150 ml ether resulted in precipitation. Yield 970 mg (90%). *Anal.* Calcd for PtC₁₂N₁₂H₈: C, 27.97; N, 32.62; H, 1.56. Found: C, 27.78; N, 32.47; H, 1.70%. ¹³C NMR (D₂O): 153.1, 124.2, 121.3, 116.4, 114.2, 108.4.

Preparation of (NBu₄)₂[Pt(CN)₂Tcbiim] 4

To a solution of 3 in water, the stoichiometric amount of NBu₄Br was added producing a white precipitate. It was recrystallized from acetone–methanol. *Anal.* Calcd for PtC₄₄N₁₂H₇₂: C, 54.80; N, 17.43; H, 7.52. Found: C, 54.72; N, 17.24; H, 7.47%.

Preparation of Pt(COD)Tcbiim 5

100 mg of Pt(COD)Cl₂ (0.27 mmol) and 63 mg of H₂Tcbiim (0.27 mmol) were refluxed in 80 ml of ethanol for 45 minutes. Upon cooling a precipitate formed which was filtered and washed with acetone and methylene chloride. Yield 122 mg (90%). *Anal.* Calcd for PtC₁₈N₈H₁₂: C, 40.38; N, 20.93; H, 2.26. Found: C, 40.16; N, 20.71; H, 2.35%.

TABLE I. Infrared Absorptions (cm⁻¹).

| | Nitrile | Cyanide |
|---------------------------------------------------------------|------------|------------|
| K ₂ [Pd(CN) ₂ Tcbiim] | 2244 | 2164, 2150 |
| K ₂ [Pt(CN) ₂ Tcbiim] | 2240 | 2162, 2143 |
| (NH ₄) ₂ [Pt(CN) ₂ Tcbiim] | 2240 | 2160, 2137 |
| (NBu ₄) ₂ [Pt(CN) ₂ Tcbiim] | 2230 | 2149, 2137 |
| Pt(COD)Tcbiim | 2240, 2229 | |

Collection and Solution of the X-ray Data

Well formed crystals of 4 were grown by slow evaporation of an acetone–methanol solution. A parallelepiped shaped crystal of dimensions 0.103 × 0.246 × 0.407 mm was mounted on a Syntex P $\bar{1}$ four circle diffractometer. Centering on 15 reflections yielded cell parameters of the triclinic system. The structure refined successfully in the space group P $\bar{1}$. A summary of crystal data is presented in Table II and programs are cited in footnote [9]. The structure was solved by conventional heavy atom and difference Fourier methods. After refining to isotropic convergence the data was corrected for absorption. Hydrogen atom positions were calculated and added to the refinement (*R* = 0.065, *R_w* = 0.079). Anisotropic refinement of the non-hydrogen atoms led to final

TABLE II. Crystal Parameters for (NBu₄)₂[Pt(CN)₂Tcbiim].

| Space group | P $\bar{1}$ |
|-----------------------------------------------------|---------------------------------------------------------------------|
| <i>a</i> , Å | 16.664(3) |
| <i>b</i> , Å | 12.422(4) |
| <i>c</i> , Å | 13.842(4) |
| α, deg. | 112.47(2) |
| β, deg. | 111.01(2) |
| γ, deg. | 89.40(2) |
| Mol. Wt. | 964.22 |
| <i>Z</i> | 2 |
| Calcd. density | 1.17 |
| Crystal Dimensions mm | (010)–0.103 (10 $\bar{1}$)–0.246 (100)–0.407 |
| Radiation Å | λ (Mo K α) 0.71069 Monochromatized from graphite crystal |
| Linear Absorption coefficient μ, | 29.28 cm ⁻¹ |
| Scan speed, deg./min. | variable 2.5–12 |
| Ratio | |
| background/peak scan | 0.8 |
| Std. reflections | (6 $\bar{3}$ 2) (82 $\bar{4}$) (4 $\bar{2}$ 0) |
| Scan range 2θ, deg. | 0–45 |
| Reflections collected | 6842 |
| Reflections observed | |
| <i>F</i> ² ≥ 3σ(<i>F</i> ²) | 5359 |


values of *R* = 0.038, *R_w* = 0.048. The final difference map showed the highest peak in the vicinity of a heavy atom with intensity 1.59 e/Å³.

Discussion


An ORTEP view of the structure of the [Pt(CN)₂Tcbiim]²⁻ anion is shown in Fig. 1. Bond distances and angles are given in Tables III and IV respectively. The complex anion is approximately planar with only the nitriles of N4 and N5 and the C12 cyanide deviating slightly from the mean plane. There is how-

TABLE III. Interatomic Distances (Esd).

| | | | |
|---------|-----------|---------|-----------|
| | | N5'–C7' | 1.151(12) |
| Pt–C11 | 1.915(9) | N4'–C6' | 1.113(10) |
| Pt–C12 | 1.918(9) | N4–C6 | 1.143(11) |
| Pt–N1' | 2.049(6) | N5–C7 | 1.134(10) |
| Pt–N1 | 2.077(6) | N9–C11 | 1.160(9) |
| N1–C5 | 1.350(9) | N10–C12 | 1.150(10) |
| N1–C2 | 1.366(9) | C5'–C4' | 1.386(12) |
| N1'–C5' | 1.353(9) | C5'–C7' | 1.399(13) |
| N1'–C2' | 1.377(10) | C4'–C6' | 1.446(12) |
| N3'–C2' | 1.319(9) | C2'–C2 | 1.442(11) |
| N3'–C4' | 1.365(11) | C4–C5 | 1.391(11) |
| N3–C2 | 1.322(9) | C4–C6 | 1.429(12) |
| N3–C4 | 1.338(9) | C5–C7 | 1.408(11) |



(a)



(b)

Fig. 1. (a) Structure of the platinum complex. (b) Stereoview of the molecule.

TABLE IV. Selected Interatomic Angles (Esd).

| | | | |
|-------------|------------|-------------|------------|
| C11-Pt-C12 | 87.71(33) | C5'-C4'-C6' | 125.23(90) |
| C11-Pt-N1' | 175.65(30) | N3'-C2'-N1' | 116.04(69) |
| C11-Pt-N1 | 97.01(27) | N3'-C2'-C2 | 128.63(79) |
| C12-Pt-N1' | 96.32(30) | N1'-C2'-C2 | 115.31(69) |
| C12-Pt-N1 | 173.61(27) | N3-C2-N1 | 114.21(66) |
| N1'-Pt-N1 | 78.85(24) | N3-C2-C2' | 130.52(75) |
| C5-N1-C2 | 104.55(63) | N1-C2-C2' | 115.26(73) |
| C5-N1-Pt | 139.75(53) | N3-C4-C5 | 111.06(69) |
| C2-N1-Pt | 114.31(49) | N3-C4-C6 | 123.27(77) |
| C5'-N1'-C2' | 103.29(64) | C5-C4-C6 | 125.63(80) |
| C5'-N1'-Pt | 141.45(58) | N1-C5-C4 | 106.56(67) |
| C2'-N1'-Pt | 115.19(48) | N1-C5-C7 | 125.47(74) |
| C2'-N3'-C4' | 101.92(70) | C4-C5-C7 | 127.96(75) |
| C2-N3-C4 | 103.58(63) | N5'-C7'-C5' | 174.7(11) |
| N1'-C5'-C4' | 107.41(75) | N4'-C6'-C4' | 179.5(34) |
| N1'-C5'-C7' | 125.26(81) | N4-C6-C4 | 177.6(10) |
| C4'-C5'-C7' | 127.32(79) | N5-C7-C5 | 174.78(96) |
| N3'-C4'-C5' | 111.31(70) | N9-C11-Pt | 176.63(75) |
| N3'-C4'-C6' | 123.39(86) | N10-C12-Pt | 177.62(84) |

ever a slight folding of the anion as shown by the 10.5 dihedral angle between the plane of the $\text{Pt}(\text{CN})_2$ moiety and the Tcbiim^{2-} ligand.

The coordination around the platinum is distorted from square by the chelating ligand which makes an angle of 78.8 at platinum. This is very similar to that found for the isoelectronic anion $[\text{Ir}(\text{CO})_2\text{Tcbiim}]^-$ [2]. The platinum-carbon distances (1.915–1.918 Å) are significantly shorter than those found in the typical salts of tetracyanoplatinates but the carbon-nitrogen distances in the cyanides are in the same range [6]. These facts suggest a slightly smaller pi accepting ability for Tcbiim^{2-} compared to CN^- although the stability of the ion indicates a comparable *trans* effect since most ligands *trans* to cyanide are quite labile.

The size of the cation apparently prevents the stacking of the planar anions, although signs of intermolecular association do not appear in concentrated solutions of $(\text{NH}_4)_2[\text{Pt}(\text{CN})_2\text{Tcbiim}]$ (which is quite soluble) as they do for salts of the iridium carbonyl analog. However if a solution of BaCl_2 is allowed to diffuse into a solution of **3**, long 'whiskers' form, which although not suitable for single crystal diffraction, do show powder X-ray patterns with strong lines corresponding to a 3.32 Å separation.

We made several attempts to prepare dimers in which Tcbiim^{2-} was tetradentate to two platinum atoms, similar to dimers we have synthesized using iridium(I) and Cu(I) [3, 10]. The mixing of *cis*- $\text{Pt}(\text{NH}_3)_2(\text{CN})_2$ in 1:1 ratio with **3** or in 2:1 ratio with H_2Tcbiim leads to the formation of $\text{Pt}(\text{CN})_2$. No reaction was observed between **3** and $\text{Pt}(\text{CN})_2$ during lengthy reflux, although $\text{Pt}(\text{CN})_2$ will slowly react with Tcbiim^{2-} producing **3** in low yield. Mixing of **3** with PtCl_4^{2-} followed by addition of two equivalents of CN^- leads to precipitation of $\text{Pt}(\text{CN})_2$. Finally, $[\text{Pt}(\text{COD})(\text{acac})](\text{BF}_4)$ was refluxed in 2:1 ratio with H_2Tcbiim in acetonitrile. This produced only **5** with no further addition of platinum complex. We conclude that the 'backside' of Tcbiim^{2-} lacks sufficient basicity to bind Pt(II) when the 'frontside' is already bound to Pt(II). It is possible that such dimers might form, as they do in the case of Cu(II), [10] when other strong donors are present *e.g.* [(dien)Cu(Tcbiim)Cu(dien)]²⁺ dien = diethylenetriamine but we do not presently know how to synthesize these analogs. It should also be noted that in the above remarks ' $\text{Pt}(\text{CN})_2$ ' does not represent a very tractable material. It is white, hard to filter, likely polymeric and is hydrated to an uncertain degree. This is quite different from the $\text{Pt}(\text{CN})_2$ used in the preparation of *cis*- $\text{Pt}(\text{NH}_3)_2(\text{CN})_2$ which is prepared from thermal decomposition of $(\text{NH}_4)_2[\text{Pt}(\text{CN})_4]$ and which may also be polymeric but is orange and anhydrous [7].

Attempts at oxidizing the salts of $[\text{Pt}(\text{CN})_2\text{Tcbiim}]^{2-}$ in analogy to the preparation of the partially oxidized, anisotropic materials, have not so

TABLE V. Atomic Coordinates.

| | | | |
|-----|--------------|--------------|--------------|
| Pt | 0.211040(18) | 0.400738(27) | 0.255731(25) |
| N1 | 0.26517(37) | 0.57426(52) | 0.36461(51) |
| N1' | 0.25353(36) | 0.45362(58) | 0.15683(50) |
| N3' | 0.32571(43) | 0.60197(64) | 0.14663(58) |
| N3 | 0.35781(42) | 0.73012(56) | 0.40334(59) |
| N5' | 0.18189(72) | 0.2171(10) | -0.11348(81) |
| N4' | 0.31132(54) | 0.50321(87) | -0.13318(72) |
| N4 | 0.42074(59) | 0.91631(79) | 0.68817(79) |
| N5 | 0.22549(64) | 0.62838(80) | 0.61187(75) |
| N9 | 0.15963(49) | 0.33383(69) | 0.41998(67) |
| N10 | 0.15054(50) | 0.14150(73) | 0.09011(69) |
| C5' | 0.25059(50) | 0.41671(77) | 0.04994(65) |
| C4' | 0.29500(53) | 0.50686(89) | 0.04494(75) |
| C2' | 0.29808(48) | 0.56629(72) | 0.20911(65) |
| C2 | 0.31000(49) | 0.62860(71) | 0.32619(65) |
| C4 | 0.34294(51) | 0.74296(68) | 0.49569(69) |
| C5 | 0.28492(48) | 0.64881(69) | 0.47355(65) |
| C7' | 0.21082(64) | 0.3063(10) | -0.03738(82) |
| C6' | 0.30461(54) | 0.50491(87) | -0.05546(82) |
| C6 | 0.38467(60) | 0.83963(81) | 0.60285(85) |
| C7 | 0.25102(58) | 0.63254(74) | 0.54748(74) |
| C11 | 0.17712(45) | 0.36136(68) | 0.35793(71) |
| C12 | 0.17201(50) | 0.23938(78) | 0.15042(71) |
| N11 | 0.89210(37) | 0.24095(52) | 0.13875(49) |
| C13 | 0.80809(63) | 0.2111(10) | 0.14883(94) |
| C14 | 0.74973(68) | 0.10292(95) | 0.0670(11) |
| C15 | 0.67538(81) | 0.0708(13) | 0.0811(14) |
| C16 | 0.61800(88) | -0.0293(14) | 0.0154(16) |
| C17 | 0.86500(64) | 0.26166(89) | 0.02887(79) |
| C18 | 0.93210(62) | 0.29942(75) | 0.00113(77) |
| C19 | 0.90180(92) | 0.3280(11) | -0.0997(11) |
| C20 | 0.95967(94) | 0.3551(14) | -0.1441(11) |
| C21 | 0.94455(63) | 0.13784(89) | 0.12616(88) |
| C22 | 0.96822(70) | 0.09672(97) | 0.21644(89) |
| C23 | 1.0284(10) | 0.0082(11) | 0.2125(13) |
| C24 | 1.0630(12) | -0.0283(14) | 0.2953(12) |
| C25 | 0.94579(73) | 0.34877(92) | 0.23948(82) |
| C26 | 0.91631(69) | 0.46293(83) | 0.26149(88) |
| C27 | 0.9679(11) | 0.56126(96) | 0.3630(11) |
| C28 | 0.9515(11) | 0.6757(10) | 0.3945(13) |
| N12 | 0.62292(38) | 0.76223(55) | 0.38369(50) |
| C29 | 0.60602(51) | 0.84006(75) | 0.48693(69) |
| C30 | 0.66548(60) | 0.83803(83) | 0.59832(71) |
| C31 | 0.64544(72) | 0.9261(11) | 0.69670(85) |
| C32 | 0.6647(11) | 1.0477(12) | 0.7208(11) |
| C33 | 0.61883(50) | 0.63268(71) | 0.36775(64) |
| C34 | 0.53276(54) | 0.57504(77) | 0.35018(71) |
| C35 | 0.54144(58) | 0.45359(84) | 0.34965(82) |
| C36 | 0.45844(69) | 0.38314(92) | 0.3212(10) |
| C37 | 0.55415(51) | 0.77770(72) | 0.28591(67) |
| C38 | 0.56066(63) | 0.71434(82) | 0.17173(73) |
| C39 | 0.49460(65) | 0.74525(88) | 0.08295(77) |
| C40 | 0.50284(82) | 0.6861(11) | -0.03084(88) |
| C41 | 0.71360(52) | 0.79652(77) | 0.39439(67) |
| C42 | 0.73393(58) | 0.92277(82) | 0.41109(78) |
| C43 | 0.82145(70) | 0.93978(97) | 0.40084(88) |
| C44 | 0.84715(82) | 1.0670(12) | 0.4263(11) |

far been successful. These efforts have included aerial oxidation, electrolysis at controlled potential, and

mixing with appropriate quantities of the oxidative addition compound formed with chlorine. Apparently the bonding strength of the stack which can form in this system is not sufficient to recoup the energy required for oxidation. This is surprising in view of the similarities of the electronic structure of Tcbiim^{2-} to cyanide and may be due to the greater ligand–ligand repulsions.

The synthetic aspects of tetracyanobiimidazole complexes of platinum are quite promising and offer potentially useful routes in several directions. The thermal decomposition of **3** analogous to the preparation of anhydrous $\text{Pt}(\text{CN})_2$ should lead to a similarly useful starting material. This method would circumvent the problems which the preparation of *cis* compounds of strong *trans* directors typically present. The formation of **5** suggests that other neutral ligands of low electronegativity would also be compatible with the $\text{Pt}(\text{Tcbiim})$ moiety and the possible preparation of $\text{Pt}(\text{L})_2\text{Tcbiim}$ derivatives by substitution reactions. These reaction pathways are under further investigation in our laboratory.

Acknowledgment

J. C. B. wishes to thank the Fulbright Commission and the U.S.A.–Spain Joint Committee for post-doctoral support. This work was aided by support from the National Science Foundation INT-8304329 and P.G.R. wishes to especially thank Prof. Andreas Ludi, Bern, Switzerland, and his students for helpful discussions.

Supplementary Information

Tables of structure factors, anisotropic thermal parameters, and least squares planes are available upon request to either of the senior authors (25 pages).

References

- 1 P. G. Rasmussen, R. L. Hough, J. E. Anderson, O. H. Bailey and J. C. Bayon, *J. Am. Chem. Soc.*, **104**, 6155 (1982).
- 2 P. G. Rasmussen, O. H. Bailey, J. C. Bayon and W. M. Butler, *Inorg. Chem.*, in press.
- 3 P. G. Rasmussen, O. H. Bailey and J. C. Bayon, *Inorg. Chem.*, in press.
- 4 P. G. Rasmussen and J. C. Bayon, *Inorg. Chim. Acta*, **81**, L15 (1984).
- 5 P. J. Nigrey, Chap. 9 in 'Extended Linear Chain Compounds', Vol. 3, ed. by J. S. Miller, Plenum: New York, (1983).
- 6 J. M. Williams, A. J. Schultz, A. E. Underhill and K. Carneiro, Chap. 3 in 'Extended Linear Chain Compounds' ed. by J. S. Miller, Vol. 1, Plenum: New York (1982).
- 7 I. B. Baranovskii and Y. Y. Kharitonov, *Doklady Akad. Nauk. SSSR*, **169**, 1335 (1966).
- 8 D. Drew and J. R. Doyle, *Inorg. Syn.*, **13**, 48 (1972).
- 9 Computer programs used during the structural analysis were SYNCOR (data reduction by W. Shmonsees), N. W. Alcock's absorption program, FORDAP (Fourier synthesis by A. Zalkin), ORFLS (full matrix least-squares refinement by Busing, Martin and Levy), ORFFE (distances, angles, and their esd's by Busing, Martin, and Levy), ORTEP (thermal ellipsoid drawings by C. K. Johnson), HATOMS (hydrogen atom positions by A. Zalkin), PLANES (least-squares by D. M. Blow), and PLUTO (crystallographic plotting program supplied by University Chemical Laboratory, Cambridge, England). Atomic scattering factors were obtained from D. T. Cromer and J. T. Waber.
- 10 P. G. Rasmussen and J. A. Anderson, *Polyhedron*, **2**, 547 (1983).