


1,2-ADDITION OF DIMETHYL(PHENYL)SILYL LITHIUM TO CYCLIC α,β -UNSATURATED KETONES AND REGIOSPECIFIC GENERATION OF CYCLIC SILYL ENOL ETHERS THROUGH BROOK REARRANGEMENT OF THE 1,2-ADDITION PRODUCTS

Masato Koreceda* and Sangho Koo

Department of Chemistry, The University of Michigan, Ann Arbor, Michigan 48109

Summary: A highly convenient two-step sequence for the regiospecific synthesis of cyclic silyl enol ethers has been developed involving the 1,2-addition of dimethyl(phenyl)silyllithium to cyclic α,β -unsaturated ketones followed by the treatment of the resulting silyl carbinols with a catalytic amount of NaH in THF at 25 °C.

Silyl enol ethers are highly versatile intermediates on which a variety of regio- and/or stereochemically controlled synthetic manipulations can be realized.¹ Accordingly, a number of efficient methods have been developed that effect the regio- and stereochemically controlled generation of silyl enol ethers.^{1,2} In 1979 - 80, Kuwajima³ and Reich⁴ independently established a novel Brook rearrangement-based method for the regiospecific synthesis of acyclic (*Z*)-silyl enol ethers (Scheme I). The origin of the exclusive formation of *Z*-isomers is attributed to the



generation of the coordination-stabilized intermediate 5 from 4. Thus, by inference this method should not be applicable to the synthesis of most cyclic silyl enol ethers. We now report that 1,2-addition products 8 of dimethyl(phenyl)silyllithium (Me₂PhSiLi) to cyclic α,β -unsaturated ketones 7 can be conveniently converted regiospecifically into the corresponding cyclic silyl enol ethers 9 (Scheme II).

Trimethylsilyllithium (TMSLi) has been reported to react with cyclic α,β -unsaturated ketones in THF/HMPA (5:1) at -78 °C to give rise to 1,4-addition products exclusively,⁵ whereas Me₂PhSiLi, which can be readily formed in THF,⁶ was found to undergo smooth 1,2-addition to cyclic α,β -unsaturated ketones at -23 °C in good to excellent

Scheme II




























Table I. 1,2-Addition of Me₂PhSiLi to Cyclic α,β-Unsaturated Ketones and Brook Rearrangement of the Adducts^a

entry	enones	silyl carbinols	% yield ^b	conditions for Brook rearrangement ^c	silyl enol ethers	% yield ^b
1			94	A		72
2			89	A	 18	73
3			89	A		76
4			74 ^{d,e}	B	 19  10	19/10 43/4
5			77	B		60
6			85 ^d	A		71
7			67 ^{e,f}	A C D	 20  11	20/11 45 ^g /15 (A) 34 ^g /21 (C) 0/23 (D)

a) Si denotes a dimethyl(phenyl)silyl group. b) Isolated yields following flash column chromatographic purification. c) Conditions: A, NaH/THF, 2 h at 25 °C; B, NaH/THF, 1 h at 25 °C; C, NaH/THF, 1 h at 55 °C; D, NaH/THF-HMPA (20/1), 1 h at 20 °C. d) 3:2 diastereomeric mixture. e) LiCl (0.30 equiv) added. f) 10:1 diastereomeric mixture. g) 2:1 diastereomeric mixture.

As an application of these dimethyl(phenyl)silyl enol ethers in synthesis, the regiospecific TiCl_4 -catalyzed aldol reactions with benzaldehyde⁸ were examined with two silyl enol ethers **18** and **19** (see eqs. 2 and 3). While these two silyl enol ethers produced the aldol products with benzaldehyde regiospecifically in comparable yields with those reported for their corresponding TMS ethers, exceptionally high erythro/threo-selectivity observed for the reaction of **18** is quite noteworthy. The erythro/threo ratio of the aldol adducts of its corresponding TMS ether with benzaldehyde was only 1:1.⁸ As in the case of its corresponding TMS ether (3:10:16:71),⁸ silyl enol ether **19** provided a mixture of four diastereomers in a 1:17:25:57 (ax. erythro/eq. threo/eq. erythro/ax. threo) ratio.



In summary, the two-step sequence delineated above offers a convenient, regiospecific means for the synthesis of dimethyl(phenyl)silyl enol ethers from their corresponding enones.⁹ These silyl enol ethers can be purified through silica gel flash column chromatography and can be used for the TiCl_4 -catalyzed aldol reaction with aldehydes.

General Procedures for the 1,2-Addition of Me_2PhSiLi to Cyclic α,β -Unsaturated Ketones **7 and Brook Rearrangement of Silyl Carbinols **8**.** **7 - 8:** To a stirred solution (-23°C) of 0.40 M Me_2PhSiLi (1.30 equiv) in THF was added a cyclic α,β -unsaturated ketone **7** (1.00 M in THF, 1.00 equiv). The mixture was stirred at -23°C for 1 h, at which point the reaction was quenched with saturated aq. NH_4Cl . The resulting mixture was extracted with ether twice and the combined organic extracts were washed first with saturated aq. NH_4Cl and then with brine. The organic layer was dried (MgSO_4), filtered, and concentrated under reduced pressure. The crude material thus obtained was purified by flash column chromatography.

8 - 9: A catalytic amount of NaH (0.50 equiv), which was obtained from 60% NaH dispersion in mineral oil by washing twice with 10 mL each of hexanes, was suspended in 10 mL of THF. To the suspension was added a cyclic silyl carbinol **8** (1.00 M in THF, 1.00 equiv) at 25°C . The mixture was stirred vigorously at that temperature for 1-2 h and then was quenched with saturated aq. NH_4Cl . The resulting mixture was worked up as above.

Acknowledgment: We are grateful for the financial support of this work by the National Institutes of Health (Grant #DK 30025). S. K. is a 1989-90 Yates Fellowship recipient.

- References:** 1) a) Colvin, E. W. *Silicon in Organic Synthesis*; Butterworth: London, 1981. Chapter 17. b) Brownbridge, P. *Synthesis* 1983, 1, 85. c) Politier, J.-M. *Org. Prep. Proced. Int.* 1988, 20, 317.
 2) For recent leading references in this area, see: a) Sakurai, H.; Miyoshi, K.; Nakadaira, Y. *Tetrahedron Lett.* 1977, 2671. b) Utimoto, K.; Obayashi, M.; Shishiyama, Y.; Inoue, M.; Nozaki, H. *Ibid.* 1980, 21, 3389. c) Negishi, E.; Chatterjee, S. *Ibid.* 1983, 24, 1341. d) Krafft, M. E.; Holton, R. A. *Ibid.* 1983, 24, 1341. e) Idem *J. Org. Chem.* 1984, 49, 3669. f) Tsuda, T.; Satomi, H.; Hayashi, T.; Saegusa, T. *Ibid.* 1987, 52, 439.
 3) a) Kuwajima, I.; Kato, M. *J. C. S. Chem. Commun.* 1979, 708. b) Kuwajima, I.; Kato, M.; Mori, A. *Tetrahedron Lett.* 1980, 21, 2745.
 4) a) Reich, H. J.; Rusek, J. J.; Olson, R. E. *J. Am. Chem. Soc.* 1979, 101, 2225. b) Reich, H. J.; Olson, R. E.; Clark, M. C. *Ibid.* 1980, 102, 1423.
 5) a) Still, W. C. *J. Org. Chem.* 1976, 41, 3063. b) Still, W. C.; Mitra, A. *Tetrahedron Lett.* 1978, 2659.
 6) Fleming, I.; Newton, T. W. *J. Chem. Soc., Perkin Trans. I* 1984, 1805.
 7) Brown, C. A.; Yamauchi, A. *J. C. S. Chem. Commun.* 1979, 100.
 8) Mukaiyama, T.; Banno, K.; Narasaka, K. *J. Am. Chem. Soc.* 1974, 96, 7503.
 9) For the alternative, transition metal-catalyzed hydrosilylation of enones, see: Ojima, I.; Kogure, T. *Organometallics* 1982, 1, 1390.