

Chern class formulas for quiver varieties

Anders Skovsted Buch¹, William Fulton²

¹Department of Mathematics, University of Chicago, Chicago, IL 60637, USA

²Department of Mathematics, University of Michigan, Ann Arbor, MI 48109, USA
(e-mail: abuch@math.uchicago.edu/wfulton@math.lsa.umich.edu)

Oblatum 6-IV-1998 & 15-V-1998 / Published online: 14 January 1999

1. Introduction

Our goal in this paper is to prove a formula for the general degeneracy locus Ω_r associated to an oriented quiver of type A_n . If we are given a sequence of vector bundles and vector bundle maps

$$E_0 \xrightarrow{\phi_1} E_1 \xrightarrow{\phi_2} E_2 \longrightarrow \cdots \longrightarrow E_{n-1} \xrightarrow{\phi_n} E_n$$

on an algebraic variety X , and a collection $r = (r_{ij})_{0 \leq i < j \leq n}$ of non-negative integers, there is a degeneracy locus $\Omega_r = \Omega_r(E_\bullet) = \Omega_r(E_\bullet, \phi_\bullet)$ defined by

$$(1.1) \quad \Omega_r = \{x \in X \mid \text{rank}(E_i(x) \rightarrow E_j(x)) \leq r_{ij} \forall i < j\} .$$

This is a closed subscheme of X ; locally, where the bundles are trivial, this is defined by vanishing of the minors of size $r_{ij} + 1$ in the product of matrices giving the map $\phi_j \circ \cdots \circ \phi_{i+1}$ from E_i to E_j , for all $i < j$.

Not all rank conditions give reasonable loci. Those that do – and the only ones we will consider – are characterized by the conditions

$$(1.2) \quad \begin{aligned} r_{ij} &\leq r_{i,j-1} \text{ and } r_{ij} \leq r_{i+1,j} && \text{for all } i < j, \text{ and} \\ r_{i+1,j-1} - r_{i,j-1} - r_{i+1,j} + r_{ij} &\geq 0 && \text{for all } i < j - 1 , \end{aligned}$$

where we set $r_{ii} = \text{rank}(E_i)$. In fact, rank conditions satisfying (1.2) are the only conditions that can actually *occur*, i.e. for which one can have equality in (1.1). When the maps are sufficiently generic, each such Ω_r is irreducible, of codimension

$$(1.3) \quad d(r) = \sum_{i < j} (r_{i,j-1} - r_{ij})(r_{i+1,j} - r_{ij}) .$$

When $n = 1$, the formula for Ω_r is the well-known Giambelli-Thom-Porteous formula, which we recall in order to introduce some notation. For a map $\phi : E \rightarrow F$ of vector bundles of ranks e and f , and a non-negative integer $r \leq \min(e, f)$, Ω_r is the locus where ϕ has rank at most r . The formula for Ω_r is the Schur polynomial

$$s_{(e-r)^{f-r}}(F - E) ,$$

which is defined as follows. Define cohomology classes h_i by the formula $\sum h_i = c(E^\vee)/c(F^\vee)$, where $c(E^\vee) = 1 - c_1(E) + c_2(E) - \dots$ is the total Chern class, and the division is carried out formally; in particular, $h_0 = 1$ and $h_i = 0$ for $i < 0$. For any sequence $\lambda = (\lambda_1, \dots, \lambda_p)$ of non-negative integers, set

$$s_\lambda(F - E) = \det(h_{\lambda_i+j-i})_{1 \leq i, j \leq p} .$$

In the Giambelli-Thom-Porteous formula, $\lambda = (e - r)^{f-r}$ denotes the sequence $e - r$ repeated $f - r$ times. In a Schur determinant $s_\lambda(F - E)$, λ will usually be a partition, i.e. a weakly decreasing sequence, but later we will also need this notation when λ is not a partition.

Our general formula for the locus Ω_r , when r is any set of rank conditions satisfying (1.2), has the form

$$\sum_{\lambda} c_\lambda(r) s_\lambda(E_\bullet) ,$$

where the sum is over sequences $\lambda = (\lambda(1), \lambda(2), \dots, \lambda(n))$, with each $\lambda(i)$ a partition. The class $s_\lambda(E_\bullet)$ is defined to be

$$s_\lambda(E_\bullet) = s_{\lambda(1)}(E_1 - E_0) \cdot s_{\lambda(2)}(E_2 - E_1) \cdot \dots \cdot s_{\lambda(n)}(E_n - E_{n-1}) .$$

The coefficients $c_\lambda(r)$ are certain integers for which we give an inductive formula.

A second purpose of this paper is to introduce these integers $c_\lambda(r)$, which we regard as generalized Littlewood-Richardson coefficients.

We have a conjectured formula for $c_\lambda(r)$ as the number of sequences (T_1, T_2, \dots, T_n) of Young tableaux, with T_i of shape $\lambda(i)$, satisfying certain conditions. This formula has been proved when the number of bundles is at most four, but it appears to be a difficult combinatorial problem to prove it in general.

In [8] a special case of this situation was studied, where the rank conditions are given by a permutation w . For maps

$$G_1 \rightarrow G_2 \rightarrow \dots \rightarrow G_m \rightarrow F_m \rightarrow F_{m-1} \rightarrow \dots \rightarrow F_1$$

with $\text{rank}(G_i) = \text{rank}(F_i) = i$, and $w \in S_{m+1}$, let

$$\Omega_w = \{x \in X \mid \text{rank}(G_q(x) \rightarrow F_p(x)) \leq r_w(p, q) \ \forall \ p, q \leq m\} \ ,$$

where $r_w(p, q) = \#\{i \leq p \mid w(i) \leq q\}$. These loci are special cases of the loci Ω_r described in this paper. The formulas given here therefore specialize to the *universal double Schubert polynomials* $\mathfrak{S}_w(c_\bullet(F_\bullet); c_\bullet(G_\bullet))$ for these loci. Since these universal Schubert polynomials specialize to quantum and double Schubert polynomials ([5], [16], [13], [7]), we derive formulas for these important polynomials. These formulas appear to be new even for the single Schubert polynomials $\mathfrak{S}_w(x)$.

Among the loci considered here are the *varieties of complexes*, which are the loci Ω_r with $r_{ij} = 0$ for $j - i \geq 2$. In this case the formula for the coefficients $c_\lambda(r)$ is particularly simple, and it agrees with our general conjectured formula.

In Section 2 we discuss the loci Ω_r in more detail, state the main theorem, and derive the main applications. This includes a precise statement of what it means for a polynomial $\sum c_\lambda(r) s_\lambda(E_\bullet)$ to give a formula for a locus Ω_r . This statement implies the assertion that if X is non-singular and Ω_r has the expected codimension $d(r)$, then

$$(1.4) \qquad [\Omega_r] = \sum_\lambda c_\lambda(r) s_\lambda(E_\bullet)$$

in the Chow group $A^{d(r)}(X)$. However, weaker assertions can be made when X is singular or the maps ϕ_i are less generic. At the end of Section 3 we sketch a generalization, which is based on explicit resolutions of singularities of these loci.

The coefficients $c_\lambda(r)$ are determined by the geometry, if this assertion is interpreted correctly. We will see in Section 2 that $c_\lambda(r)$ depends only on the differences $r_{i,j-1} - r_{ij}$ and $r_{i+1,j} - r_{ij}$. This allows the ranks of the bundles E_i to be taken large compared to the expected

codimension $d(r)$; if the Chern classes of the bundles are independent, the coefficients $c_\lambda(r)$ are then uniquely determined by (1.4).

Much of the work in a project of this kind – discovering the shape of the formula – is invisible in the final product, which has a short proof (given in Section 3). In particular, it came as a pleasant surprise to us that the polynomials for all the loci Ω_r can be written as a linear combination of the polynomials $s_\lambda(E_\bullet)$. We know of no reason for this other than the proof of the explicit formula. That the coefficients $c_\lambda(r)$ appear to be non-negative is even more surprising.

The conjectured formula for the coefficients is discussed in more detail in the final Section 4; proofs of the combinatorial assertions made there can be found in [4].

We are particularly grateful to S. Fomin, who provided an involution on pairs of tableaux which gave us the strongest evidence for the conjectured formula, and who has collaborated with us on the combinatorial aspects of this problem. Thanks also to M. Haiman and M. Shimozono for their responses to combinatorial questions. The Schubert package [12] was useful for calculations.

2. Quiver varieties; the theorem and applications

2.1. The main theorem. Given vector bundles E_0, \dots, E_n on a variety X , let H be the direct sum of the bundles $\text{Hom}(E_{i-1}, E_i)$, i.e.

$$H = \text{Hom}(E_0, E_1) \times_X \text{Hom}(E_1, E_2) \times_X \cdots \times_X \text{Hom}(E_{n-1}, E_n) .$$

Writing \tilde{E}_i for the pullback of E_i to H , we have a universal or tautological sequence of bundle maps

$$(2.1) \quad \tilde{E}_0 \xrightarrow{\Phi_1} \tilde{E}_1 \xrightarrow{\Phi_2} \tilde{E}_2 \longrightarrow \cdots \longrightarrow \tilde{E}_{n-1} \xrightarrow{\Phi_n} \tilde{E}_n$$

on H . For this universal case, it is a theorem of Lakshmibai and Magyar [14] that for r satisfying (1.2), the scheme $\tilde{\Omega}_r = \Omega_r(\tilde{E}_\bullet)$ for (2.1) is reduced and irreducible, of codimension $d(r)$. Moreover, $\tilde{\Omega}_r$ is a Cohen-Macaulay variety if X is Cohen-Macaulay. (Earlier Abeasis, del Fra, and Kraft [2] had shown, in characteristic zero, that the reduced scheme $(\tilde{\Omega}_r)_{\text{red}}$ is Cohen-Macaulay.) Note that, when the bundles are trivial, H is a Cartesian product of X and a product M of spaces of matrices, and $\tilde{\Omega}_r$ is a product of X with the corresponding locus in M ; it is this locus in M that is studied in [1], [2], and [14].

The statement that “the polynomial $P = \sum c_\lambda(r)s_\lambda(E_\bullet)$ is a formula for the locus Ω_r ,” has the usual meaning in intersection theory

(cf. [6, §14], [10, App. A]). It implies that when X is non-singular and $\text{codim}(\Omega_r, X) = d(r)$, then

$$[\Omega_r] = \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(E_{\bullet})$$

in the Chow group $A^{d(r)}X$, where $[\Omega_r]$ is the cycle defined by the scheme Ω_r . For arbitrary X and maps ϕ_i , there is a well defined cycle class Ω_r in the Chow group $A_{m-d(r)}(\Omega_r)$, where $m = \dim(X)$, whose image in $A_{m-d(r)}(X)$ is the class $\sum c_{\lambda}(r) s_{\lambda}(E_{\bullet}) \cap [X]$. Whenever Ω_r has codimension $d(r)$ in X , Ω_r is a positive cycle supported on Ω_r ; if X is Cohen-Macaulay, or more generally if $\text{depth}(\Omega_r, X) = d(r)$, this cycle is $[\Omega_r]$, but if X is not Cohen-Macaulay the coefficient of a component of Ω_r in Ω_r may be smaller than the length of Ω_r at its generic point. These classes Ω_r are compatible with the basic constructions of intersection theory, exactly as in [6, Thm. 14.3].

In fact, to give maps $\phi_i : E_{i-1} \rightarrow E_i$ for all i is the same as giving a section $s : X \rightarrow H$ of the bundle H , and $\tilde{\Omega}_r = s^{-1}(\tilde{\Omega}_r)$. The general class Ω_r is constructed by intersecting $\tilde{\Omega}_r \subset H$ with the (regular) embedding $s : X \rightarrow H$, i.e.

$$\Omega_r = s^! [\tilde{\Omega}_r] ,$$

where $s^! : A_*(\tilde{\Omega}_r) \rightarrow A_*(\Omega_r)$ is the refined intersection [6, §6]. As in [6, §14], the general properties of these classes follow from this construction. It therefore suffices to prove the corresponding formula on H , i.e. that

$$[\tilde{\Omega}_r] = \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(\tilde{E}_{\bullet}) \cap [H]$$

in $A_{N-d(r)}(H)$, where $N = \dim(H) = \dim(X) + \sum_{i=1}^n e_{i-1}e_i$, $e_i = \text{rank}(E_i)$.


It is natural to arrange the rank conditions in a triangular array:

$$\begin{array}{cccccccc}
 r_{00} & & r_{11} & & r_{22} & & r_{33} & & \cdots & & r_{nn} \\
 & r_{01} & & r_{12} & & r_{23} & & \cdots & & r_{n-1,n} & \\
 & & r_{02} & & r_{13} & & \cdots & & r_{n-2,n} & & \\
 & & & r_{03} & & \cdots & & r_{n-3,n} & & & \\
 & & & & \ddots & & & & & & \\
 & & & & & & & & & & r_{0n}
 \end{array}$$


It is useful to replace each small triangle

$$\begin{array}{cc}
 e & f \\
 & r
 \end{array}$$


occurring in this array by the rectangle of width $e - r$ and height $f - r$.




We then have the rectangular array



where R_{ij} has width $r_{i,j-1} - r_{ij}$ and height $r_{i+1,j} - r_{ij}$. Note that the expected codimension $d(r)$ is the sum of the areas of the rectangles. The condition (1.2) says that the rectangles get (weakly) shorter as one proceeds in a southeasterly direction, and they get (weakly) narrower as one travels southwest. For example, the rank conditions given in the triangular array



correspond to the rectangular array:



Our formula depends on the rectangles in this array. To be precise, it depends on the integers $r_{i,j-1} - r_{ij}$ and $r_{i+1,j} - r_{ij}$ for all $i < j$; if a width $r_{i,j-1} - r_{ij}$ is zero, we need to know the height $r_{i+1,j} - r_{ij}$, even though the rectangle R_{ij} is empty. (The conjectured formula discussed later does not have this defect.) Each R_{ij} is identified with the partition $(r_{i,j-1} - r_{ij})^{r_{i+1,j} - r_{ij}}$ for which it is the Young diagram.

At this point we need some notation. If R is a rectangle of width e and height f , and σ and τ are partitions, with the length $\ell(\sigma)$ of

σ at most f , then $(R + \sigma, \tau)$ denotes the sequence $(e + \sigma_1, e + \sigma_2, \dots, e + \sigma_f, \tau_1, \tau_2, \dots)$; this is a partition only if $e + \sigma_f \geq \tau_1$. For a partition $\lambda = (\lambda_1, \dots, \lambda_p)$, $|\lambda|$ denotes $\sum \lambda_i$, which is the number of boxes in the Young diagram of λ . For partitions λ, σ, τ with $|\sigma| + |\tau| = |\lambda|$, $c_{\sigma, \tau}^\lambda$ denotes the Littlewood-Richardson coefficient, which is the coefficient of the Schur polynomial s_λ in the expansion of $s_\sigma \cdot s_\tau$ (see [17]). We set $e_i = r_{ii} = \text{rank}(E_i)$, $r_i = r_{i-1, i}$, and $R_i = R_{i-1, i}$, so R_i has height $e_i - r_i$ and width $e_{i-1} - r_i$.

If $I = (i_1, \dots, i_p)$ is a sequence of non-negative integers that is not weakly decreasing, then $s_I(F - E)$ is either 0 or it is $\pm s_\lambda(F - E)$ for some unique partition λ and unique coefficient ± 1 . This partition and coefficient can be found by performing a sequence of moves of the type

$$(j_1, \dots, j_p) \mapsto (j_1, \dots, j_{k-1}, j_{k+1} - 1, j_k + 1, j_{k+2}, \dots, j_p)$$

if $j_{k+1} > j_k$; if one reaches a sequence (j_1, \dots, j_p) with some $j_{k+1} = j_k + 1$, then $s_I = 0$; otherwise one reaches a partition $\lambda = (\lambda_1, \dots, \lambda_p)$ in m steps, and then $s_I = (-1)^m s_\lambda$.

We now give an algorithm for constructing finite formal sums $\sum c_\lambda(r)S(\lambda)$, with λ varying over n -tuples of partitions $\lambda = (\lambda(1), \dots, \lambda(n))$. The polynomial for the degeneracy locus Ω_r will be obtained by replacing each $S(\lambda)$ by $s_\lambda(E_\bullet)$. In the algorithm we will meet symbols $S(I(1), \dots, I(n))$ where each $I(j)$ is a sequence of integers. For such symbols we imitate the above rule for Schur polynomials to write $S(I(1), \dots, I(n))$ as either zero or $\pm S(\lambda(1), \dots, \lambda(n))$ for unique partitions $\lambda(1), \dots, \lambda(n)$. If $s_{I(j)} = 0$ for any j , put $S(I(1), \dots, I(n)) = 0$; otherwise write $s_{I(j)} = \varepsilon_j s_{\lambda(j)}$ for $1 \leq j \leq n$, with $\varepsilon_j = \pm 1$, and put $S(I(1), \dots, I(n)) = (\prod \varepsilon_j) S(\lambda(1), \dots, \lambda(n))$.

We construct the polynomial $\sum c_\lambda(r)S(\lambda)$ by induction on n . For $n = 1$ we have just one rectangle $R = R_{01}$, and the polynomial is $S(R)$, which gives $s_R(E_1 - E_0)$. Given the rectangular array for r , delete the top row. This gives a smaller array, for which we have a polynomial $\sum d_\mu S(\mu)$ by induction, the sum over sequences $\mu = (\mu(1), \dots, \mu(n - 1))$ of partitions. The polynomial $\sum c_\lambda(r)S(\lambda)$ is obtained by replacing each $S(\mu)$ in $\sum d_\mu S(\mu)$ by


$$\sum \left(\prod_{i=1}^{n-1} c_{\sigma(i), \tau(i)}^{\mu(i)} \right) S(I(1), \dots, I(n)) \quad .$$

Here the sum is over all sequences $(\sigma(1), \dots, \sigma(n - 1))$ and $(\tau(1), \dots, \tau(n - 1))$ of partitions, with $|\sigma(i)| + |\tau(i)| = |\mu(i)|$, such that

the length of $\sigma(i)$ is at most the height of R_i , i.e. $\ell(\sigma(i)) \leq e_i - r_i$. Define $I(i)$ to be $(R_i + \sigma(i), \tau(i - 1))$ for $1 \leq i \leq n$, where $\tau(0)$ and $\sigma(n)$ are taken to be the empty partition. One uses the rules just given to write each $S(I(1), \dots, I(n))$ as 0 or $\pm S(\lambda)$ for a unique $\lambda = (\lambda(1), \dots, \lambda(n))$, thus arriving at a polynomial $\sum c_\lambda(r) S(\lambda)$.


Main Theorem. *The formula for Ω_r is $\sum c_\lambda(r) s_\lambda(E_\bullet)$.*

This theorem will be proved in the next section. We first interpret it in the case where the rectangular array has only two non-empty rows, i.e. $R_{ij} = \emptyset$ if $j - i > 2$. In this case the inductive polynomial $\sum d_\mu S(\mu)$ is just $S(\mu)$, for $\mu = (R_{02}, R_{13}, \dots, R_{n-2,n})$. For a rectangle shape R , the Littlewood-Richardson coefficient $c_{\sigma,\tau}^R$ vanishes unless σ and the 180° rotation of τ fit together to make R , in which case $c_{\sigma,\tau}^R = 1$.



Corollary 1. *If R_{ij} is empty for $j - i > 2$, then the formula for Ω_r is $\sum s_\lambda(E_\bullet)$, where the sum is over all $\lambda = (\lambda(1), \dots, \lambda(n))$, with $\lambda(i) = (R_i + \sigma(i), \tau(i - 1))$, such that $\sigma(i)$ and $\tau(i)$ fit together to form $R_{i-1,i+1}$ for $1 \leq i \leq n - 1$; here $\sigma(n)$ and $\tau(0)$ are empty.*


Note that, by (1.2), for any division of $R_{i-1,i+1}$ into $\sigma(i)$ and $\tau(i)$, $\sigma(i)$ always fits on the right side of R_i , and $\tau(i)$ fits below R_{i+1} , so the resulting sequences $\lambda(i) = (R_i + \sigma(i), \tau(i - 1))$ are always partitions. This formula can be remembered by the picture



The situation in the corollary covers the case of varieties of complexes, which means that $r_{ij} = 0$ for $j - i \geq 2$. In this case the triangular array is

$$\begin{array}{cccccccc}
 e_0 & e_1 & e_2 & e_3 & \cdots & e_n \\
 & r_1 & r_2 & r_3 & \cdots & r_n \\
 & & 0 & 0 & \cdots & 0
 \end{array}$$

so the array of rectangles is




P. Pragacz reports that he had known this formula for Ω_r in the case of varieties of complexes.

2.2. *Geometric description of $c_\lambda(r)$.* Although we use the notation $c_\lambda(r)$ for the coefficients, it should be emphasized that they depend only on the differences $r_{i,j-1} - r_{ij}$ and $r_{i+1,j} - r_{ij}$, not on the integers r_{ij} themselves. For example they are unchanged if the same positive integer is added to each r_{ij} .


The linear independence of ordinary Schur polynomials $s_\lambda(x_1, \dots, x_p)$ as λ varies over partitions of length at most p , implies that the polynomials $s_\lambda(E_\bullet)$ are linearly independent functions of the Chern classes of the bundles E_0, E_1, \dots, E_n , if the ranks of the bundles are suitably large (e.g. if $\ell(\lambda(i)) \leq e_i$ for $1 \leq i \leq n$).

From the preceding two paragraphs it follows that the coefficients $c_\lambda(r)$ are uniquely determined by the geometry, i.e. by the fact that $\sum c_\lambda(r) s_\lambda(E_\bullet)$ is a formula for Ω_r . To see this, one can choose the ranks e_i large, and one can find a smooth variety X on which Ω_r has the expected codimension $d(r)$, and for which the classes $s_\lambda(E_\bullet)$, for $\sum |\lambda(i)| = d(r)$, are linearly independent. For example, one can start with universal bundles E_i on large Grassmannians G_i , let $G = \prod_{i=0}^n G_i$, and set $X = \bigoplus_{i=1}^n \text{Hom}(E_{i-1}, E_i)$.

2.3. *Schubert polynomials.* In [1], the rank conditions r satisfying (1.2) are described by diagrams of dots connected by lines. One puts $e_i = r_{ii}$ dots in column i , and lines are drawn between some dots in adjacent columns. Then r_{ij} is the number of lines connecting a dot in column i to a dot in column j . The example given earlier in this section can be described by the diagram



Now fix a positive integer m . For a permutation $w \in S_{m+1}$, we form a diagram with $2m$ columns of lengths $1, 2, \dots, m, m, m - 1, \dots, 2, 1$. All possible lines are drawn among the first m columns and among the last m columns. Between the two middle columns, the i 'th dot on the right is connected to the $w(i)$ 'th dot on the left. If $w(i) = m + 1$, no connection is made. For example, if $m = 4$ and $w = 31452$, this diagram is



The number of connections between the left column with q dots and the right column with p dots is the number

$$r_w(p, q) = \#\{i \leq p \mid w(i) \leq q\} \quad .$$

There are the maximal number of connections between two columns on the left or between two on the right. This means that for a sequence E_\bullet of bundle maps

$$G_1 \rightarrow G_2 \rightarrow \dots \rightarrow G_m \rightarrow F_m \rightarrow F_{m-1} \rightarrow \dots \rightarrow F_1$$

with $\text{rank}(G_i) = \text{rank}(F_i) = i$, the locus $\Omega_r(E_\bullet)$ defined in the introduction is exactly the locus Ω_w defined in [8], with the same scheme structure. In [8] “universal Schubert polynomials” $\mathfrak{S}_w(c_\bullet(F_\bullet); c_\bullet(G_\bullet))$ were constructed, which represent the loci Ω_w . From the fact that the formula for a locus is unique, we deduce the following corollary.

Corollary 2. *With r determined by w as above,*

$$\mathfrak{S}_w(c_\bullet(F_\bullet); c_\bullet(G_\bullet)) = \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(E_\bullet) \quad .$$

When these bundle maps are specialized so that each $G_{i-1} \rightarrow G_i$ is an inclusion of bundles, and each $F_i \rightarrow F_{i-1}$ is a surjection, then $\mathfrak{S}_w(c_\bullet(F_\bullet); c_\bullet(G_\bullet))$ becomes the double Schubert polynomial

$$\mathfrak{S}_w(x_1, \dots, x_m; y_1, \dots, y_m)$$

of Lascoux and Schützenberger; here we set $x_i = c_1(\ker(F_i \rightarrow F_{i-1}))$ and $y_i = c_1(G_i/G_{i-1})$. The right side of the formula in this corollary also simplifies in this case. It follows from the definition that for a partition τ , we have

$$s_\tau(G_i - G_{i-1}) = \begin{cases} (y_i)^q & \text{if } \tau = (q), q \geq 0 \\ 0 & \text{otherwise,} \end{cases}$$

and

$$s_\tau(F_{i-1} - F_i) = \begin{cases} (-x_i)^p & \text{if } \tau = (1)^p, p \geq 0 \\ 0 & \text{otherwise.} \end{cases}$$


Thus $s_\lambda(E_\bullet) = 0$ unless $\lambda = ((q_2), (q_3), \dots, (q_m), \tau, (1)^{p_m}, \dots, (1)^{p_2})$, in which case

$$s_\lambda(E_\bullet) = (-1)^{p_2 + \dots + p_m} x_2^{p_2} \dots x_m^{p_m} y_2^{q_2} \dots y_m^{q_m} s_\tau(x/y),$$

where $s_\tau(x/y) = \det(h_{\tau_i+j-i})$, $\sum h_k = \prod(1 - y_i) / \prod(1 - x_j)$. Our formula therefore writes $\mathfrak{S}_w(x; y)$ as a signed sum of monomials in the $x_2, \dots, x_m, y_2, \dots, y_m$ times Schur polynomials $s_\tau(x/y)$. When all variables y_i are set equal to zero, we have only the terms with $q_2 = \dots = q_m = 0$, and this writes the ordinary Schubert polynomial $\mathfrak{S}_w(x) = \mathfrak{S}_w(x; 0)$ as a signed sum of monomials in x_2, \dots, x_m times (symmetric) Schur polynomials $s_\tau(x)$.

Unlike the inductive construction of Schubert polynomials from high degree to low degree, our formulas are simplest for those of low degree.

For example, for $w = 3142 \in S_4$, the corresponding array of rectangles is



Calculating $\sum c_\lambda(r)S(\lambda)$ with the algorithm of the main theorem, working from the bottom up, one has, with \emptyset the empty partition,

$$\begin{aligned}
 S(\emptyset) &\rightsquigarrow S(\emptyset, 1) \\
 &\rightsquigarrow S(\emptyset, 1, \emptyset) + S(\emptyset, \emptyset, 1) \\
 &\rightsquigarrow S(\emptyset, 2, \emptyset, \emptyset) + S(\emptyset, 1, 1, \emptyset) + S(\emptyset, 1, \emptyset, 1) \\
 &\rightsquigarrow \sum c_\lambda(r)S(\lambda) = S(\emptyset, 2, 1, \emptyset, \emptyset) + S(\emptyset, 1, 11, \emptyset, \emptyset) + \\
 &\quad S(\emptyset, 1, 2, \emptyset, \emptyset) + S(\emptyset, 1, 1, 1, \emptyset) + S(\emptyset, \emptyset, 21, \emptyset, \emptyset) + \\
 &\quad S(\emptyset, \emptyset, 11, 1, \emptyset) + S(\emptyset, 1, 1, \emptyset, 1) + S(\emptyset, \emptyset, 11, \emptyset, 1) .
 \end{aligned}$$

Substituting $s_\lambda(E_\bullet)$ for $S(\lambda)$, this is a formula for the universal double Schubert polynomial associated to $w = 3142$. It specializes to the formula

$$\begin{aligned}
 \mathfrak{S}_{3142}(x; y) &= s_{21}(x/y) + y_3s_2(x/y) + (y_3 - x_2 - x_3)s_{11}(x/y) + \\
 &\quad y_3(y_3 - x_2 - x_3)s_1(x/y)
 \end{aligned}$$

and to

$$\mathfrak{S}_{3142}(x) = s_{21}(x) - (x_2 + x_3)s_{11}(x) .$$

The rectangular array coming from a permutation $w \in S_{m+1}$ has only empty rectangles and 1×1 rectangles. In fact, this array is determined from the diagram of the permutation denoted $D'(w)$ in [8, §2]: the diagram $D'(w)$ is reflected in a vertical line, then rotated 135° clockwise, and the result fitted in the bottom of the triangle; each box in $D'(w)$ is then in the position of a non-empty rectangle of the rectangular array. In other words, the rectangle R_{ij} is non-empty iff $D'(w)$ contains a box in position $(2m - j, i + 1)$, which happens exactly when we have both $w(2m + 1 - j) \leq i + 1$ and $w^{-1}(i + 2) \leq 2m - j$.

3. Proof of the Theorem

3.1. Geometric preliminaries. It follows from the general discussion in Section 2 that it suffices to prove the formula for the universal locus $\Omega_r(E_\bullet)$ in $H = \bigoplus \text{Hom}(E_{i-1}, E_i)$. (Throughout this section we omit notation for pullbacks of bundles by canonical maps.) In particular, we know from [2] and [14] that $\Omega_r(E_\bullet)$ is reduced and irreducible of the expected codimension $d(r)$, and that $\Omega_r(E_\bullet)$ is the closure of the locus $\Omega_r^\circ(E_\bullet)$ where each of the maps $E_i(x) \rightarrow E_j(x)$ has rank equal to r_{ij} for $i < j$. We must prove that, with these assumptions,

$$[\Omega_r(E_\bullet)] = \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(E_\bullet) \cap [H]$$

in the Chow group $A_{N-d(r)}(H)$, $N = \dim(H)$.

Form the Grassmannian bundle G_0 over X with $r_i = r_{i-1,i}$ as in §2.1:

$$G_0 = \text{Gr}(r_1, E_1) \times_X \text{Gr}(r_2, E_2) \times_X \cdots \times_X \text{Gr}(r_n, E_n) .$$

Let $G = G_0 \times_X H$, with projection $\pi : G \rightarrow H$. Let $0 \rightarrow A_i \rightarrow E_i \rightarrow Q_i \rightarrow 0$ be the universal exact sequences on G_0 , and hence also on G . Let $Z \subset G$ be the intersection of the zero-schemes of the canonical maps $E_{i-1} \rightarrow E_i \rightarrow Q_i$, i.e.

$$Z = Z(E_0 \rightarrow Q_1) \cap Z(E_1 \rightarrow Q_2) \cap \cdots \cap Z(E_{n-1} \rightarrow Q_n) .$$

On Z we have maps $E_{i-1} \rightarrow A_i$ for $1 \leq i \leq n$. Composing these with the inclusions $A_{i-1} \subset E_{i-1}$ we get a sequence A_\bullet of bundles and bundle maps on Z :

$$A_1 \rightarrow A_2 \rightarrow \cdots \rightarrow A_n .$$

Let \bar{r} denote the rank conditions obtained by omitting the top row of the triangular array for r , and let $\Omega_{\bar{r}}(A_\bullet) \subset Z$ be the locus given by these maps and rank conditions. It is easy to see that $\Omega_{\bar{r}}(A_\bullet)$ is mapped into $\Omega_r(E_\bullet)$ by π .

Now Z is isomorphic to the bundle $\bigoplus_{i=1}^n \text{Hom}(E_{i-1}, A_i)$ over G_0 , and we have a canonical projection

$$\rho : Z = \bigoplus_{i=1}^n \text{Hom}(E_{i-1}, A_i) \rightarrow \bigoplus_{i=2}^n \text{Hom}(A_{i-1}, A_i) = H' .$$

Denote by Ω' the universal locus $\Omega_{\bar{r}}(A_\bullet)$ on H' . Then $\Omega_{\bar{r}}(A_\bullet)$ in Z is the inverse image of Ω' by ρ . Since the maps on H' are universal, it follows that Ω' is irreducible, and therefore $\Omega_{\bar{r}}(A_\bullet)$ is an irreducible subscheme of Z .

$$\begin{array}{ccccc}
 \Omega' & \longleftarrow & \Omega_{\bar{r}}(A_\bullet) & \longrightarrow & \Omega_r(E_\bullet) \\
 \downarrow & & \downarrow & & \downarrow \\
 H' & \xleftarrow{\rho} & Z & \xrightarrow{\quad} & G & \xrightarrow{\pi} & H \\
 & & & & \downarrow & & \downarrow \\
 & & & & G_0 & \longrightarrow & X
 \end{array}$$

Lemma 1. π maps $\Omega_{\bar{r}}(A_{\bullet})$ birationally onto $\Omega_r(E_{\bullet})$.

Proof. Let Z° denote the open subset of Z where the maps $E_{i-1} \rightarrow A_i$ are surjective. Then the schemes $\Omega_{\bar{r}}^{\circ}(A_{\bullet}) \cap Z^{\circ}$ and $\Omega_r^{\circ}(E_{\bullet})$ are isomorphic; they are both universal objects in the category of schemes Y over H , such that the pullback to Y of the sequence E_{\bullet} satisfies $\text{rank}(E_i \rightarrow E_j) = r_{ij}$ everywhere on Y . Since both $\Omega_{\bar{r}}(A_{\bullet})$ and $\Omega_r(E_{\bullet})$ are irreducible schemes, and since $\Omega_r^{\circ}(E_{\bullet}) \neq \emptyset$, the assertion follows. \square

Note that this argument gives a direct proof that the codimension of $\Omega_r(E_{\bullet})$ in H is $d(r)$. Indeed, by induction we know that Ω' has codimension $d(\bar{r})$ in H' , and $\Omega_{\bar{r}}(A_{\bullet})$ must then have the same codimension in Z . We conclude that the codimension of $\Omega_r(E_{\bullet})$ in H is

$$d(\bar{r}) + \dim(H) - \dim(Z) = d(r) .$$

3.2. *Proof of the main theorem.* By induction on n we know that $[\Omega'] = \sum c_{\mu}(\bar{r}) s_{\mu}(A_{\bullet}) \cap [H']$, so

$$[\Omega_{\bar{r}}(A_{\bullet})] = \rho^*[\Omega'] = \sum_{\mu} c_{\mu}(\bar{r}) s_{\mu}(A_{\bullet}) \cap [Z] .$$

Furthermore $[Z] = \prod_{i=1}^n c_{\text{top}}(\text{Hom}(E_{i-1}, Q_i)) \cap [G]$, so

$$[Z] = \prod_{i=1}^n s_{R'_i}(Q_i - E_{i-1}) \cap [G] ,$$

where $R'_i = (e_{i-1})^{e_i - r_i}$. Since $\pi_*[\Omega_{\bar{r}}(A_{\bullet})] = [\Omega_r(E_{\bullet})]$ by Lemma 1, we are therefore reduced to proving the identity

$$\pi_* \left(\sum_{\mu} c_{\mu}(\bar{r}) s_{\mu}(A_{\bullet}) \cdot \prod_{i=1}^n s_{R'_i}(Q_i - E_{i-1}) \cap [G] \right) = \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(E_{\bullet}) \cap [H] . \tag{3.1}$$

For this we need the following basic Gysin formula of Pragacz [18, Prop. 2.2], whose proof comes from [11], cf. [10, App. F].

Lemma 2. Let E and F be vector bundles of ranks e and f on a variety X . Let $0 \leq d \leq \min(e, f)$. Let $G = \text{Gr}(d, F)$ be the Grassmann bundle, with universal exact sequence $0 \rightarrow A \rightarrow F \rightarrow Q \rightarrow 0$ and projection

$\pi: G \rightarrow X$. Let $q = f - d$, $R = (e - d)^q$, and $R' = (e)^q$. For any partitions λ and μ , with λ of length at most q , and any $\alpha \in A_*(X)$,

$$\pi_*(s_{R'+\lambda}(Q - E)s_\mu(A - E) \cap \pi^* \alpha) = s_{R+\lambda, \mu}(F - E) \cap \alpha .$$

We also need the following special case of the factorization formula of Lascoux and Schützenberger [16] and Berele and Reger [3], cf. [18].

Lemma 3. *Let E and F be vector bundles of ranks e and f . Let $R = (e)^f$. Let λ be a partition of length at most f . Then*

$$s_\lambda(F)s_R(F - E) = s_{R+\lambda}(F - E) .$$

Note that this identity follows from Lemma 2. Finally we need the basic identity [17, §1.5]:

Lemma 4. *For bundles E_1, E_2 , and E_3 , and a partition μ ,*

$$s_\mu(E_3 - E_1) = \sum c_{\sigma\tau}^\mu s_\sigma(E_2 - E_1)s_\tau(E_3 - E_2) ,$$

the sum over partitions σ and τ with $|\sigma| + |\tau| = |\mu|$, with $c_{\sigma,\tau}^\mu$ the Littlewood-Richardson coefficient.

Now we can prove (3.1). First use Lemma 4 to replace each factor $s_{\mu(i)}(A_{i+1} - A_i)$ that occurs in each $s_\mu(A_\bullet)$ on the left side of (3.1) by the sum

$$\begin{aligned} \sum c_{\sigma(i), \tau(i)}^{\mu(i)} s_{\sigma(i)}(E_i - A_i)s_{\tau(i)}(A_{i+1} - E_i) \\ = \sum c_{\sigma(i), \tau(i)}^{\mu(i)} s_{\sigma(i)}(Q_i)s_{\tau(i)}(A_{i+1} - E_i) . \end{aligned}$$

Note that $s_{\sigma(i)}(Q_i) = 0$ if $\ell(\sigma(i)) > \text{rank}(Q_i) = e_i - r_i$. Next use Lemma 3 to replace each $s_{\sigma(i)}(Q_i) \cdot s_{R'_i}(Q_i - E_{i-1})$ in the result by $s_{R'_i + \sigma(i)}(Q_i - E_{i-1})$. The left side of (3.1) becomes

$$\begin{aligned} \sum_\mu c_\mu(\bar{r}) \sum_{\sigma(i), \tau(i)} \left(\prod_{i=1}^{n-1} c_{\sigma(i), \tau(i)}^{\mu(i)} \right) \cdot \\ \pi_* \left(\prod_{i=1}^n s_{R'_i + \sigma(i)}(Q_i - E_{i-1})s_{\tau(i-1)}(A_i - E_{i-1}) \cap [G] \right) . \end{aligned}$$

Finally, n applications of Lemma 2 yields

$$\begin{aligned} & \pi_* \left(\prod_{i=1}^n s_{R'_i + \sigma(i)}(Q_i - E_{i-1}) s_{\tau(i-1)}(A_i - E_{i-1}) \cap [G] \right) \\ &= \prod_{i=1}^n s_{R_i + \sigma(i), \tau(i-1)}(E_i - E_{i-1}) \cap [H] \ , \end{aligned}$$

and this gives the required formula

$$\begin{aligned} & \sum_{\mu} c_{\mu}(\bar{r}) \sum_{\sigma(i), \tau(i)} \left(\prod_{i=1}^{n-1} c_{\sigma(i), \tau(i)}^{\mu(i)} \right) \prod_{i=1}^n s_{R_i + \sigma(i), \tau(i-1)}(E_i - E_{i-1}) \cap [H] \\ &= \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(E_{\bullet}) \cap [H] \ . \end{aligned}$$

Although we have stated it for varieties over a field, the theorem (and its proof) extend readily to schemes of finite type over a regular base, as in [6, §20].

3.3. A generalization. There is a generalization of the theorem which may be useful in its own right, and which gives some insight into the proof. (It is not needed in this paper.) Fix E_0, \dots, E_n on X , and $r = (r_{ij})$ satisfying (1.2). Let $H = \bigoplus \text{Hom}(E_{i-1}, E_i)$, on which the tautological bundle maps are universal, and one has the universal locus $\Omega_r \subset H$.

Let $\pi : F \rightarrow H$ be the partial flag bundle parameterizing flags in each E_j of ranks $r_{0j}, r_{1j}, \dots, r_{j-1,j}$. Let $E_{0j} \subset E_{1j} \subset \dots \subset E_{j-1,j} \subset E_j$ denote the tautological flags of vector bundles on F , $1 \leq j \leq n$.

Let $Z \subset F$ be the locus on which the image of $E_{i,j-1}$ by the map $E_{j-1} \rightarrow E_j$ is contained in the subbundle E_{ij} , i.e. Z is the subscheme defined by the vanishing of all maps $E_{i,j-1} \rightarrow E_j/E_{ij}$ for $i < j$.


One sees as in Lemma 1 that π maps Z birationally onto Ω_r . In fact, if X is non-singular, this construction gives a canonical resolution of singularities of the universal locus Ω_r . It is easy to see that the class of Z is given by

$$[Z] = \prod_{i < j} z_{ij} \ ,$$

where $z_{ij} = c_{\text{top}}(\text{Hom}(E_{i,j-1}, E_{i+1,j}/E_{ij}))$.

Consider a path γ through the triangular array for r , going from r_{00} to r_{nm} . The path must be a union of line segments between


neighboring rank conditions, and it must intersect any vertical line at most once.



For each $j = 1, \dots, n$ let k_j be minimal such that γ goes through $r_{k_j,j}$. Let $F(\gamma)$ be the partial flag bundle parameterizing flags in E_j of ranks $r_{k_j,j}, r_{k_j+1,j}, \dots, r_{j-1,j}$, and let $E_{k_j,j} \subset \dots \subset E_{j-1,j} \subset E_j$ be the tautological flags on $F(\gamma)$, $1 \leq j \leq n$. If the path has m line segments, we let A_0, A_1, \dots, A_m denote the vector bundles on $F(\gamma)$ corresponding to the rank conditions passed through by the path. (In the illustration, $m = 7$, and the bundle sequence is $E_{00}, E_{01}, E_{02}, E_{12}, E_{22}, E_{23}, E_{34}, E_{44}$.)

Let $\Omega_r(\gamma) \subset F(\gamma)$ be the subscheme defined by the conditions that each map $E_{i,j-1} \rightarrow E_j/E_{ij}$ vanishes for r_{ij} on or above the path, and $\text{rank}(E_{ip} \rightarrow E_j) \leq r_{ij}$ for r_{ip} on or above the path and $p \leq j$. The canonical maps $F \rightarrow F(\gamma) \rightarrow X$ map Z birationally onto $\Omega_r(\gamma)$ which in turn is mapped birationally onto Ω_r . Our goal is to give a formula for the class of $\Omega_r(\gamma)$. To do this, we define a formal linear combination $\Phi(\gamma)$ of symbols $S(\lambda)$, where λ is a sequence of partitions, one for each line segment in γ . The formula for $\Omega_r(\gamma)$ is obtained by replacing each $S(\lambda(1), \dots, \lambda(m))$ by $\prod_{i=1}^m s_{\lambda(i)}(A_i - A_{i-1})$, and multiplying the result by $\prod z_{ij}$, the product over all i, j such that r_{ij} is on or above γ .

We define $\Phi(\gamma)$ inductively. If γ is the lowest possible path, going from r_{00} to r_{0n} to r_{nm} , then $\Phi(\gamma) = S(\emptyset, \dots, \emptyset)$, where the empty partition \emptyset is repeated $2n$ times. Otherwise, we can find a path γ' that is equal to γ except it goes lower at one place, in one of the following ways:



In Case 1, we obtain $\Phi(\gamma)$ from $\Phi(\gamma')$ by replacing each symbol $S(\dots, \mu, \dots)$ by $\sum c_{\sigma, \tau}^{\mu} S(\dots, \sigma, \tau, \dots)$. Note that in this case $F(\gamma) = F(\gamma')$ and $\Omega(\gamma) = \Omega(\gamma')$. For Case 2, each symbol $S(\dots, \mu, \nu, \dots)$ in $\Phi(\gamma')$ is replaced by the symbol $S(\dots, (R_{ij} + \nu, \mu), \dots)$. Note that the partitions $\lambda(i)$ are always empty for line segments on the left or right edges of the triangle.

The proof that this polynomial gives a formula for $[\Omega_r(\gamma)]$ is similar to that of our main theorem: one shows that $[Z]$ pushes forward to the class of this polynomial. The induction step, in either Case 1 or 2, is more transparent, as changes are made in only one segment of the formula.

If bundle maps and flags of subbundles are given on X corresponding to ranks on or above the path γ , these determine a section $s : X \rightarrow F(\gamma)$, so one has corresponding formulas for the classes $s^*[\Omega_r(\gamma)]$. When γ is the horizontal path across the top of the diagram, we recover the main theorem.

4. On the coefficients $c_{\lambda}(r)$

There are some properties of the coefficients $c_{\lambda}(r)$ that follow from geometry, i.e. from the main theorem, although they are not obvious from the algorithm defining them. We describe these first, and then discuss properties we believe for other reasons. We conclude with a comparison of the numbers $c_{\lambda}(r)$ with Littlewood-Richardson coefficients.

Consider the dual sequence

$$E_n^{\vee} \rightarrow E_{n-1}^{\vee} \rightarrow \dots \rightarrow E_0^{\vee}$$


with dual rank conditions, $\text{rank}(E_j^{\vee} \rightarrow E_i^{\vee}) \leq r_{ij}$, which we denote by r^{\vee} ; then $\Omega_{r^{\vee}}(E_{\bullet}^{\vee}) = \Omega_r(E_{\bullet})$. Note that the rectangular array for r^{\vee} is obtained by reflecting that for r in a vertical line, and replacing each rectangle by its transpose. Using the basic identity that $s_{\lambda}(F - E) = s_{\lambda'}(E^{\vee} - F^{\vee})$, where λ' is the transpose of λ , we find that

$$(4.1) \quad c_{\lambda^{\vee}}(r^{\vee}) = c_{\lambda}(r) \ ,$$

where, if $\lambda = (\lambda(1), \dots, \lambda(n))$, we put $\lambda^{\vee} = (\lambda(n)', \dots, \lambda(1)')$.

It can happen that for some k , all of the rank conditions of the form $\text{rank}(E_i \rightarrow E_k) \leq r_{ik}$ and $\text{rank}(E_k \rightarrow E_j) \leq r_{kj}$ follow from other rank conditions. This happens when, in the rectangle diagram, all the rectangles on the two 45° lines descending from position k are empty.


For the example $G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow F_3 \rightarrow F_2 \rightarrow F_1$ considered at the end of Section 2, with rank conditions r coming from $w = 3142 \in S_4$, the bundles G_1 and F_2 are inessential in this way. If an inessential bundle E_k is omitted, one has a shorter sequence $E'_\bullet : E_0 \rightarrow \cdots \rightarrow E_{k-1} \rightarrow E_{k+1} \rightarrow \cdots \rightarrow E_n$, with the map $E_{k-1} \rightarrow E_{k+1}$ being $\phi_{k+1} \circ \phi_k$, and corresponding rank conditions r' ; the array of rectangles for r' is obtained by omitting the 45° lines of empty rectangles and moving all rectangles below up a row. For example, if G_1 and F_2 are omitted from the example, one has $G_2 \rightarrow G_3 \rightarrow F_3 \rightarrow F_1$, with rectangular array



Lemma 4 can be used to expand any $s_\rho(E_{k+1} - E_{k-1})$ occurring in the formula $\sum c_\mu(r') s_\mu(E'_\bullet)$ as a sum of terms of the form $s_\sigma(E_k - E_{k-1}) \cdot s_\tau(E_{k+1} - E_k)$. Since $\Omega_{r'}(E'_\bullet) = \Omega_r(E_\bullet)$, with this interpretation we have

$$(4.2) \quad \sum_{\mu} c_{\mu}(r') s_{\mu}(E'_{\bullet}) = \sum_{\lambda} c_{\lambda}(r) s_{\lambda}(E_{\bullet}) .$$

Now we turn to our conjectured formula for the coefficients $c_{\lambda}(r)$, which interprets them by counting Young tableaux, in a way similar to and generalizing the classical Littlewood-Richardson rule. Recall that a (semistandard) *Young tableau* is a filling of the boxes in the Young diagram of a partition with integers that are weakly increasing in rows and strictly increasing down columns. Two Young tableaux P and Q can be multiplied to give another Young tableau denoted $P \cdot Q$. One way to do this is to arrange P and Q corner to corner and play the jeu de taquin, sliding inside corners through but keeping the weak and strict orderings. For example, if $P = \begin{smallmatrix} 1 & 2 \\ 3 & 4 \end{smallmatrix}$ and $Q = \begin{smallmatrix} 1 & 1 \\ 4 \end{smallmatrix}$, this can be carried out by the sequence of moves



The final tableau is $P \cdot Q$. The main fact is that this product is independent of choice of the sequence of inside corners, from which it follows that the set of tableaux form an associative monoid, called the plactic monoid. With this notion, the Littlewood-Richardson number

$c_{\mu,\nu}^\lambda$ is the number of ways a given tableau T of shape λ can be factored into a product $T = P \cdot Q$, such that P has shape μ and Q has shape ν . For proofs and relations with Schur polynomials, see [15] or [9].

Given rank conditions r (satisfying (1.2) as always), form the array of rectangles R_{ij} . We choose a fixed tableau T_{ij} on each shape R_{ij} , with the condition that each entry of T_{ij} must be strictly smaller than any entry of T_{kl} if R_{kl} lies in the wedge cut out by 45° lines below R_{ij} , i.e. if $k \leq i$ and $l \geq j$ with $(k, l) \neq (i, j)$.

From this array of rectangular tableaux we will construct a set of n -tuples of tableaux (T_1, \dots, T_n) that we call *factor sequences*. Our conjecture is that $c_\lambda(r)$ is the number of factor sequences (T_1, \dots, T_n) such that T_i has shape $\lambda(i)$ for $1 \leq i \leq n$. We first explain this for $n = 3$, where we start with an array of rectangular tableaux:

$$\begin{array}{ccc} A & B & C \\ & D & E \\ & & F \end{array}$$

Factor F into a product $F = F_1 \cdot F_2$ of tableaux. Pass F_1 up to the left, and multiply it to D from the right. Pass F_2 up to the right and multiply it to E from the left. Then factor the results:

$$D \cdot F_1 = D_1 \cdot D_2 \quad \text{and} \quad F_2 \cdot E = E_1 \cdot E_2 .$$

Pass the results up to the left and right, arriving at tableaux $A \cdot D_1$, $D_2 \cdot B \cdot E_1$, and $E_2 \cdot C$. This gives a factor sequence $(T_1, T_2, T_3) = (A \cdot D_1, D_2 \cdot B \cdot E_1, E_2 \cdot C)$.

In general one proceeds by induction. A factor sequence for the given array of rectangular tableaux is obtained by forming a factor sequence (S_1, \dots, S_{n-1}) for the array of the bottom $n - 1$ rows. Factor each S_i arbitrarily into $S_i = P_i \cdot Q_i$. Then

$$(T_1, \dots, T_n) = (T_{01} \cdot P_1, \dots, Q_{i-1} \cdot T_{i-1,i} \cdot P_i, \dots, Q_{n-1} \cdot T_{n-1,n})$$

is a factor sequence for the given array.

Conjecture. $c_\lambda(r)$ is the number of factor sequences (T_1, \dots, T_n) of shape $\lambda = (\lambda(1), \dots, \lambda(n))$ that can be made from a given array of rectangular tableaux.

The conjecture has a number of consequences:

- C1. Each $c_\lambda(r)$ is a non-negative integer.

C2. *The coefficients $c_\lambda(r)$ depend only on the rectangles R_{ij} , not on their sides.*

This means that if one of the sides of a rectangle R_{ij} is 0, the length of the other side is irrelevant. (The algorithm of the main theorem shows this when the height of a rectangle is 0, but not when the width is 0.)

Implicit in the conjecture is the assertion

C3. *The number of factor sequences of shape λ is independent of choice of fixed tableaux T_{ij} .*

Granting C3, it is not hard to see that the conjectured formula for the $c_\lambda(r)$ satisfies the duality (4.1). For this one chooses the T_{ij} so that no entry appears more than once, and uses the fact that factoring a tableau T with distinct entries into $P \cdot Q$ is equivalent to factoring its conjugate T' into $Q' \cdot P'$. It is also not difficult to verify that the conjectured formula satisfies the property (4.2) for omitting inessential bundles.

The conjecture is true for the case where R_{ij} is empty for $j - i > 2$. This follows from the description in Corollary 1 of Section 2, together with the fact that for a tableau T of rectangular shape R , for each σ and τ that fit together to form R , there is a unique factorization $T = P \cdot Q$ with P of shape σ and Q of shape τ ; conversely, any factorization of T has factors of shapes that fit together to form R .

The conjecture has been proved when $n \leq 3$. More generally, it is proved when R_{ij} is empty for $j - i > 3$ and no two non-empty rectangles in the third row are adjacent. The proof depends on a wonderful involution on pairs of tableaux produced for us by S. Fomin. This proof is given in [4].


One reason that the combinatorial formula is hard to work with is that a given factor sequence can arise in many ways by the algorithm that produces them. At first glance it would appear that to tell if some (T_1, \dots, T_n) is a factor sequence, one would have to test all possible ways of carrying out the sequence of factorings. However, there is a direct test. For this, define P_i to be the part of T_i lying to the right of the rectangle R_i , and define Q_{i-1} to be everything lying below $T_{i-1,i}$ and P_i :

$$T_i = \begin{array}{|c|c|} \hline T_{i-1,i} & P_i \\ \hline Q_{i-1} & \\ \hline \end{array}$$


Then (T_1, \dots, T_n) is a factor sequence if and only if Q_0 and P_n are empty and $(P_1 \cdot Q_1, \dots, P_{n-1} \cdot Q_{n-1})$ is a factor sequence for the lower $n - 1$ rows of the array. By induction this gives a direct algorithm to test, from the top down. Note that this algorithm, like the theorem, uses the height of a rectangle R_{ij} even if its width is zero. This criterion is proved in [4].

The full conjecture follows from an assertion that Fomin’s involution preserves factor sequences. This assertion is true for $n \leq 3$, and it has been verified in 500,000 randomly generated examples for $n \leq 10$. For a discussion of Fomin’s involution and the discussion of this, we refer again to [4].

The numbers $c_\lambda(r)$ generalize Littlewood-Richardson numbers $c_{\alpha,\beta}^\gamma$ in fact as well as in spirit. To see this, take any rectangle R containing γ , and let τ be the complement of γ in R (rotated 180°):



Form the array of rectangles



where the vertical lines are the height of R , the horizontal lines are its width, and the dot is empty. Choose r giving rise to this array. Then

$$c_{\alpha,\beta}^\gamma = c_\lambda(r) ,$$

with $\lambda = (\alpha, \beta, \tau)$. This follows easily from the theorem (and also from the conjecture).

There are analogous conjectures for the coefficients of the polynomials for the more general loci described in §3.3. In particular, all these coefficients should also be positive. Details will be given in [4].

References

- [1] S. Abeasis and A. Del Fra: Degenerations for the representations of an equioriented quiver of type A_m , *Boll. Un. Mat. Ital. Suppl. no. 2*, 157–171 (1980)
- [2] S. Abeasis, A. Del Fra, and H. Kraft: The geometry of representations of A_m , *Math. Ann.* **256**, 401–418 (1981)
- [3] A. Berele and A. Regev: Hook Young diagrams, combinatorics and representations of Lie superalgebras, *Bull. Amer. Math. Soc.* **8**, 337–339 (1983)
- [4] A. S. Buch: Combinatorics of degeneracy loci, to appear
- [5] S. Fomin, S. Gelfand, and A. Postnikov: Quantum Schubert polynomials, *J. Amer. Math. Soc.* **10**, 565–596 (1997)
- [6] W. Fulton: *Intersection Theory*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)*, vol. 2, Springer-Verlag, Berlin 1984, 1998
- [7] W. Fulton: Flags, Schubert polynomials, degeneracy loci, and determinantal formulas, *Duke Math. J.* **65**, 381–420 (1992)
- [8] W. Fulton: Universal Schubert polynomials, To appear in *Duke Math. J.*
- [9] W. Fulton: *Young Tableaux*, London Mathematical Society Student Texts, vol. 35, Cambridge University Press, 1997
- [10] W. Fulton and P. Pragacz: *Schubert varieties and degeneracy loci*, Springer Lecture Notes **1689**, 1998
- [11] T. Józefiak, A. Lascoux, and P. Pragacz: Classes of determinantal varieties associated with symmetric and skew-symmetric matrices, *Izv. Akad. Nauk SSSR Ser. Mat.* **45**, 662–673 (1981)
- [12] S. Katz and S. A. Strømme: “Schubert”, a Maple package for intersection theory and enumerative geometry, Software and documentation available at <ftp://ftp.math.okstate.edu/pub/schubert>
- [13] A. N. Kirillov and T. Maeno: Quantum double Schubert polynomials, quantum Schubert polynomials and Vafa-Intriligator formula, preprint 1996.
- [14] V. Lakshmibai and P. Magyar: Degeneracy schemes and Schubert varieties, preprint, 1997
- [15] A. Lascoux and M.-P. Schützenberger: *Le monoïde plaxique*, *Noncommutative structures in algebra and geometric combinatorics (Naples, 1978)* (Rome), *Quad, “Ricerca Sci.”* vol. **109**, CNR, pp. 129–156 (1981)
- [16] A. Lascoux and M.-P. Schützenberger: *Polynômes de Schubert*, *C. R. Acad. Sci. Paris Sér. I Math.* **294**, 447–450 (1982)
- [17] I. G. Macdonald: *Symmetric functions and Hall polynomials*, Oxford University Press, 1979, 1995
- [18] P. Pragacz: Enumerative geometry of degeneracy loci, *Ann. Sci. École Norm. Sup. (4)* **21**, 413–454 (1988)