

An Existence Theorem for Optimal Control Systems with State Variable in C , and Stochastic Control Problems¹

RICHARD F. BAUM²

Communicated by L. Cesari

Abstract. We consider an existence theorem for control systems whose state variables for every t are in C , the set of continuous functions varying over a given set I . The dependence of the state variables upon $a \in I$ is induced by their dependence upon the initial state and the state equation governing the system. In contrast, the control $u = u(t)$ is taken as a measurable function of t alone. The usual space constraints and boundary conditions are also allowed to vary over $a \in I$, and the cost functional is now taken to be a continuous functional over a suitable class of continuous functions. We also discuss an application of these results to control systems with stochastic boundary conditions.

1. Introduction

In the present paper, we prove an existence theorem for control systems whose state variables for every t are in C , the set of continuous functions varying over a given set I . The dependence of the state variable upon $a \in I$ is induced by its dependence upon the initial state and the state equation governing the system. In contrast, the admissible controls $u = u(t)$ are taken as measurable functions of t alone. The usual space constraints and boundary conditions are also allowed to vary with $a \in I$, and the cost functional is now taken to be a continuous functional over a suitable class of continuous functions. We further assume that the initial time T_1 is fixed and allow the final time $t_2(a)$ to vary with $a \in I$. Thus, the cost functional is of the form $I = K[t_2(a)$,

¹Paper received August 26, 1969. This research was accomplished under Grant No. AF-AFOSR-942-65. The author is grateful to Dr. Lamberto Cesari for his suggestions and assistance in the preparation of this paper.

²Assistant Professor, Department of Industrial Engineering, University of Michigan, Ann Arbor, Michigan.

$x(t_2(a), a)$], $a \in I$. In particular, our existence theorem applies to control systems with stochastic boundary conditions; for these, we can take as cost functional the expectation $I = \int_I g[t_2(a), x(t_2(a), a)] dP$, where g is a continuous, real-valued function and P is a probability measure over I .

The central result of this paper is the existence theorem of Section 4 for control problems with state variable in C . In turn, this theorem is based on a closure theorem with state variable in C which is new and is proved by the use of a new technique along with the general approach of Cesari for the case of state variables in E_n .

In Section 2, we formally present our control systems. In Section 3, we prove various properties of the systems; and, in Section 4, using these properties, we prove the closure theorem and the existence theorem. Finally, in Section 5, we discuss an application of our results to control systems with stochastic boundary conditions. In a future paper, we will present necessary conditions for such systems, these conditions being analogous to Pontryagin's necessary conditions for usual control systems.

2. Description of the System

We consider control systems whose initial conditions are given in terms of a known continuous function and whose state equations may also contain a known continuous function. In contrast, the control is required to be a function of time alone. More precisely, assume $I \subset E_1$, I compact, and $i(a)$ a known continuous function from I into E_n . Let $A(a)$, $a \in I$, denote the constraint sets, where $A(a)$ are compact subsets of the tx -space $E_1 \times E_n$, with t in E_1 , and $x = (x_1, \dots, x_n)$, the space variable, in E_n . We assume that $A = \bigcup_{a \in I} A(a)$ is compact. Let $T_1 < T_2$ be fixed times. For each t in $[T_1, T_2]$, let $U(t)$, the control set, be a subset of the u -space E_m , $u = (u_1, \dots, u_m)$, the control variable. Let $M(a) = \{(t, x, u) : (t, x) \in A(a), u(t) \in U(t)\}$ and $M = \bigcup_{a \in I} M(a) = \{(t, x, u) : (t, x) \in A, u \in U(t)\}$ be compact subsets of E_{1+n+m} , and let $f = (f_1, \dots, f_n)$ be a continuous vector function from M into E_n (see also Remark 2.2). Let $B(a)$, the terminal sets, $a \in I$, be closed subsets of points of E_{1+n} and let $B = \bigcup_{a \in I} B(a)$. Let T_1 denote the fixed initial time, and let τ , the set of terminal times $t_2(a)$, $a \in I$, be a given family of equicontinuous functions, $T_1 \leq t_2(a) \leq T_2$, which is closed in the uniform topology (see Remark 2.1). In order to guarantee that, for each fixed t in $[T_1, T_2]$, the state $x(t, a)$, $a \in I$, is a member of C , the set of continuous functions over I , we assume that there is a finite constant d such that

$$|f(t, x^1, u) - f(t, x^2, u)| \leq d |x^1 - x^2|$$

for all $(t, x^1, u), (t, x^2, u)$ in M , that is, f is Lipschitzian in x uniformly in t and u over the set M .

We shall now consider the class Ω of all pairs $x(t, a), u(t), T_1 \leq t \leq T_2, T_1 \leq t_2(a) \leq T_2, a \in I$, of the family of vector functions $x(t, a)$ and the vector function $u(t)$, satisfying the following conditions: (a) $x(t, a)$ is absolutely continuous (AC) in $[T_1, T_2]$ for each fixed $a \in I$; (b) $u(t)$ is measurable in $[T_1, T_2]$ (a function of t alone); (c) $(t, x(t, a)) \in A(a), T_1 \leq t \leq T_2$, for each $a \in I$; (d) $u(t) \in U(t), T_1 \leq t \leq T_2$; (e) $x(T_1, a) = i(a)$; (f) $t_2(\cdot) \in \tau$; (g) $(t_2(a), x(t_2(a), a)) \in B(a)$ for each $a \in I$; and (h) the state equation

$$dx(t, a)/dt = f(t, x(t, a), u(t)) \tag{1}$$

is satisfied a.e. in $[T_1, T_2]$ for each $a \in I$ (see also Remark 2.2).

A pair $x(t, a), u(t), T_1 \leq t \leq T_2, T_1 \leq t_2(a) \leq T_2, a \in I$, satisfying conditions (a)–(h) is said to be an admissible pair, where $x(t, a)$ is called a trajectory and $u(t)$ is called a control.

Let $K[w_1(\cdot), w_2(\cdot)]$ be a continuous functional, in the uniform topology, defined on the set of functions $W = \{(w_1(a), w_2(a)), a \in I, w_i \in C, (t_2(a), w_2(a)) \in B(a) \text{ for each } a \in I, i = 1, 2\}$. Assume that K is bounded from below on $W' = \{(w_1(a), w_2(a)), w_i \in C, (t_2(a), w_i(a)) \in B(a) \cap A(a), i = 1, 2\} \subset W$. Then, the functional

$$\begin{aligned} I[x, u] &= K[\eta(x)(a)], & a \in I \\ &= K[t_2(a), x(t_2(a), a)], & a \in I \end{aligned}$$

is called the cost functional, where $\eta(x)(a) = (t_2(a), x(t_2(a), a)), a \in I$.

We seek the absolute minimum of $I[x(\cdot, \cdot), u(\cdot)]$ in the class Ω . If $(\tilde{x}(\cdot, \cdot), \tilde{u}(\cdot))$ has the property that $I[\tilde{x}(\cdot, \cdot), \tilde{u}(\cdot)] \leq I[x(\cdot, \cdot), u(\cdot)]$ for all $(x(\cdot, \cdot), u(\cdot)) \in \Omega$, then we say that $\tilde{x}(\cdot, \cdot), \tilde{u}(\cdot)$ is an optimal pair; and we may say that $\tilde{u}(\cdot)$ is an optimal control and $\tilde{x}(\cdot, \cdot)$ is an optimal trajectory. Though the optimal pair $\tilde{x}(\cdot, \cdot), \tilde{u}(\cdot)$ may not be unique in Ω , the value of the cost functional $I[\tilde{x}(\cdot, \cdot), \tilde{u}(\cdot)]$ is the same for all optimal pairs.

Remark 2.1. In the description of Ω , the class of admissible pairs, we require, for various technical reasons, that there is a finite time T_2 such that $T_1 \leq t_2(a) \leq T_2$ for all $a \in I$ and such that the solution to the state equation exists over all of $[T_1, T_2]$. Physically, we may interpret this as demanding that a *safety margin* exists, that is, we ask that, if we ignore our boundary conditions, we can extend our trajectories past the prescribed *stopping time* $t_2(a)$ to time T_2 . Moreover, in condition (f), for the purpose of obtaining a minimizing sequence, we also require that $t_2(a), a \in I$, belongs to an equicontinuous

family τ which is closed in the uniform topology. For example, we may require that $t_2(a)$ be constant for $a \in I$ or we may take τ to be the class of all Lipschitz functions over I with a fixed Lipschitz constant. We can think of this requirement as arising from the nature of the instruments governing the stopping time.

Remark 2.2. The control systems described in this section also include systems in which the state function f contains the parameter a , that is, where Eq. (1) is replaced by

$$dx(t, a)/dt = f(t, x(t, a), u(t), a)$$

and where f is a continuous vector function from $M \times I$ into E_n . Indeed, in this case, we have only to consider the additional variables $z = (x_{n+1}, \dots, x_{n+l})$ satisfying $dx_i/dt = 0$, $x_i(T_1, a) = a$, $i = n + 1, \dots, n + l$. Then, if $\tilde{x} = (x_1, \dots, x_n, x_{n+1}, \dots, x_{n+l}) = (x, z)$ and $f_i = 0$, $i = n + 1, \dots, n + l$, we have the system

$$d\tilde{x}(t, a)/dt = \tilde{f}(t, \tilde{x}(t, a, b), u(t))$$

with $\tilde{f}_i(t, \tilde{x}, u) = f_i(t, x, z, u)$, $i = 1, \dots, n + l$, which is of the form we first presented. More generally, the control systems described in this section also include systems in which the state function f contains a *perturbation* $\eta(t, b) \in E_s$, that is, whose state equation is given by

$$dx(t, a, b)/dt = f(t, x(t, a, b), u(t), \eta(t, b))$$

where $\eta(\cdot, \cdot)$ satisfies conditions similar to those of x . In particular, assume that (i) the parameter b varies over some compact set I_0 in Euclidean space E_p , (ii) for each fixed $b \in I_0$, $\eta'(t, b)$ exists a.e. in $[T_1, T_2]$ and satisfies $\eta'(t, b) = \gamma(t, \eta, b)$ with $(t, \eta) \in A_0 \subset E_{1+s}$ compact, (iii) $\eta(T_1, b) = j(b)$, j a continuous function on I_0 , and (iv) γ is Lipschitzian in η and b , uniformly in t , over the set $A_0 \times I_0$. Such systems can now be rewritten in the previous form by introducing the variable $z \in E_p$ and setting $\tilde{x} = (x, \eta, z)$, where $z(t, b) = b$. We then obtain the system

$$\begin{aligned} d\tilde{x}(t, a, b)/dt &= (f(t, x(t, a, b), \eta(t, b), u(t)), \gamma(t, \eta(t, b), z(t, b)), 0) \\ &= \tilde{f}(t, \tilde{x}(t, a, b), u(t)) \end{aligned}$$

with initial conditions $\tilde{x}(T_1, a, b) = (i(a), j(b), b)$ and, hence, we are back to our previous form.

3. Properties of Equicontinuity of the State Variable

The following properties of the systems just described will be used in the next section.

Property 3.1. Consider a control system as described in Section 2. Then, $x(t, a)$ is uniformly continuous for $(t, a) \in [T_1, T_2] \times I$. Moreover, given a sequence $[x_k(\cdot, \cdot), u_k(\cdot)]$ of admissible pairs, then $x_k(\cdot, \cdot), k = 1, 2, \dots$, forms an equicontinuous and equibounded family of functions in $[T_1, T_2] \times I$.

Proof. Let $x(\cdot, \cdot), u(\cdot)$ be an admissible pair. By the Lipschitz assumption on f , if $a_1, a_2 \in I$,

$$\begin{aligned} |x(t, a_1) - x(t, a_2)| &\leq |i(a_1) - i(a_2)| + \int_{T_1}^t |f(s, x(s, a_1), u(s)) - f(s, x(s, a_2), u(s))| ds \\ &\leq |i(a_1) - i(a_2)| + d \int_{T_1}^{T_2} |x(s, a_1) - x(s, a_2)| ds, \quad T_1 \leq t \leq T_2 \end{aligned}$$

Hence, by Gronwall's inequality,

$$|x(t, a_1) - x(t, a_2)| \leq C |i(a_1) - i(a_2)| \tag{2}$$

$C = \exp[d(T_2 - T_1)] < \infty, T_1 \leq t \leq T_2$. Note that C does not depend upon $u(\cdot)$. Since f is continuous over the compact set M , there is a finite K , not dependent upon $u(\cdot)$, such that $|f| < K$ on M . Hence, by (2), for $(t_1, a_1), (t_2, a_2)$ in $[T_1, T_2] \times I$,

$$\begin{aligned} |x(t_1, a_1) - x(t_2, a_2)| &\leq |x(t_1, a_1) - x(t_2, a_1)| + |x(t_2, a_1) - x(t_2, a_2)| \\ &\leq K |t_1 - t_2| + C |i(a_1) - i(a_2)| \end{aligned} \tag{3}$$

Since $i(a)$ is continuous for $a \in I, I$ compact, $i(a)$ is uniformly continuous over I . Thus, from (3), it follows that $x(t, a)$ is uniformly continuous over $[T_1, T_2] \times I$. Moreover, since K and C are independent of $u(\cdot)$, if $[x_k(\cdot, \cdot), u_k(\cdot)]$ is a sequence of admissible pairs, (3) holds with x replaced by x_k , that is,

$$|x_k(t_1, a_1) - x_k(t_2, a_2)| \leq K |t_1 - t_2| + C |i(a_1) - i(a_2)|, \quad k = 1, 2, \dots$$

Hence, $x_k(t, a), (t, a) \in [T_1, T_2] \times I, k = 1, 2, \dots$, forms an equicontinuous family of functions over $[T_1, T_2] \times I$. Since A is assumed compact, this family is also equibounded. Property 3.1 is thereby proved.

Property 3.2. Consider a sequence $[x_n(t, a), T_1 \leq t \leq T_2, a \in I]$ of equibounded and equicontinuous functions of (t, a) in $[T_1, T_2] \times I$. Moreover, suppose that, for each n , there is an associated function $t_{2n}(a), a \in I, T_1 \leq t_{2n}(a) \leq T_2$, with $t_{2n}(\cdot) \in \tau, n = 1, 2, \dots$. Then, there exists at least one subsequence $[x_{n_s}(\cdot, \cdot)]$ of $[x_n(\cdot, \cdot)]$ which converges uniformly in t and a to a continuous function $x(t, a), T_1 \leq t \leq T_2, a \in I$, and for which $t_{2n_s}(a)$ converges uniformly in a to a continuous function $t_2(a), a \in I$, with $t_2(\cdot) \in \tau$ (see Ref. 1, Chapter 7).

4. Closure and Existence Theorem

We now impose further requirements on the state equation and the control set $U(t)$ and prove a closure theorem and an existence theorem for the resulting systems. The central result of this section is the closure theorem. The main step in its proof is the construction of a new control system with the state variable in E_{2m} , this system being obtained from the general system by fixing two points a_1 and a_2 in I . We then use the results of Cesari (Ref. 2) and the form of the state function f given below in assumption (α) to complete the proof. In particular, let us assume that $(\alpha) f(t, x, u) = g(t, x) + k(t, u)$ for (t, x, u) in M . We shall also require, as usual, the following assumptions: $(\beta) U(t)$ is a compact subset of E_m for t in $[T_1, T_2]$; $(\gamma) U(t)$ is an upper semicontinuous function of t in $[T_1, T_2]$; and $(\delta) k(t, U(t)) = \{y \in E_n : y = k(t, u), u \in U(t)\}$ is a convex subset of E_n for each t in $[T_1, T_2]$; consequently, $Q(t, x) = f(t, x, U(t))$ is a convex subset of E_n for each $(t, x) \in A$.

Closure Theorem. Let $A(a), a \in I$, and A be compact subsets of E_{1+n} , let $U(t)$ be a compact subset of E_m for every t in $[T_1, T_2]$, let $U(t)$ be an upper semicontinuous function of t in $[T_1, T_2]$, let $f(t, x, u) = g(t, x) + k(t, u)$ be a continuous function on M into E_n , let $M(a), a \in I$, and M be compact subsets of E_{1+n+m} , and let $k(t, U(t))$ be a convex subset of E_m for every $(t, x) \in A$. Let $x_k(t, a), T_1 \leq t \leq T_2, a \in I$, with $(t_{2k}(a), a \in I) \in \tau$, be a sequence of trajectories satisfying assumptions (a)–(f) and (h) of Section 2 and converging uniformly toward a function $x_0(t, a), T_1 \leq t \leq T_2, a \in I$, which is AC in $[T_1, T_2]$ for each $a \in I$. Also, suppose that $t_{2k}(\cdot)$ converges uniformly to $t_2(\cdot)$. Then, there exists a measurable $u_0(\cdot), u_0(t) \in U(t), T_1 \leq t \leq T_2$, such that the pair $x_0(t, a), u_0(t), T_1 \leq t \leq T_2, a \in I$, with $t_2(a), a \in I$, satisfies conditions (a)–(f) and (h) of Section 2; that is, with the exception of condition (g), $x_0(\cdot, \cdot)$ is an admissible trajectory with stopping time $t_2(\cdot) \in \tau$, and $\eta[x_k(\cdot, \cdot)](a)$ converges uniformly to $\eta[x_0(\cdot, \cdot)](a), a \in I$.

Proof. For each $a \in I$, the vector functions $x_0'(t, a)$, $x_k'(t, a) = f(t, x_k(t, a), u_k(t, a))$, $T_1 \leq t \leq T_2$, $k = 1, 2, \dots$, are defined a.e. and are L -integrable. We have to prove that, for each $a \in I$, $(t, x_0(t, a)) \in A(a)$, $T_1 \leq t \leq T_2$, that $t_2(\cdot) \in \tau$, and that there is a measurable function $u_0(t)$, $T_1 \leq t \leq T_2$, such that, for each $a \in I$,

$$x_0'(t, a) = f(t, x_0(t, a), u_0(t)), u_0(t) \in U(t) \quad \text{a.e. in } [T_1, T_2] \quad (4)$$

First, from $x_k(\cdot, \cdot) \rightrightarrows x_0(\cdot, \cdot)$ (where \rightrightarrows denotes uniform convergence) and the assumption that $(t, x_k(t, a)) \in A(a)$, $(t, a) \in [T_1, T_2] \times I$, $A(a)$ closed, it follows that $(t, x_0(t, a)) \in A(a)$, $(t, a) \in [T_1, T_2] \times I$. It is immediate from condition (f) of Section 2 that $t_2(\cdot) \in \tau$. Since f is continuous on the compact set M , there exists a finite K such that $|f| \leq K$ over M . Hence, for each $a \in I$,

$$|x_k(t', a) - x_k(t'', a)| \leq K |t' - t''| \quad (5)$$

for any t', t'' in $[T_1, T_2]$. Thus, for $a \in I$,

$$\begin{aligned} |x_k(t_{2k}(a), a) - x_0(t_2(a), a)| &\leq |x_k(t_{2k}(a), a) - x_k(t_2(a), a)| + |x_k(t_2(a), a) - x_0(t_2(a), a)| \\ &\leq K |t_{2k}(a) - t_2(a)| + |x_k(t_2(a), a) - x_0(t_2(a), a)| \end{aligned}$$

from which it follows that $x_k(t_{2k}(a), a) \rightrightarrows x_0(t_2(a), a)$ and $\eta[x_k(\cdot, \cdot)](a) \rightrightarrows \eta[x_0(\cdot, \cdot)](a)$. Hence, it only remains to find a measurable $u_0(t)$ satisfying (4).

It is immediate that, for each $a \in I$, $x_k(\cdot, a) \rightarrow x_0(\cdot, a)$, $x_0(t, a)$ AC in $[T_1, T_2]$. Moreover, for each $a \in I$, if we consider the control system with constraint set $A(a)$, control set $U(t)$, and state equation $x' = f(t, x, u)$, with $T_1 \leq t \leq T_2$, then each of the pairs $x_k(\cdot, a)$, $u_k(\cdot)$, $k = 1, 2, \dots$, belongs to $\Omega(a)$, the class of admissible pairs for this system. Hence, we can conclude from Cesari's closure theorem (see Ref. 2) that, for each $a \in I$, there exists a measurable $u_0(\cdot, a)$, $u_0(t, a) \in U(t)$, $T_1 \leq t \leq T_2$, such that

$$x_0'(t, a) = f(t, x_0(t, a), u_0(t, a)) \quad (6)$$

a.e. in $[T_1, T_2]$ and, in particular, $(x_0(\cdot, a), u_0(\cdot, a)) \in \Omega(a)$. We will show that, for any fixed $a_0 \in I$, $u_0(\cdot, a_0)$ generates all the trajectories $x_0(\cdot, a)$, $a \in I$; that is, for any $a \in I$,

$$x_0'(t, a) = f(t, x_0(t, a), u_0(t, a_0)) \quad (7)$$

a.e. in $[T_1, T_2]$. If one sets $u_0(t) = u_0(t, a_0)$, this will then complete the proof.

We now show that, for any $a_1, a_2 \in I$, $u_0(\cdot, a_1)$ generates $x_0(\cdot, a_2)$. To this end, let us consider the control system consisting of the constraint set

$\tilde{A} = \{(t, X) : (t, X_1, \dots, X_n) \in A(a_1), (t, X_{n+1}, \dots, X_{2n}) \in A(a_2)\}$, a compact subset of the tX -space $E_1 \times E_{2n}$, with t in E_1 and $X = (X_1, \dots, X_n, \dots, X_{2n})$, the state variable, in E_{2n} . Let T_1, T_2 be given as before; and, for each t in $[T_1, T_2]$, let the control set $U(t)$ be the same as before, with $u = (u_1, \dots, u_m)$ the control variable. Hence, if $\tilde{M} = \{(t, X, u) : (t, X) \in \tilde{A}, u \in U(t)\}$, then $\tilde{M} = \{(t, X, u) : (t, X_1, \dots, X_n, u) \in M(a_1), (t, X_{n+1}, \dots, X_{2n}, u) \in M(a_2)\}$ is a compact subset of E_{1+2n+m} . Let $F = (F_1, \dots, F_{2n})$ be a continuous vector function from \tilde{M} into E_{2n} , $F(t, X, u)$ given by

$$F(t, X, u) = (f(t, X_1, \dots, X_n, u), f(t, X_{n+1}, \dots, X_{2n}, u))$$

The class \tilde{Q} of admissible pairs $X(t), u(t)$, $T_1 \leq t \leq T_2$, consists of those pairs satisfying the following conditions: (A) $X(t)$ is AC in $[T_1, T_2]$; (B) $u(t)$ is measurable in $[T_1, T_2]$; (C) $(t, X(t)) \in \tilde{A}$ for $T_1 \leq t \leq T_2$; (D) $u(t) \in U(t)$ for $T_1 \leq t \leq T_2$; (E) $X(t_1) = (i(a_1), i(a_2))$; and (F) the state equation

$$dX(t)/dt = F(t, X(t), u(t))$$

is satisfied a.e. in $[T_1, T_2]$.

By the construction of this control system, it is clear that $(X_k(\cdot), u_k(\cdot)) = ((x_k(\cdot, a_1), x_k(\cdot, a_2)), u_k(\cdot)) \in \tilde{Q}$, $k = 1, 2, \dots$. We wish to show that $X_0(t) = (x_0(t, a_1), x_0(t, a_2))$, $T_1 \leq t \leq T_2$, is an admissible trajectory for this system. Again, it is immediate that $X_k(t) \rightarrow X_0(t)$, $T_1 \leq t \leq T_2$, $X_0(t)$ AC in $[T_1, T_2]$. Moreover, since $A(a_i)$ and $M(a_i)$ were assumed compact, $i = 1, 2$, \tilde{A} and \tilde{M} are compact. In addition, $\tilde{Q}(t, X) = F(t, X, U(t))$ is convex by assumption (α) since, for fixed $(t, X) \in \tilde{A}$ and any u_1, u_2 in $U(t)$, we have, for $0 \leq \alpha \leq 1$,

$$\begin{aligned} & \alpha(f(t, X_1, \dots, X_n, u_1), f(t, X_{n+1}, \dots, X_{2n}, u_1)) \\ & \quad + (1 - \alpha)(f(t, X_1, \dots, X_n, u_2), f(t, X_{n+1}, \dots, X_{2n}, u_2)) \\ & = (g(t, X_1, \dots, X_n) + \alpha k(t, u_1) + (1 - \alpha) k(t, u_2), g(t, X_{n+1}, \dots, X_{2n}) \\ & \quad + \alpha k(t, u_1) + (1 - \alpha) k(t, u_2)) \\ & = (g(t, X_1, \dots, X_n) + k(t, u_\alpha), g(t, X_{n+1}, \dots, X_{2n}) + k(t, u_\alpha)) \\ & = (f(t, X_1, \dots, X_n, u_\alpha), f(t, X_{n+1}, \dots, X_{2n}, u_\alpha)) \end{aligned}$$

where, by assumption (δ), there exists a $u_\alpha \in U(t)$ such that

$$k(t, u_\alpha) = \alpha k(t, u_1) + (1 - \alpha) k(t, u_2)$$

Hence, we can apply Cesari's closure theorems (Ref. 2) to conclude that there exists a measurable control function $u_0(t)$, $T_1 \leq t \leq T_2$, such that

$$X_0'(t) = F(t, X_0(t), u_0(t)), \quad u_0(t) \in U(t)$$

a.e. in $[T_1, T_2]$. By the construction of $X_0(t), F(t, x, u)$, and (6), it thus follows that

$$\begin{aligned} f(t, x_0(t, a_1), u_0(t, a_1)) &= f(t, x_0(t, a_1), u_0(t)) \\ f(t, x_0(t, a_2), u_0(t, a_2)) &= f(t, x_0(t, a_2), u_0(t)) \end{aligned}$$

a.e. in $[T_1, T_2]$. Hence, $k(t, u_0(t, a_1)) = k(t, u_0(t, a_2))$ a.e. in $[T_1, T_2]$ and, therefore, $f(t, x_0(t, a_2), u_0(t, a_2)) = f(t, x_0(t, a_2), u_0(t, a_1))$ a.e. in $[T_1, T_2]$, that is, $u_0(t, a_1)$ generates $x_0(t, a_2)$, $T_1 \leq t \leq T_2$. Since a_1 and a_2 were chosen arbitrarily from I , we have thereby verified relation (7); and, by previous remarks, the closure theorem is thereby proved.

Existence Theorem. Consider a control system as described in Section 2 and satisfying conditions (α)–(δ). Assume that the class of admissible pairs Ω is nonempty. Then, there exists an admissible pair $x_0(\cdot, \cdot), u_0(\cdot)$, such that $I[x_0(\cdot, \cdot), u_0(\cdot)] \leq I[x(\cdot, \cdot), u(\cdot)]$ for all admissible pairs $x(\cdot, \cdot), u(\cdot)$.

Proof. By assumption, if $(x(\cdot, \cdot), u(\cdot)) \in \Omega$, then $\eta[x(\cdot, \cdot)](\cdot) \in W'$. Since $K[\eta[x(\cdot, \cdot)](a)], a \in I$, is assumed to be bounded from below on W' , $i = \inf_{\Omega} I[x(\cdot, \cdot), u(\cdot)] > -\infty$. Since there exists at least one admissible pair by assumption, i is therefore finite. Thus, there exists a minimizing sequence of admissible pairs $x_k(t, a), u_k(t), T_1 \leq t \leq T_2, a \in I$, with $t_{2k}(\cdot) \in \tau$, such that $I[x_k(\cdot, \cdot), u_k(\cdot)] \rightarrow i$ as $k \rightarrow \infty$.

By Property 3.1, $x_k(\cdot, \cdot)$ forms an equicontinuous and equibounded family of functions in $[T_1, T_2] \times I, k = 1, 2, \dots$. By Property 3.2, there exists a subsequence, which we still call $[x_k(\cdot, \cdot)]$, which converges uniformly to a continuous function $x_0(t, a), (t, a) \in [T_1, T_2] \times I$ and for which $t_{2k}(a)$ converges uniformly to a continuous function $t_2(a) \in \tau, a \in I$. Moreover, from (5), it follows that, for any $a \in I, |x_0(t', a) - x_0(t'', a)| \leq K$ for any t', t'' in $[T_1, T_2]$, so that $x_0(\cdot, a)$ is Lipschitzian, and hence AC, in $[T_1, T_2]$, for any $a \in I$. Consequently, by the closure theorem, there exists a measurable $u_0(t), u_0(t) \in U(t), T_1 \leq t \leq T_2$, such that the pair $x_0(\cdot, \cdot), u_0(\cdot)$ satisfies assumptions (a)–(f) and (h) of Section 2 and such that $\eta[x_k(\cdot, \cdot)](a)$ converges uniformly to $\eta[x_0(\cdot, \cdot)](a), a \in I$. Since $B(a)$ is assumed closed for each $a \in I$ by assumption, it follows that $\eta[x_0(\cdot, \cdot)](a) = (t_2(a), x_0(t_2(a), a)) \in B(a)$ for each $a \in I$; that is, assumption (g) of Section 2 is also satisfied. Thus, $(x_0(\cdot, \cdot), u_0(\cdot)) \in \Omega$. Finally, since K is assumed continuous in W ,

$$I[x_0(\cdot, \cdot), u_0(\cdot)] = \lim_{k \rightarrow \infty} I[x_k(\cdot, \cdot), u_k(\cdot)] = i.$$

The existence theorem is thereby proved.

5. Stochastic Systems

Let us now consider a class of control systems with stochastic boundary conditions. Specifically, let us particularize the control systems considered thus far by making $i(a)$, $a \in I$, a random variable, and by writing the cost functional as an expectation. In particular, let us assume that we are given a probability measure P over the Borel sets restricted to I . Again, we consider a control system as described in Section 2, with the cost functional now given as follows.

Assume that $h(t, y)$ is a real-valued, continuous function on the set $B \subset E_{1+n}$. Then, the cost functional is given by

$$I[x(\cdot, \cdot), u(\cdot)] = Eh(t_2(a), x(t_2(a), a)) = \int_I h(t_2(a), x(t_2(a), a)) dP$$

where the letter E denotes the expectation with respect to the probability measure P . We also assume that the set B is closed in E_{1+n} . Thus, $h(\cdot, \cdot)$ is continuous, and hence bounded, on the compact set $B \cap A$; and it is then immediate that this cost functional has all the required properties of the functional K . Therefore, these systems are subsumed by those described in Section 2. In other words, the state equation is a differential equation with stochastic initial conditions $i(a)$, with probability distribution P on the space I of the variable a . Hence, for a given control $u(\cdot)$, we obtain, for each t , a family $x(t, a)$ of vector-valued functions of a in I . Again, the probability measure P and the function h are also used to describe explicitly the cost functional K . Consequently, the results of Section 4 are applicable to these stochastic systems.

Such systems can arise, for example, if the interval $[T_1, T_2]$ is thought of as an *interval of ignorance* for a usual control system, during which the initial conditions and the state of the system are unknown. Alternately, such a system can arise if we are forced to choose a control $u(\cdot)$ before the initial conditions are known. For a further discussion of such systems, see also Ref. 3, Chapter 2.

Remark 5.1. As shown in Remark 2.2, if we consider control systems whose state equation is of the form $x' = g(t, x, \eta) + k(t, u)$, η a random variable or a stochastic process whose sample paths possess the same differentiability properties as $x(t, a)$, then clearly we can transform such systems into systems with stochastic initial conditions; hence, the results of Section 4 pertain to such systems as well (see Example 5.3; see also Ref. 3, Section 8.6).

Example 5.1. Let us consider the control system $\dot{x}_1 = x_2, \dot{x}_2 = u$, $x_1(0, a) = 0, x_2(0, a) = a$, where $u(t) \in U = [-1 \leq u \leq 1]$ and a is any real number between 0 and 1, with probability density one. We wish to find admissible pairs which minimize $I[x(\cdot, \cdot), u(\cdot)]$ within Ω , $I[x(\cdot, \cdot), u(\cdot)] = E[x_2(T, a)]^2 = \int_0^1 [x_2(T, a)]^2 da$, with $0 \leq t \leq T, T$ fixed. If we further require that $(t, x(t, a))$ lie in some compact set A , then clearly this control system satisfies all the relevant conditions; hence, by the results of Section 4, an optimal pair exists. Since $x_2(t, a) = a + \int_0^t u(s) ds$, it can be easily shown that, if A is large enough to contain the trajectories described below, the optimal pairs can be described as follows:

Case 1: $T < \frac{1}{2}$. Set $u(t) = -1$ on $[0, T]$. Hence, $x_2(T, a) = a - T$.

Case 2: $T \geq \frac{1}{2}$. Any admissible control such that $Ex(T, a) = 0$ and such that the space constraints are satisfied is optimal. For example, we may set $u(t) = -1$ on $[0, \frac{1}{2}]$ and $u(t) = 0$ on $(\frac{1}{2}, T]$.




Fig. 1

Example 5.2. Let us consider the control system of Example 5.1, with the cost functional now given by $Eg(x_2(T, a))$, where $g(x) = 2x, x \geq 0, g(x) = -4x, x < 0$. Again, if the constraint set is large enough, the results of Section 4 apply and the optimal pairs can be described as follows:

Case 1: $T < \frac{1}{3}$. Set $u(t) = -1$ on $[0, T]$.

Case 2: $T \geq \frac{1}{3}$. Any admissible control such that $Ex_2(T, a) = \frac{1}{6}$ is optimal.

Note that, unlike Example 5.1, the optimal solution cannot be obtained by replacing x with Ex throughout the system.

Example 5.3. Let us consider the control system of Example 5.1, with the state equation now given by $\dot{x}_1 = ax_2, \dot{x}_2 = u$, with fixed initial conditions $(\xi_1, \xi_2), 0 \leq t \leq T, T$ free, and where the cost functional is given by $E[(x_1(T, a))^2 + (x_2(T, a))^2]$. From the above remarks, if the constraint set is large enough, the results of Section 4 are applicable. Moreover, the optimal pairs can be shown to be those admissible pairs with $Eax_1(T, a) = 0$ and $Ex_2(T, a) = Ex_2(T) = 0$. If we set $y_1 = Eax_1(t, a), y_2 = Ex_2(t, a)$, then $\dot{y}_1 = \frac{1}{3}y_2, \dot{y}_2 = u$, with $y_1(0) = \frac{1}{2}\xi_1, y_2(0) = \xi_2$. In particular, one set of optimal trajectories is described in Fig. 1.

References

1. KELLEY, J. L., *General Topology*, D. Van Nostrand Company, Princeton, New Jersey, 1955.
2. CESARI, L., *Existence Theorems for Weak and Usual Optimal Solutions in Lagrange Problems with Unilateral Constraints, I*, Transactions of the American Mathematical Society, Vol. 124, No. 3, 1966.
3. BAUM, R., *Optimal Control Systems with Stochastic Boundary Conditions*, University of Michigan, Ph.D. Thesis, 1969.

Additional Bibliography

- FLEMING, W. H., *Optimal Continuous-Parameter Stochastic Control*, SIAM Review, Vol. 11, No. 4, 1969.
- KUSHNER, H. J., *On the Existence of Optimal Stochastic Controls*, SIAM Journal on Control, Vol. 3, No. 3, 1965.