

Closed and Open Conformal Field Theories and Their Anomalies*

Po Hu¹, Igor Kriz²

¹ Department of Mathematics, Wayne State University, Detroit, MI 48202, USA.
E-mail: po@math.wayne.edu

² Mathematics Department, University of Michigan, Ann Arbor, MI 48109-1109, USA.
E-mail: ikriz@umich.edu

Received: 26 January 2004 / Accepted: 7 April 2004
Published online: 14 October 2004 – © Springer-Verlag 2004

Abstract: We describe a formalism allowing a completely mathematical rigorous approach to closed and open conformal field theories with general anomaly. We also propose a way of formalizing modular functors with positive and negative parts, and outline some connections with other topics, in particular elliptic cohomology.

1. Introduction

The main purpose of this paper is to give rigorous mathematical foundations for investigating closed and closed/open conformal field theories (CFT's) and their anomalies. In physics, the topic of closed/open CFT has been extensively discussed in the literature (see e.g. [8, 22, 23, 7]). Our investigation was originally inspired by two sources: Edward Witten (cf. [41]) proposed a general program for using K -theory to classify stable D -branes in string theory. On the other hand, G. Moore and G. Segal [25] obtained a mathematically rigorous approach to classifying D -branes in the case of 2-dimensional topological quantum field theory (TQFT) (see also [26] for excellent detailed lectures on this topic).

We attempted to consider a case in between, namely D -branes in conformal field theory. We point out that G. Moore in [26] has also considered this direction, and Yi-Zhi Huang and L. Kong [18] in parallel with our investigation developed a vertex operator algebra approach to open CFT, so there is overlap with existing work. However, we will end up exploring the topic in a somewhat different light, as will become apparent below. In particular, our investigation will lead us to a new approach to anomaly in rational CFT (RCFT), and D -brane categories, using 2- and 3-vector spaces. This will also lead to constructions which relate to certain mathematical topics, such as foundation of elliptic cohomology. Even in this direction, however, substantial inroads have already been made in the literature, in particular [30, 12, 13]. The main contribution of the present paper therefore is that we set out to proceed with complete mathematical rigor.

* The authors were supported by the NSF.

There are many reasons for D -branes to be easier in CFT than in string theory. First of all, one does not have to insist on anomaly cancellation, and can investigate the anomaly instead. Here by anomaly we mean a certain indeterminacy (usually finite-dimensional), allowed in the correlation functions of a CFT (much more precise discussion will be given later: this is, in some sense, the main subject of the present paper). Another reason is that CFT makes good sense even without supersymmetry, while string theory does not. (In fact, in this paper, we restrict attention to non-supersymmetric CFT, although mostly just in the interest of simplicity.) Most importantly, however, any CFT approach to string theory amounts to looking at the complicated string moduli space only through the eyes of one tangent space, which is a substantial simplification. We will make some comments on this relationship between CFT and string theory in Sect. 3 below. We should point out that in this paper, we restrict attention to CFT's defined in *oriented* surfaces. This is not a restriction on our formalism, which works in the unoriented case as well, but it simplifies the discussion somewhat. However, interesting phenomena certainly arise when considering CFT's in the non-oriented worldsheet context, even for example in the case of the critical Ising model [32, 13].

On the other hand, CFT is incomparably more complicated than 2-dimensional TQFT. Because of this, in fact, a theorem classifying D -branes (as outlined in [25] for TQFT) seems, at the present time, out of reach for CFT. However, an exact mathematical *definition* of the entire structure of closed/open CFT is a reasonable goal which we do undertake here. To this end, we use our formalism of stacks of lax commutative monoids with cancellation (SLCMC's), developed in [16], and reviewed in the Appendix.

In Sect. 2 below, we shall also describe analogues of some of the concepts of [25] for CFT's, and observe some interesting new phenomena. For example, one may ask what is the correct generalization of the category of modules over the algebra corresponding to the closed sector of 2-dimensional TQFT in defining a D -brane category. We will see fairly quickly how going in this direction leads to 2-vector spaces in the case of CFT.

In Sect. 4, we give a basic example, the free bosonic CFT (=linear σ -model), and show how to obtain the D -brane modules corresponding to Von Neumann and Dirichlet boundary conditions for open strings. As we will see, however, even in this basic case, a substantial complication is giving a mathematically rigorous treatment of the convergence issues of the CFT.

Up to this point, we suppressed the discussion of anomaly, by assuming that anomaly is 1-dimensional. However, there is an obvious suggestion: a parallel between the set of D -branes of a closed/open CFT, and the set of labels of a modular functor of RCFT, see [27–29, 35]. It therefore seems we should look for axioms for the most general possible kind of anomaly for closed/open CFT, which would include sets of both D -branes and modular functor labels.

There are, however, further clues which suggest that the notion of “sets” in this context is too restrictive. Notably, the free \mathbb{C} -vector space $\mathbb{C}S$ on the set of labels S of a modular functor is the well known Verlinde algebra [40]. But the multiplication rule of the Verlinde algebra uses only *dimensions* of vector spaces involved in the modular functor, so it seems that if one wants to consider the spaces themselves, it is that one should consider, instead of $\mathbb{C}S$, the *free 2-vector space on S* . Is it possible to axiomatize modular functors for RCFT's in a way which uses 2-vector spaces in place of sets of labels?

In Sect. 5, we answer this last question in the affirmative. This is rather interesting, because it leads to other questions: the authors [16] previously proposed RCFT as a possible tool for geometrically modelling elliptic cohomology, while Baas-Dundas-Rognes

[2] obtained a version of elliptic cohomology based on 2-vector spaces. Is there a connection? Observations made in [2] show that the right environment of such discussion would be a suitable *group completion* of the symmetric bimonoidal category \mathbb{C}_2 of vector spaces, while simultaneously noting the necessary difficulty of any such construction. Nevertheless, we propose in Sect. 6, such group completion, despite major technical difficulties. Our construction involves super-vector spaces, and thus suggests further connections with P. Deligne’s observation [19] that the modulars functor of *bc*-systems must be considered as super-vector spaces, and with the work of Stolz-Teichner [38], who, in their approach to elliptic objects, also noticed the role of fermions and, in effect, what amounts to 1-dimensional super-modular functors. In *D*-brane theory, this same construction allows axiomatization of anti-*D*-branes with 1-dimensional anomaly.

However, let us return to *D*-branes. Is it possible to formulate axioms for general anomalies of closed/open CFT’s analogous to the 2-vector space approach to modular functors? We give, again, an affirmative answer, although another surprise awaits us here: while the “set of labels” of a modular functor was naturally a 2-vector space, the “set of *D*-branes” of a closed/open CFT must be a 3-vector space! We discuss this, and propose axioms for a general anomaly of closed/open CFT in Sect. 7. An intriguing problem is to extend the group completion approach of Sect. 6 to the case of general closed/open CFT anomaly, which would give an axiomatization of anti-*D*-branes in that context.

2. Closed/Open CFT’s with 1-Dimensional Anomaly, *D*-Brane Modules and *D*-Brane Cohomology

There is substantial physical literature on the subject of *D*-branes (see e.g. [8, 22, 23, 7]). In this paper, we shall discuss a mathematically rigorous approach to *D*-branes in non-supersymmetric CFT’s. Moreover, in Sects. 2-4, we shall restrict attention to 1-dimensional anomaly allowed both on the closed CFT and the *D*-brane. More advanced settings will be left to the later sections.

We begin by defining the stack of lax commutative monoids with cancellation (SLCMC) corresponding to oriented open/closed string (more precisely conformal field) theory. SLCMC’s were introduced in [16], but to make this paper self-contained, we review all the relevant definitions in the Appendix. We consider a set *L*. This is not our set of labels, it is the set of *D*-branes. Our set of labels consists of $K' = L \times L$ which we will call *open labels* and we will put $K = K' \amalg \{1\}$, where 1 is the *closed label*.

We will now define the SLCMC \mathcal{D} of *closed/open worldsheets*, which we will need for oriented closed/open CFT. We shall first define the LCMC of its sections over a point. As usual (see [16]), the underlying lax commutative monoid is the category of finite sets labelled by a certain set *K*, not necessarily finite. Before describing the exactly correct analytic and conformal structure, we first specify that these are compact oriented surfaces (2-manifolds) *X* together with homeomorphic embeddings $c_i : S^1 \rightarrow \partial X$, $d_j : I \rightarrow \partial X$ with disjoint images. Moreover, each c_i is labelled with 1, and each d_j is labelled by one of the open labels K' . Moreover, each connected component of

$$\partial X - \bigcup \text{Im}(c_i) - \bigcup \text{Im}(d_j)$$

(which we shall call *D*-brane components) is labelled with an element of *L*, and each d_j is labelled with the pair $(\ell_1, \ell_2) \in L \times L$ of *D*-branes which the beginning point and endpoint of d_j abut. The c_i ’s and d_j ’s are considered inbound or outbound depending on the usual comparison of their orientation with the orientation of *X*.

It is now time to describe the smoothness and conformal structure on X . To this end, we simply say that X is a smooth complex 1-manifold with analytic (real) boundary and corners; this means that the interior of X is a complex 1-manifold, and the neighbourhood of a boundary point x of X is modeled by a chart whose source is either an open subset of the halfplane $\mathbb{H} = \{z \in \mathbb{C} \mid \text{Im}(z) \geq 0\}$, where 0 maps to x or an open subset of the quadrant $\mathbb{K} = \{z \in \mathbb{C} \mid \text{Im}(z) \geq 0, \text{Re}(z) \geq 0\}$, where 0 maps to x ; the transition maps are (locally) holomorphic maps which extend biholomorphically to an open neighbourhood of 0 in \mathbb{C} . The points of the boundary whose neighbourhoods are modelled by open neighborhoods of 0 in \mathbb{K} are called *corners*. Further, we specify exactly which points are the corners of an open/closed string world sheet: we require that the corners be precisely the endpoints of the images of open string boundary parametrizations. We also require that open as well as closed string parametrizations be real-analytic diffeomorphisms onto their image; this completes the definition of the objects.

An isomorphism $X \rightarrow Y$ is a diffeomorphism which preserves complex structure, D -brane labels, is smooth on the interior, and commutes with the c_i, d_j (which we shall call parametrization components - note that the set of parametrization components is not ordered, so an automorphism may switch them).

Now to define the SLCMC \mathcal{D} , the main issue is fixing the Grothendieck topology. We use simply finite-dimensional smooth manifolds with open covers. As usual, the underlying stack of lax commutative monoids is the stack of covering spaces with locally constant K -labels (analogously to [16]). Sections of \mathcal{D} over M are smooth manifolds fibered over M , where the fibers are elements of \mathcal{D} , and the structure varies smoothly in the obvious sense. It is important, however, to note that it does not seem possible to define this stack over the Grothendieck topology of *complex* manifolds and open covers; in other words, it does not appear possible to discuss *chiral* CFT's with D -branes. To see this, we consider the following

Example. The moduli space of elliptic curves E with an unparametrized hole (i.e. one closed D -brane component with a given label). It is easily seen that the moduli space of such worldsheets is the ray $(0, \infty)$, i.e. not a complex manifold. To see this, the key point is to notice that the invariant $\text{Im}(\tau)$ of the elliptic curve F obtained by attaching a unit disk to E along the D -brane component does not depend on its parametrization. Intuitively, this seems plausible since $\text{Im}(\tau)$ is the “volume”. To rigorize the argument, we first recall that if one cuts the elliptic curve along a non-separating curve, then $\text{Im}(\tau)$ can be characterized as the “thickness” of the resulting annulus (every annulus is conformally equivalent to a unique annulus of the form $S^1 \times [0, r]$ for some boundary parametrization; r is the thickness). But now any reparametrization of the D -brane component c of E is a composition of reparametrizations which are identity outside of a certain small interval $J \subset c$. Thus, it suffices to show that the invariant $\text{Im}(\tau)$ does not change under such reparametrizations. However, we can find a smooth non-separating curve $d \supset J$ in F ; then cutting F along d , the so-called change of parametrization of c becomes simply a change of parametrization of one of the boundary components of F ; we already know that does not affect thickness.

By a K -labelled closed/open CFT with 1-dimensional anomaly $(H_i)_{i \in K}$ we shall mean a CFT with 1-dimensional modular functor on the SLCMC \mathcal{D} over the stack of lax monoids S_K , with target in the SLCMC \underline{H}_K . This means a lax morphism of SLCMC's

$$\tilde{\mathcal{D}} \rightarrow (H_i)_{i \in K}$$

where $\tilde{\mathcal{D}}$ is a \mathbb{C}^\times -central extension of \mathcal{D} . These concepts were defined in [16] (see the Appendix for a review). When K is not mentioned (i.e. we speak of just a closed/open

CFT), we shall assume that there is only one D -brane, i.e. $K = \{1, m\}$, where m is the open label.

Now, however, we would like to ask a more fundamental question: what are D -branes? The answer “elements of the set L associated with a K -labelled closed/open CFT” is clearly not a satisfactory one. For one thing, D -branes should form an additive category, not a set. Thus, let us look to Moore-Segal ([25], Sect. 3) for a guideline. There, in the case of a 2-dimensional topological quantum field theory (TQFT), a candidate for an additive category of D -branes is proposed (at least in the semisimple case): the category of modules over the closed sector algebra.

Let us review these results of [25] in more detail: a (2-dimensional closed TQFT) is a Poincaré algebra \mathcal{C} , which is the same thing as a commutative Frobenius algebra. The first theorem of [25], Sect. 3 is that a closed/open 2-dimensional TQFT is equivalent to the following additional set of data: A (not necessary commutative) Frobenius algebra \mathcal{O} and a map of algebras

$$\mathcal{C} \rightarrow \mathcal{O}$$

satisfying the Cardy condition, which asserts equality between the operations corresponding to cutting an annulus which has one parametrization component and one D -brane component on each boundary component in two ways: either by an open string (curve) connecting the two D -brane components, or by a closed string (curve) separating the two boundary components. The second theorem of [25], Sect. 3 states that when \mathcal{C} is semisimple, then

$$\mathcal{O} = \text{End}_{\mathcal{C}}(M)$$

for some \mathcal{C} -module M . Therefore, the category of \mathcal{C} -modules is the correct candidate for a category of D -branes.

Which of these concepts have we extended to CFT so far? The algebra \mathcal{O} is no problem: it corresponds simply to a closed/open CFT with one D -brane. However, what is the analogue of a \mathcal{C} -module M ? We shall explain why we think it may be too naive to simply search for some suitable concept of such a \mathcal{C} -module (e.g. some modification of VOA module) which would do the job. The key point is that even in the case of TQFT, \mathcal{C} -modules are only the right answer when \mathcal{C} is semisimple. When \mathcal{C} is not semisimple, it is not obvious how to make, for a general \mathcal{C} -module M , the algebra $\text{End}_{\mathcal{C}}(M)$ into an open sector of a closed/open TQFT. It is the Cardy condition which causes trouble. When \mathcal{C} is semisimple, the simple summands themselves are open sectors of closed/open TQFT's, and moreover, they can be summed, because semisimplicity makes 0 the only possible choice for mixed sectors.

Now for a (non-chiral) CFT, semisimplicity seems like a completely unnatural assumption. For example, for a 1-dimensional free bosonic CFT (see Sect. 4 below), we have different irreducible Dirichlet branes corresponding to points of spacetime, which certainly have non-trivial, and interesting, mixed sectors. In conclusion, therefore, for a closed CFT H , there does not seem to be a satisfactory notion of “ H -module” M which would always make $(H, M \otimes M^*)$ (or some similar construction) into a closed/open CFT: the reason is that it doesn't quite work even for TQFT's, in the general case.

On the other hand, there is something we can do. Suppose (H, R) is a closed/open CFT (R is the open sector) and V is a finite-dimensional vector space over \mathbb{C} . Then there is a canonical way of making $(H, V \otimes R \otimes V^*)$ into a closed/open CFT with one D -brane: in the open CFT operations, simply insert units and traces $\mathbb{C} \rightarrow V \otimes V^*$, $V \otimes V^* \rightarrow \mathbb{C}$ wherever suggested by the diagrams. This is like putting a $|V|$ -dimensional linear Chan-Paton charge on the ends of the open strings. Therefore, if we characterized a D -brane D by the closed/open CFT (H, R) , then $(H, V \otimes R \otimes V^*)$ should represent

$V \otimes D$. This gives as an important clue: the category of D -branes should be a *lax module* over the symmetric bimonoidal category \mathbb{C}_2 of finite-dimensional complex vector spaces (see the Appendix, and also, for more detail, [9]).

At the same time, however, this presents a puzzle, namely that there is no such obvious candidate for the sum: If the closed/open CFT (H, R) represents a D -brane D and (H, S) represents E , what closed/open CFT represents $D \oplus E$? The problem is the same as above: there doesn't seem to be enough information to recover the mixed open sector of strings stretched between the D -branes D and E .

It seems at this point that we have no choice but to axiomatize the whole structure we wish to have, including a category of D -branes which is a lax module over \mathbb{C}_2 . However, even writing the axioms correctly is a challenge. We will give the answer in full generality (not assuming that the anomaly is necessarily 1-dimensional) in Sect. 7 below. But since we already gave a substantial discussion of this topic in this section, let us work out the solution here in the case of a 1-dimensional anomaly. This case is of particular interest from the point of view of string quantization, and it is also a case in which there is a substantial simplification.

So we already know that we want a “ D -brane category” \mathcal{A} which is a lax module over \mathbb{C}_2 . For simplicity, let us further require that this be a finite-dimensional free module, i.e. equivalent to a sum of finitely many copies of \mathbb{C}_2 . Now the closed sector H is a Hilbert space and the open sector R is an object of the category

$$\mathcal{A} \otimes_{\mathbb{C}_2} \mathcal{A}^* \otimes_{\mathbb{C}_2} \mathbb{C}_2^{Hilb}.$$

Here \mathbb{C}_2^{Hilb} is the symmetric bimonoidal category of Hilbert spaces (see Sect. 5 below), and $\otimes_{\mathbb{C}_2}$ denotes lax extension of scalars in the obvious sense (see [9] for reference). From this data, we can construct an SLCMC, which we denote by $C(\mathcal{A}, H, R)$ and which we will now describe.

As usual, there is no difficulty with extending the construction to families, so we will limit ourselves to the LCMC, i.e. to sections over one point. Even before getting into that, however, there is another wrinkle which we must mention (see also Sect. 7 below). To keep track of which open parametrization components share a boundary component, and in what order and orientation, one must introduce a separate SLCMC Γ of all such configurations, which we will call *incidence graphs*. Therefore, more precisely, an incidence graph will encode (1) a set of closed parametrization components together with their orientations, (2) a set of (unlabelled) closed D -brane boundary components, (3) cyclically ordered sets of open parametrization components in the same boundary component, together with their orientations. Now our construction will come with a map of SLCMC's,

$$C(\mathcal{A}, H, R) \rightarrow \Gamma.$$

To use the same notation as in Sect. 7, we will decorate sets S of boundary components with two indices, the first of which will specify closed or open label (1 or m), and the second will specify inbound or outbound orientation (*in* or *out*). Then the set of sections of $C(\mathcal{A}, H, R)$ (over a point) over a given configuration of Γ is

$$tr_{cyclic} \left(\begin{array}{cc} \hat{\otimes} R^* \hat{\otimes} & \hat{\otimes} R \\ S_{m,in} & S_{m,out} \end{array} \right) \hat{\otimes} \begin{array}{cc} \hat{\otimes} H^* \hat{\otimes} & \hat{\otimes} H \\ S_{1,in} & S_{1,out} \end{array}$$

Here tr_{cyclic} denotes the tensor product over \mathbb{C}_2 of copies of the canonical functor

$$tr : \mathcal{A}^* \otimes_{\mathbb{C}_2} \mathcal{A} \rightarrow \mathbb{C}_2$$

over all pairs of ends of open strings which share a D -brane in the given configuration of Γ (note that the orientations are arranged in such a way that one copy of \mathcal{A}^* and one copy of \mathcal{A} always arise in such case).

In view of this discussion, we can define a *closed/open CFT with D -brane category \mathcal{A} with 1-dimensional anomaly* as a (lax) map of SLCMC's over Γ

$$\tilde{\mathcal{D}}_{\{1,m\}} \rightarrow C(\mathcal{A}, H, R),$$

where $\tilde{\mathcal{D}}_{\{1,m\}}$ is a \mathbb{C}^\times -central extension of the SLCMC $\mathcal{D}_{\{1,m\}}$. We will see that the case of general anomaly (where different D -branes can have different anomalies, possibly multidimensional) is still more complicated, by yet another level. However, we leave this to Sect. 7 below.

3. Conformal Field Theory and String Theory

We will now consider the relationship between conformal field theory and string theory, and the way it reflects on our investigation. As a standing reference on string concepts, we recommend one of the standard textbooks on the subject, e.g. [14] or [31]. One added feature of string theory is, of course, supersymmetry, but we shall soon see that this turns out not to be the only complication.

We will, therefore, begin our discussion with bosonic string theory (superstrings will enter later). The essential point of string quantization is that conformal field theory quantizes parametrized strings, while physical strings should be unparametrized. Now to pass from parametrized strings to unparametrized, one needs a way to quantize the complex structure. This problem is analogous to gauge fixing in gauge theory. In fact, this is more than just an analogy: from a strictly worldsheet point of view, conformal field theory is indeed a 2-dimensional quantum field theory satisfying Schwinger axioms, and can be viewed as a gauge theory in a certain sense; however, we do not need to pursue this here. The important point is that in string theory, complex structure gauge is needed to produce a consistent theory: conformal field theory is anomalous and, in Minkowski space, contains states of negative norm.

The modern approach to gauge fixing in gauge theory, and to string quantization, is through Fadeev-Popov ghosts and BRST cohomology. In the string theory case, we start with a CFT H_m , the (*matter CFT*). In this case, BRST cohomology is essentially a semi-infinite version of Lie algebra cohomology of the complexified Witt algebra (viewed as a ‘‘Lie algebra of the semigroup of annuli’’) with coefficients in H_m . To be precise about this, we must describe the semi-infinite analogue of the complex $\Lambda(g)$ for a Lie algebra g where g is the Witt algebra. As it turns out, this semi-infinite Lie complex is also a CFT which is denoted as H_{gh} and called the Fadeev-Popov ghost CFT. A mathematical description is outlined in [35], and given in more detail in [19]. In the chiral CFT case, H_{gh} is a Hilbert completion (with a chosen Hilbert structure) of

$$\Lambda(b_n | n < 0) \otimes \Lambda(c_n, n \leq 0). \tag{1}$$

In the physical case, both chiralities are present, and H_{gh} is a Hilbert completion of the tensor product of (1) with its complex conjugate. To understand why this is a semi-infinite Lie complex of the Witt algebra, we write the generators of (1) as

$$\begin{aligned} b_n &= z^{-n+1} \frac{d}{dz}, \\ c_n &= z^{-n-2} (dz)^2. \end{aligned} \tag{2}$$

So, the b_n 's are vector fields on S^1 (elements of the Witt algebra) and the c_n 's are dual to the b_{-n} 's. An ordinary Lie complex would be the exterior algebra on the duals c_n , $n \in \mathbb{Z}$. However, a special feature of CFT is a choice of vacuum which allows us, even before gauge fixing, to define correlation functions which are finite, albeit anomalous: this is the mathematical structure known as Segal-type CFT, which we have axiomatized in [16] and here. However, this choice of vacuum of H_m is what prompts the ‘‘semi-infinite approach’’, where the exterior generators of H_{gh} are not all of the c_n 's, but half of the c_n 's and half of the b_n 's, as in (1). Now it turns out that for our further discussion, it will be important to know explicitly one part of the Virasoro action on H_{gh} , namely the conformal weights, or eigenvectors of L_0 . One may guess that b_n, c_n should be eigenvectors of conformal weight $-n$, but it turns out that one must decrease the conformal weights of the entire complex (1) by 1, so the correct conformal weight of a monomial in the b_n 's and c_n 's is $-k - 1$, where k is the sum of the subscripts of the b_n 's and c_n 's in the monomial. In fact, it turns out that the vacuum of the ghost theory, i.e. the element of H_{gh} assigned to the unit disk, is

$$b_{-1}c_0. \tag{3}$$

Now the ghost CFT has an anomaly which is described by a 1-dimensional modular functor L which has central charge -26 in the chiral case (see [35, 19]) and $(-26, -26)$ in the physical case. (In the chiral case, there is an additional complication that L must be considered a super-modular functor, see [19] and Sect. 5 below.) A CFT H_m is called *critical* if it has anomaly described by the 1-dimensional modular functor $L^{\otimes -1}$. The 26th power of the 1-dimensional free bosonic CFT described (briefly) in the next section is critical in the physical sense (with both chiralities).

Now for a critical CFT H_m , there is a certain differential Q (called the BRST differential) on the (non-anomalous) CFT

$$\mathbb{H} = H_m \hat{\otimes} H_{gh}. \tag{4}$$

In the chiral case, one has explicitly

$$Q = \sum_{r \in \mathbb{Z}} L_r^{H_m} c_{-r} - \frac{1}{2} \sum_{r,s \in \mathbb{Z}} (r - s) : c_{-r} c_{-s} b_{r+s} : - c_0 \tag{5}$$

(see [5], formula (4.59)). Here L_r^m are the Virasoro generators acting on H_m , and c_n, b_n , $n \in \mathbb{Z}$ are now understood as operators on H_{gh} in the standard way (see [5]). In the non-chiral case, one must add to (4) its complex conjugate. Q is a differential, which means that

$$QQ = 0. \tag{6}$$

The cohomological dimension is called the ghost number. The c_n 's have ghost number 1, the b_n have ghost number -1 , so the ghost number degree of Q is $+1$. We shall fix the ghost number as an algebra grading, so 1 has ghost number 0, but other conventions also exist.

What is even more interesting than (6), however, is that Q turns \mathbb{H} into a ‘‘differential graded CFT’’. If we use the usual notation where we write for a CFT, as a lax morphism of SLCMC's,

$$X \mapsto U_X,$$

then we may define a differential graded CFT by the relation

$$\sum (1 \otimes \dots \otimes Q \otimes \dots 1) U_X = 0, \tag{7}$$

with the correct sign convention. For simplicity, we assumed in (7) that all boundary components of X are outbound, the adjoint operator Q^* is used on inbound boundary components. Physically, (7) is due to the fact that Q is a conserved charge corresponding to the Noether current of a supersymmetry (called BRST symmetry) of the Lagrangian of \mathbb{H} (see [5]).

In any case, we now see that the BRST cohomology

$$H = H^*(\mathbb{H}, Q) \tag{8}$$

is a non-anomalous CFT. Infinitesimally, this implies that

$$[Q, L_n] = 0,$$

where L_n are the standard Virasoro algebra (in our case in fact Witt algebra) generators. However, more is true. In fact, one has

$$L_n = [Q, b_n],$$

so

$$Qx = 0 \Rightarrow L_n x \in \text{Im}(Q). \tag{9}$$

Because of (9), L_n actually act trivially on H , so H is in fact a TQFT (which means that U_X only depends on the topological type of X). Therefore, our machinery would certainly seem to apply to H , in fact so would that of Moore-Segal [25]. There are, however, two difficulties.

First of all, H_m may not actually be a CFT as we defined it because of convergence problems. For example, when H_m is the free bosonic CFT on the (25, 1)-dimensional Minkowski space, the inner product on the space H_m is indefinite, so this space cannot be Hilbert-completed with respect to its inner product. This is more than a technical difficulty: in physical language, this is the cause of the 1-loop divergence of bosonic string theory. In our language, this means that the state space of our would-be TQFT is infinite-dimensional, so U_E for an elliptic curve E is infinity, or more precisely undefined. So there isn't, in fact, any variant of the (25, 1)-dimensional free bosonic CFT for which the machinery outlined above would work mathematically and produce a true TQFT.

In physics, this is an argument why the free bosonic string theory is not physical, and one must consider superstring theory. Our definition of CFT works on the SLCMC of superconformal surfaces, but the convergence problems persist, i.e. again, for the free (9, 1)-dimensional super-CFT, the BRST cohomology would be TQFT is infinite-dimensional. Physicists argue that the (infinite) even and odd parts of the TQFT are “of equal dimension”, and thus the 1-loop amplitude vanishes (a part of the “non-renormalization theorem”). However, we do not know how to make this precise mathematically.

There is another, more interesting caveat, namely that H is actually not exactly the object one wants to consider as the physical spectrum of string theory. Working, for simplicity, in the bosonic case, one usually restricts to states of ghost number 0, which, at least in the free case, is isomorphic to the quotient H_0 of the submodule $Z_0 \subset H_m$ of states $x \in H_m$ satisfying $L_n x = 0$ for $n > 0$ and $L_0 x = x$, by the submodule B_0

of states of null norm (the Goddard-Thorn no ghost theorem). In bosonic string theory, the vacuum of H_m is in H_0 , but this is a tachyon, which is not the vacuum of H : the vacuum in H is b_{-1} , as remarked above, and has ghost number -1 . In superstring theory, the tachyon is factored out by the so-called GSO projection, while the vacuum of course persists, and has also ghost number -1 , but a different name, due to the different structure of the theory, which we have no time to discuss here.

One may ask what mathematical structure there is on H_0 itself. Here the answer depends strongly on whether we work chirally or not. In the chiral case, Borchers [4] noticed that H_0 is a Lie algebra. The Lie algebra structure comes from $[u, v] = u_0 v$ where u_0 is the residue of the vertex operator $Y(u, z)$. The Jacobi identity follows immediately from the vertex operator algebra Jacobi identity. From CFT point of view, this operation is analogous to the Lie bracket in Batalin-Vilkovisky algebras.

However, when both chiralities are present (which is the case we are interested in), the rabbit hole goes deeper than that. First note that the CFT vertex operator is not holomorphic, and curve integrals do not seem to be the right operations to consider. Instead, elements of Z_0 are operator-valued $(1, 1)$ -forms, and therefore can be naturally integrated over worldsheets. Indeed, one can see that integration of an element of Z_0 over worldsheets produces an *infinitesimal deformation* of CFT. Elements of B_0 also deform the CFT, but only by a gauge transformation, so elements of H_0 give rise to infinitesimal deformations of string theory. We may therefore (despite potentially serious convergence problems) wish to consider a *moduli space* M of string theories, to which H_0 is a tangent space at one point. In fact, points of the curved space M should be the true states of string theory, while the points of the tangent space H_0 are only an approximation. In the physical theory, one conjectures that the space M contains all of the 5 original superstring theories as states, and a continuum of states in between. As seen even by studying the basic example of toroidal spacetime, some states in M differ only by “boundary conditions on open strings”, and such conditions are called D -branes. When there is a well defined spacetime manifold X , D -branes as a rule are associated with submanifolds of X with some additional structure. These, however, are classical and not quantum objects (cf. Polchinski [31]), so that approach also has its drawbacks. While rigorous mathematical attempts to define and investigate D -branes from the manifold point of view have (with some success) also been made in the literature exist, the “tangent” CFT approximation which we consider here is, in some sense, complementary. Finding a mathematical theory which would unify both points of view is an even much more complex task, which we do not undertake here.

4. An Example: The 1-Dimensional Free Scalar CFT

We shall now give the standard examples of D -branes in the free bosonic CFT in dimension 1 (which is the CFT description of the linear σ -model). Unfortunately, even for this most basic CFT, a mathematically rigorous description of its convergence issues is nowhere to be found in the literature. The best outline we know of is given in [35].

A good first guess for the free (bosonic) field theory state space is, analogously with the lattice theories (see [16])

$$H = L^2(\mathbb{R}, \mathbb{C}) \hat{\otimes} \hat{S}ym < z^n, \bar{z}^n | n > 0 > . \quad (10)$$

Here $L^2(\mathbb{R}, \mathbb{C})$ denotes L^2 -functions with respect to the Gaussian measure. The quantum number associated with this space is the momentum.

To be more precise, (10) should be a Heisenberg representation of a certain infinite-dimensional Heisenberg group. To describe it, we start with the topological vector space V of all harmonic functions on S^1 or, more precisely, harmonic functions on small open sets in \mathbb{C} which contain S^1 and, say, the topology of uniform convergence in an open set containing S^1 . Thus, V is topologically generated by the functions

$$z^n, \bar{z}^n, n \in \mathbb{Z}$$

and

$$\ln ||z||.$$

To define the Heisenberg group, we would like to find a \mathbb{C}^\times -valued cocycle on V which would be invariant under the action of $Diff^+(S^1)$. However, similarly as in the case of lattice theories, we do not know any such cocycle. Instead, one considers the space U of harmonic \mathbb{C} -valued functions on (an open domain containing) the unit interval I . The point is that the harmonic functions on I break up into holomorphic and antiholomorphic parts; a topological basis of the holomorphic part is given by the elements

$$z^n, n \in \mathbb{Z}, \ln(z),$$

and a topological basis of the antiholomorphic parts is given by their complex conjugates. Therefore, the holomorphic and antiholomorphic parts U_+ and U_- of U have well defined winding numbers which can be added to a total winding number; let $V' \subset U$ be the set of functions of total winding number 0. Then the map $exp(?) = e^{2\pi i?}$ gives a projection

$$V' \rightarrow V$$

whose kernel consists of the constant functions. Now to get the free field theory, one proceeds analogously to lattice theories (see [16]), specifying a cocycle on U . We shall specify separately cocycles on both U_+ and U_- . However, because the integrality condition is replaced by equality of winding numbers on U_+ and U_- , we have more freedom in choosing the cocycle. For example, we can put, on both U_+ and U_- ,

$$c(f, g) = exp \frac{1}{2} \left(\int_{S^1} f dg - \Delta_f g(0) + \frac{1}{2} \Delta_f \Delta_g \right), \tag{11}$$

(where Δ_f is the winding number). The effect of this is that if we apply the cocycle to lifts of two harmonic functions f, g on a worldsheet to its universal cover, whose restriction to boundary components are f_i, g_i (as is done in [16] for the lattice theories), the Greene formula implies that

$$\begin{aligned} c(f, g) &= exp \frac{1}{4} \left(\sum_{i < j} (\Delta_{f_i} \Delta_{g_j} + \Delta_{g_i} \Delta_{f_j}) + \sum_{i=1}^n \Delta_{f_i} \Delta_{g_i} \right) \\ &= exp \frac{1}{4} \left(\sum_{i=1}^n \Delta_{f_i} \sum_{i=1}^n \Delta_{g_i} \right) = 1. \end{aligned} \tag{12}$$

Thus, the situation is simpler than in the case of lattice theory. Now similarly as in the case of lattice theory, the cocycle c we have constructed, when restricted to V' , is trivial on the constant functions, so we get a canonical map

$$\mathbb{C} \rightarrow \tilde{V}' \tag{13}$$

(where $\tilde{?}$ denotes the Heisenberg group with respect to a given cocycle). Similarly as in the case of lattice theories, in fact, $c(f, g) = 0$ for $f, g \in V'$, f constant, so the subgroup (13) is normal, so the desired Heisenberg group can be defined by

$$\tilde{V} = \tilde{V}'/\mathbb{C}. \tag{14}$$

Then H should be the Heisenberg representation of the right *real form* of (14). Note, however, that the above construction comes with no obvious natural choice of real form. Let us postpone the discussion of this issue, as we shall see it is related to the convergence issues of the CFT. Now, conformal field theory structure is specified as usual: looking at the Heisenberg representation $H_{\partial X}$ of the central extension $\tilde{V}_{\partial X}$ of the space $V_{\partial X}$ of harmonic functions on the boundary of a worldsheet X , we have already constructed a canonical splitting of the pullback of the central extension to the subspace V_X of harmonic functions on X ; we would like to define the field theory operator associated with X as the vector space of invariants of $H_{\partial X}$ with respect to V_X .

A usual “density argument” (cf. [35, 16, 33]) shows that the invariant vector space H_X is always at most 1-dimensional. In more detail, if we denote by $Harm(X)$ the space of harmonic functions on X and by $Harm(\partial X)$ the space of harmonic functions on a small neighborhood of ∂X , and also by D the unit disk, then, by restriction, we may form the double coset space

$$Harm(X) \backslash Harm(\partial X) / \prod_{\partial} Harm(D) \tag{15}$$

(the product is over boundary components of X). Then (15) is isomorphic to $H^1(\bar{X}, Harm) \cong \mathbb{C}$, where $Harm$ is the sheaf of harmonic functions and \bar{X} is the worldsheet obtained from X by gluing unit disks on the boundary components. Since H can be interpreted as a (completed) space of functions on

$$Harm(\partial X) / \prod_{\partial} Harm(D), \tag{16}$$

the identification of (15) shows that only functions on the orbits \mathbb{C} have a chance to be $Harm(X)$ -fixed points. However, studying further the constant functions in $Harm(X)$, we see that only functions supported on $\{0\} \subset \mathbb{C}$ have a chance of being fixed points.

These observations also point to a difficulty with a Hilbert space model for H . What kind of reasonable Hilbert space functions on \mathbb{R} contain distributions supported on a single point? Now recall the reason why the Hilbert space is not yet fixed: we haven’t fixed the real structure on V . One clue for such real structure is that, from the desired interpretation of H as functions on (16),

$$A = Harm(D) \subset V$$

should be our “Lagrangian subspace” so that $H = \widehat{Sym}(A)$, (cf. [33], Section 9.5). Thus, we can define the real structure on V by specifying the inner product on A . The choice enjoying the desired invariances is

$$\langle f, g \rangle = \int \bar{f} dg. \tag{17}$$

But then this is only a semidefinite Hermitian product on A , where constants have norm 0! The above discussion shows that this is more than a technical difficulty. To get the field operator U_X to converge, we must take the inner product (17), which leads to

$$H = \prod_{k \in \mathbb{R}} \text{Sym} \langle z^n, \bar{z}^n | n > 0 \rangle$$

(\prod is the Cartesian product). Physically, the quantum number k is the momentum. Then the notion of ‘‘Hilbert product’’ of copies of H and ‘‘trace’’ must be adjusted. For more details, see the Appendix. With these choices, convergence of U_X can be proven similarly as in [16] for lattice theories (e.g. using boson-fermion correspondence at 0 momentum), so the free bosonic CFT is rigorous. This convergence problem does not arise if we consider the σ -model on a compact torus instead of flat Euclidean space.

For completeness, we note that we haven’t discussed inbound boundary components, and closed worldsheets. The former topic offers no new phenomena and can be treated simply by reversing the sign of the cocycle. Discussing closed worldsheets amounts really to discussing in detail the anomaly, which is $H^1(X, \underline{Harm})$, analogously to the fermionic case treated in [19].

Now we want to give examples of simple elementary D -branes in the free CFT. The above discussion shows that we would have to work in compact spacetime (a torus) to make the examples fit the scheme proposed in Sect. 2 literally. However, we elect instead to stick to the flat spacetime \mathbb{R} , where the situation seems more fundamental. It must be then understood, however, that the notion of closed/open CFT in this situation must also be generalized in a way analogous to closed CFT (as discussed in the Appendix), to solve the convergence issue.

Consider the $1/2$ -disk B consisting of elements of D with non-negative imaginary part. We consider B an open string worldsheet where the real boundary elements are the D -brane component, and the open string component is parametrized by the map $e^{\pi i t}$. Then we can consider the space of all harmonic functions on the boundary of B which obey a suitable boundary condition on the D -brane component. The boundary conditions allowable first of all must be conformally invariant. The most obvious such condition is that the derivative of the function in question be 0 in the direction of a certain vector $u \in S^1$, $Im(u) > 0$ or $u = 1$. A priori, all of those conditions are allowable. However, if we want to follow the methods we used above to describe closed free CFT, additional conditions are needed. Namely, we need the vector space of functions satisfying the condition to have a central extension which is a Heisenberg group. Moreover, to get consistency of open and closed CFT, we need the Heisenberg group to be obtained by restriction of the cocycle (11), and the real structure must also be induced from the closed CFT real structure.

We will see however that the real structure is incompatible with the closed CFT real structure unless $u = 1$ or $u = i$. In effect, let f be a holomorphic function which sends \mathbb{R} to \mathbb{R} (e.g. a polynomial with real coefficients). Then $uf - \bar{u}\bar{f}$ has zero derivative in the direction u on \mathbb{R} (as $z \mapsto uf(z/u) - \bar{u}\bar{f}(z/u)$ has zero derivative in the direction of u on the line $z \in u\mathbb{R}$). Here by $\bar{\cdot}$ we mean the usual complex conjugation. But now recall that in the real structure involved in defining closed CFT, the complex conjugate of z^n is z^{-n} , so the complex conjugate of $uz^n - \bar{u}\bar{z}^n$ is $\bar{u}z^{-n} - u\bar{z}^{-n}$, which is not of the form $uf - \bar{u}\bar{f}$ for a holomorphic function f sending \mathbb{R} to \mathbb{R} , unless $u = (\pm)1$ or $u = (\pm)i$.

Now for $u = i$, the functions on B which we get are

$$z^n + \bar{z}^n, \quad n \in \mathbb{Z},$$

$$\ln ||z||. \tag{18}$$

(Note the singularities on the D -brane components of ∂B - those needn't bother us.) The cocycle is defined by the same formula as (11), although one integrates over the parametrization component only (the resulting factor of $1/2$ is related to the squaring relation between the open and closed string couplings). The corresponding Heisenberg representation (with the same discussion of convergence at momentum quantum numbers as in the closed case) is the von Neumann open string sector, or the 1-dimensional “ $D0$ -brane” of the 1-dimensional free bosonic CFT. Together with the closed sector, this defines (up to the convergence-related modifications) a closed/open CFT with 1-dimensional anomaly in the sense of Sect. 2 (the vacua are obtained completely analogously as in the closed case).

When $u = 1$, the functions satisfying the boundary condition (which is the Dirichlet condition) are

$$z^n - \bar{z}^n, \quad n \in \mathbb{Z} - \{0\},$$

$$1,$$

$$arg(z) = \frac{1}{2} \ln \frac{z}{\bar{z}}.$$

Although at first everything looks analogous as in the von Neumann case, there is an important difference: if we define again the cocycle by (11) integrated over the parametrization component of B (which we must to get consistency), this time the cocycle is degenerate with kernel

$$\langle 1, arg(z) \rangle.$$

This is a case of spontaneous symmetry breaking: we get a 2-dimensional continuum of irreducible representations, one for each weight of 1 and $arg(z)$ (which are independent real numbers).

To name these sectors, we must figure out what these quantum numbers mean. To this end, in turn, we must review our recipe for defining a closed/open CFT from these Heisenberg representations. To get consistency, the recipe must be the same as in the closed case: for a closed/open worldsheet X , take the tensor product $H_{\partial X}$ of state spaces corresponding to the parametrization components of X (suitably completed, as above). By the representations we constructed, the group $Harm(X)$ of harmonic functions on X acts on $H_{\partial X}$ (the central extension given by our cocycle splits on $Harm(X)$ canonically, as above). The scalar multiples of the vacuum in $H_{\partial X}$ now form the space of invariants $H_{\partial X}^{Harm(X)}$. The discussion is analogous to the closed case, and we omit the details.

However, the sector numbers we are interested in correspond to weights, on each open parametrization component c of X , of functions which have given values α, β on the endpoints of c (this is linear in α, β). Note that we can get any such pair of numbers α, β by taking a linear combination of 1 and $arg(z)$. Now consider, on X , harmonic functions which are constant on the D -brane components of X : such functions must act by identity (have weight 0). We see from this that the fixed point space will be 0 unless weights corresponding to the two endpoints of each open D -brane component coincide. Therefore, these weights can be interpreted as position coordinates of the two endpoints of the open string, in other words position coordinates of the (instanton) D -branes. The

2-dimensional continuum of state spaces we got describes all the possible mixed sectors H_{ab} of open strings ending on any two of these D -branes with positions $a, b \in \mathbb{R}$. Convergence issues work out after a discussion analogous to the closed case.

To build a closed/open CFT with D -brane category in the sense of Sect. 2 from the 1-dimensional free boson, note that we have restricted attention to D -brane categories which are finite-dimensional free lax \mathbb{C}_2 -modules. Therefore, we must select *finitely many* positions of instanton D -branes a_1, \dots, a_N (and the $D0$ -brane, if we wish). The D -branes can then be summed using the mixed sectors in the obvious way. The mixed sectors between the $D0$ and $D(-1)$ -branes are 0. As in the closed case, the Hilbert tensor product and trace must be modified to get proper convergence behavior. We have not axiomatized anti- D -branes here, but that can be done using the formalism of Sect. 6, at the cost of increasing technical difficulty substantially.

5. CFT Anomaly via 2-Vector Spaces and Elliptic Cohomology

In this section, we give a new definition of modular functor which generalizes the definition given in [16]. Consider a free finitely generated lax module \mathcal{M} over the lax semiring \mathbb{C}_2 (the category of finite-dimensional \mathbb{C} -vector spaces; it is convenient to let the morphisms of \mathbb{C}_2 be all linear maps; thereby, \mathbb{C}_2 is not a groupoid, and we have to use the version of lax algebra theory which works over categories - see the appendix; of course, it is possible to consider the subcategory \mathbb{C}_2^\times of \mathbb{C}_2 whose morphisms are isomorphisms). Consider further \mathbb{C}_2^{Hilb} , the \mathbb{C}_2 -algebra of Hilbert spaces with the Hilbert tensor product. We put $\mathcal{M}^{Hilb} = \mathcal{M} \otimes_{\mathbb{C}_2} \mathbb{C}_2^{Hilb}$. Now consider $H \in_1 \mathcal{M}^{Hilb}$ (the symbol \in_1 means a map of lax \mathbb{C}_2 -modules $\mathbb{C}_2 \rightarrow ?$; in our case, this is the same thing as $H \in Obj(\mathcal{M}^{Hilb})$). We shall define two LCMC's $C(\mathcal{M})$ and $C(\mathcal{M}, H)$ (underlying LCM of sets) which are, in standard ways, extended into SLCMC's over the Grothendieck category of finite-dimensional smooth complex manifolds.

The LCMC's are constructed as follows: the objects of $C(\mathcal{M})$ over the pair of finite sets (S, T) are 1-elements of $\mathcal{M}^{\otimes S} \otimes \mathcal{M}^{*\otimes T}$; the morphisms are 2-isomorphisms. Here \mathcal{M}^* is the dual lax \mathbb{C}_2 -module of \mathcal{M} , whose objects are (lax) morphisms of lax \mathbb{C}_2 -modules $\mathcal{M} \rightarrow \mathbb{C}_2$ and morphisms are natural isomorphisms compatible with the operations. The gluing maps are given by trace over \mathbb{C}_2 , i.e. the evaluation morphism $\mathcal{M} \otimes \mathcal{M}^* \rightarrow \mathbb{C}_2$. An object of $C(\mathcal{M}, H)$ over (S, T) consists of an object M of $C(\mathcal{M})$, and 2-morphism $u : M \rightarrow_2 H^{\hat{\otimes} S} \hat{\otimes} H^{*\hat{\otimes} T}$ whose image consists of trace class elements: (Choosing a basis of \mathcal{M} , a 1-element of \mathcal{M}^{Hilb} becomes a collection of Hilbert spaces, so $\hat{\otimes}$ -powers of H are collections of $\hat{\otimes}$ -powers; an element is trace class if each of its components is trace class. For generalizations beyond the trace class context, see Remarks in the Appendix.) Here $H^* \in_1 \mathcal{M}^{* Hilb}$ is defined by putting, for $V \in_1 \mathcal{M}$, $H^*(V) = Hom_{\mathcal{M}^{Hilb}}(H, V)$. Morphisms are commutative diagrams of the obvious sort. To define gluing operations, note that

$$u : M \rightarrow_2 H^{\otimes(S+U)} \hat{\otimes} H^{*\hat{\otimes}(T+U)}$$

induces a 2-morphism $tr(u) : (1 \otimes tr_1)M \rightarrow_2 H^{\hat{\otimes} S} \hat{\otimes} H^{*\hat{\otimes} T}$, where $tr_1 : \mathcal{M}^{\otimes U} \otimes \mathcal{M}^{*\otimes U} \rightarrow \mathbb{C}_2$ is the evaluation morphism; it is defined by using the canonical morphism $tr_2 : tr_1(H \otimes H^*) \rightarrow \mathbb{C}$.

To define the corresponding SLCMC's, (which we denote by the same symbols), as usual, it suffices to define sections over pairs of constant covering spaces $(U \times S, U \times T)$ of a complex manifold U . For defining $C(\mathcal{M})$, we need a concept of a holomorphically

varying 1-element of \mathcal{M} . To this end, we denote by $Hol(U, \mathbb{C}_2)$ the lax commutative monoid of finite-dimensional holomorphic bundles on U . Then the concept we need is

$$M_U \in_1 \mathcal{M}_U := \mathcal{M} \otimes_{\mathbb{C}_2} Hol(U, \mathbb{C}_2).$$

This determines the SLCMC $C(\mathcal{M})$. To define $C(\mathcal{M}, H)$, note that H does not depend on U , so in the case of the constant covering space described above, we simply need a

$$u_U : M_U \rightarrow_2 H_U^{\hat{\otimes} S} \hat{\otimes} H_U^{*\hat{\otimes} T},$$

where H_U is the constant H -bundle on U , and the 2-morphism means a morphism of holomorphic bundles.

Definition. A modular functor on an SLCMC C with labels \mathcal{M} is a (lax) morphism of SLCMC's $\phi : C \rightarrow C(\mathcal{M})$. A CFT on C with modular functor on labels \mathcal{M} with state space H is a (lax) morphism of SLCMC's $\Phi : C \rightarrow C(\mathcal{M}, H)$.

For a very detailed discussion of issues related to laxness, see [9]. Now note that $C(?)$ is a 2-functor from the 2-category $\mathbb{C}_2 - mod$ of lax \mathbb{C}_2 -modules (1-morphisms are equivalences of \mathbb{C}_2 -modules, and 2-morphisms are natural isomorphisms compatible with \mathbb{C}_2 -module structure) into the 2-category of SLCMC's. Similarly, $C(?, ?)$ is a 2-functor from the 2-category $\mathbb{C}_2 - mod_*$ of pairs \mathcal{M}, H , where \mathcal{M} is a \mathbb{C}_2 -module, and $H \in_1 \mathcal{M}^{Hilb}$. Here 1-morphisms in $\mathbb{C}_2 - mod, \mathbb{C}_2 - mod_*$ are equivalences of lax \mathbb{C}_2 -modules, 2-morphisms are natural isomorphisms compatible with \mathbb{C}_2 -module structure. (In $\mathbb{C}_2 - mod_*$, 1-morphisms $(\mathcal{M}, H) \rightarrow (\mathcal{N}, K)$ are 1-morphisms $\phi : \mathcal{M} \rightarrow \mathcal{N}$ in $\mathbb{C}_2 - mod$ together with a 2-isomorphism $\lambda : \phi(H) \rightarrow K$; a 2-morphism $\phi \rightarrow \psi$ is a 2-morphism in $\mathbb{C}_2 - mod$ which induces an isomorphism $\phi(H) \rightarrow \psi(H)$ commuting with the λ 's.)

We use this to build a 2-category Γ of CFT's as a "comma 2-category". The objects are tuples \mathcal{M}, H, Φ where, Φ is a CFT on C with labels \mathcal{M} and state space H , 1-morphisms are tuples Φ, Ψ, f, ι , where Φ, Ψ are CFT's with labels \mathcal{M}, \mathcal{N} and state spaces H, K , f is a $\mathbb{C}_2 - mod_*$ -1-morphism $(\mathcal{M}, H) \rightarrow (\mathcal{N}, K)$ and ι is a natural isomorphism $f(M_X) \rightarrow_2 N_X$, where M_X, N_X are the 1-elements of \mathcal{M}, \mathcal{N} assigned to $X \in Obj C$ by Φ, Ψ which commutes with SLCMC structure maps and the u 's assigned by Φ, Ψ (we have used the notation of sections over a point, but we mean this in the stack sense for sections over any complex manifold U). 2-morphisms $\Phi, \Psi, f, \iota \rightarrow \Phi, \Psi, g, \kappa$ are given by 2-isomorphisms $f \rightarrow g$ in $\mathbb{C}_2 - mod_*$ which commute with ι, κ (hence the u 's).

Next, we shall show that Γ is a symmetric monoidal 2-category. This means that we have a lax 2-functor \oplus with the same coherence 1-isomorphisms as in a symmetric monoidal category, but coherence diagrams commute up to 2-cells; the 2-cells, in turn, are required to satisfy all commutations valid for the trivial 2-cells of coherence diagrams in an ordinary symmetric monoidal category. Thus, the main point is to construct the 2-functor \oplus . Suppose we have two objects \mathcal{M}, H, Φ and \mathcal{N}, K, Ψ of Γ . Then their sum is $\mathcal{M} \oplus \mathcal{N}, H \oplus K, \Phi \oplus \Psi$: The first component is the direct sum in $\mathbb{C}_2 - mod$. $H \oplus K$ is the direct sum induced by that functor on 1-morphisms. The symbol $\Phi \oplus \Psi$, however, has to be defined explicitly. For simplicity, we shall restrict to sections over a point. Then, the data which remains to be defined, for an object X of the source SLCMC, is

$$M \oplus N, \tag{19}$$

and

$$u : M \oplus N \rightarrow_2 (H \oplus K)^{\otimes S} \otimes (H \oplus K)^{* \otimes T}. \tag{20}$$

Of this data, for X connected, (19) is, again, the direct sum induced by the direct sum of \mathbb{C}_2 -modules on 1-morphisms, composed with the canonical map

$$\begin{aligned} & \mathcal{M}^{\otimes S} \otimes \mathcal{M}^{* \otimes T} \oplus \mathcal{N}^{\otimes S} \otimes \mathcal{N}^{* \otimes T} \\ & \rightarrow (\mathcal{M} \oplus \mathcal{N})^{\otimes S} \otimes (\mathcal{M} \oplus \mathcal{N})^{* \otimes T}. \end{aligned} \tag{21}$$

Analogously, (20) is given by the \oplus of \mathbb{C}_2 -modules on 2-morphisms, composed with (21). For X non-connected, note that we are forced to define all data by applying the tensor product to the data on connected components. This definition extends to 1-morphisms and 2-morphisms in a standard way to produce a symmetric monoidal 2-category of CFT's Γ .

Note that we may not always wish to work with the whole Γ , but with some symmetric monoidal sub-2-category Λ ; for example, we may take direct \oplus -sums of copies of a given CFT.

Now there is an infinite loop space machine for 2-categories: for example, Segal's machine. Segal's machine is supposed to construct an \mathcal{F} -space, which is a functor from the category \mathcal{F} of finite sets with base point $*$ and based maps into spaces (alternately, one can think of this as a category of partial maps); it is also required that the functor (called \mathcal{F} -space) B be *special*, which means that the product map from $B(n)$ to the product of copies of $B(1)$ by the maps which send all numbers in $\{1, \dots, n\}$ except i into the basepoint be an equivalence.

Now to produce a special \mathcal{F} -space from a symmetric monoidal category C , simply consider the category $C(n)$ which is a category of diagrams, whose objects are tuples (x_T) of objects of C indexed by non-empty subsets of S , together with isomorphisms

$$\bigoplus_{i \in T} x_{(i)} \cong x_T. \tag{22}$$

Morphisms are commutative diagrams of the obvious kind (see [37]). Now $C(?)$ is a functor from \mathcal{F} into categories, so applying the classifying space gives the requisite \mathcal{F} -space. It is special by basic theorems about the homotopy types of classifying spaces.

However, now note that the same definition (22) in the case of a symmetric monoidal 2-category C gives a 2-category $C(n)$. The only difference is that on 1-morphisms, we do not consider merely diagrams commutative on the nose, but up to 2-cells and 2-morphisms are systems of 2-cells which further commute with the 2-cells thus introduced. With that, however, $C(?)$ becomes a (strict) functor from \mathcal{F} into 2-categories.

So, we are done if we can produce a functorial classifying space construction B_2 on 2-categories, and show that the \mathcal{F} -space $B_2 C(?)$ is special. The latter is a straightforward exercise which we omit. For the former, however, we remark that to define a classifying space of a 2-category C , we can first form the bar construction $B_1 = B(\text{Mor}(C))$, i.e. the bar construction on 2-morphisms. However, if C is lax, then B_1 is not a category, but composition is defined with respect to a contractible operad (without permutations). The operad D is as follows: the space $D(n)$ is the standard $(n - 1)$ -simplex and the composition is given by joining:

$$D(k) \times D(n_1) \times \dots \times D(n_k) \rightarrow D(n_1) * \dots * D(n_k) = D(n_1 + \dots + n_k).$$

Nevertheless, it is well known that such A_∞ -categories still have a classifying space functor (for example, one can “rectify” them by push-forward change of operads to the one point operad without permutations, which encodes associativity).

Thus, we have produced a symmetric monoidal 2-category of CFT’s, and an infinite loop space machine for such a case. Therefore, we have an infinite loop space E . This is related to the kind of construction used in [16] to give a candidate for an elliptic-like cohomology theory and seems like an improvement in the sense that it gives a model for the *additive* infinite loop structure (a “free” construction was used in [16]).

However, Baas-Dundas-Rognes [2] point out that this kind of construction is naive. The problem is that there are not enough isomorphisms of free lax \mathbb{C}_2 -modules: they are essentially just permutation matrices composed with diagonal matrices of line bundles. In [2], a solution to this problem is proposed, conjecturally calculating the algebraic K -theory of \mathbb{C}_2 . The point is to consider, instead of invertible matrices of finite-dimensional vector spaces, numerically invertible matrices, which means that the corresponding matrix of dimensions of the entry vector spaces is invertible.

Unfortunately, this approach does not seem satisfactory for the purposes of CFT: along with an iso $f : \mathcal{M} \rightarrow \mathcal{N}$, we need to consider also the inverse $\mathcal{M}^* \rightarrow \mathcal{N}^*$ of the adjoint morphism $\mathcal{N}^* \rightarrow \mathcal{M}^*$; there is no candidate for such inverse when f is only numerically invertible.

Another clue that something else is needed is the following example of *bc*-systems, whose anomaly, it seems, can only be expressed by considering “modular functors with positive and negative parts”.

Example. Consider the chiral *bc*-system of Ω^α -forms, $\alpha \in \mathbb{Z}$ (see also Sect. 3 above). The *bc*-system was first considered mathematically by Segal [35], but the observation that the super-modular functor formalism is needed to capture its properties is due to P. Deligne ([19]). In the case, the state space of the *bc*-system is the “fermionic Fock space”

$$\mathcal{F}_\alpha = \hat{\Lambda}(H_+ \oplus \overline{H}_-), \tag{23}$$

where $H = \langle z^n dz^\alpha | n \in \mathbb{Z} \rangle$ and H_+ is, say, the subspace $\langle z^n dz^\alpha | n \geq 0 \rangle$. We select some real form to make this a positive definite Hilbert space (cf. [16], Chap. 2). Then the modular functor is 1-dimensional, and is given by the determinant line of $\Omega^\alpha X$, the space of holomorphic α -forms on a worldsheet X . The reason why a super-modular functor is needed here is that we are dealing with Grassmannians, and signs must be introduced when permuting odd-degree variables for the CFT to be consistent; no such signs, however, occur in CFT’s with 1-dimensional anomaly as considered above (see [19] for more details).

Thus, it seems that \mathbb{C}_2 in our definition of modular functors and CFT’s should be replaced by some sort of “group completion” which would involve $\mathbb{Z}/2$ -graded vector spaces. A candidate for such a construction is given in the next section, although we will see that this comes at the price of substantially increasing technical difficulty.

6. The Group Completion of \mathbb{C}_2

As argued above, it would be desirable to have a group completion $\hat{\mathbb{C}}_2$ of \mathbb{C}_2 over which we could do the analogues of all of our constructions as suggested by Baas-Dundas-Rognes [2]: this would give approaches to axiomatizing CFT with positive and

negative-dimensional modular functors, as well as anti- D -branes with 1-dimensional anomaly. In this section, we propose such a construction. However, as also pointed out in [2], any such construction is necessarily accompanied by substantial difficulties. The first problem is to even define what we mean by “group completion”. It is easy to show that for a lax commutative ring R , BR is always an Eilenberg-MacLane space (and hence cannot be used for our purposes), but there is strong evidence that a large class of weaker categorical notions of “weakly group-complete” lax commutative semirings suffer from the same problem [39].

We take an alternate approach of introducing topology into the picture. This means we construct a model of a topological lax commutative semiring $\hat{\mathcal{C}}_2$ where there is an object -1 so that $1 \oplus (-1)$ is in the same connected component of 0 . While this approach does seem to lead to a viable definition, one must overcome a variety of technical difficulties caused by the additional topology.

The first issue is what is the appropriate 2-category $TCat$ of topological categories? The point is that requiring functors to be continuous on objects appears to restrict too much the notion of equivalence of topological categories, and consequently alter their lax colimits. To remedy this situation, we define 1-morphisms $\mathcal{C} \rightarrow \mathcal{D}$ in $TCat$ to be of the form

$$\begin{array}{ccc} \mathcal{C}' & \xrightarrow{F} & \mathcal{D} \\ G \downarrow & & \\ \mathcal{C} & & \end{array} \tag{24}$$

where F is a continuous functor and G is a *partition* which we define as follows: A partition is given by a topological space X and a continuous map

$$f : X \rightarrow \text{Obj}(\mathcal{C})$$

such that the topology on $\text{Obj}(\mathcal{C})$ is induced by f (we work in the category of weakly Hausdorff compactly generated topological spaces). Then we have $\text{Obj}(\mathcal{C}') = X$, $\text{Mor}(\mathcal{C}')$ is a pullback of the form

$$\begin{array}{ccc} \text{Mor}(\mathcal{C}') & \longrightarrow & \text{Mor}(\mathcal{C}) \\ \downarrow & & \downarrow s \times T \\ X \times X & \xrightarrow{f \times f} & \text{Obj}(\mathcal{C}) \times \text{Obj}(\mathcal{C}). \end{array}$$

Two functors F_1, F_2 as in (24) are considered *equal* if they coincide on a common partition, i.e. we have a commutative diagram

$$\begin{array}{ccccc} & & \mathcal{C}'_2 & & \\ & G_2 \swarrow & \uparrow H_2 & \searrow F_2 & \\ & \mathcal{C} & \mathcal{C}' & \xrightarrow{F} & \mathcal{D} \\ & \swarrow G_1 & \downarrow H_1 & \nearrow F_1 & \\ & & \mathcal{C}'_1 & & \end{array}$$

where H_1, H_2 are partitions. Composition is defined by pullback in the usual way, using the following

Lemma 1. *If we have a pullback*

$$\begin{array}{ccc}
 Y' & \xrightarrow{j} & X' \\
 h \downarrow & & \downarrow f \\
 Y & \xrightarrow{g} & X
 \end{array} \tag{25}$$

in the category of compactly generated weakly Hausdorff spaces such that f induces the (compactly generated) topology on X , then h induces the (compactly generated) topology on Y .

Proof. Suppose $V \subset Y$, $v \in (Y - V) \cap Cl(V)$, $h^{-1}(V)$ closed. Then there exists K compact where v is a limit point of $K \cap V$. So, we may replace V by $K \cap V$ and assume $Cl(V) = K = Y$. Next, let $Z = g(K)$, so Z is compact.

Case 1. $g(V) \neq Z$. Then there exists $T \subset X'$ compact, $T \cap f^{-1}(g(V))$ not closed. Consider the pullback

$$\begin{array}{ccc}
 T' & \longrightarrow & T \\
 \downarrow & & \downarrow \\
 g^{-1}f(T) & \longrightarrow & f(T).
 \end{array}$$

Then T' is compact since g is proper. But then $j(T' \cap h^{-1}(V)) = f^{-1}(g(V)) \cap T$, so $T' \cap h^{-1}(V)$ cannot be closed in T' (since $j|_{T'}$ is closed, T' being compact). This is a contradiction.

Case 2. $g(V) = Z$. Then in particular, there exists $z' \in V$, $g(v) = g(z') =: z$. Now we may assume that

$$v \text{ is a limit point of } V \cap g^{-1}(\{z\}). \tag{26}$$

Indeed, otherwise, since $K = Y$ is compact weakly Hausdorff, it is normal, hence regular and there exist U, W open in K , $U \cap W = \emptyset$, $v \in U$, $Cl(V \cap g^{-1}(\{z\})) \subset W$. But then we may replace Y by $Y - V$ (and X by $g(Y - V)$), and we are back to Case 1. So we may assume (26). But then we may replace X by $\{z\}$ and Y by $g^{-1}(\{z\})$. But then (25) is a product, in which case the statement of the lemma is obviously true (a product projection induces the topology on its target in the compactly generated weakly Hausdorff category). Thus, we have a contradiction again. \square

Now 2-morphisms in $TCat$ are defined as follows: we can assume we have two 1-morphisms F_1, F_2 given as

$$\begin{array}{ccc}
 \mathcal{C}' & \xrightarrow{F_1, F_2} & \mathcal{D} \\
 G \downarrow & & \\
 \mathcal{C} & &
 \end{array}$$

where G is a partition. Then a 2-morphism is given by a partition

$$\mathcal{C}'' \xrightarrow{G'} \mathcal{C}'$$

and a continuous natural transformation

$$F_1 G' \rightarrow F_2 G'.$$

Again, two 2-morphisms are identified if they coincide after pullback via a partition, similarly as above.

This completes the definition of the 2-category $TCat$. This 2-category is defined in such a way that it has lax limits defined in the same way as in Cat [9]. (Lax limit is given as the category whose objects and morphisms are lax cones from a point or an arrow to a diagram with the topology induced from the product; by “lax”, we always mean “up to coherences which are iso”.)

Next, one may discuss lax monads in $TCat$, which, in our definition, are lax functors $C : TCat \rightarrow TCat$ with lax natural transformations

$$\mu : CC \rightarrow C, \quad \eta : Id \rightarrow C$$

which are associative and unital up to coherence isos with commutative coherence diagram the same as for lax monoids. For a lax monad C in $TCat$ we then have a category of lax C -algebras whose objects are objects M of $TCat$ together with a functor

$$\theta : CM \rightarrow C$$

satisfying associativity and unitality up to coherence isos with commutative coherence diagrams of the same form as those for categories with lax action of a lax monoid.

Then lax algebras over a lax monad in $TCat$ form a 2-category which has lax limits created by the forgetful functor to $TCat$. We may be interested in lax algebras over a strict monad, for example the monad associated with a theory T . One example of a lax monad whose lax algebras we are interested in is gotten from a 2-theory (Θ, T) and a lax T -algebra I . Then we can define a lax monad $C_{\Theta, I}$ not over $TCat$, but over the category $Tcat^{I^k}$ of strict functors $I^k \rightarrow TCat$. In effect, $C_{\Theta, I}(X)$ has

$$C_{\Theta, I}(X)_i = \coprod \Theta(m)((\gamma_1, \dots, \gamma_p); \gamma) \times \prod_{i=1}^p X\gamma_i(j_1, \dots, j_m),$$

where the coproduct is indexed over $m, (j_1, \dots, j_m) \in I, \gamma \in T(m)^k, \gamma(j_1, \dots, j_m) = i, \gamma_1, \dots, \gamma_p \in T(m)^k$. Then lax $C_{\Theta, I}$ -algebras are precisely lax algebras over (Θ, T) with underlying lax T -algebra I .

We are now ready to describe a topological lax semiring $\hat{\mathbb{C}}_2$ with an object -1 such that $1 \oplus (-1)$ is in the same connected component as 0 . First consider the lax semiring $s\mathbb{C}_2$ of pairs $(V_+, V_-), V_+, V_- \in Obj(\mathbb{C}_2)$ with the lax \mathbb{C}_2 -module structure given by $\mathbb{C}_2 \oplus \mathbb{C}_2$ and multiplication

$$(V_+, V_-) \otimes (W_+, W_-) = (V_+ \otimes W_+ \oplus V_- \otimes W_-, V_+ \otimes W_- \oplus V_- \otimes W_+).$$

Now in $s\mathbb{C}_2$, consider the full subcategory J on pairs (V_+, V_-) , where $dim(V_+) = dim(V_-)$. Then J is a lax $s\mathbb{C}_2$ -module, and $J \oplus \mathbb{C}_2$ is a lax commutative \mathbb{C}_2 -algebra with a lax commutative \mathbb{C}_2 -algebra morphisms $J \oplus \mathbb{C}_2 \rightarrow \mathbb{C}_2$ (an augmentation) and $J \oplus \mathbb{C}_2 \rightarrow s\mathbb{C}_2$ (the inclusion). Thus, we have a lax simplicial commutative \mathbb{C}_2 -algebra (=lax functor $\Delta^{Op} \rightarrow$ lax commutative \mathbb{C}_2 -algebras)

$$B_{\mathbb{C}_2}(\mathbb{C}_2, J \oplus \mathbb{C}_2, s\mathbb{C}_2). \tag{27}$$

We propose $\hat{\mathbb{C}}_2$ to be the realization of (27). This needs some explaining, namely we must define realization. We shall describe a lax realization functor

$$\mathcal{A} \rightarrow |\mathcal{A}|$$

from lax simplicial commutative \mathbb{C}_2 -algebras to topological commutative \mathbb{C}_2 -algebras (i.e. commutative \mathbb{C}_2 -algebras in $TCat$). We want to mimic as closely as possible the strict construction. This means that we will define (27) as the lax simplicial realization in the 2-category of lax \mathbb{C}_2 -modules, which we must define. First, let, for a space X , $\mathbb{C}_2 X$ be the free lax \mathbb{C}_2 -module on X (objects and morphisms are finite formal linear combinations with coefficients in objects and morphisms of \mathbb{C}_2 , and the topology is induced from the topologies of finite powers of X). We want the realization $|\mathcal{A}|$ to be the lax coequalizer of

$$\bigoplus_{m,n} (\mathbb{C}_2 \Delta_m \otimes \mathcal{A}_n) \begin{matrix} \xrightarrow{\quad} \\ \rightrightarrows \\ \xrightarrow{\quad} \end{matrix} \bigoplus_n (\mathbb{C}_2 \Delta_n \otimes \mathcal{A}_n) \tag{28}$$

(the arrows are the usual two arrows coming from lax simplicial structure, \oplus, \otimes are over \mathbb{C}_2). To construct the lax coequalizer (28), we can take the objects of

$$\bigoplus_{m,n} (\mathbb{C}_2 \Delta_m \otimes \mathcal{A}_n) \oplus \bigoplus_n (\mathbb{C}_2 \Delta_n \otimes \mathcal{A}_n). \tag{29}$$

To get morphisms, we take the morphisms of (29), and adjoin isomorphisms between all source and target objects of the arrows in (28). Take the free topological category spanned by these morphisms, modulo the obvious commutative diagrams required. This gives us a category with the lax \mathbb{C}_2 -module (29) as a subcategory. The free construction we must then perform is applying the *strict* left adjoint to the forgetful functor from the category of lax \mathbb{C}_2 modules with lax submodule (29) on the same set of objects (taking only functors which are identity on objects) to the category of categories with subcategory (29) on the same set of objects (taking only functors which are identity on objects). As usual, the functors are strict because objects and coherences are already specified.

This completes the construction of the lax simplicial realization (27). One must still prove that this is a lax \mathbb{C}_2 -algebra, but this is accomplished analogously as in the strict case, using the shuffle map (and the morphism definition (24) to assure continuity).

Now topological SLCMC's $C(\mathcal{M}), C(\mathcal{M}, H)$ for a finitely generated free topological lax $\hat{\mathbb{C}}_2$ -module \mathcal{M} are defined analogously as over \mathbb{C}_2 . (Since the underlying stack of covering spaces $I = Set$ does not change, LCMC's can be described as lax algebras over a lax monad in $TCat$ as above, and therefore stacks over a Grothendieck topology can be defined to be, as usual, contravariant functors which take Grothendieck covers to lax limits.)

However, the topology would be of little use if we simply took for our definition of modular functor a lax morphism of SLCMC's from \mathcal{C} to $C(\mathcal{M})$ (similarly for CFT's). Instead, the corresponding "derived notion" is appropriate. This means that we should consider lax morphisms of topological SLCMC's

$$B(C_{\Theta,S}, C_{\Theta,S}, \mathcal{C}) \rightarrow C(\mathcal{M}), \tag{30}$$

where Θ denotes the 2-theory of LCMC's, and S the lax commutative monoid of finite sets, as above. The left hand side is obtained by taking the bar construction section-wise and then applying the lax left adjoint to the forgetful functor from SLCMC's to pre-stacks of LCMC's.

It still remains to define a realization functor from lax simplicial lax $C_{\Theta,S}$ -algebras to topological lax $C_{\Theta,S}$ -algebras. Analogously as in the case of B_{C_2} , however, we may simply lax-realize in the 2-category $(TCat)^{Set^2}$ (where it is easy to construct lax colimits, cf. [9]), and use the lax analogue of Milnor's map $|A| \times |B| \rightarrow |A \times B|$ to obtain lax $C_{\Theta,S}$ -algebra structure on the realization. We omit the details.

7. The General Anomaly for Open-Closed CFT

In this section, we shall apply the principles of Sect. 5 to propose a general definition of open-closed CFT with both multiple D -branes and multi-dimensional conformal anomaly (although without any group completion). We shall see, however, that this is necessarily even much more complicated than what we have done in Sect. 5. We have already argued that neither the "set of D -branes" nor the "set of labels" should be sets. Rather, they should be higher vector spaces. However, on a boundary component of the worldsheet where several open parametrization components are present, we need to take traces of "sets of labels" over "sets of D -branes". This suggests that our model of "set of D -branes" must be one categorical level above our notion of "set of labels".

Therefore, we propose that the "set of D -branes" be a 3-vector space \mathcal{A} . When dealing with 3-vector spaces, note that they are 2-categories. 3-vector spaces are, by definition, 2-lax modules over the 2-lax commutative semiring \mathbb{C}_3 . We must, of course, define these notions. On generalizing from lax to 2-lax structures, we find it easiest to follow the approach of [17]. Let T be a theory. Then let $Th(T)$ be the free theory on T , with the canonical projection of theories $\phi : Th(T) \rightarrow T$. Let G be a groupoid with objects $Th(T)$ and one isomorphism $x \rightarrow y$ for every $x, y \in Th(T)$ which satisfy $\phi(x) = \phi(y)$. Then $(Th(T), G)$ is a theory (strictly) enriched over categories and a lax T -algebra is the same thing as a strict $(Th(T), G)$ -algebra enriched over categories.

Now to go to the next level, consider the forgetful functor

$$U : \text{Theories enriched over groupoids} \rightarrow \mathcal{P},$$

where \mathcal{P} is the (strict) category of pairs (T, G) where T is a theory, G is a graph with objects T . Then let F be the left adjoint of U . Notice that F is the identity on objects T , so we may write

$$F(T, G) = (T, F(G)).$$

Now we have a map of theories enriched over groupoids:

$$\psi : (Th(T), F(G)) \rightarrow (Th(T), G).$$

Therefore, we may create a 2-category $(Th(T), F(G), H)$ by putting exactly one 2-isomorphism between every $\alpha, \beta \in F(G)$ with $\psi(\alpha) = \psi(\beta)$. Then the 2-category $(Th(T), F(G), H)$ is naturally a theory (strictly) enriched over 2-categories, and a 2-lax T -algebra is a 2-category which is a strict $(Th(T), F(G), H)$ -algebra enriched over 2-categories. (Obviously, one may proceed further in the same way to define even higher laxness, but we shall not need that here.)

One remark to be made is that theories, strictly speaking, do not model universal algebras which are modelled on more than one set, such as a module over a ring (which is

modelled over two sets). However, algebras modelled over k sets can be easily included in the formalism by modifying the concept of theory to a category with objects \mathbb{N}^k (i.e. k -tuples of natural numbers) with the axiom that for all $a, b \in \mathbb{N}^k$, $a + b$ is the categorical product of a, b . This is the concept of *multisorted* theories (see [9] for more details and references). All of our constructions generalize to this context.

Now to identify the categorical levels with the levels we considered before, we will denote objects of 3-vector spaces by \in_0 and morphisms of 3-vector spaces by \rightarrow_0 . Thus, a 0-morphism $\mathbb{C}_3 \rightarrow \mathbb{C}_3$ (where \mathbb{C}_3 is the lax symmetric monoidal category of 2-vector spaces) is a 2-vector space, and the notations $\rightarrow_1, \rightarrow_2$ of such 2-vector spaces will coincide with the notations we used above.

Now given the 3-vector space \mathcal{A} (“the set of D -branes”), we must introduce the “set of labels” for anomalies. The “set of closed labels” will be, as before, a 2-vector space, which we will denote by \mathcal{C} . The “set of open labels” will be an object of the form

$$\mathcal{O} \in_0 \mathcal{A} \otimes_{\mathbb{C}_3} \mathcal{A}^*.$$

(We remark here that \mathcal{A}^* for a 3-vector space \mathcal{A} is defined analogously as in the case of 2-vector spaces.) Now we would like to define an SLCMC $C(\mathcal{A}; \mathcal{C}, \mathcal{O})$. All our SLCMC’s in this section shall have two labels, 1 and m (closed and open). However, note that there is another subtlety we must provide for, namely that the set Γ of all possible *incidence graphs* whose vertices are open and closed parametrization and D -brane components and edges describe their incidence relations with the obvious conditions (e.g. all vertices have degree 2, etc.) is itself an SLCMC, and in order to correctly keep track of incidences on the boundary, we must consider SLCMC’s over Γ (see also the end of Sect. 2 above).

We shall only describe sections of $C(\mathcal{A}; \mathcal{C}, \mathcal{O})$ over a given object G of Γ over a single point, over four given sets $S_{1,in}, S_{1,out}, S_{m,in}, S_{m,out}$ of inbound and outbound closed and open “components”. Let P denote the set of closed D -brane components of G . Before making the definition, note that we have canonical dual 0-morphisms

$$\mathbb{C}_3 \xrightarrow{\eta} \mathcal{A} \otimes_{\mathbb{C}_3} \mathcal{A}^* \xrightarrow{\epsilon} \mathbb{C}_3. \tag{31}$$

(If no further discussion is made, (31) requires a finiteness assumption about \mathcal{A} .) Their composition is a 2-vector space which we shall denote by $tr_0\mathcal{A}$. The set of sections of $C(\mathcal{A}; \mathcal{C}, \mathcal{O})$ are 1-elements

$$M \in_1 \bigotimes_{S_{1,in}} \mathcal{C}^* \otimes \bigotimes_{S_{1,out}} \mathcal{C} \otimes \bigotimes_P tr_0\mathcal{A} \otimes tr_{0,cyclic} \left(\bigotimes_{S_{m,in}} \mathcal{O}^* \otimes \bigotimes_{S_{m,out}} \mathcal{O} \right). \tag{32}$$

Here the tensor products are over \mathbb{C}_2 , and $tr_{0,cyclic}$ denotes composition with the tensor product of the appropriate number of ϵ ’s; note that although not explicitly written, the definition of $tr_{0,cyclic}$ makes use of all of the structure of G . Now in order to give (32) a structure of LCMC, one must show an appropriate gluing property, but this is analogous to our discussion for closed CFT’s.

Now let, as above, \mathcal{D} be the SLCMC of closed-open worldsheet with one closed and one open label. Then an open-closed CFT anomaly (modular functor) is a map of SLCMC’s over Γ

$$\mathcal{D} \rightarrow C(\mathcal{A}; \mathcal{C}, \mathcal{O}).$$

Now to define open-closed CFT, we must add the ‘‘Hilbert spaces’’. The ‘‘closed Hilbert space’’ is, as above,

$$H \in_1 \mathcal{C}^{Hilb}.$$

The ‘‘open Hilbert space’’ is a 1-morphism

$$K : \eta \rightarrow_1 \mathcal{O}^{Hilb},$$

where η is as in (31). We shall now define an SLCMC $C(\mathcal{A}; \mathcal{O}, \mathcal{C}; H, K)$ over Γ . As above, we shall specialize to sections over a single point and single object of Γ , with the same notation as above. Then a section consists of a section (32) of $C(\mathcal{A}; \mathcal{C}, \mathcal{O})$ and a 2-morphism

$$M \rightarrow_2 \underset{S_{1,in}}{\hat{\otimes}} H^* \underset{S_{1,out}}{\hat{\otimes}} \underset{P}{\hat{\otimes}} H \underset{P}{\hat{\otimes}} \eta_{tr_0 \mathcal{A}}$$

$$\hat{\otimes} tr_{1,cyclic} \left(\underset{S_{m,in}}{\hat{\otimes}} K^* \underset{S_{m,out}}{\hat{\otimes}} \underset{S_{m,out}}{\hat{\otimes}} K \right).$$

Here $tr_{1,cyclic}$ is given by the structure of 2-category, and $\eta_{tr_0 \mathcal{A}}$ is the canonical ‘‘unit’’ 1-element of $tr_0 \mathcal{A}$. To be more precise, write, in (31),

$$\eta(\mathbb{C}_2) = \bigoplus_{i=1}^n \mathcal{V}_i \otimes_{\mathbb{C}_2} \phi_i,$$

so

$$tr_0 \mathcal{A} = \bigoplus_{i=1}^n \phi_i \mathcal{V}_i.$$

But then one can show

$$\phi_i \mathcal{V}_j = \delta_i^j \mathbb{C}_2,$$

so we have

$$tr_0 \mathcal{A} = \bigoplus_{i=1}^n \mathbb{C}_2,$$

and we can write

$$\eta_{tr_0 \mathcal{A}} = \bigoplus_{i=1}^n \mathbb{C} \in_1 \bigoplus_{i=1}^n \mathbb{C}_2.$$

Of course, such discussion reveals the weaknesses of the higher vector space formalism, and the desirability to really work, again, in a suitable higher group completion. However, we do not work out that approach here.

8. Appendix: Stacks of Lax Commutative Monoids with Cancellation

To make this paper self-contained, we review here the basic definitions [16] related to stacks of lax commutative monoids with cancellation (SLCMC’s). We must begin by defining lax algebras. The formalism we use is theories according to Lawvere, and their extension which we call 2-theories. Recall first that a *theory* according to Lawvere [20]

is a category T with objects \mathbb{N} (the set of all natural numbers $0, 1, 2, \dots$) such that n is the product of n copies of 1 . Categories of algebraic structures given by a set of operations and relations on one set X can be encoded by a theory, where $T_n = \text{Hom}(n, 1)$ is the set of all n -ary operations of the algebraic structure (including all possible compositions, repetitions of one or more variables, etc.).

Definition. A 2-theory consists of a natural number k , a theory T and a (strict) contra-variant functor Θ from T to the category of categories (and functors) with the following properties. Let T^k be a category with the same objects as T , and $\text{Hom}_{T^k}(m, n) = \text{Hom}_T(m, n)^{\times k}$ (obvious composition). Then

$$\text{Obj}(\Theta(m)) = \coprod_n \text{Hom}_{T^k}(m, n),$$

on morphisms, Θ is given by precomposition on $\text{Obj}(\Theta(m))$, and $\gamma \in \text{Hom}_{T^k}(m, n)$ is the product, in $\Theta(m)$, of the n -tuple $\gamma_1, \dots, \gamma_n \in \text{Hom}_T(m, 1)$ with which it is identified by the fact that T is a theory. We also speak of a 2-theory fibered over the theory T .

The example relevant to CFT is the 2-theory of commutative monoids with cancellation. T is the theory of commutative monoids with an operation $+$, and $k = 2$. The 2-theory Θ has three generating operations, addition (or disjoint union) $+$: $X_{a,c} \times X_{b,d} \rightarrow X_{a+b,c+d}$, unit $0 \in X_{0,0}$ and cancellation (or gluing) \checkmark : $X_{a+c,b+c} \rightarrow X_{a,b}$. The axioms are commutativity, associativity and unitality for $+$, 0 , transitivity for \checkmark and distributivity of \checkmark under $+$.

To get further, one needs to define algebras and lax algebras over theories and 2-theories. An algebra over a theory T is a set I together with, for each $\gamma \in T_n$, a map $\bar{\gamma} : I^{\times n} \rightarrow I$, satisfying appropriate axioms. These axioms can be written out explicitly, but a quick way to encode them is to notice that for a set I , we have the endomorphism theory $\text{End}(I)$, where $\text{End}(I)(n) = \text{Map}(I^{\times n}, I)$, and we may simply say that a structure of T -algebra on I is given by a map of theories $T \rightarrow \text{End}(I)$.

To define an algebra over a 2-theory Θ fibered over a theory T , we must first have an algebra I over the theory T (the ‘indexing theory’). This gives us, for $\gamma \in \text{Hom}_{T^k}(m, 1)$, a k -tuple of maps $\bar{\gamma} : I^{\times m} \rightarrow I$. In an algebra over the 2-theory, we have, in addition, a map

$$X : I^{\times k} \rightarrow \text{Sets}. \tag{33}$$

For a morphism in $\phi \in \text{Mor}(\Theta)$ from $(\gamma_1, \dots, \gamma_n) \in \text{Hom}_{T^k}(m, n)$ to $\gamma \in \text{Hom}_{T^k}(m, 1)$, we have, for each choice i_1, \dots, i_m of elements of I , maps

$$\bar{\phi} : X(\bar{w}_1(i_1, \dots, i_m)) \times \dots \times X(\bar{w}_n(i_1, \dots, i_m)) \rightarrow X(\bar{w}(i_1, \dots, i_m)), \tag{34}$$

satisfying appropriate axioms. Once again, we can avoid writing them down explicitly by defining the endomorphism 2-theory. Consider a set I and a map

$$X : I^k \rightarrow \text{Sets}.$$

To such data there is assigned a 2-theory $\text{End}(X)$ fibered over the theory $\text{End}(I)$: let

$$\Theta(w; w_1, \dots, w_n)$$

consist of the set of all possible simultaneous choices of maps

$$X(w_1(i_1, \dots, i_m)) \times \dots \times X(w_n(i_1, \dots, i_m)) \rightarrow X(w(i_1, \dots, i_m)), \quad (35)$$

where i_j range over elements of I . A structure of an algebra over the 2-theory Θ fibered over T is given by a morphism of 2-theories

$$(\Theta, T) \rightarrow (End(X), End(I)).$$

A lax algebra over a theory is a category I , with maps $\bar{\gamma}$ which are functors. We do not, however, require that these maps define a strict morphism from T to the endomorphism theory of I . Instead, this is only true up to certain natural isomorphisms, which we call coherence isomorphisms, which in turn are required to satisfy certain commutative diagrams, which are called coherence diagrams. This is, of course, always the case when defining lax algebras of any kind. But now the benefit of introducing theories is that the coherences and coherence diagrams always have the same shape. To be more precise, recall that the notion of theory itself is an algebraic structure which can be encoded by the sequence of sets $T(n)$, and certain operations on these sets satisfying certain identities. Denoting the set of operations defining theories by G (for ‘generators’), and identities by R (for ‘relations’), we observe that the set of coherence isomorphisms we must require for lax T -algebras is always in bijective correspondence with G , while the set of coherence diagrams needed is in bijective correspondence with R !

The concept of lax algebra over a 2-theory is defined in a similar fashion, but one important point is that one doesn’t want to consider the most general possible type of laxness (since that would lead to a 3-category). Rather, one starts with a lax algebra I over the indexing theory, and a strict functor

$$X : I^{\times k} \rightarrow Categories;$$

appropriate coherence isomorphisms and diagrams then follow in the same way as in the case of lax algebras over a theory (are indexed by operations and identities of 2-theories interpreted as a ‘universal algebras’ – see [16]).

The lax commutative monoid we most frequently consider is the groupoid S of finite sets and isomorphisms (the operation is disjoint union). More generally, we often consider a set of labels K and the lax commutative monoid of S_K of sets A labelled by K , i.e. maps $A \rightarrow K$. Again, the operation is disjoint union. The example of lax commutative monoid with cancellation fibered over S considered in [16] is the groupoid \mathcal{C} of *worldsheets* or rigged surfaces. These are 2-dimensional smooth manifolds with smooth boundary; further, each boundary component is parametrized by a smooth diffeomorphism with S^1 , and the surface has a complex structure with respect to which the boundary parametrization is analytic. Morphisms are biholomorphic diffeomorphisms commuting with boundary parametrization. Addition is disjoint union, and cancellation is gluing of boundary components. Similarly, again, one can consider the LCMC \mathcal{C}_K of worldsheets with K -labelled boundary components, which is an LCMC over S_K .

To complete the picture, one needs to consider stacks. We note that lax algebras over a theory and lax algebras (in our sense) over a 2-theory form 2-categories which have lax limits of strict diagrams (see Fiore [9]). For older references, which however work in slightly different contexts (and with different terminology), see Borceux [3] or [9]. For the 2-category structure, 1-morphisms are lax morphisms of lax algebras (functors such that there is a natural coherence isomorphism for every element of G), and 2-morphisms are natural isomorphisms which commute with the operations given by the theory (or

2-theory). Now for any 2-category \mathcal{C} with lax limits, and every Grothendieck topology \mathcal{B} , we can define \mathcal{B} -stacks over \mathcal{C} : they are simply contravariant functors $\mathcal{B} \rightarrow \mathcal{C}$ which turn Grothendieck covers into lax limits. Note that such stacks then themselves form a 2-category with respect to stack versions of the same 1-morphisms and 2-morphisms.

Now to turn \mathcal{C} into a stack of lax commutative monoids with cancellation, we must first specify the Grothendieck topology. Note that there are two choices of the topology, either just (finite-dimensional) smooth manifolds and open covers (non-chiral setting) or finite-dimensional complex manifolds and open covers (chiral setting). As remarked in Sect. 2 above, however, D -branes can only be considered in the non-chiral setting. To define the stack, one must first define the underlying stack of lax commutative monoids; the answer is simply the stack of covering spaces with finitely many sheets. Now one must define smooth or holomorphic families of worldsheets. We shall only make the definition in the holomorphic case, the smooth case is analogous. The most convenient way to make this precise is to consider, for a worldsheet X , the complex manifold Y obtained by gluing, locally, solid cylinders to the boundary components of X . Then, a holomorphic family of rigged surfaces X over a finite dimensional complex manifold B is a holomorphic map

$$q : Y \rightarrow B$$

transverse to every point, such that $\dim(Y) = \dim(B) + 1$ and B is covered by open sets U_i for each of which there are given holomorphic regular inclusions

$$s_{i,c} : D \times U_i \rightarrow Y$$

with

$$q \circ s_{i,c} = Id_{U_i},$$

where c runs through some indexing set C_i . Further, if $U_i \cap U_j \neq \emptyset$, we require that there be a bijection $\iota : C_i \rightarrow C_j$ such that

$$s_{i,c}|_{D \times (U_i \cap U_j)} = s_{j,\iota(c)}|_{D \times (U_i \cap U_j)}.$$

Then we let

$$X = Y - \left(\bigcup_i \bigcup_{c \in C_i} s_{i,c}((D - S^1) \times U_i) \right).$$

Then the fiber of X over each $b \in B$ is a rigged surface, which vary holomorphically in b , in the sense we want. (Note that the reason the maps s_c cannot be defined globally in B is that it is possible for a non-trivial loop in $\pi_1(B)$ to permute the boundary components of X .) The treatment of \mathcal{C}_K is analogous. As a rule, we shall use the same symbol for the SLCMC's $\mathcal{C}, \mathcal{C}_K$ as for the corresponding LCMC's (their sections over a point).

We are done with the review of SLCMC's, but we shall still briefly cover CFT's, as defined in [16]. Although this definition is subsumed by Sect. 5 above, the reader might still find the more elementary definition useful while reading the earlier sections. Let $\mathcal{H}_1, \dots, \mathcal{H}_n$ be complex (separable) Hilbert spaces. Then on $\mathcal{H}_1 \otimes \dots \otimes \mathcal{H}_n$, there is a natural inner product

$$\langle a_1 \otimes \dots \otimes a_n, b_1 \otimes \dots \otimes b_n \rangle = \langle a_1, b_1 \rangle \langle a_2, b_2 \rangle \dots \langle a_n, b_n \rangle.$$

The Hilbert completion of this inner product space is called the *Hilbert tensor product*

$$\mathcal{H}_1 \hat{\otimes} \dots \hat{\otimes} \mathcal{H}_n. \tag{36}$$

Now an element of (36) is called *trace class* if there exist unit vectors $e_{ij} \in \mathcal{H}$, where $j = 1, \dots, n$ and i runs through some countable indexing set I such that

$$x = \sum_{i \in I} \mu_i (e_{i1} \otimes \dots \otimes e_{in})$$

and

$$\sum_{i \in I} |\mu_i| < \infty.$$

The vector subspace of (36) of vectors of trace class will be denoted by

$$\mathcal{H}_1 \boxtimes \dots \boxtimes \mathcal{H}_n. \tag{37}$$

Note that (37) is not a Hilbert space. We have, however, canonical maps

$$\boxtimes : (\mathcal{H}_1 \boxtimes \dots \boxtimes \mathcal{H}_n) \otimes (\mathcal{H}_{n+1} \boxtimes \dots \boxtimes \mathcal{H}_{m+n}) \rightarrow \mathcal{H}_1 \boxtimes \dots \boxtimes \mathcal{H}_{m+n}$$

and, if \mathcal{H}^* denotes the dual Hilbert space to a complex Hilbert space \mathcal{H} ,

$$tr : \mathcal{H} \boxtimes \mathcal{H}^* \boxtimes \mathcal{H}_1 \boxtimes \dots \boxtimes \mathcal{H}_n \rightarrow \mathcal{H}_1 \boxtimes \dots \boxtimes \mathcal{H}_n.$$

This allows us to define a particular example of stack of LCMC's based on \mathcal{H} , which we will call $\underline{\mathcal{H}}$. The underlying stack of lax commutative monoids (T -algebras) is S . Now let $B \in \mathcal{B}$. Let s, t be sections of the stack S over B , i.e. covering spaces of B with finitely many sheets. Then we have an infinite-dimensional holomorphic bundle over B ,

$$(\mathcal{H}^*)^{\boxtimes s} \boxtimes \mathcal{H}^{\boxtimes t}. \tag{38}$$

What we mean by that is that there is a well defined sheaf of holomorphic sections of (38) (note that it suffices to understand the case when s, t are constant covering spaces, which is obvious). Now a section of $\underline{\mathcal{H}}$ over a pair of sections s, t of S is a *global* section of (38) over b ; the only automorphisms of these sections covering $Id_s \times Id_t$ are identities. The operation $+$, \boxtimes are given by the operations \boxtimes, tr (see above).

We can also define a variation of this LCMC for the case of labels indexed over a finite set K . We need a collection of Hilbert spaces

$$\mathcal{H}_K = \{\mathcal{H}_k | k \in K\}.$$

Then we shall define a stack of LCMC's $\underline{\mathcal{H}}_K$. The underlying stack of T -algebras (commutative monoids) is S_K . Let s, t be sections of S_K over $B \in \mathcal{B}$. The place of (38) is taken by

$$(\mathcal{H}_K^*)^{\boxtimes s} \boxtimes \mathcal{H}_K^{\boxtimes t}. \tag{39}$$

By the sheaf of holomorphic section of (39) when B is a point we mean that \boxtimes -powers of \mathcal{H}_k (or \mathcal{H}_k^*) for each label $k \in K$ are taken according to the number of points of $\Gamma(t)$ (resp. $\Gamma(s)$); when s and t are constant covering spaces B , the space of sections of (39) is simply the set of holomorphically varied elements of the spaces of sections over points of B (which are identified). This is generalized to the case of general s, t in the obvious way (using functoriality with respect to permutations of coordinates). As above, the only automorphisms of these sections covering $Id_s \times Id_t$ are identities.

Remark. For technical reasons (different types of convergence), the above setup involving Hilbert spaces and trace class elements is sometimes insufficient (see Example below and Sect. 4 above). Because of that, it is beneficial to generalize to a context where $\underline{\mathcal{H}}$ simply means any SLCMC over S (resp. S_K in the labelled case) whose spaces of sections are vector spaces, the operation $\check{?}$ is linear, and the operation $+$ is bilinear. We shall further assume that $\underline{\mathcal{H}}$ is a *sheaf* in the sense that the only endomorphisms over the identity in S (resp. S_K) is the identity.

Example. Consider the free bosonic CFT (10) discussed in Sect. 4. As remarked above, the description 10 is actually already not quite right: the vacuum state is to be an eigenstate of momentum 0, but there is no non-zero function in $L^2(\mathbb{R}, \mathbb{C})$ with support in the set $\{0\}$. For the same reason, we also find that the operator U_{A_q} associated with the standard annulus A_q is not trace class as defined (since, for example, 1 is a limit point of the spectrum of U_{A_q}). This is the usual problem in quantum mechanics. In the present setting, a solution along the lines of the Remark can be obtained as follows: Let \mathcal{F} be the bosonic Fock space, i.e.

$$\mathcal{F} = \widehat{Sym}\langle z^n, \bar{z}^n, n > 0 \rangle.$$

Then the sections of $\underline{\mathcal{H}}$ over (s, t) (over a point) are elements

$$f \in \prod_{k \in \mathbb{R}^{|s|+|t|}} \mathcal{F}^{\hat{\otimes}|t|} \hat{\otimes} \mathcal{F}^{*\hat{\otimes}|s|}$$

(the product is a categorical product of vector spaces) which have the property that for every pair of injections $i : u \rightarrow s, j : u \rightarrow t$ and every map

$$k : (s - i(u)) \coprod (t - j(u)) \rightarrow \mathbb{R},$$

we have

$$\int_{x \in \mathbb{R}^u} \mu(\phi_{i,j,k}(x)) < \infty,$$

where $\phi_{i,j,k}(x) = f(y), y \in \mathbb{R}^s \coprod \mathbb{R}^t$ is defined by $y(i(r)) = y(j(r)) = x(r), y(r) = k(r)$ for $r \in (s - i(u)) \coprod (t - j(u))$ and, for $z \in \mathcal{F}^{\hat{\otimes}|t|} \hat{\otimes} \mathcal{F}^{*\hat{\otimes}|s|}$,

$$\mu(z) = inf(\{ \sum |a_i| \mid z = \sum_{x \in s \coprod t} a_i \otimes e_x, \|e_x\| = 1 \text{ for all } x \},$$

and gluing along i, j is defined by

$$\check{f}(k) = \int_{x \in \mathbb{R}^u} tr(\phi_{i,j,k}(x)). \tag{40}$$

The expression (40) is always defined because of the condition imposed, and the condition is preserved by the gluing operation by Fubini's theorem.

Now we can define an *abstract CFT* based on an SLCMC \mathcal{D} with underlying stack of lax commutative monoids (SLCM) S simply as a 1-morphism of SLCMC's, over Id_S ,

$$\mathcal{D} \rightarrow \underline{\mathcal{H}}.$$

A similar definition applies if \mathcal{D} has underlying SLCM S_K , with $\underline{\mathcal{H}}$ replaced by $\underline{\mathcal{H}}_K$.

However, this notion still is not definitive in the sense that it does not capture anomaly. In Sect. 5 above, we give the most general definition of modular functor, but it is useful to review a direct definition from [16] at least in one special case, namely the 1-dimensional anomaly. To this end, we give the definition of \mathbb{C}^\times -central extension (or, equivalently, 1-dimensional modular functor) on an LCMC \mathcal{D} . This is a strict morphism of stacks of LCMC's

$$\psi : \tilde{\mathcal{D}} \rightarrow \mathcal{D} \tag{41}$$

over Id on the underlying stacks of LCM's with the following additional structure (for simplicity, let us just work in the holomorphic (chiral) setting): For each object B of \mathcal{B} , and each pair of sections s, t of S over B , and each section α of \mathcal{D} over $s, t, B, B' \rightarrow B$,

$$\psi^{-1}(\alpha|_{B'}) \tag{42}$$

with varying B' is the space of sections of a complex holomorphic line bundle over B . Furthermore, functoriality maps supplied by the structure of a stack of LCMC's on $\tilde{\mathcal{D}}$ are linear maps on these holomorphic line bundles. Regarding the operation $+$, we require that the map induced by $+$

$$\psi^{-1}(\alpha|_{B'}) \times \psi^{-1}(\beta|_{B'}) \rightarrow \psi^{-1}((\alpha + \beta)|_{B'}) \tag{43}$$

be a bilinear map, which induces an isomorphism of holomorphic line bundles

$$\psi^{-1}(\alpha|_{B'}) \otimes_{\mathcal{O}_{B'}} \psi^{-1}(\beta|_{B'}) \rightarrow \psi^{-1}((\alpha + \beta)|_{B'}) \tag{44}$$

(\mathcal{O}_B is the holomorphic structure sheaf on B).

Regarding the operation $\check{?}$, we simply require that if α is a section of \mathcal{D} over $s+u, t+u, B$, where u is another section of S over B , and $\check{\alpha}$ is the section over s, t, B which is obtained by applying the operation $\check{?}$ to α , then the map of holomorphic line bundles coming from LCMC structure

$$\psi^{-1}(\alpha|_{B'}) \rightarrow \psi^{-1}(\check{\alpha}|_{B'}) \tag{45}$$

($B' \rightarrow B$) is an isomorphism of holomorphic line bundles.

By a CFT with 1-dimensional modular functor over \mathcal{D} with underlying stack S we shall mean a CFT

$$\phi : \tilde{\mathcal{D}} \rightarrow \underline{\mathcal{H}} \tag{46}$$

(where $\tilde{\mathcal{D}}$ is a \mathbb{C}^\times -central extension of \mathcal{D} which has the property that ϕ is a linear map on the spaces of sections (42). Similarly in case \mathcal{D} has underlying stack S_K , we simply replace S by S_K everywhere throughout the definition.

It is appropriate to comment on a weaker kind of morphism of lax algebras where we do not require that the coherence maps be iso. By a *pseudomorphism* of lax T -algebras (and similarly in the cases of lax \mathbb{E} , T -algebra and their stacks) we shall mean a functor

$$f : X \rightarrow Y$$

together with morphisms (called cross-morphism, not necessarily iso)

$$\gamma(f, \dots, f) \rightarrow f\gamma \tag{47}$$

which commute with all the coherences in the lax T -algebra sense (we shall refer to these required commutative diagrams as cross-diagrams).

Remark. Although the anomaly of the linear σ -model considered in Sect. 4 is 1-dimensional, in Sect. 5 we considered higher-dimensional anomalies. It is therefore appropriate to reconcile the above remark concerning generalizing the SLCMC \mathcal{H} to cases when the Hilbert/trace class model fails due to non-convergence with our discussion of higher-dimensional modular functors via 2-vector spaces. In other words, what is the right generalization of $H \in \text{Obj}(\mathcal{M}^{\text{Hilb}})$ in $C(\mathcal{M}, H)$? The main point is that the Hilbert tensor powers $H^{\hat{\otimes} t} \hat{\otimes} H^{*\hat{\otimes} s}$ should be replaced by a “vector space indexed over \mathcal{M} ” which depends only on s, t and have appropriate designated “trace maps”.

The category $\mathcal{M}^{\text{Vect}}$ of *vector spaces indexed over \mathcal{M}* is defined as

$$\mathcal{M} \otimes_{\mathbb{C}_2} \mathbb{C}_2^{\text{Vect}},$$

where $\mathbb{C}_2^{\text{Vect}}$ is the lax commutative semiring of \mathbb{C} -vector spaces (not necessarily finitely dimensional).

Now $H \in \text{Obj}(\mathcal{M}^{\text{Vect}})$, we may consider a pseudomorphism of SLCMC’s over S (see above)

$$h : S^2 \rightarrow C(\mathcal{M}^{\text{Vect}}).$$

Here by $C(\mathcal{M}^{\text{Vect}})$ we mean the analogous construction as $C(\mathcal{M})$, but with the duals taken over \mathbb{C}^{Vect} , so $(\mathcal{M}^{\text{Vect}})^* =_{\text{def}} (\mathcal{M}^*)^{\text{Vect}}$. Then, an SLCMC $C(\mathcal{M}, h)$ is defined as follows. Sections over a point over $\sigma \in \text{Obj}(S)^2$ consist of a section M of $C(\mathcal{M})$ over σ and a 2-morphism

$$M \rightarrow h(\sigma).$$

Stacking, and the necessary verifications, are completed in the usual way.

Acknowledgement. We thank G. Moore for pointing out to us his lectures [26]. We would also like to thank N. A. Baas and J. Rognes for their helpful mathematical comments, and for detecting mistakes in the original version of this paper.

References

1. Ando, M.: Power operations in elliptic cohomology and representations of loop groups. *Trans. AMS* **352**, 5619–5666 (2000)
2. Baas, N.A., Dundas, B.I., Rognes, J.: Two-vector bundles and forms of elliptic cohomology. To appear in *Segal Proceedings*, Cambridge University Press
3. Borceux, F.: *Handbook of categorical algebra 1–2*, Encyclopedia of Mathematics and its Applications, Cambridge: Cambridge University Press, pp.50–52
4. Borchers, R.E.: Monstrous moonshine and monstrous Lie superalgebra. *Invent. Math.* **109**, 405–444 (1992)
5. d’Hoker, E.: String theory, In: *Quantum fields and strings: a course for mathematicians*, Vol. **2**, Providence RI: AMS and IAS, 1999, pp. 807–1012
6. Deligne, P., Freed, D.: Notes on Supersymmetry (following J. Bernstein) In: *Quantum fields and strings, a course for mathematicians*, Vol. **1**, Providence RI: AMS, 1999, pp. 41–98
7. Diaconescu, D.E.: Enhanced D -brane categories from string field theory. *JHEP* **0106**, 16 (2001)
8. Douglas, M.R.: D -branes, categories and $N = 1$ SUSY. *J. Math. Phys.* **42**, 2818–2843 (2001)
9. Fiore, T.: *Lax limits, lax adjoints and lax algebras: the categorical foundations of conformal field theory*. To appear
10. Frenkel, I.: Vertex algebras and algebraic curves. *Seminaire Bourbaki 1999–2000, Asterisque* **276**, 299–339 (2002)
11. Frenkel, I., Lepowsky, J., Meurman, A.: Vertex operator algebras and the monster. *Pure and applied Mathematics*, Vol. **134**, London–NewYork: Academic Press, 1999
12. Fuchs, J., Runkel, I., Schweigert, C.: TFT construction of RCFT correlators I: Partition functions. *Nucl. Phys. B* **646**, 353 (2002)

13. Fuchs, J., Runkel, I., Schweigert, C.: TFT construction of RCFT correlators II: Unoriented world sheets. *Nucl. Phys. B* **678**, 511 (2004)
14. Green, M.B., Schwartz, J.H., Witten, E.: *Superstring theory*. Vol. **1,2**, Cambridge: Cambridge University Press, 1988
15. Horava, P.: Equivariant Topological Sigma Models. *Nucl. Phys. B* **418**, 571–602 (1994)
16. Hu, P., Kriz, I.: Conformal field theory and elliptic cohomology. *Advances in Mathematics* **189**(2), 325–412 (2004)
17. Hu, P., Kriz, I., Voronov, A.A.: *On Kontsevich's Hochschild cohomology conjecture*. <http://arxiv.org/abs/amth.AT/0309369>, 2003
18. Huang, Y.Z., Kong, L.: Open-string vertex algebras, tensor categories and operads. *Comm. Math. Phys.* **250**, 433–471 (2004)
19. Kriz, I.: On spin and modularity in conformal field theory. *Ann. Sci. de ENS* **36**, 57–112 (2003)
20. Lawvere, W.F.: Functorial semantics of algebraic theories. *Proc. Nat. Acad. Sci. U.S.A.* **50**, 869–87 (1963)
21. Lazaroiu, C.I.: On the structure of open-closed topological field theory in two-dimensions. *Nucl. Phys. B* **603**, 497–530 (2001)
22. Lazaroiu, C.I.: Generalized complexes and string field theory. *JHEP* **06**, 52 (2001)
23. Lazaroiu, C.I.: Unitarity, D -brane dynamics and D -brane categories. *JHEP* **12**, 31 (2001)
24. Lewellen, D.: Sewing constraints for conformal field theories on surfaces with boundaries. *Nucl. Phys. B* **372**, 654 (1992)
25. Moore, G.: Some Comments on Branes, G -flux, and K -theory. *Int. J. Mod. Phys. A* **16**, 936–944 (2001)
26. Moore, G.: *Lectures on branes, K-theory and RR-charges*. <http://www.physics.rutgers.edu/~gmoore/day1/12.html>
27. Moore, G., Seiberg, N.: Classical and Quantum Conformal Field Theory. *Commun. Math. Phys.* **123**, 177–254 (1989)
28. Moore, G., Seiberg, N.: Taming the conformal ZOO. *Phys. Lett. B* **220**, 422–430 (1989)
29. Moore, G., Seiberg, N.: Lectures on RCFT. In: H.C. Lee (ed.), *Physics, Geometry and Topology*, RiverEdge, World Scientific, 1990, pp. 263–361
30. Ostrik, V.: Module categories, weak Hopf algebras and modular invariants. *Transform. Groups* **8**, 177 (2003)
31. Polchinski, J.: *String theory*. Vols. **1,2**, Cambridge: Cambridge Univ. Press, 1999
32. Pradisi, G., Sagnotti, A., Stanev, Y.A.: Planar duality in $SU(2)$ WZW models. *Phys. Lett. B* **354**, 279 (1995)
33. Pressley, A., Segal, G.: *Loop groups*. Oxford: Oxford University Press, 1986
34. Segal, G.: Elliptic cohomology. *Seminaire Bourbaki 1987/88, Asterisque* **161–162**, Exp. No. 695, (1988) 4, 187–201 (1989)
35. Segal, G.: *The definition of conformal field theory*. Preprint, 1987
36. Segal, G.: *ITP lectures*. <http://doug-pc.itp.ucsb.edu/online/geom99/>, 1999
37. Segal, G.: Categories and cohomology theories. *Topology* **13**, 293–312 (1974)
38. Stolz, S., Teichner, P.: What is an elliptic object?, In: U. Tillmann (ed.), *Proc. of 2002 Oxford Symp. in Honour of G.Segal*, Cambridge: Cambridge Univ. Press, 2004
39. Thomason, R.W.: Beware the phony multiplication on Quillen's $\mathcal{A}^{-1}A$. *Proc. AMS* **80**(4), 569–573 (1980)
40. Verlinde, E.: Fusion rules and modular transformations in $2D$ conformal field theory. *Nucl. Phys. B* **300**, 360–376 (1988)
41. Witten, E.: Overview of K -theory applied to strings. *Int. J. Mod. Phys. A* **16**, 693–706 (2001)

Communicated by M.R. Douglas