

Automorphic vector bundles on connected Shimura varieties*

J.S. Milne

Mathematics Department, University of Michigan, Ann Arbor, MI 48109, USA

Introduction	91
0. Review of terminology concerning Shimura varieties	93
1. The Taniyama group, the period torsor, and conjugates of Shimura varieties	95
2. The compact dual symmetric Hermitian space and its conjugates	101
3. The principal bundle $Y^0(G, X)$; statement of the first main theorem	105
4. Automorphic vector bundles	112
5. Conjugates of automorphic vector bundles	115
6. Proof of Theorem 3.10 for the symplectic group	118
7. Proof of Theorem 3.10 for connected Shimura varieties of abelian type	121
8. First completion of the proof of Theorem 3.10	123
9. Second completion of the proof of Theorem 3.10	124
Appendix: Pairs defining connected and nonconnected Shimura varieties	126
Bibliography	128

Introduction

A connected Shimura variety $S^0(G, X)$ is defined by a semisimple group G over \mathbb{Q} and a symmetric Hermitian domain X . For any automorphism τ of \mathbb{C} (as an abstract field), it is known that the conjugate $\tau S^0(G, X)$ of $S^0(G, X)$ has a canonical realization as a connected Shimura variety $S^0(G', X')$, and that the pair (G', X') defining the second Shimura variety can be constructed from the first pair by using the Taniyama group. In more down-to-earth terms, we can say that with an automorphic function f on X and a special point x of X , it is possible to associate a new automorphic function ${}^\tau f$ on a different domain X' ; the associated $f \mapsto {}^\tau f$ commutes with the Hecke operators, and $\tau(f(x)) = {}^\tau f(x')$ for an explicitly defined special point x' on X' . (A proof of this result for most connected Shimura varieties can be found in Milne and Shih (1982b) and for the remaining varieties in Milne (1983); see also Borovoi (1983/4). It is the analogue for connected Shimura varieties of a conjecture of Langlands (1979).)

* Parts of this work have been supported by the National Science Foundation; it was completed while the author was at the Mathematical Sciences Research Institute, Berkeley

The purpose of the present paper is to extend these statements to holomorphic automorphic forms. Such forms of a fixed type are sections of a vector bundle over a connected Shimura variety. The vector bundles for whose sections our results hold arise from equivariant vector bundles on the compact dual of X ; we call them *automorphic vector bundles* (the importance of these vector bundles has been emphasized by P. Deligne; the name was suggested to me by M. Harris). Holomorphic automorphic forms in the classical sense corresponding to automorphy factors for the full group arise as sections of automorphic vector bundles. We show in §5 that the conjugate $\tau\mathcal{V}$ of an automorphic vector bundle can be canonically realized as an automorphic vector bundle on $S^0(G', X')$. More precisely, we show (Theorem 5.2) that there is an automorphic vector bundle \mathcal{V}' on $S^0(G', X')$ and a canonical isomorphism $\tau\mathcal{V} \xrightarrow{\sim} \mathcal{V}'$ lifting the isomorphism $\tau S^0(G, X) \xrightarrow{\sim} S^0(G', X')$ and commuting with the Hecke operators; the data defining \mathcal{V}' is constructed from the that defining \mathcal{V} by using the Taniyama group and the period torsor. In more down-to-earth terms, we can say that with an automorphic form f and a special point x of X , it is possible to associate a new automorphic form ${}^{\tau}x f$ on the domain X' ; the association $f \mapsto {}^{\tau}x f$ commutes with the Hecke operators, and $\tau(f(x))$ can often be related to ${}^{\tau}x f(x')$ where x' is the same special point of X' as above.

In a sequel to this paper, these results will be used to obtain similar results for automorphic forms on nonconnected Shimura varieties. In this way, we shall obtain an analogue for automorphic vector bundles over Shimura varieties of Langlands's conjecture on the conjugates of Shimura varieties. In particular, this will allow us (without any assumptions on the underlying Shimura variety) to define *canonical* models of automorphic vector bundles, and to give a definitive definition of what it means for a holomorphic automorphic form to be rational over a number field.

The theorem on automorphic vector bundles is obtained as a rather direct consequence of a theorem (again a generalization of Langlands's conjecture) concerning a certain principal bundle $Y^0(G, X)$ over $S^0(G, X)$. This theorem is stated in §3 and proved in §6 and §7 for connected Shimura varieties of abelian type. In §8 and §9 we give two methods of extending the result to all connected Shimura varieties. The first, which is the shorter, uses a statement (Borovoi (1983/4), 3.21) for which no proof is currently available; the second makes use of an idea from Harris (1985). Automorphic vector bundles are defined in §4 and their conjugates are described in §5. The first two sections contain preliminary material on the Taniyama group, the period torsor, conjugates of connected Shimura varieties, and conjugates of the compact duals of Hermitian symmetric domains.

Lacking at this point are theorems describing how automorphisms of \mathbb{C} act on the Fourier-Jacobi series of automorphic forms (or even a general algebraic definition of such series) and on the Eisenstein series associated with cusp forms on boundary components. It is however possible to give precise conjectures, again in terms of the Taniyama group and the period torsor, and I hope to take up these questions in future papers.

A discussion of the relation of these results to those of other authors will be given in the sequel to this paper. Here we mention only that the results were suggested by those of Harris (1984, 1985), which in turn were suggested by questions of Shimura (1980) and Deligne.

0. Review of terminology concerning Shimura varieties

A reductive group is always assumed to be connected. When G is an algebraic group, G^{der} and G^{ad} are the associated derived and adjoint groups of G , and $Z(G)$ is the centre of G . The action of G on itself by inner automorphisms factors through a homomorphism $\text{ad}: G^{\text{ad}} \rightarrow \text{Aut}(G)$. The simply connected covering group of a semisimple group G is denoted by \tilde{G} .

When G is an algebraic group over \mathbb{R} , $G(\mathbb{R})^+$ is the identity component of $G(\mathbb{R})$ (for the real topology), and $G(\mathbb{R})_+$ is the inverse image of $G^{\text{ad}}(\mathbb{R})^+$ in $G(\mathbb{R})$. In the case that G is defined over \mathbb{Q} , we write $G(\mathbb{Q})^+$ for $G(\mathbb{Q}) \cap G(\mathbb{R})^+$ and $G(\mathbb{Q})_+$ for $G(\mathbb{Q}) \cap G(\mathbb{R})_+$. The symbol $\bar{}$ denotes closure in $G(\mathbb{A}^f)$ where $\mathbb{A}^f \stackrel{\text{df}}{=} \hat{\mathbb{Z}} \otimes \mathbb{Q}$ is the ring of finite adèles. We write \mathbb{A} for $\mathbb{R} \times \mathbb{A}^f$ and \mathbb{A}' for $\mathbb{C} \times \mathbb{A}^f$.

The real torus $\text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_m$ is denoted by \mathbf{S} ; thus $\mathbf{S}(\mathbb{R}) = \mathbb{C}^\times$ and $\mathbf{S}(\mathbb{C}) = \mathbb{C}^\times \times \mathbb{C}^\times$ (the projections onto the two factors correspond respectively to the identity automorphism of \mathbb{C} and to complex conjugation). Associated with any homomorphism $h: \mathbf{S} \rightarrow G$, there is a *weight map*

$$w_h: \mathbb{G}_m \rightarrow G, \quad r \mapsto h(r)^{-1}, \quad \text{all } r \in \mathbb{R}^\times \subset \mathbf{S}(\mathbb{R}),$$

and a cocharacter

$$\mu_h: \mathbb{G}_m \rightarrow G_{\mathbb{C}}, \quad z \mapsto h_{\mathbb{C}}(z, 1), \quad \text{all } z \in \mathbb{C}^\times.$$

By a *pair* (G, X) defining a connected Shimura variety we mean a semisimple algebraic group G over \mathbb{Q} and a $G^{\text{ad}}(\mathbb{R})^+$ -conjugacy class X of homomorphisms $\mathbf{S} \rightarrow G_{\mathbb{R}}^{\text{ad}}$ satisfying the conditions (2.1.1)–(2.1.3) of Deligne (1979). Then X has a canonical structure of a Hermitian symmetric domain, and we write x for a point of X when we are regarding it in this way and $h_x: \mathbf{S} \rightarrow G_{\mathbb{R}}^{\text{ad}}$ and $\mu_x: \mathbb{G}_m \rightarrow G_{\mathbb{C}}^{\text{ad}}$ for the homomorphism and cocharacter associated with x . The connected Shimura variety $S^0(G, X)$ is defined to be the projective system $(\Gamma \backslash X)_{\Gamma \in \Sigma(G)}$ (or its limit), where $\Sigma(G)$ is the set of net arithmetic subgroups of $G^{\text{ad}}(\mathbb{Q})$ containing the image of a congruence subgroup in $G(\mathbb{Q})$. Each complex manifold $\Gamma \backslash X$ has a unique structure as an algebraic variety, and the morphisms in the projective system are algebraic. It is sometimes also convenient to regard $S^0(G, X)$ as being the projective system of varieties $(\Gamma \backslash X)$ with Γ running over the set $\tilde{\Sigma}(G)$ of net congruence subgroups of $G(\mathbb{Q})$. We often write $S^0_{\Gamma}(G, X)$ (or $S^0_K(G, X)$ when $\Gamma = G(\mathbb{Q}) \cap K$, K compact an open in $G(\mathbb{A}^f)$) for the algebraic variety $\Gamma \backslash X$. When G is simply connected,

$$S^0(G, X)(\mathbb{C}) \stackrel{\text{df}}{=} \varprojlim_{\Gamma} S^0_{\Gamma}(G, X)(\mathbb{C}) = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}^f).$$

In the last term, $q \in G(\mathbb{Q})$ acts on $(x, g) \in X \times G(\mathbb{A}^f)$ according to the rule:

$$q(x, g) = (qx, qg).$$

For x in X , x_Γ denotes the image of x in $\Gamma \backslash X$, and $[x] = (x_\Gamma)_{\Gamma \in \Sigma(G)}$ denotes its image in $S^0(G, X)$.

The action of $G^{\text{ad}}(\mathbb{Q})^+$ on $S^0(G, X)$,

$$g: \Gamma \backslash X \rightarrow \Gamma' \backslash X, \quad x_\Gamma \mapsto (gx)_{\Gamma'}, \quad \Gamma' = \text{ad}(g)\Gamma$$

extends by continuity to the completion $G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ of $G^{\text{ad}}(\mathbb{Q})^+$ for the topology defined by the subgroups in $\Sigma(G)$. The map $G(\mathbb{Q})_+ \rightarrow G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ extends by continuity to the closure $G(\mathbb{Q})_+^-$ of $G(\mathbb{Q})_+$ in $G(\mathbb{A}^f)$, and $G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ is generated by $G^{\text{ad}}(\mathbb{Q})^+$ and the image $G(\mathbb{Q})_+^-/Z(\mathbb{Q})$ of $G(\mathbb{Q})_+^-$; more precisely,

$$G^{\text{ad}}(\mathbb{Q})^{+\wedge} = G(\mathbb{Q})_{+\star G(\mathbb{Q})}^- \cdot G^{\text{ad}}(\mathbb{Q})^+$$

(Deligne (1979), 2.1.6.2). In fact $G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ is also generated by $G^{\text{ad}}(\mathbb{Q})^+$ and the image of $\tilde{G}(\mathbb{A}^f)$. In the case that G is simply connected, $G(\mathbb{Q})_+^- = G(\mathbb{A}^f)$, and the actions of $G(\mathbb{A}^f)$ and $G^{\text{ad}}(\mathbb{Q})^+$ on $S^0(G, X)$ are given by

$$\begin{aligned} a[x, g] &= [x, ga^{-1}], & x \in X, \quad a, g \in G(\mathbb{A}^f); \\ q[x, g] &= [gx, \text{ad}(q)g], & x \in X, \quad q \in G^{\text{ad}}(\mathbb{Q})^+, \quad g \in G(\mathbb{A}^f). \end{aligned}$$

We write (g) for the automorphism of $S^0(G, X)$ defined by $g \in G^{\text{ad}}(\mathbb{Q})^{+\wedge}$.

By a *morphism* $f: (G, X) \rightarrow (G', X')$ of pairs defining connected Shimura varieties we mean a homomorphism $f: G \rightarrow G'$ of algebraic groups over \mathbb{Q} carrying the conjugacy class X into X' . The map $h \mapsto \text{ad}(f) \circ h: X \rightarrow X'$ automatically sends special points of X to special points of X' . Such an f defines a morphism $S^0(f): S^0(G, X) \rightarrow S^0(G', X')$ of connected Shimura varieties taking the action of $g \in G(\mathbb{A}^f)$ into that of $f(g) \in G'(\mathbb{A}^f)$. We say that f is an *embedding* if $f: G \rightarrow G'$ is injective. In this case $S^0(f)$ is a projective system of closed immersions (see Deligne (1971), 1.15).

By a *pair* (G_1, X_1) defining a Shimura variety we mean a reductive group G_1 over \mathbb{Q} and a $G_1(\mathbb{R})$ -conjugacy class X_1 of homomorphisms $\mathbf{S} \rightarrow G_{1\mathbb{R}}$ satisfying the conditions (2.1.1)–(2.1.3) of (Deligne (1979)). For such a pair (G_1, X_1) , a connected component X_1^+ of X_1 can be identified with a $G_1^{\text{ad}}(\mathbb{R})^+$ -conjugacy class of maps $\mathbf{S} \rightarrow G_{1\mathbb{R}}^{\text{ad}}$, and $(G, X) = (G_1^{\text{der}}, X_1^+)$ is a pair defining a connected Shimura variety; in this situation, we write $(G, X) = (G_1, X_1)^+$. The connected component of $S(G_1, X_1)$ containing the image of X_1^+ can be identified with $S^0(G, X)$. The action of $G_1(\mathbb{A}^f)$ on $S(G_1, X_1)$ factors through $G_1(\mathbb{A}^f)/Z_1(\mathbb{Q})^-$, where Z_1 is the centre of G_1 , and the stabilizer of $S^0(G, X)$ in $S(G_1, X_1)$ is $G_1(\mathbb{Q})_+^-/Z_1(\mathbb{Q})^-$ (here $-$ denotes closure in $G_1(\mathbb{A}^f)$). If we assume that $H^1(\mathbb{Q}, Z_1) = 0$, then $G^{\text{ad}}(\mathbb{Q}) = G_1(\mathbb{Q})/Z_1(\mathbb{Q}) \subset G_1(\mathbb{A}^f)/Z_1(\mathbb{Q})^-$, and the stabilizer of $S^0(G, X)$ in $G_1(\mathbb{A}^f)/Z_1(\mathbb{Q})^-$ is the closure of $G^{\text{ad}}(\mathbb{Q})^+$ in $G_1(\mathbb{A}^f)/Z_1(\mathbb{Q})^-$. This closure can be identified with $G^{\text{ad}}(\mathbb{Q})^{+\wedge}$, and its action on $S^0(G, X)$, when converted into a left action, agrees with that defined above [ibid. 2.1.16].

All vector spaces and vector bundles are of finite dimension. The category of representations of an algebraic group G on k -vector spaces is denoted by $\text{Rep}_k(G)$.

Motives are always meant in the sense of (absolute) Hodge cycles (see (Deligne and Milne (1982), § 6)).

We rarely distinguish a vector bundle from its associated locally free sheaf of sections. By a *variation of real Hodge structures* on a complex manifold X , we mean a local system of real vector spaces \mathbf{V} on X together with, at each point x of X , a real Hodge structure on the fibre \mathbf{V}_x at x ; these Hodge structures are required to vary continuously in x , and the associated Hodge filtrations F^* on the fibres of $\mathcal{O}_X \otimes \mathbf{V}$ are required to vary holomorphically and satisfy the axiom of transversality: $\forall F^p \subset \Omega_X^1 \otimes F^{p-1}$ (Deligne 1979), 1.1.7). To define a variation of *rational Hodge structures* replace “real” with “rational” in the preceding definition.

The algebraic closure of \mathbb{Q} in \mathbb{C} is denoted by $\bar{\mathbb{Q}}$. When necessary, we denote the inclusion $\bar{\mathbb{Q}} \hookrightarrow \mathbb{C}$ by ι . We often use $=$ to denote a canonical isomorphism. The equivalence class containing $*$ is often written $[*]$.

1. The Taniyama group, the period torsor, and conjugates of Shimura varieties

Recall (Milne and Shih 1982a) that the *Serre group* is a pair $(\mathfrak{S}, h_{\text{can}})$ consisting of a proalgebraic torus \mathfrak{S} over \mathbb{Q} and a homomorphism $h_{\text{can}}: \mathbf{S} \rightarrow \mathfrak{S}_{\mathbf{R}}$ whose weight is defined over \mathbb{Q} . The pair is universal in the following sense: for any torus T over \mathbb{Q} and homomorphism $h: \mathbf{S} \rightarrow T_{\mathbf{R}}$ whose cocharacter μ is defined over a *CM* field and whose weight w is defined over \mathbb{Q} , there is a unique \mathbb{Q} -rational homomorphism $\rho: \mathfrak{S} \rightarrow T$ such that $\rho_{\mathbf{R}} \circ h_{\text{can}} = h$.

The *Taniyama group* [ibid. § 3] is an extension

$$1 \rightarrow \mathfrak{S} \rightarrow \mathfrak{I} \xrightarrow{\pi} \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow 1$$

of pro-algebraic groups together with a continuous section $\text{sp}: \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q}) \rightarrow \mathfrak{I}(\mathbb{A}^f)$.

For any τ in $\text{Aut}(\mathbb{C})$, ${}^{\tau}\mathfrak{S} = \pi^{-1}(\tau|\bar{\mathbb{Q}})$ is a right \mathfrak{S} -torsor with a distinguished \mathbb{A}^f -point $\text{sp}(\tau)$.

Let $\text{CM}_{\bar{\mathbb{Q}}}$ be the category of motives over $\bar{\mathbb{Q}}$ generated by abelian varieties of *CM*-type over $\bar{\mathbb{Q}}$ and the Tate motive, and let $\text{CM}_{\mathbb{Q}}$ be the category of motives over \mathbb{Q} generated by the abelian varieties over \mathbb{Q} of potential *CM*-type, the Tate motive, and the Artin motives. The objects of these categories will be called *CM*-motives over $\bar{\mathbb{Q}}$ and \mathbb{Q} respectively. Both categories are Tannakian and have natural \mathbb{Q} -linear fibre functors sending a motive M to the Betti cohomology group of ιM , and it is known that \mathfrak{S} and \mathfrak{I} are the pro-algebraic groups associated with $\text{CM}_{\bar{\mathbb{Q}}}$ and $\text{CM}_{\mathbb{Q}}$ (see Deligne (1982b) for \mathfrak{I}). In particular, this means that with each \mathbb{Q} -linear representation (r, V) of \mathfrak{I} there is associated a *CM*-motive M over \mathbb{Q} , well-defined up to a unique isomorphism. The Betti cohomology group $H_B(\iota M) = V$, and the Hodge structure on $H_B(\iota M)$ is defined by $r \circ h_{\text{can}}$.

Let M be a CM motive over \mathbb{Q} . A tensor t of $H_B(\iota M)$ will be called a Hodge cycle if there is a Hodge cycle (t_{dR}, t_{et}) of $H_{dR}(M) \times H_{et}(M)$ on M relative ι (in the sense of Deligne (1982a), p. 28) such that t and (t_{dR}, t_{et}) have the same image as a tensor of $H_{dR}(\iota M) \times H_{et}(\iota M)$. For any $\tau \in \text{Gal}(\mathbb{Q}/\mathbb{Q})$, Deligne's theorem [ibid., 2.11] implies that there exists a Hodge cycle ${}^{\tau}t$ of $H_B(\tau M)$ corresponding to the tensor $(\tau t_{dR}, \tau t_{et})$ of $H_{dR}(\tau M) \times H_{et}(\tau M)$. For example, when t is the class of an algebraic cycle Z on an abelian variety A , ${}^{\tau}t$ is the class of τZ on τA .

The functor H_{dR} is a second fibre functor on $\mathbf{CM}_{\mathbb{Q}}$ with values in the category of vector spaces over \mathbb{Q} . Therefore $\mathcal{P} = \mathcal{H}om^{\otimes}(H_B, H_{dR})$ is a torsor for \mathfrak{I} (see Deligne and Milne (1982), 3.2), which we call the *period torsor*. The comparison isomorphisms $c(M): H_B(M) \otimes \mathbb{C} \xrightarrow{\sim} H_{dR}(M_{\mathbb{C}})$ define a canonical element c in $\mathcal{P}(\mathbb{C})$. The \mathbb{Q} -structures $H_B(M)$ and $H_{dR}(M)$ determine actions of $\text{Aut}(\mathbb{C})$ on $H_B(M) \otimes \mathbb{C}$ and $H_{dR}(M_{\mathbb{C}})$, and for $\tau \in \text{Aut}(\mathbb{C})$, we write

$$z_{\infty}(\tau) = c^{-1} \circ \tau(c) \in \mathfrak{I}(\mathbb{C}).$$

The map $z_{\infty}: \text{Aut}(\mathbb{C}) \rightarrow \mathfrak{I}(\mathbb{C})$ is a one cocycle:

$$z_{\infty}(\sigma\tau) = z_{\infty}(\sigma) \cdot \sigma z_{\infty}(\tau), \sigma, \tau \in \text{Aut}(\mathbb{C}).$$

Note that z_{∞} does not factor through the quotient $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ of $\text{Aut}(\mathbb{C})$.

Proposition 1.1. *The element $z_{\infty}(\tau)$ lies in ${}^{\tau}\mathfrak{S}(\mathbb{C})$.*

Proof. We have to show that $z_{\infty}(\tau)$ maps to τ in $\text{Gal}(\mathbb{Q}/\mathbb{Q})$. Recall that $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ is the group associated with the Tannakian category $\mathbf{Art}_{\mathbb{Q}}$ of Artin motives over \mathbb{Q} , and that the map $\pi: \mathfrak{I} \rightarrow \text{Gal}(\mathbb{Q}/\mathbb{Q})$ corresponds to the inclusion of $\mathbf{Art}_{\mathbb{Q}}$ into $\mathbf{CM}_{\mathbb{Q}}$. It therefore suffices to show that, for all Artin motives M , $z_{\infty}(\tau)(M): H_B(M_{\mathbb{C}}) \otimes \mathbb{C} \rightarrow H_B(M) \otimes \mathbb{C}$ is simply $\tau \otimes 1$.

Every Artin motive M is a direct factor of a motive of the form $h(X)$ with X a finite scheme over \mathbb{Q} (see Deligne and Milne (1982), p. 211), and so we may suppose that $M = h(X)$. Then $H_B(M) = \text{Hom}(X(\mathbb{Q}), \mathbb{Q})$ and M corresponds to the representation of $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ on $H_B(M)$ induced by its action on $X(\mathbb{Q})$. For any \mathbb{Q} -algebra R , $H_{dR}(M_R) = A \otimes R$, where $A = \Gamma(X, \mathcal{O}_X)$, and $A \otimes \mathbb{Q} = \text{Hom}(X(\mathbb{Q}), \mathbb{Q})$ with $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ acting on the second term through its action on $X(\mathbb{Q})$ and \mathbb{Q} . In summary:

- (a) $H_B(M) = \text{Hom}(X(\mathbb{Q}), \mathbb{Q})$ with $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ acting through its action on $X(\mathbb{Q})$;
- (b) $H_B(M) \otimes \mathbb{C} = \text{Hom}(X(\mathbb{Q}), \mathbb{C})$ with $\text{Aut}(\mathbb{C})$ acting through its action on \mathbb{C} ;
- (c) $H_{dR}(M_{\mathbb{C}}) = \text{Hom}(X(\mathbb{Q}), \mathbb{C})$ with $\text{Aut}(\mathbb{C})$ acting through its action on $X(\mathbb{Q})$ and \mathbb{C} . All actions are on the left. With these identifications, c becomes the identity map, and for $\lambda \in \text{Hom}(X(\mathbb{Q}), \mathbb{C}) = H_B(M) \otimes \mathbb{C}$,

$$\begin{aligned} z_{\infty}(\tau)(\lambda) &\stackrel{df}{=} (c^{-1} \circ \tau c)(\lambda) \stackrel{df}{=} (c^{-1} \circ \tau \circ c \circ \tau^{-1})(\lambda) \\ &\stackrel{(b)}{=} (c^{-1} \circ \tau)(\tau^{-1} \circ \lambda) \stackrel{(c)}{=} \tau \circ (\tau^{-1} \circ \lambda) \circ \tau^{-1} = \lambda \circ \tau^{-1} \stackrel{(a)}{=} (\tau \otimes 1)(\lambda), \end{aligned}$$

as required.

Remark 1.2. (a) We leave open the question of giving a description of \mathcal{P} and its canonical \mathbb{C} -valued point (equivalently of $z_\infty(\tau)$) in the spirit of Langlands's definition of the Taniyama group (in particular, a description that avoids mentioning CM motives). As Deligne pointed out to me, $\pi_*(\mathcal{P})$ is the $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -torsor $\text{Spec}(\bar{\mathbb{Q}})$.

(b) It is possible to give a slightly different interpretation of $z_\infty(\tau)$. For a CM -motive M over $\bar{\mathbb{Q}}$, write $H_\tau(M) = H_B(\tau \iota M)$. Then $H_\tau(M)$ is a fibre functor on $\mathbf{CM}_{\bar{\mathbb{Q}}}$, and for any \mathbb{Q} -algebra R , ${}^t\mathfrak{S}(R) = \mathcal{H}om^\otimes(H_{id} \otimes R, H_\tau \otimes R)$ (see Deligne and Milne (1982), 6.23). An element of ${}^t\mathfrak{S}(\mathbb{C})$ is therefore a \mathbb{C} -linear functorial isomorphism $H_B(\iota M) \otimes \mathbb{C} \xrightarrow{\sim} H_B(\tau \iota M) \otimes \mathbb{C}$, compatible with tensor products, and such that $t \otimes 1$ corresponds to ${}^t t \otimes 1$ for all Hodge cycles t of $H_B(\iota M)$. Consider the maps

$$\begin{aligned} H_B(\iota M) \otimes \mathbb{C} &\xrightarrow{1 \otimes \tau^{-1}} H_B(\iota M) \otimes \mathbb{C} \xrightarrow{c(\iota M)} H_{dR}(M) \otimes_{\mathbb{Q}, \iota} \mathbb{C} \\ &\xrightarrow{1 \otimes \tau} H_{dR}(M) \otimes_{\mathbb{Q}, \tau \iota} \mathbb{C} \xrightarrow{c(\tau \iota M)^{-1}} H_B(\tau \iota M) \otimes \mathbb{C}. \end{aligned}$$

Obviously the composite is \mathbb{C} -linear, and for any Hodge cycle t of $H_B(\iota M)$,

$$t \otimes 1 \leftrightarrow t \otimes 1 \leftrightarrow t_{dR} \otimes_{\mathbb{Q}, \iota} 1 \leftrightarrow t_{dR} \otimes_{\mathbb{Q}, \tau \iota} 1 \leftrightarrow {}^t t \otimes 1.$$

Since the maps are functorial and compatible with tensor products, they define an element of ${}^t\mathfrak{S}(\mathbb{C})$, which is clearly $z_\infty(\tau)$.

Let (G, X) be a pair defining a connected Shimura variety, and let x be a special point of X . By definition, this means that there is a (maximal) \mathbb{Q} -rational torus T in G such that h_x factors through $(T/Z)(\mathbb{R})$, $Z = Z(G)$. From the universal property of \mathfrak{S} , we know that there is a unique \mathbb{Q} -rational homomorphism $\rho_x: \mathfrak{S} \rightarrow T/Z$ such that $h_x = (\rho_x)_{\mathbb{R}} \circ h_{\text{can}}$. The map $\rho_x: \mathfrak{S} \rightarrow G^{\text{ad}}$ defines an action of \mathfrak{S} on G , and the \mathfrak{S} -torsor ${}^t\mathfrak{S}$ can be used to twist G (or any other covering group of G^{ad}). Thus we obtain an algebraic group ${}^{\tau, x}G = {}^t\mathfrak{S} \times_{\mathfrak{S}, \rho_x} G$ over \mathbb{Q} such that (as a $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$ -set)

$${}^{\tau, x}G(\bar{\mathbb{Q}}) = \{s \cdot g \mid s \in {}^t\mathfrak{S}(\bar{\mathbb{Q}}), g \in G(\bar{\mathbb{Q}})\} / \mathfrak{S}(\bar{\mathbb{Q}})$$

where $s_1 \in \mathfrak{S}(\bar{\mathbb{Q}})$ acts according to the rule

$$(s \cdot g)s_1 = s s_1 \cdot \text{ad}(\rho_x(s_1^{-1}))g.$$

Then ${}^{\tau, x}G$ is a semisimple group having ${}^t\mathfrak{S} \times_{\mathfrak{S}, \rho_x} T = T$ as a subtorus.

Let $q \in {}^{\tau, x}G^{\text{ad}}(\bar{\mathbb{Q}})$, and let $s \in \rho_{x*}({}^t\mathfrak{S})(\bar{\mathbb{Q}})$; then q can be written $[s \cdot q_s]$ for a unique element $q_s \in G^{\text{ad}}(\bar{\mathbb{Q}})$, and $s \mapsto s q_s$ defines a \mathbb{Q} -rational automorphism (q) of $\rho_{x*}({}^t\mathfrak{S})$ (as a G^{ad} -torsor). In this way, ${}^{\tau, x}G$ can be identified with the group of automorphisms of the G -torsor $\rho_{x*}({}^t\mathfrak{S})$ (acting on the right).

The point $\text{sp}(\tau)$ in ${}^t\mathfrak{S}(\mathbb{A}^f)$ defines a canonical continuous isomorphism

$$g \mapsto {}^{\tau, x}g = [\text{sp}(\tau) \cdot g]: \tilde{G}(\mathbb{A}^f) \rightarrow {}^{\tau, x}\tilde{G}(\mathbb{A}^f).$$

In (Milne and Shih (1982b), 8.2) it is shown how to construct a canonical isomorphism

$$g \mapsto {}^{\tau,x}g: G^{\text{ad}}(\mathbb{Q})^{+\wedge} \rightarrow {}^{\tau,x}G^{\text{ad}}(\mathbb{Q})^{+\wedge}$$

compatible with the preceding isomorphism (cf. (3.7) below). Define ${}^{\tau}h$ to be the homomorphism $\mathbf{S} \rightarrow {}^{\tau,x}G_{\mathbb{R}}^{\text{ad}}$ associated with the cocharacter $\tau\mu_x$ of $T/Z \subset {}^{\tau,x}G^{\text{ad}}$. If we let ${}^{\tau,x}X$ denote the ${}^{\tau,x}G^{\text{ad}}(\mathbb{R})^+$ -conjugacy class of maps $\mathbf{S} \rightarrow {}^{\tau,x}G_{\mathbb{R}}^{\text{ad}}$ containing ${}^{\tau}h$, then the pair $({}^{\tau,x}G, {}^{\tau,x}X)$ again defines a connected Shimura variety. We write ${}^{\tau}x$ for ${}^{\tau}h$ regarded as a point of ${}^{\tau,x}X$.

Proposition 1.3. *For any special points x and x' of X , there is a canonical isomorphism*

$$\varphi^0(\tau; x', x): S^0({}^{\tau,x}G, {}^{\tau,x}X) \rightarrow S^0({}^{\tau,x'}G, {}^{\tau,x'}X)$$

such that $\varphi^0(\tau; x', x) \circ ({}^{\tau,x}g) = ({}^{\tau,x'}g) \circ \varphi^0(\tau; x', x)$ for all $g \in G^{\text{ad}}(\mathbb{Q})^{+\wedge}$.

Proof. Because $\tilde{G}^{\text{ad}}(\mathbb{Q})^{+\wedge} \rightarrow G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ is surjective, and $S^0(G, X)$ is the quotient of $S^0(\tilde{G}, X)$ by the kernel of this map, it suffices to prove the proposition for (\tilde{G}, X) . Thus we may assume that G is simply connected.

Lemma 1.4. *Let (G, X) and (G', X') be pairs defining connected Shimura varieties. Let f and f' be isomorphisms*

$$f: G \xrightarrow{\sim} G', \quad f': G^{\text{ad}}(\mathbb{Q})^{+\wedge} \xrightarrow{\sim} G'^{\text{ad}}(\mathbb{Q})^{+\wedge},$$

and let $\gamma \in G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ be such that, when f is extended by continuity to an isomorphism $\hat{f}: G^{\text{ad}}(\mathbb{Q})^{+\wedge} \xrightarrow{\sim} G'^{\text{ad}}(\mathbb{Q})^{+\wedge}$, then $\hat{f} \circ \text{ad}(\gamma) = f'$. Under these conditions

$$\varphi = S^0(f_1) \circ (\gamma): S^0(G, X) \rightarrow S^0(G', X')$$

is an isomorphism such that $\varphi \circ (a) = (f'(a)) \circ \varphi$ for all $a \in G^{\text{ad}}(\mathbb{Q})^{+\wedge}$; moreover, if f is replaced with $f \circ \text{ad}(q)$, $q \in G^{\text{ad}}(\mathbb{Q})^+$, and γ with $q^{-1}\gamma$, then φ is unchanged.

Proof. Straightforward.

Thus we must find a pair (f, γ) , well-defined up to replacement by $(f \circ \text{ad}(q), q^{-1}\gamma)$ with $q \in {}^{\tau,x}G^{\text{ad}}(\mathbb{Q})^+$, such that

- (i) f is an isomorphism ${}^{\tau,x}G \rightarrow {}^{\tau,x'}G$ sending ${}^{\tau,x}X$ into ${}^{\tau,x'}X$;
- (ii) $\gamma \in {}^{\tau,x}G^{\text{ad}}(\mathbb{Q})^{+\wedge}$ satisfies $f(\text{ad}(\gamma)({}^{\tau,x}a)) = {}^{\tau,x'}a$ for all $a \in G^{\text{ad}}(\mathbb{Q})^{+\wedge}$.

Let c be the class of $\tau\mathfrak{S}$ in $H^1(\mathbb{Q}, \mathfrak{S})$. The existence of the section sp shows that c maps to zero in $H^1(\mathbb{Q}_l, \mathfrak{S})$ for all primes l . Consider $\rho_x(c)$ and $\rho_{x'}(c) \in H^1(\mathbb{Q}, G^{\text{ad}})$. These elements have the same image in $H^1(\mathbb{R}, G^{\text{ad}})$; this can be proved by a direct calculation, which is carried out in (Milne and Shih (1982b), pp. 315–316), or by noting that (Deligne (1979), 1.2.2) implies that ${}^{\tau,x}G_{\mathbb{R}}^{\text{ad}} \approx {}^{\tau,x'}G_{\mathbb{R}}^{\text{ad}}$. Since the Hasse principle holds for G^{ad} (see for example (Milne (1986), I.9.9), it follows that $\rho_x(c)$ and $\rho_{x'}(c)$ are equal.

Choose a pair (G_1, X_1) as in (A.4) of the appendix. The constructions reviewed above for (G, X) have analogues for (G_1, X_1) . We write x_1 and x'_1 for x and x' regarded as points of X_1 when it is necessary to make this distinction. They are special, and (A.4b) implies they give rise to homomorphisms $\rho_{x_1}, \rho_{x'_1} : \mathfrak{S} \rightarrow G_1$ lifting ρ_x and $\rho_{x'}$. Because of (A.4c), the map $H^1(\mathbb{Q}, G_1) \rightarrow H^1(\mathbb{Q}, G_1^{\text{ad}})$ is injective, and so $\rho_{x_1*}({}^t\mathfrak{S})$ and $\rho_{x'_1*}({}^t\mathfrak{S})$ are isomorphic G_1 -torsors. The choice of an isomorphism $t: \rho_{x_1*}({}^t\mathfrak{S}) \xrightarrow{\sim} \rho_{x'_1*}({}^t\mathfrak{S})$ determines an isomorphism $f_1: {}^{\tau, x}G_1 \xrightarrow{\sim} {}^{\tau, x'}G_1$, and it is known (Langlands (1979), p. 232) that f_1 maps ${}^{\tau, x}X_1$ into ${}^{\tau, x'}X_1$. From the discussion preceding the statement of the proposition, we know that t is determined up to replacement by an element $t \circ (q)$, $q \in {}^{\tau, x}G_1(\mathbb{Q})$. Since ${}^{\tau, x}G_1(\mathbb{Q})$ is dense in ${}^{\tau, x}G_1(\mathbb{R})$, we can modify t so that f_1 maps the component ${}^{\tau, x}X$ of ${}^{\tau, x}X_1$ into the component ${}^{\tau, x'}X$ of ${}^{\tau, x'}X_1$. Then f_1 restricts to a morphism $f: ({}^{\tau, x}G, {}^{\tau, x}X) \rightarrow ({}^{\tau, x'}G, {}^{\tau, x'}X)$, and when t is replaced by $t \circ (q)$, $q \in {}^{\tau, x}G_1(\mathbb{Q})_+$, then f is replaced by $f \circ \text{ad}(\bar{q})$ where \bar{q} is the image of q in ${}^{\tau, x}G^{\text{ad}}(\mathbb{Q})^+$.

If s and s' denote the images of $\text{sp}(\tau)$ in $\rho_{x_1*}({}^t\mathfrak{S})(\mathbb{A}^f)$ and $\rho_{x'_1*}({}^t\mathfrak{S})(\mathbb{A}^f)$ respectively, then there exists an element $\gamma_1 \in {}^{\tau, x}G_1(\mathbb{A}^f)$ such that $t(s\gamma_1) = s'$. Note that γ_1 is uniquely determined by the choice of t , and that if t is replaced by $t \circ (q)$, $q \in {}^{\tau, x}G_1(\mathbb{Q})_+$, then γ_1 is replaced by $q^{-1}\gamma_1$. Note also that $f_1 \circ \text{ad}(\gamma_1)({}^{\tau, x}a) = {}^{\tau, x'}a$ for all $a \in G_1(\mathbb{A}^f)$. We shall show that the image γ of γ_1 in ${}^{\tau, x}G_1(\mathbb{A}^f)/Z_1(\mathbb{Q})^-$ lies in the subgroup ${}^{\tau, x}G^{\text{ad}}(\mathbb{Q})^{+ \wedge}$ (cf. §0). Since the maps $a \mapsto {}^{\tau, x}a$ and $a \mapsto {}^{\tau, x'}a$ on $G_1(\mathbb{A}^f)$ induce the maps of the same name on the subquotient $G^{\text{ad}}(\mathbb{Q})^{+ \wedge}$ of $G_1(\mathbb{A}^f)$ (Milne and Shih (1982b), 8.2), the lemma will then show that the pair (f, γ) defines a map $\varphi^0(\tau; x', x)$ having the desired properties. It is independent of the choice of t , and the usual argument [ibid. p. 339–340] shows that it is independent of the choice of (G_1, X_1) .

Write G'_1 for ${}^{\tau, x}G_1$. We shall show that γ_1 lies in the closure of $G'_1(\mathbb{Q})^{+ -}$ of $G'_1(\mathbb{Q})^+$ in $G'_1(\mathbb{A}^f)$. Recall (Deligne (1979), 2.5.1) that $G'_1(\mathbb{Q})^{+ -}$ is the fibre over 1 of $G'_1(\mathbb{A}^f) \rightarrow \pi_0(G'_1(\mathbb{Q}) \backslash G'_1(\mathbb{A}))$, and (Deligne (1971), 2.4) that

$\pi_0(G'_1(\mathbb{Q}) \backslash G'_1(\mathbb{A})) \xrightarrow{\sim} \pi_0(H_1(\mathbb{Q}) \backslash H_1(\mathbb{A}))$ where $H_1 = G'_1/G_1^{\text{der}} = G_1/G_1^{\text{der}}$. The maps ρ_{x_1} and $\rho_{x'_1}$ become equal when composed with $\nu: G_1 \rightarrow H_1$, and so, when the above constructions are carried out with G_1 replaced with H_1 , one sees immediately that the image of γ_1 in $H_1(\mathbb{A}^f)$ lies in $H_1(\mathbb{Q})$. Therefore γ lies in $G_1(\mathbb{Q})^{+ -}$.

Remark 1.5. In §3 we shall need to use a slight strengthening of some of the above arguments. Let $H_1(\mathbb{R})_+$ be the image of $Z_1(\mathbb{R})$ in $H_1(\mathbb{R})$ and let $H_1(\mathbb{Q})_+ = H_1(\mathbb{Q}) \cap H_1(\mathbb{R})_+$. Write ρ for $\nu \circ \rho_{x_1} = \nu \circ \rho_{x'_1}$. An isomorphism

$t: \rho_{x_1*}({}^t\mathfrak{S}) \xrightarrow{\sim} \rho_{x'_1*}({}^t\mathfrak{S})$ induces an automorphism of $\rho_*({}^t\mathfrak{S})$, and hence defines an element $q \in H_1(\mathbb{Q})$. If t is replaced by $t \circ (g)$, $g \in {}^{\tau, x}G_1(\mathbb{Q})_+$, then q is replaced by $q \cdot \nu(g)$. Since $\nu(G_1(\mathbb{R})_+) \subset \nu(Z_1(\mathbb{R}))$, we see that we obtain an element $q_1 \in H_1(\mathbb{Q})/H_1(\mathbb{Q})_+$ which is independent of the choice of t . Probably it is possible to show in general that this is 1 by refining the proof of the “Second Lemma

of Comparison” in Langlands (1979), p. 232, but we shall use a different argument to prove this in the cases of immediate interest to us.

The construction in (A.2) leads to a pair (G_1, X_1) such that the weight is in fact zero. Let G_0 be the subgroup of G_1 constructed in (A.5), and let $H_0 = G_0/G$. Write x_0 and x'_0 for x and x' regarded as homomorphisms $\mathbb{S} \rightarrow G_{0,\mathbb{R}}$. Then $\rho_{x_0,*}({}^r\mathfrak{S})$ and $\rho_{x'_0,*}({}^r\mathfrak{S})$ are isomorphic G_0 -torsors (see the argument in (Milne and Shih (1982b), pp. 315–316)). This shows that there is an element $q_0 \in H_0(\mathbb{Q})/H_0(\mathbb{Q})_+$ mapping to $q_1 \in H_1(\mathbb{Q})/H_1(\mathbb{Q})_+$. But $H_{0\mathbb{R}/w}(\mathbb{G}_m)$ is anisotropic, and is therefore connected. Since $w(\mathbb{G}_m) = 1$, $H_0(\mathbb{R})$ is itself connected, and so $v(Z_0(\mathbb{R})) = H_0(\mathbb{R})$. Therefore that $q_0 = 1$, as required.

Theorem 1.6. *For each $\tau \in \text{Aut}(\mathbb{C})$, there is a unique isomorphism*

$$\varphi_{\tau,x}^0: \tau S^0(G, X) \rightarrow S^0({}^{\tau,x}G, {}^{\tau,x}X)$$

such that

- (i) *the point $\tau[x]$ is mapped to $[{}^{\tau,x}]$;*
- (ii) *$\varphi_{\tau,x}^0 \circ \tau(g) = ({}^{\tau,x}g) \circ \varphi_{\tau,x}^0$ for all $g \in G^{\text{ad}}(\mathbb{Q})^+ \wedge$.*

Moreover, if x' is a second special point of X , then

$$\varphi^0(\tau; x', x) \circ \varphi_{\tau,x}^0 = \varphi_{\tau,x'}^0.$$

Proof. See Milne and Shih (1982b) for Shimura varieties of abelian type and Milne (1983) for the general case.

Remark 1.7. Theorem 1.6 has a down-to-earth interpretation. Let (G, X) be a pair defining a connected Shimura variety, and let f be an automorphic function on X relative to a congruence subgroup Γ of $G(\mathbb{Q})$. We can regard f as an algebraic function on $\Gamma \backslash X$. By definition, $\Gamma = K \cap G(\mathbb{Q})$ for some compact open subgroup K in $G(\mathbb{A}^f)$, and we define the congruence subgroup ${}^{\tau,x}\Gamma$ of ${}^{\tau,x}G(\mathbb{Q})$ to be ${}^{\tau,x}K \cap {}^{\tau,x}G(\mathbb{Q})$ where ${}^{\tau,x}K$ is the image of K under the isomorphism $g \mapsto {}^{\tau,x}g: G(\mathbb{A}^f) \xrightarrow{\sim} {}^{\tau,x}G(\mathbb{A}^f)$. Let τ be an automorphism of \mathbb{C} , and let x be a special point of X . Then the theorem defines an isomorphism

$$(\varphi_{\tau,x}^0)_\Gamma: \tau(\Gamma \backslash X) \rightarrow {}^{\tau,x}\Gamma \backslash {}^{\tau,x}X.$$

It therefore associates with f an automorphic function ${}^{\tau,x}f = \tau f \circ (\varphi_{\tau,x}^0)_\Gamma^{-1}$ on the Hermitian symmetric domain ${}^{\tau,x}X$ relative to ${}^{\tau,x}\Gamma$ such that

$${}^{\tau,x}f([{}^{\tau,x}]) = \tau(f([x])) \text{ (for the chosen special point } x).$$

Moreover, ${}^{\tau,x}(f \circ (g)) = {}^{\tau,x}f \circ ({}^{\tau,x}g)$ for all $g \in G^{\text{ad}}(\mathbb{Q})^+ \wedge$.

Let G be a reductive algebraic group over \mathbb{Q} , and let ρ be a homomorphism $\mathfrak{S} \rightarrow G$. It is possible to give a motivic description of ${}^{\tau,\rho}G \stackrel{df}{=} {}^{\tau}\mathfrak{S} \times_{\mathfrak{S},\rho} G$. Choose a faithful \mathbb{Q} -linear representation $r: G \hookrightarrow GL(V)$ of G . There exists a family of tensors $(t_\alpha)_{\alpha \in A}$ of V (that is, elements of vector spaces of the form $T_j^i V = V^{\otimes i} \otimes (V^\vee)^{\otimes j}$) such that G is the subgroup of $GL(V)$ fixing the t_α . If M is the CM-motive over \mathbb{Q} corresponding to the representation $r \circ \rho$, then $H_B(M) = V$

and the t_α are Hodge cycles on M . Write ${}^{\tau,\rho}V = {}^\tau\mathfrak{S} \times_{\mathfrak{S},\rho} V$, so that (as $\text{Gal}(\mathbb{Q}/\mathbb{Q})$ -modules)

$${}^{\tau,\rho}V \otimes \mathbb{Q} = \{s \cdot v \mid s \in {}^\tau\mathfrak{S}(\mathbb{Q}), g \in V \otimes \mathbb{Q}_1\} / \mathfrak{S}(\mathbb{Q})$$

where $s_1 \in \mathfrak{S}(\mathbb{Q})$ acts according to the rule

$$(s \cdot v)s_1 = s s_1 \cdot \rho(s_1^{-1})v.$$

Proposition 1.8. *There is a canonical isomorphism ${}^{\tau,\rho}V \approx H_B(\tau M)$, and ${}^{\tau,\rho}G$ is the subgroup of $GL(H_B(\tau M))$ fixing the Hodge cycles ${}^\tau t_\alpha$ on τM .*

Proof. Let $\tilde{\mathfrak{S}}$ be the image of \mathfrak{S} in $GL(H_B(M))$, and let ${}^\tau\tilde{\mathfrak{S}}$ be the $\tilde{\mathfrak{S}}$ -torsor defined by ${}^\tau\mathfrak{S}$. For any \mathbb{Q} -algebra R , the R -valued points of ${}^\tau\tilde{\mathfrak{S}}$ can be identified with the set of R -linear maps $H_B(M) \otimes R \rightarrow H_B(\tau M) \otimes R$ taking t to ${}^\tau t$ for all Hodge cycles t on M (see Deligne and Milne (1982), especially (6.23 a)). Clearly ${}^{\tau,\rho}V = {}^\tau\tilde{\mathfrak{S}} \times_{\mathfrak{S},\rho} V$, and the map $[s \cdot v] \rightarrow s(v)$ identifies this with $H_B(\tau M)$. Choose a point $s_0 \in {}^\tau\tilde{\mathfrak{S}}(\mathbb{Q})$, and consider the map

$$\text{Aut}(H_B(\tau M) \otimes \mathbb{Q}, ({}^\tau t_\alpha)_{\alpha \in A}) \rightarrow {}^{\tau,\rho}G(\mathbb{Q}), \quad a \mapsto [s_0 \cdot \text{ad}(s_0^{-1})a].$$

One checks immediately that this is independent of the choice of s_0 , that it defines an isomorphism, and that it commutes with the action of $\text{Gal}(\mathbb{Q}/\mathbb{Q})$. It therefore defines an isomorphism of algebraic groups over \mathbb{Q} .

2. The compact dual symmetric Hermitian space and its conjugates

We first review Deligne’s interpretation (Deligne 1979) of a Hermitian symmetric domain X and its compact dual \check{X} as parameter spaces for Hodge structures and filtrations. Then we prove analogues of (1.3) and (1.6) for \check{X} .

Let V be a vector space over a field k of characteristic zero. A homomorphism $\mu: \mathbb{G}_m \rightarrow GL(V)$ defines a decomposition

$$V = \bigoplus V^i, \quad V^i = \{v \in V \mid \mu(z)v = z^i v, \text{ all } z \in k^\times\}$$

and a decreasing filtration of V

$$\dots F^p V \supset F^{p+1} V \dots, \quad F^p V = \bigoplus_{i \geq p} V^i.$$

Let G be a reductive group over k . A homomorphism $\mu: \mathbb{G}_m \rightarrow G$ defines a filtration F^* on V for each representation (r, V) of G , namely, that corresponding to $r \circ \mu$. These filtrations are compatible with the formation of tensor products and duals. Conversely, any functor $(r, V) \mapsto (F^*, V)$ from representations of G to filtrations compatible with tensor products and duals arises from a homomorphism $\mu: \mathbb{G}_m \rightarrow G$. We call such a functor a *filtration* of $\text{Rep}_k(G)$, and we write $\text{Filt}(\mu)$ for the filtration defined by μ .

Proposition 2.1. *Let G be a reductive group over a field k of characteristic zero, and let μ be a cocharacter of G . From the adjoint action of G on $\mathfrak{g} = \text{Lie}(G)$, we acquire a filtration F^* of \mathfrak{g} .*

(a) The subalgebra $F^0 \mathfrak{g}$ is the Lie algebra of a parabolic subgroup $F^0 G$ of G ; moreover, $F^0 G$ is the subgroup of G respecting the filtration (defined by μ) on each representation of G .

(b) The subalgebra $F^1 \mathfrak{g}$ is the Lie algebra of the unipotent radical $F^1 G$ of $F^0 G$; moreover $F^1 G$ is the subgroup of $F^0 G$ that acts trivially on the graded module $\otimes (F^p V/F^{p+1} V)$ associated with each representation (r, V) of G .

(c) The centralizer $Z(\mu)$ of μ is a Levi subgroup of $F^0 G$; in particular, $Z(\mu) \xrightarrow{\sim} F^0 G/F^1 G$, and the composite $\bar{\mu}$ of μ with $F^0 G \rightarrow F^0 G/F^1 G$ is central. Two cocharacters of G define the same filtration of G if and only if they define the same group $F^0 G$ and induce the same map $\mathbf{G}_m \rightarrow F^0 G/F^1 G$.

Proof. See (Saavedra (1972), especially IV.2.2.5).

Let G be a reductive group over \mathbf{C} , and let $\mu_0: \mathbf{G}_m \rightarrow G$ be a cocharacter of G . We let \check{X} be the set of filtrations of $\mathbf{Rep}_{\mathbf{C}}(G)$ that are $G(\mathbf{C})$ -conjugate to $\text{Filt}(\mu_0)$.

Proposition 2.2. *The action*

$$G(\mathbf{C}) \times \check{X} \rightarrow \check{X}, \quad (g, \mu) \mapsto \text{Filt}(\text{ad}(g) \circ \mu)$$

defines a bijection $G(\mathbf{C})/P_0(\mathbf{C}) \xrightarrow{\sim} \check{X}$, where P_0 is the parabolic subgroup $F^0 G$ of G .

Proof. We have to show that $P_0(\mathbf{C})$ is the subgroup fixing $\text{Filt}(\mu_0)$ under the above action, but the filtration on a vector space V defined by $\text{ad}(g) \circ \mu_0$ is obtained from the filtration defined by μ_0 by applying g , and so this follows from (2.1 a).

Remark 2.3. (a) The bijection in (2.2) endows \check{X} with a complex structure. In fact, because P_0 is parabolic, \check{X} has the structure of a smooth projective variety over \mathbf{C} .

(b) According to (2.1 c), the points of \check{X} can be identified with the set of equivalence classes $[P, \mu]$, where P is a parabolic subgroup of G and μ is a cocharacter μ of P such that (P, μ) is conjugate under $G(\mathbf{C})$ to (P_0, μ_0) ; the classes $[P, \mu]$ and $[P', \mu']$ are equal if and only if $P=P'$ and μ and μ' define the same cocharacter of $P/R_u P$.

(c) For any faithful representation (r, V) of G it is obvious that \check{X} can be identified with the set of filtrations of V conjugate to that defined by μ_0 . Slightly less obviously, \check{X} can also be identified with the set of filtrations of $\text{Lie}(G)$ conjugate to that defined by $\text{ad} \circ \mu_0$.

By a global tensor of a sheaf \mathcal{V} of \mathcal{O}_S -modules, we mean an element of $\Gamma(S, T_j^i \mathcal{V})$ for some i and j , where $T_j^i \mathcal{V} = \mathcal{V}^{\otimes i} \otimes \check{\mathcal{V}}^{\otimes j}$.

Proposition 2.4. *Let μ_0 be a cocharacter of a reductive group G over \mathbf{C} , and let $\check{X} = G(\mathbf{C})/P_0(\mathbf{C})$ be the corresponding space of filtrations; let $r: G \rightarrow \text{GL}(V)$ be a faithful representation of G , and let \mathcal{V} be the constant vector bundle $\check{X} \times V$ on \check{X} .*

(a) *There is a unique filtration of \mathcal{V} by algebraic subbundles such that the filtration on each fibre $\mathcal{V}_{\{P, \mu\}}$ is that defined by μ .*

(b) *Let (t_α) be a family of tensors of V such that G is the subgroup of $GL(V)$ fixing the t_α . Let F^* be a filtration (by algebraic subbundles) of the constant vector bundle $\mathcal{V}' = S \times V$ on a smooth complex variety S , and let t'_s be the global tensor $(1, t_\alpha)$ of \mathcal{V}' . If for each s in S , there is an isomorphism $\mathcal{V}'_s \xrightarrow{\sim} V$ of filtered vector spaces mapping each $t'_{\alpha, s}$ to t_α , then there is a unique morphism $\gamma: S \rightarrow \check{X}$ such that when we identify $\gamma^* \mathcal{V}$ with $S \times V = \mathcal{V}'$ in the obvious way, the filtrations on $\gamma^* \mathcal{V}$ and \mathcal{V}' agree.*

Proof. (a) Let $\mathfrak{F}(V)$ be the flag variety of filtrations on V conjugate under $GL(V)$ to that defined by μ_0 . Then the map sending $\text{Filt}(\mu)$ to the filtration on V defined by μ is a closed immersion of \check{X} into $\mathfrak{F}(V)$, and the pullback of the universal bundle on $\mathfrak{F}(V)$ is \mathcal{V} . Clearly the filtration on the universal bundle induces the correct filtration on \mathcal{V} .

(b) For $y \in \mathfrak{F}(V)$, let F_y be the corresponding filtration of V . Then y is in the image of \check{X} in $\mathfrak{F}(V)$ if and only if there is an isomorphism of filtered vector spaces $(V, F_0) \xrightarrow{\sim} (V, F_y)$ fixing each t_α because such isomorphisms are defined by elements of $G(\mathbb{C})$. Therefore the map $S \rightarrow \mathfrak{F}(V)$ defined by \mathcal{V} and the universal property of $\mathfrak{F}(V)$ factors through \check{X} , and the resulting map $S \rightarrow \check{X}$ has the correct properties.

Now let (G, X) be a pair defining a connected Shimura variety.

Proposition 2.5. *Let $r: G_{\mathbb{R}}^{\text{ad}} \hookrightarrow GL(V)$ be a faithful real representation of G^{ad} , and let \mathbf{V} be the constant sheaf on X defined by V .*

(b) *For each x in X , $r \circ h_x$ is a Hodge structure on the stalk $\mathbf{V}_x (= V)$ of \mathbf{V} ; \mathbf{V} , together with these Hodge structures, is a variation of real Hodge structures on X .*

(b) *There exists a bilinear form $\psi: V \times V \rightarrow \mathbb{R}$ defining a polarization of the real Hodge structure $(r \circ h_x, V)$ for all x .*

(c) *Let $\{t_\alpha\}$ be a family of tensors of V such that $G_{\mathbb{R}}^{\text{ad}}$ is the subgroup of $GL(V)$ fixing the t_α . Then as x runs through the points of X , the Hodge structures \mathbf{V}_x run through a connected component of the set of all Hodge structures on V for which the tensors t_α are all of type $(0, 0)$.*

Proof. See Deligne (1979), 1.1.14.

We now apply (2.2) and the preceding discussion to the group G^{ad} and the cocharacter μ_0 corresponding to a point $o \in X$. In particular, we define the dual space \check{X} to be the set of filtrations of $\text{Rep}_{\mathbb{C}}(G^{\text{ad}})$ conjugate under $G^{\text{ad}}(\mathbb{C})$ to μ_0 (and hence to all μ_x for x in X). It is the compact dual symmetric Hermitian space of X in the usual sense (Helgason 1978, V.2). Note that $G(\mathbb{C})$ acts on \check{X} through $G(\mathbb{C}) \rightarrow G^{\text{ad}}(\mathbb{C})$.

Proposition 2.6. *The map $\beta: X \rightarrow \check{X}$ sending a point x in X to the filtration of $\text{Rep}_{\mathbb{C}}(G^{\text{ad}})$ defined by μ_x embeds X as an open complex submanifold of \check{X} . For $o \in X$, let K_0 be the isotropy group at o in $G(\mathbb{R})^+$, and let P_0 be the isotropy group at $o \in \check{X}$ in $G(\mathbb{C})$; then $K_0 = P_0 \cap G(\mathbb{R})^+$, and the inclusion of K_0 into P_0*

identifies $(K_0)_{\mathbb{C}}$ with a Levi subgroup of P_0 ; there is an equivariant commutative diagram

$$\begin{array}{ccc} G^{\text{ad}}(\mathbb{R})_+/K_0 & \hookrightarrow & G^{\text{ad}}(\mathbb{C})/P_0(\mathbb{C}) \\ \downarrow \approx & & \downarrow \approx \\ X & \hookrightarrow & \check{X}. \end{array}$$

Proof. This is proved in Deligne (1979), 1.1.14. We merely note that the injectivity of $X \rightarrow \check{X}$ follows from the fact that the Hodge filtration determines the Hodge decomposition.

The map β is the *Borel embedding* of X into \check{X} . Since \check{X} is an algebraic variety, we can speak of the variety $\tau\check{X}$ for τ an automorphism of \mathbb{C} . We shall show that $\tau\check{X}$ has a natural realization as the dual of a Hermitian symmetric domain.

Recall that for all automorphisms τ of \mathbb{C} , we have a canonical element $z_{\infty}(\tau) \in {}^{\tau}\mathfrak{G}(\mathbb{C})$. This gives rise to a canonical map

$$g \mapsto {}^{\tau,x}g = [z_{\infty}(\tau) \cdot g]: G'(\mathbb{C}) \xrightarrow{\approx} {}^{\tau,x}G'(\mathbb{C})$$

for central extension G' of G^{ad} .

Proposition 2.7. *Let ${}^{\tau,x}\check{X}$ be the dual Hermitian symmetric space associated with $({}^{\tau,x}G, {}^{\tau,x}X)$. For any special point of X , there is a unique isomorphism (of algebraic varieties) $\varphi_{\tau,x}^{\vee}: \tau\check{X} \rightarrow {}^{\tau,x}\check{X}$ such that*

- (i) *the point τx is mapped to ${}^{\tau,x}x$, and*
- (ii) *$\varphi_{\tau,x}^{\vee} \circ \tau(g) = ({}^{\tau,x}g) \circ \varphi_{\tau,x}^{\vee}$, for all g in $G^{\text{ad}}(\mathbb{C})$.*

Proof. The uniqueness is obvious. For the existence, note that there is a commutative diagram

$$\begin{array}{ccccc} G^{\text{ad}}(\mathbb{C}) & \xrightarrow{\tau} & G^{\text{ad}}(\mathbb{C}) & \longrightarrow & {}^{\tau,x}G^{\text{ad}}(\mathbb{C}) \\ \uparrow & & \uparrow & & \uparrow \\ T(\mathbb{C}) & \xrightarrow{\tau} & T(\mathbb{C}) & \xrightarrow{\text{id}} & T(\mathbb{C}) \\ \uparrow \mu_x & & \uparrow \tau\mu_x & \nearrow \mu_{\tau x} & \\ \mathbb{G}_m(\mathbb{C}) & \xrightarrow{\tau} & \mathbb{G}_m(\mathbb{C}) & & \end{array}$$

in which the unmarked arrow on the top row is $g \mapsto {}^{\tau,x}g$ and T is a suitable subtorus of G^{ad} . Let P_x, P' , and P'' be the subgroups of $G_{\mathbb{C}}^{\text{ad}}, G_{\mathbb{C}}^{\text{ad}}$, and ${}^{\tau,x}G_{\mathbb{C}}^{\text{ad}}$ respectively fixing the filtrations defined by $\mu_x, \tau\mu_x$, and $\mu_{\tau x}$. Then $P' = \tau P_x$ and P'' is a subgroup of ${}^{\tau,x}G_{\mathbb{C}}^{\text{ad}}$ such that $P''(\mathbb{C}) = \{{}^{\tau,x}p \mid \tau^{-1}p \in P(\mathbb{C})\}$. On passing to the quotients, we obtain maps $\check{X} \xrightarrow{\tau} \tau\check{X} \rightarrow {}^{\tau,x}\check{X}$, and the second of these obviously satisfies (i) and (ii).

Remark 2.8. (a) The subgroup P of G^{ad} is defined over the reflex field $E = E(G^{\text{ad}}, X)$, and ρ_x factors through $T_E \subset P$. The subgroup τP of G^{ad} still contains T because T is defined over \mathbb{Q} , and so we can use ρ_x and ${}^\tau\mathfrak{S}$ to twist τP . Clearly ${}^{\tau,x}(\tau P) = P''$.

(b) Let x' be a second special point of X . Then ${}^{\tau,x}g \mapsto {}^{\tau,x'}g$ is an isomorphism ${}^{\tau,x}G^{\text{ad}}(\mathbb{C}) \rightarrow {}^{\tau,x'}G^{\text{ad}}(\mathbb{C})$ giving rise to a commutative diagram

$$\begin{array}{ccccc} G^{\text{ad}}(\mathbb{C}) & \xrightarrow{\tau} & G^{\text{ad}}(\mathbb{C}) & \rightarrow & {}^{\tau,x}G^{\text{ad}}(\mathbb{C}) & g \mapsto {}^{\tau,x}g. \\ & & & \searrow & \downarrow & \searrow \downarrow \\ & & & & {}^{\tau,x'}G^{\text{ad}}(\mathbb{C}) & {}^{\tau,x'}g \end{array}$$

On passing to the quotients, we obtain a commutative diagram

$$\begin{array}{ccccc} X & \xrightarrow{\tau} & \tau X & \xrightarrow{\varphi_{\tau,x}^{\vee}} & {}^{\tau,x}\check{X} \\ & & \searrow \varphi_{\tau,x'}^{\vee} & & \downarrow \hat{\varphi}(\tau;x',x) \\ & & & & {}^{\tau,x'}\check{X}. \end{array}$$

3. The principal bundle $Y^0(G, X)$; statement of the first main theorem

We begin by reviewing some elementary constructions from complex differential geometry (see for example Kobayashi and Nomizu (1963/69)).

Let S be a connected complex manifold, and let $\pi_1 = \pi_1(S, s)$ be the fundamental group of S regarded as the group of covering transformations of the universal covering space \tilde{S} of S (acting on the left). A homomorphism $r: \pi_1(S, s) \rightarrow G$ from $\pi_1(S, s)$ into a complex Lie group G gives rise to a (right) principal G -bundle

$$Y(r) = \sim \backslash \tilde{S} \times G, \quad (\gamma s, r(\gamma)g) \sim (s, g)$$

on S , and there is a canonical flat connection on $Y(r)$. Every principal G -bundle Y over S admitting a flat connection arises in this way.

Let V be a complex vector space. A homomorphism $r: \pi_1(S, s) \rightarrow GL(V)$ gives rise to a vector bundle

$$\mathcal{V}(r) = \sim \backslash \tilde{S} \times V, \quad (\gamma s, r(\gamma)v) \sim (s, v)$$

on S , and there is a canonical flat connection on \mathcal{V} . The sections of $Y(r)$ over any open subset U of S can be identified with the isomorphisms

$$a: \mathcal{O}_U \otimes V \xrightarrow{\sim} \mathcal{V}|U \text{ (trivializations of } \mathcal{V} \text{ over } U).$$

Now suppose that r factors through a reductive algebraic subgroup G of $GL(V)$. There will exist a finite family of tensors (t_α) of V such that G is the subgroup fixing the t_α . Each t_α gives rise to a global tensor t'_α of \mathcal{V} , and the sections of $Y(r)$ over any open subset U can then be identified with the isomorphisms $a: \mathcal{O}_U \otimes V \xrightarrow{\sim} \mathcal{V}|U$ under which each $1 \otimes t_\alpha$ corresponds to $t'_\alpha|U$ (trivializations of \mathcal{V} over U respecting the t_α).

Proposition 3.1. *In addition to the hypotheses in the last paragraph, assume that S is algebraic, that \mathcal{V} is an algebraic vector bundle on S , and that the tensors t'_α are algebraic sections of the $T^1_j \mathcal{V}$. Then $Y(r)$ is algebraic, and it represents the functor of S -varieties whose value on $\pi: T \rightarrow S$ is the set of isomorphisms*

$$a: \mathcal{O}_T \otimes V \xrightarrow{\cong} \pi^* \mathcal{V} \text{ such that } 1 \otimes t_\alpha \text{ corresponds to } \pi^*(t'_\alpha) \text{ for all } \alpha.$$

Proof. Suppose first that $G = GL(V)$ (and there are no tensors). If \mathcal{V} is trivial, i.e., $\mathcal{V} = \mathcal{O}_S \otimes V$, then the functor is represented by G_S . Since \mathcal{V} is locally trivial for the Zariski topology, it follows easily that the functor is represented by a G_S -torsor Y , and it is obvious from the discussion preceding the proposition that the analytic space associated with Y is $Y(r)$. In the general case, the tensors t_α define a $G_{\mathbb{C}}$ -subtorsor of the $GL(V)_{\mathbb{C}}$ -torsor Y whose associated analytic space is again $Y(r)$.

We apply these remarks to a pair (G, X) defining a connected Shimura variety. For each $\Gamma \in \tilde{\Sigma}(G)$, Γ is the fundamental group of S^0_Γ and (by definition) $\Gamma \subset G(\mathbb{Q}) \subset G(\mathbb{C})$. The above construction gives us a principal $G(\mathbb{C})$ -bundle $Y^0_\Gamma(G, X) = \Gamma \backslash X \times G(\mathbb{C})$ over $S^0_\Gamma(G, X)$. Because of our conventions we are forced to turn this into a left principal bundle (by making $g \in G(\mathbb{C})$ act as g^{-1} in the natural action; that is, $g[x, c] = [x, cg^{-1}]$). For varying Γ , these bundles form a projective system $Y^0(G, X)$, which can be regarded as a principal $G(\mathbb{C})$ -bundle over $S^0(G, X)$. In the case that G is simply connected,

$$Y^0(G, X) = G(\mathbb{Q}) \backslash X \times G(\mathbb{C}) \times G(\mathbb{A}^f)$$

with $q \in G(\mathbb{Q})$ acting on $(x, c, a) \in X \times G(\mathbb{C}) \times G(\mathbb{A}^f)$ according to the rule

$$q(x, c, a) = (qx, qc, qa).$$

There is an action of $G^{\text{ad}}(\mathbb{Q})^+$ on $Y^0(G, X)$: $g \in G^{\text{ad}}(\mathbb{Q})^+$ maps $[x, c]_{\Gamma \in \Gamma \backslash X \times G(\mathbb{C})}$ to $[gx, \text{ad}(g)c]_{\Gamma' \in \Gamma' \backslash X \times G(\mathbb{C})}$, where $\Gamma' = \text{ad}(g)\Gamma$. We let $G(\mathbb{Q})$ act on $S^0(G, X)$ through the map $G(\mathbb{Q}) \rightarrow G^{\text{ad}}(\mathbb{Q})$; this extends by continuity to an action of the closure $G(\mathbb{Q})^+_{\mathbb{C}}$ of $G(\mathbb{Q})_+$ in $G(\mathbb{A}^f)$. Therefore $G(\mathbb{C}) \times G(\mathbb{Q}^+_{\mathbb{C}})$ and $G^{\text{ad}}(\mathbb{Q})^+$ both act on $Y^0(G, X)$. If $q \in G(\mathbb{Q})^+_{\mathbb{C}}$, then $(q, q) \in G(\mathbb{C}) \times G(\mathbb{Q}^+_{\mathbb{C}})$ and the image of q in $G^{\text{ad}}(\mathbb{Q})^+$ have the same action, and consequently we obtain an action of

$$\mathcal{G}(G) = (G(\mathbb{C}) \times G(\mathbb{Q}^+_{\mathbb{C}})) *_{G(\mathbb{Q})^+} G^{\text{ad}}(\mathbb{Q})^+$$

on $Y^0(G, X)$. There are obvious homomorphisms $G(\mathbb{C}) \rightarrow \mathcal{G}(G) \rightarrow G^{\text{ad}}(\mathbb{Q})^+_{\mathbb{C}}$, and the action of $\mathcal{G}(G)$ on $Y^0(G, X)$ is compatible via these maps with the actions of $G(\mathbb{C})$ and $G^{\text{ad}}(\mathbb{Q})^+_{\mathbb{C}}$ on $Y^0(G, X)$ and $S^0(G, X)$ respectively.

In the case that G is simply connected, $G(\mathbb{Q}^+_{\mathbb{C}}) = G(\mathbb{A}^f)$, and the actions are given by

$$\begin{aligned} a[x, z, g] &= [x, z, ga^{-1}], & x \in X, z \in G(\mathbb{C}), a, g \in G(\mathbb{A}^f), \\ c[x, z, g] &= [x, zc^{-1}, g], & x \in X, c, z \in G(\mathbb{C}), g \in G(\mathbb{A}^f). \end{aligned}$$

In the general case, $\mathcal{G}(G)$ is generated by the images of $G(\mathbb{C})$, $\tilde{G}(\mathbb{A}^f)$, and $G^{\text{ad}}(\mathbb{Q})^+_{\mathbb{C}}$.

Now let $r: G_{\mathbb{C}} \rightarrow GL(V)$ be a representation of $G_{\mathbb{C}}$. In this case we obtain a vector bundle $\mathcal{V}(r)$ on $S^0(G, X)$ together with an action of $G(\mathbb{Q})^{\pm}$ on $\mathcal{V}(r)$.

Proposition 3.2. *The principal $G(\mathbb{C})$ -bundle $Y^0(G, X)$ is algebraic, and the elements of $\mathcal{G}(G)$ act algebraically on it.*

Proof. We shall need to use the following result.

Lemma 3.3. *Let S be an algebraic variety embedded as an open subvariety of a complete algebraic variety \bar{S} .*

(a) *If $\bar{S} - S$ has codimension ≥ 3 , then the functor $\mathcal{F} \mapsto \mathcal{F}^{\text{an}}$ taking an algebraic vector bundle on S to its associated analytic vector bundle defines an equivalence of categories.*

(b) *If $\bar{S} - S$ has codimension ≥ 2 , then $\mathcal{F} \mapsto \mathcal{F}^{\text{an}}$ is fully faithful, and $\Gamma(S, \mathcal{F}) = \Gamma(S, \mathcal{F}^{\text{an}})$.*

Proof. This follows from theorems of Serre, Grothendieck, Siu, and Trautmann; see Hartshorne (1970), p. 222.

We first prove the proposition under the assumption that the boundary of $S_F^0(G, X)$ in its Baily-Borel compactification has codimension ≥ 3 . Choose a faithful representation $r: G \rightarrow GL(V)$ of G , and let (t_{α}) be a finite family of tensors of V such that G is the subgroup of $GL(V)$ fixing the t_{α} . Then (3.3a) shows that the sheaf \mathcal{V}_r on $S_F^0(G, X)$ corresponding to V is algebraic, and (3.3b) shows that the global tensors t'_{α} are algebraic. It now follows from Proposition 3.1 that $Y_F^0(G, X)$ is algebraic. An element of $G(\mathbb{Q})^{\pm}$ acts as an algebraic morphism on the family $(Y_F^0(G, X))$ because (3.3a) shows that it does so on the family (\mathcal{V}_r) . On applying this to G^{ad} we find that an element of $G^{\text{ad}}(\mathbb{Q})^{\pm}$ acts algebraically on $Y_F^0(G^{\text{ad}}, X)$, and therefore on $Y_F^0(G, X)$, because $Y_F^0(G, X)$ is a finite covering of $Y_F^0(G^{\text{ad}}, X)$. Finally an element of $G(\mathbb{C})$ acts algebraically on $Y^0(G, X)$ because it defines a morphism of the functor that $Y^0(G, X)$ represents.

Next we assume that the boundary has codimension ≥ 2 . In this case, there will be a totally real field F and a pair (G_*, X_*) with $G_* = \text{Res}_{F/\mathbb{Q}} G$ such that the boundary of $S_F^0(G_*, X_*)$ has codimension ≥ 3 and the natural map $G \hookrightarrow G_*$ sends X into X_* . Choose a faithful representation r_* of G_* , and let r be its restriction to G . Then the sheaf \mathcal{V} on $S^0(G, X)$ defined by r is obtained by restriction from the similar sheaf on $S^0(G_*, X_*)$, and so it is algebraic. Moreover the global tensors of \mathcal{V} are again algebraic, and so the same argument as before applies.

In the only remaining case, the boundary has codimension = 1. But then G is SL_2 or PGL_2 , and the result is easy to prove, for example by making use of the universal elliptic curve (cf. § 6).

In summary, with a pair (G, X) defining a connected Shimura variety, we have associated a $G_{\mathbb{C}}$ -torsor $Y^0(G, X)$ over $S^0(G, X)$, a flat connection ∇ on $Y^0(G, X)$, and an action a of $\mathcal{G}(G)$ on $(Y^0(G, X), \nabla)$; moreover, a special point x on X defines a point $y \in Y^0(G, X)$ lying over $[x]$, namely, its image under the map $X \rightarrow Y^0(G, X)$. The association is functorial: a morphism $(G, X) \rightarrow (G', X')$ carrying a special point x of X to a special point x' of X'

gives rise to a morphism of quadruples

$$(Y^0(G, X), \mathcal{V}, a, y) \rightarrow (Y^0(G', X'), \mathcal{V}', a', y').$$

We need an adèlic version of (3.1). Write $p: Y^0(G, X) \rightarrow S^0(G, X)$ for the structure map, and let $r: G \hookrightarrow GL(V)$ be a faithful representation of G . Let (t_α) be a finite family of tensors such that G is the subgroup of $GL(V)$ fixing the t_α , and let t'_α be the global tensor corresponding to t_α . Then there is a canonical equivariant trivialization $a_0: \mathcal{O}_{Y_0} \otimes V \xrightarrow{\sim} p^* \mathcal{V}(r)$, and under the trivialization, $1 \otimes t_\alpha$ corresponds to $p^*(t'_\alpha)$ for each α . The next proposition will show that $Y^0(G, X)$ is universal for this property.

Proposition 3.4. *Let $r: G \hookrightarrow GL(V)$, (t_α) , and (t'_α) be as above. For any morphism $\pi: T \rightarrow S^0(G, X)$ of $\mathcal{G}(G)$ -varieties and equivariant isomorphism $a: \mathcal{O}_T \otimes V \xrightarrow{\sim} \pi^* \mathcal{V}(r)$ such that $1 \otimes t_\alpha \leftarrow \pi^*(t'_\alpha)$ for all α , there is a unique $\mathcal{G}(G)$ -equivariant S^0 -morphism $\psi: T \rightarrow Y^0(G, X)$ such that a is the inverse image of a_0 .*

Proof. This follows easily from (3.1).

Proposition 3.5. *There is a canonical $G(\mathbb{C})$ -equivariant map*

$$\gamma(G, X): Y^0(G, X) \rightarrow \check{X}.$$

Proof. As there is a canonical map $Y^0(G, X) \rightarrow Y^0(G^{ad}, X)$, and \check{X} is unchanged when G is replaced by G^{ad} , we may assume that $G = G^{ad}$. Let $r: G \rightarrow GL(V)$ be a faithful representation of G such that the corresponding vector bundle \mathcal{V} on $S^0(G, X)$ is algebraic, and consider the constant vector bundle $\mathcal{V}_\Gamma = Y^0_\Gamma(G, X) \times V$. According to (3.1), an element $y \in Y^0_\Gamma$ can be identified with an isomorphism $a_y: V \xrightarrow{\sim} (\mathcal{V}_\Gamma)_y$. Endow $(\mathcal{V}_\Gamma)_y$ with the Hodge filtration defined by the Hodge structure $a_y \circ r \circ h_x: \mathbf{S} \rightarrow GL((\mathcal{V}_\Gamma)_y)$, where x_Γ is the image of y in $S^0_\Gamma(G, X)$. Then (2.4b) shows that there is a unique morphism of algebraic varieties $\gamma_\Gamma: Y_\Gamma(G, X) \rightarrow \check{X}$ such that $\gamma^*_\Gamma(\mathcal{V}) \approx \mathcal{V}_\Gamma$ as filtered vector bundles. For varying Γ , the γ_Γ form a projective system, that is, a morphism $\gamma: Y^0(G^{ad}, X) \rightarrow \check{X}$. The map commutes with the actions of $G(\mathbb{C})$.

Remark 3.6. (a) The composite of the canonical map $X \rightarrow Y^0(G, X)$ with $\gamma(G, X)$ is the Borel embedding β .

(b) When G is simply connected, γ is the map $[x, c, a] \mapsto c^{-1} \beta(x)$, $x \in X$, $c \in G(\mathbb{C})$, $a \in G(\mathbb{A}^f)$.

Let τ be an automorphism of \mathbb{C} , and let x be a special point of X . The point $(z_\infty(\tau), \text{sp}(\tau)) \in {}^t\mathfrak{S}(\mathbb{A}')$ defines an isomorphism

$$g \mapsto {}^t, x g: G(\mathbb{A}') \rightarrow {}^t, x G(\mathbb{A}').$$

(Recall that $\mathbb{A}' = \mathbb{C} \times \mathbb{A}^f$.)

Proposition 3.7. *There is a canonical map*

$$g \mapsto {}^{\tau, x}g: \mathcal{G}(G) \rightarrow \mathcal{G}({}^{\tau, x}G)$$

compatible with the above map.

Proof. For simplicity, we first prove this under the assumption that G is simply connected. Choose a pair (G_1, X_1) as in (A.4) of the Appendix. Let H be the torus G_1/G , and let v be the map $G_1 \rightarrow H$. As in (1.5), we write $H(\mathbb{R})_+$ for $v(Z_1(\mathbb{R})) \subset H(\mathbb{R})$ and $H(\mathbb{Q})_+$ for $H(\mathbb{Q}) \cap H(\mathbb{R})_+$. Because of (A.4c), any $q \in G^{\text{ad}}(\mathbb{Q})^+$ lifts to an element $\tilde{q} \in G_1(\mathbb{Q})_+$, and one checks immediately that $(c, a)^*q \mapsto [c\tilde{q}, a\tilde{q}]$ is a well-defined homomorphism $\mathcal{G}(G) \rightarrow G_1(\mathbb{A}')/Z_1(\mathbb{Q})$.

Lemma 3.8. *There is an exact sequence*

$$0 \rightarrow \mathcal{G}(G) \rightarrow G_1(\mathbb{A}')/Z_1(\mathbb{Q}) \xrightarrow{v} H(\mathbb{A}')/H(\mathbb{Q})_+ \rightarrow 0$$

Proof. We first show that the final map is surjective. For this it suffices to show that v defines a surjection $G_1(\mathbb{A}^f) \rightarrow H(\mathbb{A}^f)$. Choose \mathbb{Z} -structures on G_1 and H . Then Lang’s lemma applied to the reduction of G shows that $G_1(\mathbb{Z}_p) \rightarrow H(\mathbb{Z}_p)$ is surjective for almost all p . It therefore suffices to show that $G_1(\mathbb{Q}_p) \rightarrow H(\mathbb{Q}_p)$ is surjective for all p , but this follows from the fact that $H^1(\mathbb{Q}_p, G) = 0$ for all finite primes p .

Next we prove the exactness at the middle term. From the diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & Z_1(\mathbb{R}) & \rightarrow & G_1(\mathbb{R}) & \rightarrow & G^{\text{ad}}(\mathbb{R}) & \rightarrow & 0 \\ & & \parallel & & \downarrow^v & & \downarrow & & \\ & & Z_1(\mathbb{R}) & \rightarrow & H(\mathbb{R}) & \rightarrow & H(\mathbb{R})/v(Z_1(\mathbb{R})) & \rightarrow & 0 \end{array}$$

we see that $G^{\text{ad}}(\mathbb{R})^+$ maps into the identity component of $H(\mathbb{R})/v(Z_1(\mathbb{R}))$. But $H(\mathbb{R})/v(Z_1(\mathbb{R}))$ is discrete, and so $G^{\text{ad}}(\mathbb{R})^+$ maps into $v(Z_1(\mathbb{R}))$. It follows from this that $\mathcal{G}(G)$ maps to zero under the composite of the two maps.

Suppose that $v(c, a) \in H(\mathbb{Q})_+ \times H(\mathbb{Q})_+$ for some $(c, a) \in G_1(\mathbb{C}) \times G_1(\mathbb{A}^f)$, say $v(c, a) = (q, q)$. Consider the diagram

$$\begin{array}{ccccc} G_1(\mathbb{Q}) & \rightarrow & H(\mathbb{Q}) & \rightarrow & H^1(\mathbb{Q}, G) \\ \downarrow & & \downarrow & & \downarrow \\ G_1(\mathbb{R}) & \rightarrow & H(\mathbb{R}) & \rightarrow & H^1(\mathbb{R}, G). \end{array}$$

The image of q in $H(\mathbb{R})$ lifts to $Z_1(\mathbb{R}) \subset G_1(\mathbb{R})$ because it lies in $H(\mathbb{R})_+$. Therefore the image of q in $H^1(\mathbb{Q}, G)$ maps to zero in $H^1(\mathbb{R}, G)$, which implies that it is zero by the Hasse principle. Hence q lifts to an element \tilde{q} in $G_1(\mathbb{Q})$. Now

$v(c\tilde{q}^{-1}, a\tilde{q}^{-1}) = 1$, and so $c' \stackrel{df}{=} c\tilde{q}^{-1}$ and $a' \stackrel{df}{=} a\tilde{q}^{-1}$ lie in $G(\mathbb{C})$ and $G(\mathbb{A}^f)$ respectively. Let \bar{q} be the image of \tilde{q} in $G^{\text{ad}}(\mathbb{Q})$. Then $(c', a')^*\bar{q} \in \mathcal{G}(G)$, and it maps to $(c, a) \bmod (Z_1(\mathbb{Q}))$.

Finally we prove the exactness at the first term. Let $(c, a)^*q \in G^{\text{ad}}(\mathbb{Q})^{\wedge}$, and choose a lifting \tilde{q} of q to $G_1(\mathbb{Q})$. If $(c, a)^*q$ maps to zero in $G_1(\mathbb{A}')/Z_1(\mathbb{Q})$,

then there exists a $z \in Z_1(\mathbb{Q})$ such that $(c\tilde{q}, a\tilde{q}) = (z, z)$. Now $c = z\tilde{q}^{-1} \in G(\mathbb{C}) \cap G_1(\mathbb{Q}) = G(\mathbb{Q})$; write it q' . Then $q'\tilde{q} \in Z_1(\mathbb{Q})$, and so the image of q' in $G^{\text{ad}}(\mathbb{Q})$ is q^{-1} . Consequently, $(q', q') * q = 1$, but $(q', q') * q = (c, a) * q$.

We now return to the proof of the proposition. If $z \in Z_1(\mathbb{Q})$ then the element $[s \cdot z]$ of ${}^{\tau, x}G_1(\mathbb{Q})$ is independent of s . Therefore $z \mapsto [s \cdot z]$ is an inclusion $Z \hookrightarrow {}^{\tau, x}G_1$, and the map

$$g \mapsto {}^{\tau, x}g = [(z_{\infty}(\tau), \text{sp}(\tau)) \cdot g]: G_1(\mathbb{A}') \rightarrow {}^{\tau, x}G_1(\mathbb{A}')$$

induces the identity map on $Z_1(\mathbb{A}')$. In particular it maps $Z_1(\mathbb{Q}) \subset G_1(\mathbb{A}')$ into $Z_1(\mathbb{Q}) \subset {}^{\tau, x}G_1(\mathbb{A}')$, and so we have a map $\psi: G_1(\mathbb{A}')/Z_1(\mathbb{Q}) \rightarrow {}^{\tau, x}G_1(\mathbb{A}')/Z_1(\mathbb{Q})$. Since $g \mapsto {}^{\tau, x}g: G_1(\mathbb{A}') \rightarrow {}^{\tau, x}G_1(\mathbb{A}')$ induces the identity map on $H(\mathbb{A}')$, ψ maps $\mathcal{G}(G)$ into $\mathcal{G}({}^{\tau, x}G)$, and this is the map we want.

When G is not simply connected, then $\mathcal{G}(G)$ is the quotient of $\mathcal{G}(\tilde{G})$ by

$$\mathcal{G}(\tilde{G}, G) \stackrel{\text{df}}{=} \text{Ker}(\tilde{Z}(\mathbb{A}')/\tilde{Z}(\mathbb{Q}) \rightarrow Z(\mathbb{A}')/Z(\mathbb{Q})),$$

where \tilde{Z} and Z are the centres of \tilde{G} and G respectively. There is a canonical isomorphism $\mathcal{G}(\tilde{G}, G) \xrightarrow{\sim} \mathcal{G}({}^{\tau, x}\tilde{G}, {}^{\tau, x}G)$ and $g \mapsto {}^{\tau, x}g: \mathcal{G}(\tilde{G}) \rightarrow \mathcal{G}({}^{\tau, x}\tilde{G})$ is compatible with this isomorphism (cf. the above discussion involving Z_1). Hence the map $g \mapsto {}^{\tau, x}G$ on $\mathcal{G}(G)$ can be defined by passing to the quotient with the map $g \mapsto {}^{\tau, x}g$ on $\mathcal{G}(\tilde{G})$.

From the pair $({}^{\tau, x}G, {}^{\tau, x}X)$ and the special point ${}^{\tau, x}x$ of ${}^{\tau, x}X$, we can construct a ${}^{\tau, x}G(\mathbb{C})$ -torsor $Y^0({}^{\tau, x}G, {}^{\tau, x}X)$, a flat connection ${}^{\tau, x}\mathcal{V}$, an action ${}^{\tau, x}a$ of $\mathcal{G}(G)$ on $(Y^0({}^{\tau, x}G, {}^{\tau, x}X), {}^{\tau, x}\mathcal{V})$, and a point ${}^{\tau, x}y$ on $Y^0({}^{\tau, x}G, {}^{\tau, x}X)$.

Proposition 3.9. *Let x' be a second special point. There is a canonical isomorphism $\phi^Y(\tau; x', x): Y^0({}^{\tau, x}G, {}^{\tau, x}X) \rightarrow Y^0({}^{\tau, x'}G, {}^{\tau, x'}X)$ such that*

- (i) $\phi^Y(\tau; x', x)$ is compatible with the flat connections;
- (ii) $\phi^Y(\tau; x', x) \circ ({}^{\tau, x}g) = ({}^{\tau, x'}g) \circ \phi^Y(\tau; x', x)$ for all $g \in \mathcal{G}(G)$;
- (iii) the following diagram commutes

$$\begin{array}{ccc} \phi^{\vee}(\tau; x', x): & {}^{\tau, x}\check{X} & \rightarrow & {}^{\tau, x'}\check{X} \\ & \uparrow \gamma & & \uparrow \gamma \\ \phi^Y(\tau; x', x): & Y^0({}^{\tau, x}G, {}^{\tau, x}X) & \rightarrow & Y^0({}^{\tau, x'}G, {}^{\tau, x'}X) \\ & \downarrow & & \downarrow \\ \phi^0(\tau; x', x): & S^0({}^{\tau, x}G, {}^{\tau, x}X) & \rightarrow & S^0({}^{\tau, x'}G, {}^{\tau, x'}X). \end{array}$$

Proof. The proof of (1.3) applies with minor modifications. For simplicity, we first assume that G is simply connected. Choose a pair (G_1, X_1) (with the weight equal to zero) and an isomorphism $t: \rho_{x_1, *}({}^{\tau}\mathfrak{S}) \xrightarrow{\sim} \rho_{x'_1, *}({}^{\tau}\mathfrak{S})$, as in the proof of (1.3). Again t defines an isomorphism $f_1: {}^{\tau, x}G_1 \xrightarrow{\sim} {}^{\tau, x'}G_1$. Let s and s' be the images of $(z_{\infty}(\tau), \text{sp}(\tau))$ in $\rho_{x_1, *}({}^{\tau}\mathfrak{S})(\mathbb{A}')$ and $\rho_{x'_1, *}({}^{\tau}\mathfrak{S})(\mathbb{A}')$ respectively. Then there is an element $\gamma_1 \in {}^{\tau, x}G_1(\mathbb{A}')$ such that $t_1(s\gamma_1) = s'$. The same argument as in the proof of (1.3) shows that the image of γ_1 in $H(\mathbb{A}')$ lies in its subgroup $H(\mathbb{Q})$, and the Remark 1.5 shows that it lies in $H(\mathbb{Q})_+ \subset H(\mathbb{Q})$. Therefore, accord-

ing to (3.8), the image γ of γ_1 in ${}^{\tau,x}G_1(\mathbb{A}')/{}^{\tau,x}Z_1(\mathbb{Q})$ lies in $\mathcal{G}({}^{\tau,x}G)$. Now $\varphi^0(\tau; x', x)$ can be defined to be $Y^0(f) \circ (\gamma)$ where f is the restriction of f_1 to ${}^{\tau,x}G$.

In the case that G is not simply connected, first construct the diagram for (\tilde{G}, X) , and then pass to the appropriate quotients.

Theorem 3.10. *Let (G, X) be a pair defining a connected Shimura variety, and let x be a special point of X . For any automorphism τ of \mathbb{C} , there is a unique isomorphism $\varphi_{\tau,x}^Y: \tau Y^0(G, X) \rightarrow Y^0({}^{\tau,x}G, {}^{\tau,x}X)$ lying over $\varphi_{\tau,x}^0$ and such that*

- (i) τy is mapped to ${}^{\tau}y$;
- (ii) $\varphi_{\tau,x}^Y \circ \tau(g) = ({}^{\tau,x}g) \circ \varphi_{\tau,x}^Y$ for all $g \in \mathcal{G}(G)$.

Moreover,

(iii) $\varphi_{\tau,x}^Y$ is compatible with the flat connections on $\tau Y^0(G, X)$ and $Y^0({}^{\tau,x}G, {}^{\tau,x}X)$. If x' is a second special point, then $\varphi^Y(\tau; x', x) \circ \varphi_{\tau,x}^Y = \varphi_{\tau,x'}^Y$.

Proof. We prove only the uniqueness of $\varphi_{\tau,x}^Y$ here; the proof of the existence will occupy § 6–§ 9.

Let φ_1 and φ_2 be two maps $\tau Y^0(G, X) \rightarrow Y^0({}^{\tau,x}G, {}^{\tau,x}X)$ satisfying the conditions (i) and (ii), and let Z be the closed subset of $\tau Y^0(G, X)$ on which the two maps agree. Then (ii) shows that Z contains a fibre of the map $\pi: \tau Y^0(G, X) \rightarrow \tau S^0(G, X)$ whenever it contains a single point of the fibre. Therefore $Z = \pi^{-1}(Z')$ for some subset Z' of $\tau S^0(G, X)$, and because Z is closed, so also must be Z' . Condition (i) shows that Z' contains $\tau[x]$, and (ii) then shows that it contains all translate of $\tau[x]$ by elements of $G^{\text{ad}}(\mathbb{Q})^{+\wedge}$. As these points make up a dense subset of $S^0(G, X)$ for the Zariski topology, it follows that $Z' = S^0(G, X)$ and $Z = Y^0(G, X)$.

Corollary 3.11. *Assume that there is a map $\varphi_{\tau,x}^Y$ as in the theorem; then there a commutative diagram:*

$$\begin{array}{ccc}
 \tau \tilde{X} & \xrightarrow{\varphi_{\tau,x}^{\tilde{X}}} & {}^{\tau,x} \tilde{X} \\
 \uparrow \tau \gamma & & \uparrow \gamma \\
 \tau Y^0(G, X) & \xrightarrow{\varphi_{\tau,x}^Y} & Y^0({}^{\tau,x}G, {}^{\tau,x}X) \\
 \downarrow & & \downarrow \\
 \tau S^0(G, X) & \xrightarrow{\varphi_{\tau,x}^0} & S^0({}^{\tau,x}G, {}^{\tau,x}X).
 \end{array}$$

The two upper maps are compatible with the map $g \mapsto {}^{\tau,x}g: G(\mathbb{C}) \rightarrow {}^{\tau,x}G(\mathbb{C})$.

Proof. It remains to prove that the upper square commutes, i.e., that the maps $\varphi_{\tau,x}^Y \circ \tau \gamma(G, X)$ and $\gamma({}^{\tau,x}G, {}^{\tau,x}X) \circ \varphi_{\tau,x}^Y$ are equal. They are maps $\tau Y^0(G, X) \rightarrow {}^{\tau,x} \tilde{X}$, and we know that they agree at τy . Both maps are constant on the orbits of $G(\mathbb{Q})_+^-$ in $\tau Y^0(G, X)$, and so they agree on all translates of y . Finally, both maps are compatible with the map $g \mapsto {}^{\tau,x}g: G(\mathbb{C}) \rightarrow {}^{\tau,x}G(\mathbb{C})$, and so they agree on a fibre of $\pi: \tau Y^0(G, X) \rightarrow \tau S^0(G, X)$ when they agree at a single point. The argument in the preceding proof now shows that the closed subset where the maps agree is the whole of $\tau Y^0(G, X)$.

4. Automorphic vector bundles

In this section we define automorphic vector bundles, and discuss some of their structure. For a similar discussion in the case of nonconnected Shimura varieties, see Harris (1985), § 3.

Let (G, X) be a pair defining a connected Shimura variety, and let $\beta: X \hookrightarrow \check{X}$ be the Borel embedding. The action of $G(\mathbb{R})_+$ on X extends to a transitive action of $G(\mathbb{C})$ on \check{X} . Since \check{X} is a projective algebraic variety, every holomorphic vector bundle on \check{X} is algebraic. By a $G_{\mathbb{C}}$ -vector bundle on \check{X} , we mean a vector bundle (\mathcal{J}, p) on \check{X} together with an action of $G_{\mathbb{C}}$ on \mathcal{J} (as an algebraic variety) such that

- (a) $p(g \cdot w) = g \cdot p(w)$ for all $g \in G(\mathbb{C})$, $w \in \mathcal{J}$;
- (b) the maps $g: \mathcal{J}_x \rightarrow \mathcal{J}_{gx}$ are linear for all $g \in G(\mathbb{C})$ and $x \in \check{X}$.

Such a vector bundle restricts to a $G(\mathbb{R})_+$ -vector bundle $\tilde{\mathcal{V}} = \beta^* \mathcal{J}$ on X , and

for each $\Gamma \in \tilde{\Sigma}(G)$, $\mathcal{V}_{\Gamma} \stackrel{df}{=} \Gamma \backslash \tilde{\mathcal{V}}$ is a vector bundle on $\Gamma \backslash X = S^0_{\Gamma}(G, X)$. The family $\mathcal{V} = (\mathcal{V}_{\Gamma})$ forms a projective system, and there is a natural action of $G(\mathbb{Q})_+$ on \mathcal{V}

$$g: \Gamma \backslash \tilde{\mathcal{V}} \rightarrow g \Gamma g^{-1} \backslash \tilde{\mathcal{V}}, \quad v(\text{mod } \Gamma) \mapsto gv(\text{mod } g \Gamma g^{-1}),$$

which extends by continuity to the closure $G(\mathbb{Q})_+^-$ of $G(\mathbb{Q})_+$ in $G(\mathbb{A}^f)$. A $G(\mathbb{Q})_+^-$ -vector bundle \mathcal{V} on $S^0(G, X)$ arising in this way from a $G(\mathbb{C})$ -vector bundle \mathcal{J} on \check{X} will be called an *automorphic vector bundle*. We sometimes write $\mathcal{V} = \mathcal{V}(\mathcal{J})$. For each x in X , the fibre $(\mathcal{V}_{\Gamma})_{x_{\Gamma}} = \tilde{\mathcal{V}}_x = \mathcal{J}_{\beta(x)}$, and so

$$\mathcal{V}_{[x]} \stackrel{df}{=} \lim_{\leftarrow} (\mathcal{V}_{\Gamma})_{x_{\Gamma}} = \mathcal{J}_{\beta(x)}.$$

When G is simply connected, \mathcal{V} is the $G(\mathbb{A}^f)$ -vector bundle

$$\mathcal{V} = G(\mathbb{Q}) \backslash \tilde{\mathcal{V}} \times G(\mathbb{A}^f)$$

on $S^0(G, X) = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}^f)$, and the action of $a \in G(\mathbb{A}^f)$ is given by

$$a[v, g] = [v, ga^{-1}].$$

On the other hand, when the action of $G(\mathbb{C})$ on \mathcal{J} factors through $G^{\text{ad}}(\mathbb{C})$, then we can regard \mathcal{V} as being the projective system $(\Gamma \backslash \tilde{\mathcal{V}})$, $\Gamma \in \Sigma(G)$, and the action of $G^{\text{ad}}(\mathbb{Q})_{\uparrow}$ on $S^0(G, X)$ lifts to \mathcal{J} .

We wish to show that automorphic vector bundles are algebraic, but first we need a lemma.

Lemma 4.1. *Let G be an affine algebraic group over a field k , and let $\pi: Y \rightarrow S$ be a torsor for G_S over an algebraic variety S .*

(a) *The functor $\mathcal{V} \mapsto \pi^* \mathcal{V}$ defines an equivalence between the category of vector bundles on S and the category of G -vector bundles on Y .*

(b) *If Y has a flat G -connection, then to give a (flat) connection on \mathcal{V} is the same as to give a (flat) G -connection on $\pi^* \mathcal{V}$.*

Proof. (a) This follows from descent theory: the map π is faithfully flat, and because Y is a G -torsor, to give a descent datum on an \mathcal{O}_Y -module \mathcal{V}' is the same as to give a G -action on it. That $\pi^*(\mathcal{V})$ is locally free if and only if \mathcal{V} is locally free follows from the fact that a coherent module is locally free if and only if it is flat.

(b) Again this is a straightforward application of descent theory.

Proposition 4.2. *The vector bundle $\mathcal{V}(\mathcal{J})$ is obtained by descent from $\gamma^*(\mathcal{J})$; hence \mathcal{V} and the action of $G(\mathbb{Q})_+^-$ on it are algebraic.*

Proof. The first statement is obvious from the commutative diagram:

$$\begin{array}{ccccc} X & \rightarrow & Y^0(G, X) & \xrightarrow{\gamma} & \check{X} \\ & \searrow & \downarrow & & \\ & & S^0(G, X) & & \end{array}$$

The second statement follows from the first (because of (3.2)).

Obviously, a section of \mathcal{V}_Γ over S_Γ^0 gives rise to a section of $\check{\mathcal{V}}$ over X . We define an *automorphic form of type \mathcal{J} and level Γ* to be a section of (the analytic vector bundle) $\check{\mathcal{V}}$ over X that arises from a section of (the algebraic vector bundle) \mathcal{V}_Γ . Thus an automorphic form of type \mathcal{J} and any level is an element of the union $\cup \Gamma(S_\Gamma^0, \mathcal{V}_\Gamma)'$, where $\Gamma(S_\Gamma^0, \mathcal{V}_\Gamma)'$ denotes the image of $\Gamma(S_\Gamma^0, \mathcal{V}_\Gamma)$ in $\Gamma(X, \check{\mathcal{V}})$. When G^{ad} has no \mathbb{Q} -rational factors isomorphic to PGL_2 , then the boundary of S_Γ^0 in its Baily-Borel compactification has codimension ≥ 2 , and so $\Gamma(S_\Gamma^0, \mathcal{V}_\Gamma) = \Gamma(S_\Gamma^0, \mathcal{V}_\Gamma^{\text{an}})$ (see 3.3). Therefore, in this case, an automorphic form of type \mathcal{J} and level Γ is simply a holomorphic section of $\check{\mathcal{V}}$ over X fixed under the action of Γ .

Let o be a point of X , and regard it also as a point of \check{X} . The isotropy group at o in $G_{\mathbb{C}}$ is a parabolic subgroup P_0 of $G_{\mathbb{C}}$, and the action of $G_{\mathbb{C}}$ on \mathcal{J} induces a linear action of P_0 on the fibre \mathcal{J}_o of \mathcal{J} .

Proposition 4.3. *The map $\mathcal{J} \mapsto \mathcal{J}_o$ defines an equivalence between the category of $G_{\mathbb{C}}$ -vector bundles \mathcal{J} on \check{X} and $\mathbf{Rep}_{\mathbb{C}}(P_0)$.*

Proof. This is standard; the vector bundle corresponding to a representation (r, V) of P_0 is $\mathcal{J} = G_{\mathbb{C}} \times V/P_0$.

Remark 4.4. Let $V = \mathcal{J}_o$. If $r: P_0 \rightarrow GL(V)$ extends to a representation (r, V) of G , then the map

$$(g, v) \mapsto [g, r(g)^{-1}v]: G_{\mathbb{C}} \times V \rightarrow \mathcal{J}$$

induces an isomorphism $\check{X} \times V \xrightarrow{\cong} \mathcal{J}$, and so $\mathcal{V}(\mathcal{J})$ is the vector bundle associated with (r, V) (as in § 3).

Recall (2.6) that there is an equivariant commutative diagram

$$\begin{array}{ccc} G(\mathbb{R})_+/K_0 & \hookrightarrow & G(\mathbb{C})/P_0(\mathbb{C}) \\ \downarrow & & \downarrow \\ X & \hookrightarrow & \check{X} \end{array}$$

with K_0 a maximal compact subgroup of $G(\mathbb{R})_+$. If we let $R_u P_0$ denote the unipotent radical of P_0 , then $K_0 \hookrightarrow P_0$ defines an isomorphism $(K_0)_{\mathbb{C}} \xrightarrow{\sim} P_0/R_u P_0$. Now (4.3) has the following corollary.

Corollary 4.5. *There are natural one-to-one correspondences between isomorphism classes of the following objects:*

- (a) $G_{\mathbb{C}}$ -vector bundles \mathcal{F} such that $R_u P_0$ acts trivially on \mathcal{F}_0 ;
- (b) semisimple complex representations of P_0 ;
- (c) representations of $K_{0\mathbb{C}}$ on complex vector spaces;
- (d) representations of K_0 on complex vector spaces.

Proof. A representation of P_0 is semisimple if and only if it is trivial on $R_u P_0$. Therefore the correspondence between (a) and (b) follows from (4.3). We saw above that $K_{0\mathbb{C}} = P_0/R_u P_0$, which proves the correspondence between the objects in (c) and those in (a) and (b). Since K_0 is the compact form of $K_{0\mathbb{C}}$, the correspondence between (c) and (d) is part of Weyl’s unitary trick.

In the next two remarks we show that certain automorphic vector bundles have additional structures.

Remark 4.6. Suppose that \mathcal{F} is a $G_{\mathbb{C}}^{\text{ad}}$ -vector bundle on \check{X} (rather than a $G_{\mathbb{C}}$ -vector bundle). Then at each point $[P_x, \mu_x]$ of \check{X} , μ_x defines a filtration on the fibre \mathcal{F}_x . These filtrations vary holomorphically and define a filtration of $F^p \mathcal{F}$ of \mathcal{F} by $G(\mathbb{C})$ -stable subbundles,

$$\dots F^p \mathcal{F} \supset F^{p+1} \mathcal{F} \dots$$

For each p , $F^p \mathcal{F}$ is the $G^{\text{ad}}(\mathbb{C})$ -vector bundle defined by the subrepresentation of P_0 on $(F^p \mathcal{F})_0$. This filtration induces a holomorphic filtration of $\check{\mathcal{V}}$ by $G^{\text{ad}}(\mathbb{R})^+$ -stable subbundles, and a filtration of \mathcal{V} by $G^{\text{ad}}(\mathbb{Q})^+ \wedge$ -stable algebraic subbundles $F^p \mathcal{V}$. For each x in X , the action of h_x on $\check{\mathcal{V}}_x$ defines a splitting of the filtration,

$$F^p \check{\mathcal{V}}_x = \bigoplus (\check{\mathcal{V}})^{p',q'}, \quad (\check{\mathcal{V}})^{p',q'} = \{v \mid h(z)v = z^{-p'} \bar{z}^{-q'} v\},$$

and these splittings define a decomposition of the C^∞ -bundle $\check{\mathcal{V}}^\infty \stackrel{df}{=} \mathcal{O}_{X^\infty} \otimes \mathcal{V}$,

$$\check{\mathcal{V}}^\infty \xrightarrow{\sim} \bigoplus (\check{\mathcal{V}})^{p,q}.$$

(The splitting is C^∞ because $(\check{\mathcal{V}})^{p,q} = F^p \cap \bar{F}^q$, and the filtration \bar{F}^\bullet is anti-holomorphic.)

Remark 4.7. In the case that \mathcal{F} arises from a representation of G (rather than of P_0), it is possible to define local systems underlying the automorphic vector bundles. Before explaining this, we recall the correspondence between local systems and vector bundles with flat connection (see for example Deligne (1970), I.2). Let S be a connected complex manifold and consider the tensor category of pairs (\mathcal{V}, ∇) with \mathcal{V} an analytic vector bundle on S and ∇ a flat connection.

The map $(\mathcal{V}, \nabla) \mapsto \mathbf{V} \stackrel{df}{=} \mathcal{V}^\nabla$ defines an equivalence between this category and

the tensor category of local systems of complex vector spaces on the manifold; it has quasi-inverse $V \mapsto (\mathcal{O}_S \otimes_{\mathbb{C}} V, \nabla_{\text{can}})$.

A complex representation $r: G_{\mathbb{C}} \rightarrow GL(V)$ of G defines a $G(\mathbb{C})$ -vector bundle $\mathcal{J} = V \times \tilde{X}$ over \tilde{X} with a canonical flat connection ∇ . The flat connection defines a similar connection on $\tilde{\mathcal{V}} = \beta^* \mathcal{J} (= V \times X)$ and on the automorphic vector bundle \mathcal{V} . In particular, we get a local system of complex vector spaces $\mathbf{V} (= \mathcal{V}^{\nabla})$ on $S^0(G, X)$, stable under the action of $G(\mathbb{Q})_{\pm}$ on \mathcal{V} , and such that $\mathcal{O}_{S^0} \otimes_{\mathbb{C}} \mathbf{V} = \mathcal{V}$.

Suppose \mathcal{J} is defined by a real representation (r, V) of $G_{\mathbb{R}}$. Then for each $\Gamma \in \tilde{\Sigma}(G, X)$, the representation $(r|_{\Gamma}, V)$ defines a local system of real vector spaces \mathbf{V}_{Γ} on $\Gamma \backslash X$ such that $\mathcal{O} \otimes_{\mathbb{R}} \mathbf{V}_{\Gamma} = \mathcal{V}^{\nabla}$. The action of $G(\mathbb{Q})_{\pm}$ on $S^0(G, X)$ lifts to an action on the projective system $\mathbf{V} = (\mathbf{V}_{\Gamma})$. When r factors through G^{ad} , the constant sheaf defined by V on X is (in a natural way) a polarizable variation of real Hodge structures spaces (see (2.5)), and so \mathbf{V} acquires a similar structure.

Finally if \mathcal{J} arises from a representation of G^{ad} on a \mathbb{Q} -vector space V , then we get a variation of rational Hodge structures \mathbf{V} on $S^0(G, X)$ such that $\mathbf{V} \otimes \mathbb{C} = \mathcal{V}^{\nabla}$. In this case we can also define, for each prime l , a local system of \mathbb{Q}_l -vector spaces $\mathbf{V}_l = (\mathbf{V}_{l,\Gamma})$ for the étale topology: $\mathbf{V}_{l,\Gamma}$ is the locally constant sheaf on S^0_{Γ} associated with the representation of Γ on $\mathbb{Q}_l \otimes V$ (see for example Milne (1980), p. 165). Symbolically, we may write $\mathbf{V}_l = \mathbb{Q}_l \otimes \mathbf{V}$. There is again a natural action of $G^{\text{ad}}(\mathbb{Q})^+ \wedge$ on \mathbf{V} and \mathbf{V}_l .

Remark 4.8. The construction of \mathcal{V} (and the extra structure on it) is functorial in (G, X, \mathcal{J}) . Moreover, if \mathcal{J} and \mathcal{J}' are two $G_{\mathbb{C}}$ -equivariant vector bundles on \tilde{X} , then an equivariant differential operator $\delta: \mathcal{J} \rightarrow \mathcal{J}'$ gives rise (in a canonical way) to an equivariant differential operator $\gamma: \mathcal{V} \rightarrow \mathcal{V}'$. (See Grothendieck (1967), IV.16.8) for the definition and basic formalism of differential operators on sheaves of \mathcal{O}_S -modules for S a scheme).

Remark 4.9. We outline the relation between the above definition of automorphic forms and the more usual definition in terms of automorphy factors.

Fix a point $o \in X$, and let $\tilde{\mathcal{V}}$ be the $G(\mathbb{R})_+$ -vector bundle on X defined by a $G_{\mathbb{C}}$ -vector bundle \mathcal{J} on \tilde{X} . Choose a trivialization $\alpha: V \times X \xrightarrow{\sim} \tilde{\mathcal{V}}$, and write $\gamma(\alpha(v, x)) = \alpha(j(\gamma, x)v, \gamma x)$ for $\gamma \in G(\mathbb{R})_+$, $v \in V$, and $x \in X$. Then $j: G(\mathbb{R})_+ \times X \rightarrow GL(V)$ is a holomorphic automorphy factor for (G, X) with values in V , and an automorphic form of level Γ and type \mathcal{J} can be identified (through α) with an automorphic form of level Γ for j .

Conversely, let j be a holomorphic automorphy factor for (G, X) with values in V . The map $r: K_0 \rightarrow GL(V)$, $kt \rightarrow j(k, o)$, is a representation of K on V , which (see 4.5) defines an automorphic vector bundle \mathcal{J} on \tilde{X} . If r is irreducible, then it is known that automorphic forms of level Γ for j correspond to automorphic forms of level Γ and type \mathcal{J} (see Murakami (1966), p. 137).

5. Conjugates of automorphic vector bundles

In this section, we state the main theorem for automorphic vector bundles and show how to deduce it from Theorem 3.10.

Let \mathcal{J} be a $G_{\mathbb{Q}}$ -vector bundle on \check{X} , and fix a special point x of X . The $G_{\mathbb{Q}}$ -vector bundle $\tau\mathcal{J}$ on $\tau\check{X}$ corresponds under the isomorphism $\varphi_{\tau,x}^{\check{X}}: \tau\check{X} \rightarrow {}^{\tau,x}\check{X}$ of (2.7) to a ${}^{\tau,x}G_{\mathbb{Q}}$ -vector bundle ${}^{\tau,x}\mathcal{J}$ on ${}^{\tau,x}\check{X}$, and ${}^{\tau,x}\mathcal{J}$ defines an automorphic vector bundle ${}^{\tau,x}\mathcal{V}$ on $S^0({}^{\tau,x}G, {}^{\tau,x}X)$.

Lemma 5.1. *If x' is a second special point of X , then there is a canonical isomorphism $\varphi^{\mathcal{V}}(\tau; x', x): {}^{\tau,x'}\mathcal{V} \rightarrow {}^{\tau,x}\mathcal{V}$ lying over $\varphi^0(\tau; x', x)$ and such that*

$$\varphi^{\mathcal{V}}(\tau; x', x) \circ ({}^{\tau,x}g) = ({}^{\tau,x'}g) \circ \varphi^{\mathcal{V}}(\tau; x', x), \quad \text{all } g \in {}^{\tau,x}\tilde{G}(\mathbb{A}^f).$$

Proof. From the commutative diagram in (2.8) we see that there is a canonical isomorphism ${}^{\tau,x}\mathcal{J} \xrightarrow{\approx} {}^{\tau,x'}\mathcal{J}$ lying over $\check{\varphi}(\tau; x', x): {}^{\tau,x}\check{X} \rightarrow {}^{\tau,x'}\check{X}$, and (4.2) and (3.9) show that this gives rise to the required isomorphism.

Theorem 5.2. *Let \mathcal{V} be an automorphic vector bundle on $S^0(G, X)$. There is a canonical isomorphism $\varphi_{\tau,x}^{\mathcal{V}}: \tau\mathcal{V} \rightarrow {}^{\tau,x}\mathcal{V}$ such that*

$$\begin{array}{ccc} \varphi_{\tau,x}^{\mathcal{V}}: & \tau\mathcal{V} & \rightarrow & {}^{\tau,x}\mathcal{V} \\ & \downarrow & & \downarrow \\ \varphi_{\tau,x}^0: & \tau S^0(G, X) & \rightarrow & S^0({}^{\tau,x}G, {}^{\tau,x}X) \end{array}$$

commutes and $\varphi_{\tau,x}^{\mathcal{V}} \circ \tau(g) = ({}^{\tau,x}g) \circ \varphi_{\tau,x}^{\mathcal{V}}$ for all $g \in G(\mathbb{Q})^-$; moreover, if x' is a second special point of \check{X} , then $\varphi^{\mathcal{V}}(\tau; x', x) \circ \varphi_{\tau,x}^{\mathcal{V}} = \varphi_{\tau,x'}^{\mathcal{V}}$.

Proof. From (3.11) we have a commutative diagram

$$\begin{array}{ccc} \tau\check{X} & \xrightarrow{\varphi_{\tau,x}^{\check{X}}} & {}^{\tau,x}\check{X} \\ \uparrow_{\tau\gamma} & & \uparrow_{\gamma} \\ \tau Y^0(G, X) & \xrightarrow{\varphi_{\tau,x}^{\check{Y}}} & Y^0({}^{\tau,x}G, {}^{\tau,x}X) \\ \downarrow & & \downarrow \\ \tau S^0(G, X) & \xrightarrow{\varphi_{\tau,x}^0} & S^0({}^{\tau,x}G, {}^{\tau,x}X), \end{array}$$

and from the very definition of ${}^{\tau,x}\mathcal{J}$, there is a commutative diagram

$$\begin{array}{ccc} \tau\mathcal{J} & \xrightarrow{\approx} & {}^{\tau,x}\mathcal{J} \\ \downarrow & & \downarrow \\ \tau\check{X} & \xrightarrow{\approx} & {}^{\tau,x}\check{X}. \end{array}$$

On pulling this back, we obtain a similar diagram

$$\begin{array}{ccc} \tau\gamma^*(\mathcal{J}) & \xrightarrow{\approx} & \gamma^*({}^{\tau,x}\mathcal{J}) \\ \downarrow & & \downarrow \\ \tau Y^0(G, X) & \xrightarrow{\approx} & Y^0({}^{\tau,x}G, {}^{\tau,x}X). \end{array}$$

Now (4.2) proves that this gives rise to canonical commutative diagram as in the statement of the theorem. Moreover it is clear from (3.10(ii)) that the map $\varphi_{\tau,x}^{\mathcal{V}}$ commutes with the actions of the Hecke operators, and it follows from (3.11) that $\varphi^{\mathcal{V}}(\tau; x', x) \circ \varphi_{\tau,x}^{\mathcal{V}} = \varphi_{\tau,x'}^{\mathcal{V}}$.

Remark 5.3. The map $\varphi_{\tau,x}^{\mathcal{V}}$ is functorial: suppose we are given compatible maps $(G, X) \rightarrow (G', X')$ and $\mathcal{J} \rightarrow \mathcal{J}'$; if a special point x of X is mapped to a special point x' of X' , then we obtain a commutative diagram

$$\begin{array}{ccc} \varphi_{\tau,x}^{\mathcal{V}}: & \tau\mathcal{V} & \rightarrow & {}^{\tau,x}\mathcal{V} \\ & \downarrow & & \downarrow \\ \varphi_{\tau,x'}^{\mathcal{V}}: & \tau\mathcal{V}' & \rightarrow & {}^{\tau,x'}\mathcal{V}' \end{array}$$

Remark 5.4. When \mathcal{J} is a $G_{\mathbb{C}}^{\text{ad}}$ -vector bundle on \check{X} (rather than a $G_{\mathbb{C}}$ -vector bundle), it is possible to give a more direct definition of ${}^{\tau,x}\mathcal{J}$ (hence of ${}^{\tau,x}\mathcal{V}$) and a characterization of $\varphi_{\tau,x}^{\mathcal{V}}$.

(a) The vector bundle \mathcal{J} corresponds (by 4.3) to a representation (r, V) of P_x , the subgroup of $G_{\mathbb{C}}^{\text{ad}}$ fixing $\beta(x)$. On applying τ we obtain a representation τr of τP_x on V . As in (2.8a), we can use ${}^{\tau}\mathfrak{S}$ to twist $\tau r: \tau P_x \rightarrow GL(V)$ and obtain a representation ${}^{\tau,x}r: {}^{\tau,x}P_x \rightarrow GL({}^{\tau,x}V)$ of ${}^{\tau,x}P_x$. But (see 2.8a), ${}^{\tau,x}P_x$ is the subgroup of ${}^{\tau,x}G(\mathbb{C})$ fixing ${}^{\tau,x}$, and so ${}^{\tau,x}r$ gives rise to a ${}^{\tau,x}G_{\mathbb{C}}$ -vector bundle \mathcal{J}' on \check{X} . Clearly $\mathcal{J}' = {}^{\tau,x}\mathcal{J}$.

(b) Recall that $\mathcal{V}_{[x]} = \mathcal{J}_x$. The map $\varphi_{\tau,x}^{\mathcal{V}}$ is the unique isomorphism $\tau\mathcal{V} \rightarrow {}^{\tau,x}\mathcal{V}$ lying over $\varphi_{\tau,x}$ and such that

(i) the action of $\varphi_{\tau,x}^{\mathcal{V}}$ on the fibre over $\tau[x]$ can be identified with the map $\tau(V \otimes \mathbb{C}) \rightarrow {}^{\tau,x}V \otimes \mathbb{C}$ defined by $z_{\infty}(\tau)$;

(ii) $\varphi_{\tau,x}^{\mathcal{V}} \circ \tau(g) = ({}^{\tau,x}g) \circ \varphi_{\tau,x}^{\mathcal{V}}$ for all $g \in G(\mathbb{Q})_{\bar{\cdot}}$.

Moreover $\varphi_{\tau,x}^{\mathcal{V}}$ carries the natural filtration on $\tau\mathcal{V}$ (see 4.6) into the natural filtration on ${}^{\tau,x}\mathcal{V}$.

We clarify the condition (i). The representation $r \circ \rho_x$ on V defines a CM motive M over $\bar{\mathbb{Q}}$, and $H_B(M) = V$. The composite $c(\tau\iota M)^{-1} \circ (1 \otimes \tau) \circ c(\iota M)$ is a τ -linear map $H_B(\iota M) \otimes \mathbb{C} \rightarrow H_B(\tau\iota M) \otimes \mathbb{C}$ (see 1.2b), and so it defines a linear map $\tau(H_B(M) \otimes \mathbb{C}) \rightarrow H_B(\tau\iota M) \otimes \mathbb{C}$, which we know is $z_{\infty}(\tau)$. But $\tau(H_B(M) \otimes \mathbb{C}) = \tau(V \otimes \mathbb{C})$ and (see 1.8) $H_B(\tau\iota M) = {}^{\tau,x}V$, and so $z_{\infty}(\tau)$ can be regarded as a map

$$\tau(\mathcal{V}_x) = \tau(V \otimes \mathbb{C}) \rightarrow {}^{\tau,x}V \otimes \mathbb{C} = {}^{\tau,x}\mathcal{V}_{\tau x}.$$

[It would be possible to state similar improvements of (5.2) for all automorphic vector bundles if we had a satisfactory theory of *fractional CM-motives* over \mathbb{Q} .]

Remark 5.4. For some automorphic vector bundles $\mathcal{V}(\mathcal{J})$ it is possible to prove (5.2) without using (3.10). For example, let \mathcal{J} be $\mathcal{T}_{\check{X}}$, the tangent bundle to \check{X} ; for any $o \in \check{X}$, \mathcal{J} corresponds to the adjoint representation of P_o on $\text{Lie}(P_o)$. Then $\mathcal{V}(\mathcal{J})$ is the tangent bundle to $S^0(G, X)$, and ${}^{\tau,x}\mathcal{J}$ is the tangent bundle to ${}^{\tau,x}\check{X}$. Therefore the isomorphism of pro-varieties $\varphi_{\tau,x}: \tau S^0(G, X) \rightarrow S^0({}^{\tau,x}G, {}^{\tau,x}X)$ defines an equivariant isomorphism $\mathcal{T}(\varphi_{\tau,x}): \tau\mathcal{V}(\mathcal{J}) \xrightarrow{\sim} \mathcal{V}({}^{\tau,x}\mathcal{J})$. To show

that this is the same as the map in (5.2), it suffices to show that it satisfies conditions (i) and (ii) of (5.4). Condition (ii) is obvious, and (i) can be shown by unwinding the various definitions, because $\mathcal{T}_{\bar{x}}$ is the G -vector bundle associated with the adjoint representation of P_0 on $\text{Lie}(G_{\mathbb{C}})/\text{Lie}(P_0)$.

A similar remark applies if \mathcal{J} is the bundle of n -jets of $\mathcal{T}_{\bar{x}}$.

Remark 5.6. When \mathcal{J} arises from a representation $r: G_{\mathbb{C}} \rightarrow GL(V)$ of $G_{\mathbb{C}}$ (rather than of P_0), then $\varphi_{\tau,x}^{\mathcal{V}}$ carries the flat connection $\tau\nabla$ on $\tau\mathcal{V}$ (see 4.7) into the natural flat connection ${}^{\tau,x}\nabla$ on ${}^{\tau,x}\mathcal{V}$. It therefore defines an isomorphism of the local system $\tau\nabla$ on $\tau S^0(G, X)$ (defined by the pair $(\tau\mathcal{V}, \tau\nabla)$) with the local system ${}^{\tau,x}\nabla$ on $S^0({}^{\tau,x}G, {}^{\tau,x}X)$ (defined by the pair $({}^{\tau,x}\mathcal{V}, {}^{\tau,x}\nabla)$).

When \mathcal{J} is defined by a real representation (r, V) of $G_{\mathbb{R}}$, $\varphi_{\tau,x}^{\mathcal{V}}$ defines an isomorphism of variations of real polarizable Hodge structures.

Finally, when \mathcal{J} is defined by a rational representation (r, V) of G , $\varphi_{\tau,x}^{\mathcal{V}}$ defines an isomorphism of variations of rational Hodge structures; moreover, it defines for each prime l an isomorphism of l -adic sheaves.

Remark 5.7. Theorem 5.2 has a more down-to-earth interpretation. Let $\mathcal{V} = \mathcal{V}(\mathcal{J})$ be an automorphic vector bundle on $S^0(G, X)$, and let f be an automorphic form on X of type \mathcal{J} and level Γ with $\Gamma \in \tilde{\Sigma}(G)$. We can regard f as an algebraic section of \mathcal{V}_{Γ} over $S^0_{\Gamma}(G, X)$. Define the congruence subgroup ${}^{\tau,x}\Gamma$ of ${}^{\tau,x}G(\mathbb{Q})$ as in (1.7). Let τ be an automorphism of \mathbb{C} , and let x be a special point of X . Then the theorem associates with f an automorphic form ${}^{\tau,x}f = \tau f \circ (\varphi_{\tau,x}^{\mathcal{V}})^{-1}$ on the Hermitian symmetric domain ${}^{\tau,x}X$ of type ${}^{\tau,x}\mathcal{J}$ and level ${}^{\tau,x}\Gamma$; moreover, ${}^{\tau,x}(f \circ (g)) = {}^{\tau,x}f \circ ({}^{\tau,x}g)$ for all $g \in G(\mathbb{Q})_{\bar{\tau}}$, and ${}^{\tau,x}f([\bar{x}])$ can be related to $\tau(f([\bar{x}]))$ (by means of $z_{\infty}(\tau)$) when \mathcal{J} is a $G_{\mathbb{C}}^{\text{ad}}$ -vector bundle.

6. Proof of Theorem 3.10 for the symplectic group

Let V be a vector space over \mathbb{Q} , and let ψ_0 be a nondegenerate skew-symmetric form on V . The group $G = \text{Sp}(V, \psi_0)$ of automorphisms of V preserving ψ_0 is semisimple, and there is a unique conjugacy class S^+ of homomorphisms $\mathbf{S} \rightarrow G^{\text{ad}}(\mathbb{R})^+$ such that

- (a) (G, S^+) is a pair defining a Shimura variety;
- (b) each h in S^+ is of weight -1 ;
- (c) the symmetric form $\psi_0(v, h(i)v')$ is positive definite for all $h \in S^+$.

Our goal in this section is to prove (3.10) for the pair (G, S^+) , but first we shall prove (5.2) for the automorphic vector bundle defined by the representation of G on V .

Let G_1 be the group of symplectic similitudes $G\text{Sp}(V, \psi_0)$ of (V, ψ_0) , and let S^{\pm} be the Siegel double space in the sense of Deligne (1979), 1.3.1. Then (G_1, S^{\pm}) is a pair defining a Shimura variety, and $(G, S^+) = (G_1, S^{\pm})^+$.

For an abelian variety A , we set $TA = \lim_{\leftarrow} A_m$ (limit over all positive integers m ordered by division), and we set $V^{\mathcal{J}}A = TA \otimes \mathbb{Q}$. Note that $V^{\mathcal{J}}A$ depends only on the isogeny class of A . Consider triples (A, ψ, k) consisting of an abelian

variety A over \mathbb{C} (defined up to isogeny), a polarization ψ of A , and an isomorphism $k: V^f(A) \xrightarrow{\sim} V(\mathbb{A}^f)$ carrying ψ to ψ_0 . We define $\mathfrak{U}(V, \psi_0)$ to be the set of isomorphism classes of triples of this form for which there is an isomorphism of symplectic spaces $(H_1(A, \mathbb{Q}), \psi) \xrightarrow{\sim} (V, \psi_0)$. The group $G_1(\mathbb{A}^f)$ acts on $\mathfrak{U}(V, \psi_0)$ according to the rule:

$$[A, \psi, k]g = [A, \psi, g^{-1}k], \quad g \in G_1(\mathbb{A}^f).$$

Lemma 6.1. *There is a bijection $S(G_1, S^\pm)(\mathbb{C}) \xrightarrow{\sim} \mathfrak{U}(V, \psi_0)$ commuting with the actions of $G(\mathbb{A}^f)$.*

Proof. Corresponding to

$$[x, g] \in S(G_1, S^\pm)(\mathbb{C}) = G_1(\mathbb{Q}) \backslash X \times G(\mathbb{A}^f)$$

we choose A to be the abelian variety (defined up to isogeny) associated with the rational Hodge structure (V, h_x) . Then ψ_0 defines a polarization ψ of A , and we define k to be the composite $V^f(A) \xrightarrow{\sim} V \otimes \mathbb{A}^f \xrightarrow{g^{-1}} V(\mathbb{A}^f)$. (See for example Milne and Shih (1982b), 2.3.)

Let x be a special point of S^+ . Then $h_x: \mathbb{S} \rightarrow G_{1\mathbb{R}}$ defines a map $\rho_x: \mathfrak{S} \rightarrow G_1 = \mathrm{GSp}(V, \psi_0)$, and we can use ρ_x and ${}^v\mathfrak{S}$ to twist (V, ψ_0) . In this way we obtain a new pair $({}^{v,x}V, {}^{v,x}\psi_0)$ with $\mathrm{GSp}({}^{v,x}V, {}^{v,x}\psi_0) = {}^{v,x}G_1$. Let (A, Ψ, k) be the triple associated with $[x, g]$. Then $H_1(\tau A, \mathbb{Q}) = {}^{v,x}V$ and $\tau\psi = {}^{v,x}\psi_0$ (cf. 1.8). The map $\varphi_{\tau,x}: \tau S^0(G, X) \rightarrow S^0({}^{v,x}G, {}^{v,x}X)$ corresponds under the bijective in (6.1) to $\tau[A, \psi, k] \mapsto [\tau A, \tau\psi, {}^v k]$ where ${}^v k$ is the composite of the maps

$$V^f(\tau A) \xrightarrow{\tau^{-1}} V^f(A) \xrightarrow{k} V(\mathbb{A}^f) \xrightarrow{\mathrm{sp}(\tau)} {}^{v,x}V(\mathbb{A}^f).$$

See [ibid., 7.16].

Just as in (4.7), the pair (V, ψ_0) defines a polarized variation of rational Hodge structures (\mathbf{V}, Ψ_0) on $S(G_1, S^\pm)$. We use v to denote a linear dual.

Proposition 6.2. *There is a canonical abelian scheme $\pi: \mathcal{A} \rightarrow S(G_1, S^\pm)$ over $S(G_1, S^\pm)$ and a polarization Ψ of \mathcal{A} such that $((R^1\pi_* \mathbb{Q})^v, \Psi) = (\mathbf{V}, \Psi_0)$.*

Proof. Choose a lattice $V_0 \subset V$ such that ψ_0 is integral on V_0 , and for any integer $N \geq 3$, define $K(N)$ to be the subgroup of $G(\mathbb{A}^f)$ stabilizing $V_0 \otimes \mathbb{A}^f$ and acting as the identity map on V_0/NV_0 . Then $S_{K(N)}(G_1, S^\pm)$ is the moduli scheme over \mathbb{C} for polarized abelian varieties of dimension $\dim(V)/2$ and degree the discriminant of ψ_0 on V_0 , and we can take $\mathcal{A}_{K(N)}$ to be the universal abelian scheme over $S_{K(N)}(G_1, S^\pm)$ (Mumford (1965), 7.9). The subgroups $K(N)$ are cofinal among compact-open subgroups of $G_1(\mathbb{A}^f)$, and we define \mathcal{A} to be the inverse limit, $\lim_{\leftarrow} \mathcal{A}_{K(N)}$.

Remark 6.3. (a) For each point s of $S(G_1, S^\pm)$, the fibre \mathcal{A}_s is the abelian variety attached to s in (6.1).

(b) From \mathcal{A} we get an isomorphism $k_l: (R^1\pi_*\mathbb{Q}_l)^\vee \xrightarrow{\sim} \mathbf{V}(\mathbb{Q}_l)$ for each l , where \mathbf{V} is the rational local system on $S(G_1, S^\pm)$ defined by V . Write $V_l(\mathcal{A})$ for $(R^1\pi_*\mathbb{Q}_l)^\vee$; then we can think of the family $k=(k_l)$, $k_l: V_l(\mathcal{A}) \rightarrow \mathbf{V}(\mathbb{Q}_l)$, as being a level structure on \mathcal{A} . The triple (\mathcal{A}, Ψ, k) has the following universal property: for any $G_1(\mathbb{A}^f)$ -scheme S over \mathbb{C} and triple $(\mathcal{A}', \Psi', k')$ over S such that $(\mathcal{A}', \Psi', k')_s \in \mathfrak{A}(V, \Psi_0)$ for all $s \in S(\mathbb{C})$, there exists a unique isomorphism $\alpha: S \rightarrow S(G_1, S^\pm)$ of $G(\mathbb{A}^f)$ -schemes such that $\alpha^*(\mathcal{A}, \Psi, k) = (\mathcal{A}', \Psi', k')$.

This can be proved easily using the universal property of each $\mathcal{A}_{K(N)}$.

When we apply the construction in (6.2) and (6.3b) to $({}^{\tau,x}V, {}^{\tau,x}\psi)$, we get a polarized abelian scheme $({}^{\tau,x}\mathcal{A}, {}^{\tau,x}\Psi)$ over $S({}^{\tau,x}G_1, S^\pm)$ with a level structure ${}^{\tau,x}k$, where ${}^{\tau,x}k_l$ is an isomorphism $V_l(\mathcal{A}) \xrightarrow{\sim} {}^{\tau,x}\mathbf{V}(\mathbb{Q})$.

Proposition 6.4. *There is a unique isomorphism $\varphi_{\tau,x}^{\mathcal{A}}: \tau\mathcal{A} \rightarrow {}^{\tau,x}\mathcal{A}$ such that*
 (a) *the following diagram commutes*

$$\begin{array}{ccc} \tau\mathcal{A} & \xrightarrow{\varphi_{\tau,x}^{\mathcal{A}}} & {}^{\tau,x}\mathcal{A} \\ \downarrow & & \downarrow \\ \tau S(G_1, S^\pm) & \xrightarrow{\varphi_{\tau,x}} & S({}^{\tau,x}G_1, S^\pm); \end{array}$$

- (b) $\varphi_{\tau,x}^{\mathcal{A}}$ sends $\tau\Psi$ to ${}^{\tau,x}\Psi$;
- (c) *the following diagram commutes*

$$\begin{array}{ccccc} V^f(\tau\mathcal{A}) & \xrightarrow{\tau^{-1}} & V^f(\mathcal{A}) & \xrightarrow{k} & \mathbf{V}(\mathbb{A}^f) \\ \downarrow \varphi_{\tau,x}^{\mathcal{A}} & & & & \downarrow \text{sp}(\tau) \\ V^f({}^{\tau,x}\mathcal{A}) & \xrightarrow{{}^{\tau,x}k} & & & {}^{\tau,x}\mathbf{V}(\mathbb{A}^f). \end{array}$$

Proof. Apply the universal property of ${}^{\tau,x}\mathcal{A}$ (and use that $V \otimes \mathbb{A}^f = {}^{\tau,x}V \otimes \mathbb{A}^f$).

Corollary 6.5. *There is a unique isomorphism $\varphi_{\tau,x}^{\mathcal{V}}: \tau\mathcal{V} \xrightarrow{\sim} {}^{\tau,x}\mathcal{V}$ lying over $\varphi_{\tau,x}$ and such that*

- (i) *the restriction of $\varphi_{\tau,x}^{\mathcal{V}}$ to the fibre over $[x, 1]$ can be identified with the map $\tau V(\mathbb{C}) \rightarrow {}^{\tau,x}V(\mathbb{C})$ defined by $z_\infty(\tau)$;*
- (ii) $\varphi_{\tau,x}^{\mathcal{V}} \circ \tau(g) = ({}^{\tau,x}g) \circ \varphi_{\tau,x}^{\mathcal{V}}$ for all $g \in G_1(\mathbb{A}^f)$.

Proof. The uniqueness is obvious. For the existence, note that $\mathcal{V} = \mathcal{H}_1^{dR}(\tau\mathcal{A})$ and ${}^{\tau,x}\mathcal{V} = \mathcal{H}_1^{dR}({}^{\tau,x}\mathcal{A})$, and so $\varphi_{\tau,x}^{\mathcal{V}}$ can be taken to be $\mathcal{H}_1^{dR}(\varphi_{\tau,x}^{\mathcal{A}})$.

Corollary 6.6. *Theorem 5.2 is true for the automorphic vector bundle \mathcal{V} defined by the representation of G on V .*

Proof. The restriction of $\varphi_{\tau,x}^{\mathcal{V}}$ to $\tau S^0(G, X)$ has the correct properties. For a second special point x' , it is not difficult to trace through the various constructions and see that $\varphi^{\mathcal{V}}(\tau; x', x) \circ \varphi_{\tau,x}^{\mathcal{V}} = \varphi_{\tau,x'}$.

Corollary 6.7. *Theorem 3.10 is true for (G, S^+) .*

Proof. This follows from (6.6) because (3.4) allows us to identify $Y^0(G, S^+)$ and $Y^0({}^{\tau,x}G, {}^{\tau,x}S^+)$ respectively with the spaces of trivializations of \mathcal{V} and ${}^{\tau,x}\mathcal{V}$ preserving the polarizations.

7. Proof of Theorem 3.10 for connected Shimura varieties of abelian type

Lemma 7.1. *Let $f: (G, X) \rightarrow (G', X')$ be an embedding of pairs defining connected Shimura varieties. If Theorem 3.10 is true for (G', X') , then it is also true for (G, X) .*

Proof. Let Y be the inverse image of $\tau Y^0({}^{\tau,x}G, {}^{\tau,x}X)$ under $\phi_{\tau,x}^{Y'}$,

$$\begin{array}{ccc} \tau Y^0(G, X) & \hookrightarrow & Y^0(G', X') \\ & & \downarrow \phi_{\tau,x}^{Y'} \\ \tau Y^0({}^{\tau,x}G, {}^{\tau,x}X) & \hookrightarrow & Y^0({}^{\tau,x}G', {}^{\tau,x}X'). \end{array}$$

Then Y is a closed subset containing τy , and it contains a fibre of the map $\pi: \tau Y^0(G, X) \rightarrow \tau S^0(G, X)$ whenever it contains a single point; moreover, its image in $\tau S^0(G, X)$ is stable under the action of $\mathcal{G}(G)$. The same argument as in the proof of the uniqueness statement in (3.10) now shows that $Y = \tau Y^0(G, X)$. Therefore the restriction of $\phi_{\tau,x}^{Y'}$ to $\tau Y^0(G, X)$ is an isomorphism satisfying the conditions (i)–(iii) of the theorem.

If x' is a second special point of X , then the restriction of $\phi^{Y'}(\tau; x', x)$ to $Y^0({}^{\tau,x}G, {}^{\tau,x}X)$ is $\phi^Y(\tau; x', x)$, and so the equality $\phi^Y(\tau; x', x) \circ \phi_{\tau,x}^Y = \phi_{\tau,x'}^Y$ is implied by the similar equality for $Y(G', X')$.

Recall (Milne and Shih (1982b), § 1, especially 1.3) that if (G, X) is primitive of abelian type, then there is an embedding $(G, X) \hookrightarrow (\mathrm{Sp}(V, \psi), S^+)$ for some symplectic space (V, ψ) . Therefore (6.7) and the lemma and show that Theorem 3.10 holds for every pair (G, X) that is primitive of abelian type. By definition, for every pair (G, X) of abelian type, there exists a family of pairs (G_i, X_i) , primitive of abelian type, and a morphism $(\Pi G_i, \Pi X_i) \rightarrow (G, X)$ with $\Pi G_i \rightarrow G$ an isogeny [ibid. p. 293]. Therefore, the next two lemmas complete the proof of Theorem 3.10 for pairs (G, X) of abelian type.

Lemma 7.2. *Let $(G', X') \rightarrow (G, X)$ be a morphism such that $G' \rightarrow G$ is an isogeny. If Theorem 3.10 is true for (G', X') , then it is also true for (G, X) .*

Proof. Note that in this case $G'^{\mathrm{ad}} = G^{\mathrm{ad}}$ and $X = X'$. Moreover, $S^0(G, X)$ is the quotient of $S^0(G', X)$ by the kernel $\mathcal{G}(G', G)$ of the map $G'^{\mathrm{ad}}(\mathbb{Q})^{+\wedge} \rightarrow G^{\mathrm{ad}}(\mathbb{Q})^{+\wedge}$. Similarly, $Y^0(G, X)$ is the quotient of $Y^0(G', X)$ by the kernel of the surjective homomorphism $\mathcal{G}(G') \rightarrow \mathcal{G}(G)$. The map $\phi_{\tau,x}^{Y'}$ for (G, X) can therefore be obtained from the corresponding map $\phi_{\tau,x}^{Y'}$ for (G', X') by passing to the quotient.

Lemma 7.3. *For $i = 1, \dots, n$, let (G_i, X_i) be a pair defining a connected Shimura variety, and let $G = \Pi G_i$ and $X = \Pi X_i$. If Theorem 3.10 is true for each pair (G_i, X_i) then it is true for (G, X) .*

Proof. This is obvious.

Proposition 7.4. *Theorem 3.10 is true for all connected Shimura varieties of abelian type.*

Proof. We have already noted that this follows from (6.7) and the preceding lemmas.

In fact, in the proof of the general case of (3.10) we shall need to use (7.4) only for connected Shimura varieties associated with groups G of type A_1 (then $\tilde{G} = SL_1(B)$ for B a quaternion algebra defined over a totally real field F).

We now make some remarks that will assist us in the next two sections in the proof of the general case.

Lemmas 7.2 and 7.3 show that it suffices to prove Theorem 3.10 for (G, X) with G simply connected and G^{ad} \mathbb{Q} -simple.

For the remainder of this section, we assume that G is simply connected. Let x be a special point of X . For each homomorphism $r: G(\mathbb{Q}) \rightarrow G(\mathbb{C})$, we define the principal $G_{\mathbb{C}}$ -bundle $Y(r)$ on $S^0(G, X)$ to be

$$Y(r) = G(\mathbb{Q}) \backslash X \times G(\mathbb{C}) \times G(\mathbb{A}^f)$$

where $q \in G(\mathbb{Q})$ acts on $(x, c, a) \in X \times G(\mathbb{C}) \times G(\mathbb{A}^f)$ according to the rule

$$q(x, c, a) = (qx, r(q) \cdot c, qa).$$

There is an obvious flat connection $\nabla(r)$ on $Y(r)$, and an action $a(r)$ of $G(\mathbb{A}^f)$ on $(Y(r), \nabla(r))$. Let $y(r)$ be the point $[x, 1, 1]$ on $Y(r)$.

Our next lemma was suggested by a similar result in Harris (1986), 3.6.

Lemma 7.5. *The map $r \mapsto (Y(r), \nabla(r), a(r), y(r))$ gives a bijection between the set of homomorphisms $r: G(\mathbb{Q}) \rightarrow G(\mathbb{C})$ and the set of isomorphism classes of quadruples (Y, ∇, a, y) consisting of a principal $G_{\mathbb{C}}$ -bundle Y , a flat connection ∇ on Y , an action a of $G(\mathbb{A}^f)$ on (Y, ∇) , and a point y lying over $[x]$.*

Proof. Suppose we are given a quadruple (Y, ∇, a, y) . Because X is simply connected, there is a map $\psi: X \times G(\mathbb{C}) \times G(\mathbb{A}^f) \rightarrow Y$ compatible with the projections to $S^0(G, X)$ (\mathbb{C}), the flat structures, and the actions of $G(\mathbb{A}^f)$ and $G(\mathbb{C})$. When we normalize ψ by requiring that $\psi(x, 1, 1) = y$, then it is uniquely determined. The map r corresponding to (Y, ∇, a, y) is determined by the rule:

$$\psi(qx, 1, q) = r(q)^{-1}y, \quad \text{all } q \in G(\mathbb{Q}).$$

Remark 7.6. Let x be a special point of X . Corresponding to the pair $({}^r x G, {}^r x X)$ and the special point ${}^r x$ of ${}^r x X$, we have a quadruple $(Y^0({}^r x G, {}^r x X), {}^r x \nabla, {}^r x a, {}^r y)$. On pulling back by $\varphi_{r,x}^0$, applying τ^{-1} , and finally pulling back again relative to the map $g \mapsto {}^r x g: G(\mathbb{C}) \rightarrow {}^r x G(\mathbb{C})$, we obtain a similar quadruple on $S^0(G, X)$, which we denote by $({}^x Y, {}^x \nabla, {}^x a, {}^x y)$. According to (7.5), this corresponds to a homomorphism $r_x: G(\mathbb{Q}) \rightarrow G(\mathbb{C})$, and to prove the existence of an isomorphism $\varphi_{r_x, x}^Y$ satisfying (i)–(iii) of Theorem 3.10 it suffices to show that this homomorphism is the natural inclusion.

8. First completion of the proof of Theorem 3.10

Let (G, X) be a pair defining a connected Shimura variety, and assume that G is simply connected and that G^{ad} is \mathbb{Q} -simple. Then $G = \text{Res}_{F/\mathbb{Q}} H$ with H an absolutely almost simple group over a totally real number field F . Let x be a special point of X , and let T be a maximal torus of G such that h_x factors through $(T/Z)(\mathbb{R})$. There is a maximal torus T' of H such that $T = \text{Res}_{F/\mathbb{Q}} T'$.

For any totally real number field F' containing F and such that $T'_{F'}$ splits over a quadratic imaginary extension L of F' , we can write

$$\text{Lie}(H_L) = \text{Lie}(T'_L) \otimes \left(\bigotimes_{\alpha \in R} \text{Lie}(H_L)_\alpha \right)$$

where $R = R(H_{\mathbb{C}}, T'_L)$. Recall that a root α is said to be *totally compact* if it is a compact root of $(H_{F'} \otimes_{F', \sigma} \mathbb{R})$ for all embeddings $\sigma: F' \hookrightarrow \mathbb{R}$. For each root α that is not totally compact, the subgroup H_α is defined to be the connected subgroup of $H_{F'}$ such that

$$\text{Lie}(H_\alpha)_L = \text{Lie}(T'_L) \oplus \text{Lie}(H_L)_\alpha \oplus \text{Lie}(H_L)_{-\alpha}.$$

Proposition 8.1. *If F' is chosen to be sufficiently large, then $H(F)$ is contained in the subgroup of $H(\mathbb{Q})$ generated by $\bigcup H_\alpha(F')$, where the union is taken over all nontotally compact roots α .*

Proof. This is stated (without proof) in Borovoi (1983/84), 3.21.

Choose a field F' as in the proposition, and let $G_* = \text{Res}_{F'/\mathbb{Q}} H$ and, for each nontotally compact root α of G , let $G_\alpha = \text{Res}_{F'/\mathbb{Q}} H_\alpha$. There are natural maps $(G, X) \hookrightarrow (G_*, X_*)$ and $(G_\alpha, X_\alpha) \hookrightarrow (G_*, X_*)$, each α . Write x_* for the image of x in X_* , and write x_α for it regarded as an element of X_α . Then the construction in (7.6) applied to the triples (G_*, X_*, x_*) and $(G_\alpha, X_\alpha, x_\alpha)$ gives us homomorphisms $r_{x_*}: G_*(\mathbb{Q}) \rightarrow G_*(\mathbb{C})$ and $r_{x_\alpha}: G_\alpha(\mathbb{Q}) \rightarrow G_\alpha(\mathbb{C})$. Clearly the maps r_x, r_{x_*} , and r_{x_α} are compatible with the inclusions $G \hookrightarrow G_*$ and $G_\alpha \hookrightarrow G_*$; that is, the following diagram commutes,

$$\begin{array}{ccc} r_x: & G(\mathbb{Q}) & \rightarrow & G(\mathbb{C}) \\ & \downarrow & & \downarrow \\ r_{x_*}: & G_*(\mathbb{Q}) & \rightarrow & G_*(\mathbb{C}) \\ & \uparrow & & \uparrow \\ r_{x_\alpha}: & G_\alpha(\mathbb{Q}) & \rightarrow & G_\alpha(\mathbb{C}) \end{array}$$

Since we know Theorem 3.10 for groups of type A_1 , each map r_{x_α} is the natural inclusion. The proposition shows that $G(\mathbb{Q}) \subset \bigcup G_\alpha(\mathbb{Q})$, and so r_x must also be the natural inclusion.

To complete the proof of (3.10), we have to show that if x' is a second special point, then $\varphi^Y(\tau; x', x) \circ \varphi^Y_{\tau, x} = \varphi^Y_{\tau, x'}$. Each is a map $\tau Y^0(G, X) \rightarrow Y^0(\tau, x' G, \tau, x' X)$ compatible with the homomorphism $g \mapsto \tau, x' g: G(\mathbb{A}') \rightarrow \tau, x' G(\mathbb{A}')$. To prove that the maps are equal, it suffices to show that they agree on a single point (compare

the proof of the uniqueness assertion of (3.10)). If x and x' both lie in X_α for some noncompact root α as above, then the maps agree on the whole of $Y^0(G_\alpha, X_\alpha)$, and so they are equal. If $x' = gx$ for some $g \in G(\mathbb{Q})$, then one can show directly that the two maps agree on the τy , where y is the image of x in $Y^0(G, X)$. To complete the proof, we need to recall one final result.

Proposition 8.2. *Let x and x' be special points of X . If F' is chosen large enough, then there exists a sequence $(x =)x_0, \dots, x_m(=x')$ of special points of X_* and an element $g \in G_*(\mathbb{Q})$ such that*

- (a) *for each positive $i < m$, there is a noncompact root α such that x_{i-1} and x_i lie in X_α ;*
- (b) $g x_{m-1} = x_m$.

Proof. See Borovoi (1983/84), 1.18 or Milne (1983), pp. 260–261.

We now complete the proof for a general pair of special points (x, x') . After replacing (G, X) with a suitable pair (G_*, X_*) , we can assume that there exist points x_0, \dots, x_m and an element g as in the statement of the proposition. The remarks preceding the proposition show that

$$\varphi^Y(\tau; x_i, x_{i-1}) \circ \varphi^Y_{\tau, x_{i-1}} = \varphi^Y_{\tau, x_i} \quad \text{for } i = 1, \dots, m.$$

On multiplying these equalities, we find that

$$\varphi^Y_{\tau, x'} = \varphi^Y(\tau; x_m, x_{m-1}) \circ \dots \circ \varphi^Y(\tau; x_1, x_0) \circ \varphi^Y_{\tau, x}.$$

It is clear from the definition of the maps $\varphi^Y(\tau; \dots)$ (see 3.7) that

$$\varphi^Y(\tau; x_m, x_{m-1}) \circ \dots \circ \varphi^Y(\tau; x_1, x_0) = \varphi^Y(\tau; x_m, x_0)$$

and so this completes the proof of Theorem 3.10 (assuming 8.1).

9. Second completion of the proof of Theorem 3.10

Again we assume that (G, X) is a pair defining a connected Shimura variety with G simply connected and G^{ad} \mathbb{Q} -simple. Let x be a special point X , and let $r_x: G(\mathbb{Q}) \rightarrow G(\mathbb{C})$ be the map defined in (7.6). If we can show that r_x is the natural inclusion map, then the same argument as in the previous section will complete the proof of Theorem 3.10.

Now let $r: G^{\text{ad}} \rightarrow GL(V)$ be a representation of G^{ad} . We can twist this to obtain a representation ${}^{\tau, x}r: {}^{\tau, x}G^{\text{ad}} \rightarrow GL({}^{\tau, x}V)$. This defines the automorphic vector bundle ${}^{\tau, x}\mathcal{V}$ on $S^0({}^{\tau, x}G, {}^{\tau, x}X)$ together with its flat connection ${}^{\tau, x}\nabla$ and action ${}^{\tau, x}a$ of ${}^{\tau, x}G(\mathbb{A}^f)$. On pulling this back by $\varphi_{\tau, x}^0$ to $\tau S^0(G, X)$ and applying τ^{-1} , we obtain a vector bundle ${}^x\mathcal{V}$ on $S^0(G, X)$ together with a flat connection ${}^x\nabla$ and action ${}^x a$ of $G(\mathbb{A}^f)$. From the triple $({}^x\mathcal{V}, {}^x\nabla, {}^x a)$ we obtain a representation $r_x(V)$ of $G(\mathbb{Q})$ on $V \otimes \mathbb{C}$. Note that Theorem 5.2 would imply that $r_x(V)$ is simply the map induced by r .

Lemma 9.1. *Let $r: G \rightarrow GL(V)$ be a representation of G factoring through $G_{\mathbb{C}}^{\text{ad}}$. With the above notations, there is a commutative diagram*

$$\begin{array}{ccc} G(\mathbb{Q}) & \xrightarrow{r_x} & G(\mathbb{C}) \\ r_x(V) & \searrow & \downarrow r \\ & & GL(V \otimes \mathbb{C}). \end{array}$$

Proof. This simply says that the above construction of $r_x(V)$ is compatible with the construction in (7.6) of r_x .

Lemma 9.2. *Consider the representation r of G^{ad} on $\mathfrak{g} \stackrel{\text{df}}{=} \text{Lie}(G)$. If for all triples (G, X, x) , $r_x(\mathfrak{g})$ is the restriction of r to $G^{\text{ad}}(\mathbb{Q})$, then $r_x: G(\mathbb{Q}) \rightarrow G(\mathbb{C})$ is the natural inclusion.*

Proof. From the preceding lemma, we get a commutative diagram

$$\begin{array}{ccc} G(\mathbb{Q}) & \xrightarrow{r_x} & G(\mathbb{C}) \\ \downarrow & & \downarrow \\ G^{\text{ad}}(\mathbb{Q}) & \xrightarrow{\text{nat}} & G^{\text{ad}}(\mathbb{C}) \end{array}$$

in which the map in the lower row is the natural inclusion. This implies that the map $\gamma: \mathfrak{g} \rightarrow r_x(\mathfrak{g}) \cdot \mathfrak{g}^{-1}$ has image contained in the centre $Z(\mathbb{C})$ of $G(\mathbb{C})$. It is moreover a homomorphism. If we knew that $G(\mathbb{Q})/Z(\mathbb{Q})$ was a simple group the proof would now be complete because we would have that $\gamma(G(\mathbb{Q})) = \{1\}$. Since we do not know it, we instead must argue as in Milne (1983), p. 251–252. Fix a $g \in G(\mathbb{Q})$. Because $G(\mathbb{Q})$ is generated by the groups $T_{x'}(\mathbb{Q})$ with x' running through the special points of X [ibid. 3.10], we can assume that g lies in some $T_{x'}(\mathbb{Q})$. If we write $G = \text{Res}_{F/\mathbb{Q}} H$ and pass to a larger totally real field F' , then we obtain inclusions $(G, X) \hookrightarrow (G_*, X_*)$ and $G_x \hookrightarrow G_*$ with G_x a reductive group of type A_1 and $T_{x'} \subset G_x$. Moreover, after possibly enlarging the totally real field F again, we can assume that $G_x \otimes \mathbb{Q}_l$ is isotropic. On applying the above construction to the pair (G_*, X_*) we obtain a map $\gamma_*: G_*(\mathbb{Q}) \rightarrow Z_*(\mathbb{C})$ whose restriction to $G(\mathbb{Q})$ is γ . According to Platonov and Rapincuk (1979), $G_x(\mathbb{Q})/Z_x(\mathbb{Q})$ is simple, and so γ_* is trivial on $G_x(\mathbb{Q})$. As $g \in G_x(\mathbb{Q})$, this shows that $\gamma(g) = 1$.

It remains to verify that $r_x(\mathfrak{g}): G^{\text{ad}}(\mathbb{Q}) \rightarrow GL(\mathfrak{g})$ is given by the adjoint representation. Recall that $S^0(G, X) = G(\mathbb{Q}) \backslash X \times G(\mathbb{A}^f)$, and write π for the projection $X \times G(\mathbb{A}^f) \rightarrow S^0(G, X)$. For any $G(\mathbb{A}^f)$ -local system \mathbf{V} on $S^0(G, X)$, $\pi^*(\mathbf{V})$ is spanned by its space of $G(\mathbb{A}^f)$ -invariant global sections, and this last space can be identified with \mathbf{V}_x for any $x \in X$. Therefore, from any such local system and point x , we obtain a representation $G(\mathbb{Q}) \rightarrow GL(\mathbf{V}_x)$. Because of the dictionary recalled in (4.7), a $G(\mathbb{A}^f)$ -vector bundle with flat connection (\mathcal{V}, ∇) also gives rise to a representation of $G(\mathbb{Q})$ on the fibre \mathcal{V}_x .

Lemma 9.3. *Let (\mathcal{V}, ∇) be a $G(\mathbb{A}^f)$ -vector bundle with flat connection on $S^0(G, X)$, and assume that \mathcal{V} is a $G(\mathbb{A}^f)$ -subbundle of an automorphic vector bundle $\mathcal{V}(\mathcal{J})$ on $S^0(G, X)$. Then the representation $r: G(\mathbb{Q}) \rightarrow GL(\mathcal{V}_x)$ is continuous for the complex topology; if it is trivial on $Z(\mathbb{Q})$, then it is induced by a morphism of algebraic groups $G_{\mathbb{Q}}^{\text{ad}} \rightarrow GL(\mathcal{V}_x)$.*

Proof. The first assertion follows from the fact that the representation r of $G(\mathbb{Q})$ on \mathcal{V}_x can be realized as a subrepresentation of the representation of $G(\mathbb{R})$ on $\Gamma(X, \beta^*(\mathcal{J}))$, which is obviously continuous. For the second, note that the homomorphism $r^{\text{ad}}: G(\mathbb{Q})/Z(\mathbb{Q}) \rightarrow GL(\mathcal{V}_x)$ extends by continuity to a homomorphism $\hat{r}: G^{\text{ad}}(\mathbb{R})^+ \rightarrow GL(\mathcal{V}_x)$. Let r' be the homomorphism of algebraic groups $G_{\mathbb{R}}^{\text{ad}} \rightarrow GL(\mathcal{V}_x)$ such that $\text{Lie}(r') = \text{Lie}(\hat{r})$. Then $r'(\mathbb{R})$ and \hat{r} agree on an open subgroup of $G^{\text{ad}}(\mathbb{R})^+$, and so they agree on the whole of $G^{\text{ad}}(\mathbb{R})^+$.

For a vector bundle \mathcal{V} on a space, write $\mathcal{J}el^n(\mathcal{V})$ for the bundle of jets of length n of \mathcal{V} (see Grothendieck (1967), IV.16.8).

Lemma 9.4. *Let \mathcal{J} be the $G_{\mathbb{Q}}$ -vector bundle on \check{X} corresponding to the adjoint representation of G on $\text{Lie}(G)$. Then there is an equivariant embedding $\mathcal{J} \hookrightarrow \mathcal{J}el^2(T_{\check{X}})$.*

Proof. See Harris (1985), p. 172.

We now complete the proof of the theorem. Let x be a special point of X . After replacing (G, X) by a pair (G_*, X_*) and applying (7.1), we may assume that T_x splits over a quadratic extension of F . On applying (9.1) and (9.3) to the automorphic vector bundle \mathcal{V} defined by the adjoint representation, we find that the map $r_x(\mathfrak{g}): G(\mathbb{Q}) \rightarrow GL(\mathfrak{g})$ is defined by an algebraic map ρ . Since we know the theorem for each H_{α} , the restriction of ρ to the algebraic subgroup H_{α} of G is given by the adjoint representation, and the next lemma shows that this implies that ρ itself is given by the adjoint representation.

Lemma 9.5. *The group G is generated (as an algebraic group) by the subgroups H_{α} .*

Proof. This follows from the fact that, for any embedding σ of F into \mathbb{R} , $[\mathfrak{p}_{\sigma}, \mathfrak{p}_{\sigma}] = \mathfrak{k}_{\sigma}$, where $\mathfrak{g}_{\sigma} = \mathfrak{k}_{\sigma} \oplus \mathfrak{p}_{\sigma}$ is a Cartan decomposition of $\mathfrak{g}_{\sigma} \stackrel{\text{def}}{=} \text{Lie}(G \otimes_{F, \sigma} \mathbb{R})$.

This completes the second proof of Theorem 3.10.

Appendix: Pairs defining connected and nonconnected Shimura varieties

In this section, we review some results that allow one to modify the centres of the groups defining Shimura varieties. Throughout, ι denotes complex conjugation.

Lemma A.1. *Let G be a semisimple adjoint group over a field of characteristic zero, and let L be a finite Galois extension of k that splits G . For any finite central covering $G' \rightarrow G$ of G , there exists a central extension defined over k*

$$1 \rightarrow N \rightarrow G_1 \rightarrow G \rightarrow 1$$

such that $G_1^{\text{der}} = G'$ and N is equal to a product of copies of $\text{Res}_{L/k} \mathbb{G}_m$.

Proof. See for example Milne and Shih (1982b), 3.1.

Proposition A.2. *For any pair (G, X) defining a connected Shimura variety, there exists a pair (G_1, X_1) defining a Shimura variety and such that*

- (a) $(G_1, X_1)^+ = (G, X)$;
- (b) the weight w_X of any $h \in X_1$ is defined over \mathbb{Q} ;
- (c) the centre $Z(G_1)$ of G_1 is a product of copies of $\text{Res}_{L/\mathbb{Q}} \mathbb{G}_m$ for some CM-field L Galois over \mathbb{Q} .

Proof. Let x be a special point of X , and let T be a maximal torus of G^{ad} such that h factors through $T_{\mathbb{R}}$. Then $T_{\mathbb{R}}$ is anisotropic, and so ι acts as -1 on $X^*(T)$. It follows that, for any $\tau \in \text{Aut}(\mathbb{C})$, $\tau \iota$ and $\iota \tau$ have the same action on $X^*(T)$, and so T splits over a CM-field L , which can be chosen to be Galois over \mathbb{Q} . Construct G_1 as in the lemma with $G_1^{\text{der}} = G$ and $G_1/N = G^{\text{ad}}$. The inverse image T_1 of T in G_1 is a maximal torus. Choose $\mu_1 \in X_*(T_1)$ to lift $\mu \in X_*(T)$. The weight $w_1 \stackrel{\text{df}}{=} -\mu_1 - \iota \mu_1$ of μ_1 lies in $X_*(Z_1)$, where $Z_1 = Z(G_1) = N$. Clearly $\iota w_1 = w_1$ and so, as $H^1(\mathbb{R}, X_*(Z_1)) = 0$, there exists a $\mu_0 \in X_*(Z_1)$ such that $(\iota + 1)\mu_0 = w_1$. When we replace μ_1 by $\mu_1 - \mu_0$ we find that $w_1 = 0$; in particular, w_1 is defined over \mathbb{Q} . Let $h_1: \mathbb{S} \rightarrow G_1$ correspond to μ_1 , and let X_1 be the conjugacy class of h_1 . Then (G_1, X_1) fulfills the requirements.

Now let (G, X) be a pair defining a Shimura variety. Recall that $x \in X$ is special if h_x factors through a \mathbb{Q} -rational torus $T \subset G$. We say that x is a CM-point if there exists a \mathbb{Q} -rational homomorphism $\rho_x: \mathbb{S} \rightarrow G$ such that $\mu_{\text{can}} \circ \rho_x = \mu_x$. Clearly a CM-point is special, and a special point is CM if and only if

$$(\tau - 1)(\iota + 1)\mu_x = 0 = (\iota + 1)(\tau - 1)\mu_x$$

all $\tau \in \text{Aut}(\mathbb{C})$. (Note, *pace* Deligne (1979), 2.2.4, it is important to distinguish these notions.)

Proposition A.3. *Every special point of X is a CM-point when (G, X) satisfies the conditions:*

- (a) the weight w_X is defined over \mathbb{Q} ;
- (b) the centre Z of G is split by a CM-field.

Proof. The conditions say respectively that

$$\begin{aligned} (\tau - 1)(\iota + 1)\mu_x &= 0, & \text{for all } \tau \in \text{Aut}(\mathbb{C}), \quad x \in X; \\ \tau \iota \mu &= \iota \tau \mu, & \text{for all } \tau \in \text{Aut}(\mathbb{C}), \quad \mu \in X_*(Z). \end{aligned}$$

Let $x \in X$ be a special, and let T be a maximal torus through which h_x factors. The argument in the proof of (A.2) shows that $\tau \iota \mu = \iota \tau \mu$ for $\mu \in X_*(T/Z)$, and since

$$X_*(T) \otimes \mathbb{Q} = X_*(Z) \otimes \mathbb{Q} \oplus X_*(T/Z) \otimes \mathbb{Q}$$

we see that the same equation holds for $\mu \in X_*(T)$. Therefore $(\iota + 1)(\tau - 1)\mu_x = (\tau - 1)(\iota + 1)\mu_x$, and we have already observed that this is zero.

Corollary A.4. *For any pair (G, X) defining a connected Shimura variety, there exists a pair (G_1, X_1) defining a Shimura variety and such that*

- (a) $(G_1, X_1)^+ = (G, X)$;
- (b) every special point of X_1 is CM;
- (c) $H^1(k, Z(G_1)) = 0$ for all fields k of characteristic zero.

Proof. Combine the last two results.

Proposition A.5. *Assume that (G, X) satisfies the conditions of (A.3). There then exists a \mathbb{Q} -rational reductive group $G_0 \subset G$ such that*

- (a) all $h \in X$ factor through $G_{0, \mathbb{R}}$, and
- (b) $\text{ad } h(\iota)$ is a Cartan involution on $G_0/w_X(\mathbb{G}_m)$.

Proof. We assume that no proper \mathbb{Q} -rational reductive subgroup of G satisfies (a) and show that G then satisfies (b). Let $H = G/(G^{\text{der}}, w(\mathbb{G}_m))$. Then $X^*(Z/w(\mathbb{G}_m)) \otimes \mathbb{Q} = X^*(H) \otimes \mathbb{Q}$, and so (A.3b) implies that $\iota \tau = \tau \iota$ on $X^*(H)$ for all $\tau \in \text{Aut}(\mathbb{C})$. Let V^+ be the maximal subspace of $X^*(H) \otimes \mathbb{Q}$ on which ι acts as $+1$. The last statement shows that V^+ is stable under $\text{Aut}(\mathbb{C})$. There therefore

exists a quotient torus H^+ of H such that $V^+ = X^*(H^+) \otimes \mathbb{Q}$. For any $x \in X$, $\mathbf{S}/\mathbf{G}_m \xrightarrow{h_x} G_{\mathbb{R}}/w(\mathbf{G}_m) \rightarrow H_{\mathbb{R}}^+$ is trivial because ι acts as -1 on $X^*(\mathbf{S}/\mathbf{G}_m)$, and so h_x factors through $\text{Ker}(G_{\mathbb{R}} \rightarrow H_{\mathbb{R}}^+)$; we must have $H^+ = 0$. It follows now that $(Z/w(\mathbf{G}_m))_{\mathbb{R}}$ is anisotropic, and (b) holds.

Remark A.6. For the group G_0 constructed in (A.4), $Z(G_0)(\mathbb{Q})$ is discrete in $Z(G_0)(\mathbb{A}^f)$ (cf. Deligne (1979), 2.1.11).

Bibliography

- Borovoi, M.: Langlands's conjecture concerning conjugation of Shimura varieties. *Sel. Math. Sov.* **3**, 3–39 (1983/84)
- Deligne, P.: *Equations Différentielles à Points Singuliers Réguliers*. (Lect. Notes Math., vol. 163) Berlin Heidelberg New York: Springer 1970
- Deligne, P.: *Travaux de Shimura*. Sémin. Bourbaki, Fév. 1971, Exposé 389, Lect. Notes Math. vol. 244. Berlin Heidelberg New York: Springer 1971
- Deligne, P.: Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques. *Proc. Symp. Pure Math., A.M.S.*, **33**, part 2, 247–290 (1979)
- Deligne, P.: Hodge cycles on abelian varieties (Notes by J.S. Milne). In: *Hodge Cycles, Motives, and Shimura Varieties* (Lect. Notes Math., vol. 900, Berlin Heidelberg New York: Springer 1982a, pp. 9–100)
- Deligne, P.: *Motifs et groupes de Taniyama*. In: *Hodge Cycles, Motives, and Shimura Varieties*, (Lect. Notes Math., vol. 900) Berlin Heidelberg New York: Springer 1982b, pp. 261–279
- Deligne, P., Milne, J.: *Tannakian categories*. In: *Hodge Cycles, Motives, and Shimura Varieties*. (Lect. Notes Math., vol. 900) Berlin Heidelberg New York: Springer 1982, pp. 101–228
- Grothendieck, A.: *Etude locale des schémas et des morphismes de schémas*. *Inst. Hautes Etudes Sci., Publ. Math.* **32**, 5–361 (1967)
- Harris, M.: *Arithmetic vector bundles on Shimura varieties*. In: (Satake I., Morita, Y. (ed.)) *Automorphic Forms of Several Variables* *Prog. Math.* **46**, Birkhäuser, Boston, 1984, pp. 138–159
- Harris, M.: *Arithmetic vector bundles and automorphic forms on Shimura varieties*. I. *Invent. math.* **82**, 151–189 (1985)
- Hartshorne, R.: *Ample Subvarieties of Algebraic Varieties*. Lect. Notes Math., vol. 156) Berlin Heidelberg New York: Springer 1970
- Helgason, S.: *Differential Geometry, Lie Groups, and Symmetric Spaces*. New York: Academic Press, 1978
- Kobayashi, S., Nomizu, K.: *Foundations of Differential Geometry I, II*, New York: Wiley (Interscience) 1963/69
- Langlands, R.: *Automorphic representations, Shimura varieties, and motives*. *Ein Märchen*. *Proc. Symp. Pure Math., A.M.S.*, **33**, 205–246 (1979)
- Milne, J.: *Etale Cohomology*, Princeton: Princeton University Press 1980
- Milne, J.: *The action of an automorphism of \mathbb{C} on a Shimura variety and its special points*. *Prog. Math.* Vol. 35, Boston: Birkhäuser, 1983, pp. 239–265
- Milne, J.: *Arithmetic Duality Theorems, Perspectives in Math.*, 1, Boston: Academic Press 1986
- Milne, J., Shih, K.-y.: *Langlands's construction of the Taniyama group*. In: *Hodge Cycles, Motives, and Shimura Varieties* (Lect. Notes Math., vol. 900, Berlin Heidelberg New York: Springer 1982a, pp. 229–260)
- Milne, J., Shih, K.-y.: *Conjugates of Shimura varieties*. In: *Hodge Cycles, Motives, and Shimura Varieties* (Lect. Notes Math., vol. 900, Berlin Heidelberg New York: Springer 1982b, pp. 280–356)
- Mumford, D.: *Geometric Invariant Theory*. Berlin Heidelberg New York: Springer 1965
- Murakami, S.: *Cohomology Groups of Vector-valued Forms on Symmetric Spaces*, mimeographed notes, University of Chicago, 1966
- Platonov, V., Rapinčuk, A.: *On the group of rational points of three-dimensional groups*. *Sov. Math. Dokl.* **20**, 693–697 (1979)
- Saavedra Rivano, N.: *Catégories Tannakiennes*. (Lect. Notes Math., vol. 265) Berlin Heidelberg New York: Springer 1972
- Shimura, G.: *On some problems of algebraicity*. *Proc. Int. Cong. Math.* 1978, Vol I, pp. 373–379