AN ANNOTATED LIST OF THE CARICES OF THE DOUGLAS LAKE REGION

BOTANY # 296

Submitted by Margaret D. Feigley August 12, 1948

AN ANNOTATED LIST OF THE CARICES OF THE DOUGLAS LAKE REGION

This list of the Carices of the Douglas Lake Region is based upon specimens collected during the summers of 1947 and 1948 in the vicinity of the University of Michigan Biological Station, situated on Douglas Lake, Cheboygan County, Michigan.

The work was undertaken as a special problem in Botany.course number 296.

- 1. Carex albursina Sheldon
 Collected in the hardwoods two miles west of Mickinaw City by Agnes Kugel.
- 2. Carex anceps Muhl. (Carex laxiflora var. patulifolia(Dewey) Carey Collected in the Colonial Point Hardwoods.
- 3. Carex aurea Nutt.
 Wolfs Bog along an old logging road. Common along margins of beach pools and in boggy ground.
- 4. Carex Bebbii Olney (Carex tribuloides var. Bebbii (Olney) var. Bailey Common in wet soil, on beaches, in bogs and along roadside ditches:
- 5. Carex brunnescens (Pers.) Poir. (C. brunnescens var. gracilior Britt.)
 In Thuja bog near mouth of Carp Creek.
- 6. Carex Buxbaumii Wahl.
 Invading a mat at a little shallow bog within Thuja woods, west of Mackinaw City; collected by Dr. Frank C. Gates.
- 7. Carex castanea Wahl.
 In very wet muck beside the stream in the Gorge; near the Pellston road.
- 8. Carex comosa Boott.
 Fontinalis Run beside the water's edge; rather common in low wet places.
- 9. Carex convoluta Mack.
 Collected in the hardwoods two miles west of Mackinaw City by Agnes Kugel.
- 10. Carex crinita Lam.
 Abundant along Carp Creek in the Gorge.
- 11. Carex communis Bailey
 Collected in maple woods Riggsville Corners; rather frequent.

- 12. Carex debilis Michx. var. Rutdgei Bailey
 In moist rich humus in the Gorge near P ellston road.
- 13. Carex Dewyana Schwein.

 In maple woods at Riggsville Corners and in the hardwoods two miles west of Mackinaw City.
- 14. Carex diandra Schrank.

 Margin of a beach Pool, Cross Fillage.
- 15. Carex disperma Dewey
 Very common in bogs throughout the area.
- 16. Carex eburnea Boott.

 Very common on moist sandy soil beneath the conifers, Wilderness Park.
- 17. Carex flava L.

 Very common along the margins of veach pools and in wet boggy places.
- 18. Carex Garberi Fern. (C. Bicolor of American authors, not All.; C. Hassie of recent authors, not Bailey)
- 19. Carex gracilima Schwein.
 Common in hardwoods.
- 20. Carex gynocrates Wormsk.

 Reeses Bog in dense Thuja stands.
- 21. Carex hystricina Muhl.

 Very common in wet situations of all sorts.
- 22. Carex interior Bailey (C. Scirpoides Schkuhr. in part)
 Found in a wide variety of habitats; extremely variable.
- 25. Carex intumescens Rudge Fairly common in wet woods.
- 24. Carex lasiocarpa Ehrh.
 Common in bogs and in most wet situations; forms the mat at Smiths Bog.
- 25. Carex laxiflora Lam.
 Common in moist hardwoods.
- 26. Carex leptalea Wahl.
 Collected in Reeses Bog and in Mud Lake Bog in Thujas.

- 27. Carex lupulina Muhl. (C. lupulina var. pedunculata Gray)
 Collected on Burt Lake near the mouth of Carp Creek.
- 28. Carex oligosperma Michx.
 Collected in Reeses Bog: not common.
- 29. Carex pauciflora Michx. Locally common in bogs.
- 50. Carex Peckii Howe
 Collected on the gravelly shores of Hat Island by Dr. Frank C. Gates
- 31. Carex pennsylvanica Lam.
 Collected by Dr. Frank C. Gates in a Piney- aspen second growth
 on old gravel beaches just north of French Farm Lake.
- 32. Carex Pseudo-cyperus L.

 Along a roadside ditch about one mile south of Riggsville Corners.
- 53. Carex retrorsa Schwein.
 Fairly common in wet situations; collected at Fontinalis Run and at Nigger Creek.
- 34. Carex riparia var. lacustris(Willd.) Kükenth.
 Collected beside a roadside ditch about one mile south of Riggsville
 Corners; very infrequent.
- 35. Carex rostrata Stokes
 Frequent in boggy situations throughout the area.
- 36. Carex scabrata Schwein.
 Collected by Cecilia Reikert in Wolfs Bog; infrequent.
- 37. Carex Schweinitzii Dewey
 Frequent in wet situations such as margins of streams.
- 38. Carex scoparia Schkuhr.
 Common along beaches and in bogs.
- 39. Carex stipata Muhl.

 Very common along wet ditches, margins of streams, and in bogs.
- 40. Carex stricta Lam. (C. stricta var. angustata(Boott.) Bailey Collected at Nigger Creek and at Fontinalis Run.
- 411 Carex substricts (Kükenth.) Mack. (C. aquatilis of most authors)
 Frequent along margins of beach pools, Cross Village.

- 42. Carex tenera Dewey
 Wolfs Bog along an old logging road.
- 45. Carex trisperma Dewey Frequent in bogs.
- 44. Carex Tuckermani Boott.
 Collected by Dr. Frank C. Gates in open thickets in marginal fosse of Vestal Bog; rare.
- 45. Carex vesicaria L. Very common along streams and in boggy places; forms large hummocks at Nichols and Smiths Bogs.
- 46. Carex viridula Michx. (C. Oederl var. pumila (Cos. & Germ.) Fern. Common around beach pools along the shore of Lake Michigan.
- 47. Carex vulpinoidea Michx, Collected frequently in all sorts of moist situations; beside roadside ditches, margins of streams, and in bogs.

ADDITIONS

- 48. Carex angustior Mack. (stellulata var. angustata Carey) Collected in the Gorge.
- 49. Carex aenea Fern. (C. foenea of Michigan authors, not Willd.) Collected along a logging road in Luce County, Michigan, near Newberry.
- 50. Carex paupercula Michx. Rather frequent in bogs.
- 51. Carex muskingumensis Schwein.
 Collected along a bog stream about two miles northwest of Mattoon,
 Wisconsin; Langlade County.

SPECIES COLLECTED IN THE VARIOUS AREAS

Big Stone Bay:

- C. Bebbii Olney
- C. hystricina Muhl.

Carp Creek(where it enters Burt Lake):

Carex brunnescens (Pers.) Poir.

- C. comosa Boott.
- C. disperma Dewey
- C. hystricina Muhl.
- C. Deweyana Schwein.
- C. interior Bailey
- C. intumescens Rudge
- C. lasiocarpa Ehrh.
- C. retrorsa Schwein.
- C. scoparia Schkuhr.
- C. stipata Muhl.
- C. trisperma Dewey

Cecil Bay:

Carex Bebbii Olney

- C. hystricina Muhl.
- C. lasiocarpa Ehrh.
- C. scoparia Schkuhr.

Cross Village:

Carex communis Bailey

- C. __diandra Schrank.
- C. hystricina Muhl.
- C. interior Bailey
- C. stipata Muhl.
- C. stricta Lam.
- C. wiridula Michx.

Bryants Bog:

- C. disperma Dewey
- C. pauciflora Michx.
- C. trisperma Dewey

Fontinalis Run:

Carex comosa Boott.

- C. flava L.
- C. hystricina Muhl.
- C. retrorsa Schwein.
- C. stipata Muhl.
- C. vesicaria L.
- C. vulpinoidea Michx.
- C. substricta Kükenth Mack.

rench Farm Lake:

Carex pennsylvanica Lam.

Hat Island: Carex Peckii Howe

Livingstons Bog:

Carex brunnescens (Pers.) Pois.

- C. intumescens Rudge
- C. pauciflora Michx.
- C. trisperma Dewey

Luce County, Michigan (small area near Newberry):

Carex brunnescens (Pers.) Poir.

- C. Buxbaumii Wahl.
- C. oligosperma Michx.
- C. scoparia Schkuhr.
- C. trisperma Dewey
- C. aeneaFern.

Mackinaw City (hardwoods two miles west):

Carex albursina

- C. convoluta
- C. Buxbaumii

Maple River (near bridge on the Pellston road):

- C. Bebbii Olney
- C. stipata Muhl.
- C. stricta Lam.
- C. vulpinoidea

Mud Lake Bog:

Carex gynocrates Wormsk.

- C. Leptalea Wahl.
- C. pauciflora Michx.

Mullet Lake:

Carex Bebbii Olney

Nichols Bog:

Carex Bebbii Olney

- C. lasiocarpa Ehrh.
- C. vesicaria L.

Nigger Creek:

Carex lupulina Muhl.

- C. stipata Muhl.
- C. vulpinoidea Michx.

```
Reeses Bog:
  Carex Bebbii Olney
  C. Cflava L. >
  C. gynocrates Wormsk.
  C. hystricina Muhl.
  C. interior Bailey
  C. Schweinitzii Dewey
  C. vulpinoidea Michx. -
The Gorge (between the Pellston Road and the iron bridge on the hogback
  Carex angustior Mack.
  C. Garberi Fern.
  C. disperma Dewey
  C. aurea Nutt.
  C. gracillima Schwein.
  C. hystricina Muhl.
  C. laxiflora Lam.
  C. lupulina Muhl.
  C. paupercula Michx.
  C. Pseudo-cyperus L.
  C. Schweinitzii Dewey
  C. scoparia Schjuhr.
  C. Stipata Muhl.
  C. vesicaria L.
  C. debilis Michx. var . Rutdgei Bailey
Roadside ditch about one mile south of Riggsville Corners on an east-
  west road:
  Carex comosa Boott.
  C. crinita Lam.
  C. flava L.
  C. hystricina Muhl.
   C. intumescens Rudge
   C. lupulina Muhl.
   C. Pseudo-cyperus L.
   C. riparia var lacustris Willd. Kükenth.
   C. retrorsa Schwein.
  .C. vesicaria L.
```

Road to Smiths Bog(bridge on Nigger Creek about three fourths of a mile beyond the turn to the bog.)

Carex castanea Wahl.

C. vulpinoidea Michx.

- C. flava L.
- C. interior Bailey
- C. Schweinitzii Dewey
- C. vulpinoidea Michx.

Riggsville Corners (hardwoods):

Carex communis Bailey

C. Deweyana Schwein.

C. disperma Dewey

C. gracillima Schwein.

C. gyhocrates Wormsk.

C. interior Bailey

C. laxiflora Lam.

Smiths Bog

C. Bebbii Olney

C. lasiocarpa Ehrh.

C. vesicaria L.

Wolfs Bog

Carex Bebbii Olney

Carex aurea Nutt.

C. flava L.

C. hystricina Muhl.

C. interior Bailer

C. paupercula Michx.

C. tenera Dewey

C. scabrata Schwein.