



## COMPARISON OF MACROINVERTEBRATE-DERIVED STREAM QUALITY METRICS BETWEEN SNAG AND RIFFLE HABITATS<sup>1</sup>

Kristine F. Stepenuck, Ronald L. Crunkilton, Michael A. Bozek, and Lizhu Wang<sup>2</sup>

**ABSTRACT:** We compared benthic macroinvertebrate assemblage structure at snag and riffle habitats in 43 Wisconsin streams across a range of watershed urbanization using a variety of stream quality metrics. Discriminant analysis indicated that dominant taxa at riffles and snags differed; Hydropsychid caddisflies (*Hydropsyche betteni* and *Cheumatopsyche* spp.) and elmids (*Optioservus* spp. and *Stenemlis* spp.) typified riffles, whereas isopods (*Asellus intermedius*) and amphipods (*Hyaella azteca* and *Gammarus pseudolimnaeus*) predominated in snags. Analysis of covariance indicated that samples from snag and riffle habitats differed significantly in their response to the urbanization gradient for the Hilsenhoff biotic index (BI), Shannon's diversity index, and percent of filterers, shredders, and pollution intolerant Ephemeroptera, Plecoptera, and Trichoptera (EPT) at each stream site ( $p \leq 0.10$ ). These differences suggest that although macroinvertebrate assemblages present in either habitat type are sensitive to detecting the effects of urbanization, metrics derived from different habitats should not be intermixed when assessing stream quality through biomonitoring. This can be a limitation to resource managers who wish to compare water quality among streams where the same habitat type is not available at all stream locations, or where a specific habitat type (i.e., a riffle) is required to determine a metric value (i.e., BI). To account for differences in stream quality at sites lacking riffle habitat, snag-derived metric values can be adjusted based on those obtained from riffles that have been exposed to the same level of urbanization. Comparison of nonlinear regression equations that related stream quality metric values from the two habitat types to percent watershed urbanization indicated that snag habitats had on average 30.2 fewer percent EPT individuals, a lower diversity index value than riffles, and a BI value of 0.29 greater than riffles.

(KEY TERMS: land use; macroinvertebrate community assemblages; streams; riffles; snags; urbanization; discriminant analysis; imperviousness.)

Stepenuck, Kristine F., Ronald L. Crunkilton, Michael A. Bozek, and Lizhu Wang, 2008. Comparison of Macroinvertebrate-Derived Stream Quality Metrics Between Snag and Riffle Habitats. *Journal of the American Water Resources Association* (JAWRA) 44(3):670-678. DOI: 10.1111/j.1752-1688.2008.00197.x

<sup>1</sup>Paper No. J05136 of the *Journal of the American Water Resources Association* (JAWRA). Received September 27, 2005; accepted December 6, 2007. © 2008 American Water Resources Association. **Discussions are open until December 1, 2008.**

<sup>2</sup>Respectively, Graduate Student, Wisconsin Cooperative Fishery Research Unit, U.S. Geological Survey, College of Natural Resources, University of Wisconsin-Stevens Point at the time this paper was prepared, now Volunteer Stream Monitoring Program Coordinator, University of Wisconsin-Extension and Wisconsin Department of Natural Resources, 445 Henry Mall, Room 202, Madison, Wisconsin 53706; Professor, College of Natural Resources, University of Wisconsin-Stevens Point; Unit Leader, Wisconsin Cooperative Fishery Research Unit, USGS, College of Natural Resources, University of Wisconsin-Stevens Point; Research Scientist, Wisconsin Department of Natural Resources, Bureau of Science Services at the time this article was prepared, now Director, Institute for Fisheries Research, Michigan Department of Natural Resources, University of Michigan, Ann Arbor, Michigan 48109 (E-mail/Stepenuck: kfstepenuck@wisc.edu).

## INTRODUCTION

Anthropogenic disturbances in a watershed, particularly those associated with urban development, are known to affect macroinvertebrate community assemblages. Shifts in structure and function, including changes in composition, richness, abundance, functional feeding groups, and diversity have been reported from streams subject to such disturbances (Pitt and Bozeman, 1983; Pedersen and Perkins, 1986; Jones and Clark, 1987; Richards and Host, 1994; Lenat and Crawford, 1994; Elliott *et al.*, 1997). As such, benthic macroinvertebrates have been shown to be useful in assessing the health of aquatic systems (Kaesler and Cairns, 1972). Not only are these organisms ubiquitous, not very mobile, and abundant, making them ideal bioindicators, but most or a portion of their life cycle is aquatic and they can be found in a wide variety of habitat types. These properties offer the opportunity to assess temporal changes in stream quality rather than a snapshot estimate which is typically provided by chemical or physical assessments (Rosenberg and Resh, 1993).

Numerous metrics have been developed to describe stream quality based on abundance and composition of macroinvertebrate populations (Shannon, 1948; Margalef, 1958; Chutter, 1972; Hilsenhoff, 1977; Chesters, 1980; Plafkin *et al.*, 1989; Rosenberg and Resh, 1993). However, a variety of site-specific attributes may affect these stream quality metrics because of their influence on macroinvertebrate communities. These attributes include substrate size and type (Hynes, 1970a,b; Allan, 1975; Minshall and Minshall, 1977; Reice, 1980; Williams, 1980; Erman and Erman, 1984; Clements, 1987), velocity (Minshall and Minshall, 1977; Williams, 1980; Erman and Erman, 1984), habitat type (Minshall and Minshall, 1977), and physical and chemical properties (Baker and Sharp, 1998).

Understanding how site-specific habitat attributes affect macroinvertebrate assemblages and their calculated stream quality metrics is crucial in biomonitoring. This is particularly important where habitat differences cannot be standardized because of inherent differences in geomorphic or watershed characteristics or where the type of available habitat is limited or altered as a result of development or disturbance in a watershed (Malmqvist and Rundle, 2002). A better understanding of these differences may enable adjustment of the metric values across habitat types to maximize their potential use. This would benefit resource managers in their decision making processes by allowing them to compare water quality among streams where the same habitat type is not available at all stream locations, or where a specific habitat

type (i.e., a riffle) is required to determine a metric value [i.e., biotic index (BI)].

Stream quality assessments using macroinvertebrates often rely on metrics based on data from standardized samples collected from riffles (Plafkin *et al.*, 1989). However, riffles are not always present, thus investigators often substitute other habitat types such as snags (defined herein as branches or other debris onto which organic material has collected). Although differences in macroinvertebrate composition between riffle and pool habitats or among different rocky substrate sizes have been documented (e.g., Minshall and Minshall, 1977; King, 1981; McCulloch, 1986; Quinn and Hickey, 1990; Brown and Brussock, 1991; Payne and Miller, 1991), only limited studies have evaluated differences in macroinvertebrate collections between riffle and snag habitats (Hooper, 1993; Wang *et al.*, 2006). More studies are especially needed to evaluate the influences of such different habitats on the macroinvertebrate metrics and how to adjust for these differences in water quality assessments. To address this, our first objective was to confirm differences in macroinvertebrate taxonomic assemblage between habitat types at our study sites. Our second objective was to assess differences in stream quality metrics between snag and riffle habitats in these southeastern Wisconsin streams subject to a gradient of watershed urbanization. These were a subset of those streams studied by Wang *et al.* (2006). Our third objective was to determine adjusted snag metric scores that could be used to evaluate stream quality in systems without riffles as compared to those with riffles that were subject to the same level of urbanization (or vice versa). These adjusted scores can be valuable to resource managers who wish to make comparisons of stream quality assessments between sites with different available habitat types.

*Study Area*

We assessed watershed land use and macroinvertebrate assemblages at 43 stream sites in Southeastern Wisconsin (Figure 1). All streams were located in unique subwatersheds in the Southeastern Wisconsin Till Plains Ecoregion (Omernik and Gallant, 1989). Most of this area has low relief and slopes are predominantly level to slightly rolling. The stream drainage system is poorly developed and undrained depressions are common. Wang *et al.* (2000) provided a detailed description of each study stream. Because of nutrient rich soils and low relief, this ecoregion was historically an important agricultural area, but by 1990 about two million people lived in this region, and agricultural land use had decreased.




FIGURE 1. Location of Forty-Three Stream Sites in Southeastern Wisconsin.

Three criteria were used to select the study sites. First, to minimize natural variation in biotic communities, sites with similar watershed soil types, stream size and slope, and natural hydrological and temperature regimes were selected. They represented a range of watershed urban development (as measured by imperviousness) and environmental quality, from least impacted regional reference conditions to heavily degraded conditions (Figure 2). Second, sites were chosen in watersheds for which detailed land use data could be obtained. Finally, sites with historical fish population data were selected to meet requirements of a concurrent study about the effects of urbanization on fish communities (Wang *et al.*, 2000, 2001).

Subwatershed areas ranged between 5.55 and 101.73 km<sup>2</sup>, with a mean of 28.33 km<sup>2</sup>. Impervious




FIGURE 2. Distribution of Sampling Sites Across a Gradient of Percent Watershed Imperviousness.

surfaces within subwatersheds (a measure of perturbation in urban watersheds; Schueler, 1994) ranged from 1.9 to 46.3% with a mean of 12.0%. To minimize variation in potential natural biological attributes, all sampling sites were warmwater (summer maximum daily mean temperature exceeded 24°C), second to third order in size, and had low to moderate gradients (< 6 m/km). These stream sites, in the absence of differing watershed land uses, were expected to have similar habitat and biological communities (Lyons, 1996).

## METHODS

### *Macroinvertebrates*

In October 1997, we sampled aquatic macroinvertebrates from riffle and snag habitats separately using a 600  $\mu$ m mesh D-frame kick net. Riffles were defined as areas where water velocity was greatest (preferably 0.3 m/s or greater) and the surface was turbulent because of a decrease in depth. Bottom substrates were composed of cobble, gravel, small rocks, or other stable substrate of fairly uniform size. Snags were defined as partially or fully submerged accumulations of vegetative debris (e.g., leaves, grasses) on branches, logs or other objects within the stream channel.

During sampling, we approached each habitat type from downstream to minimize disturbance prior to positioning the D-frame net. At riffles, we positioned the D-frame net on the stream bottom and vigorously disturbed the substrate just upstream from the net by kicking. This process dislodged macroinvertebrates, allowing them to be washed into the net (Hilsenhoff, 1987). At snags, we positioned the D-frame net in the water column where it would collect most of the dislodged debris. Depending on water depth, we disturbed the snag using either hands or feet. If the net became clogged, we cleared it and continued the collection process.

At each site, we repeated this process at three locations for each habitat type. We sampled riffles across a transect perpendicular to flow. If the stream was too narrow for three successive samples to be collected, we collected them by moving to an upstream riffle. We collected snag samples in areas of highest water velocity with first consideration given to largest snags. We sampled additional snags with similar consideration given to size of snag and water velocity.

For each habitat type, we combined the three samples to ensure that at least 125 organisms with assigned BI tolerance values were obtained

(Hilsenhoff, 1987). This number of organisms is usually present when enough debris to fill an 8-ounce jar has been collected (Hilsenhoff, 1987). We visually inspected debris in the field to assess general number of organisms collected. At highly degraded sites where macroinvertebrates or appropriate sampling habitat types were rare, a more extensive sampling effort was required. In such cases, regardless of the number of macroinvertebrates collected, we stopped sampling after one hour or after a standard habitat sampling reach (35 times mean stream width; Lyons, 1992) had been assessed.

By hand or using forceps, we removed organisms found clinging to the net or to large organic debris and placed them in sample collection jars. We then discarded the large organic debris at the field site. We preserved macroinvertebrate samples in 80% isopropyl alcohol, and transferred the samples to fresh alcohol after 24 and 48 hours. We later transported samples to the laboratory for sorting and identification.

In the laboratory, we placed macroinvertebrates and debris in a Pyrex<sup>®</sup> plate positioned over a grid of twelve 6.5 cm<sup>2</sup> squares. Using a random number table, we chose a grid square from which to remove invertebrates. With the aid of an illuminated magnifying lamp, we removed all invertebrates from the grid square. Organisms that had BI tolerance values were enumerated. We repeated this process in additional, consecutively numbered grid squares until a minimum of 125 organisms with BI tolerance values had been enumerated or until the entire sample had been sorted.

We identified macroinvertebrate taxa to the lowest taxonomic level for which a BI tolerance value was defined (Hilsenhoff, 1987). We identified organisms without defined BI tolerance values to the lowest level of taxonomic resolution possible. We slide-mounted chironomid larvae in CMC-10<sup>®</sup> mounting media (Master's Chemical Company, Inc., Bensenville, IL) to increase clarity of specimens, and allowed them to clear for two days prior to identifying them.

#### *Watershed Land Use*

We used percent watershed imperviousness as a surrogate for degree of urbanization using a Geographic Information System database (developed by the Southeastern Wisconsin Regional Planning Commission, 1990). The database was developed from 1:4,800 air photos. Nine major land use categories were defined: residential, commercial, industrial, transportation, communication and utilities, government and institutional, recreational, agricultural, and open lands. Each of these categories included sub-categories, for a total of 63 possible land use types. Each

land use type was assigned a percent effective connected imperviousness based on surfaces such as roads, rooftops, and parking lots. These surfaces were typically connected to surface waterways by some direct route, such as a storm sewer, drainage pipe, or surface drainage way (Booth and Jackson, 1997). Watershed imperviousness was calculated for each land use type, summed and standardized by watershed area. Watershed boundaries and upstream land area were determined following methods described in Wang *et al.* (2000).

#### *Analyses*

For our assessments, we calculated several stream quality metrics (Plafkin *et al.*, 1989) based on macroinvertebrate data, including Hilsenhoff's BI (Hilsenhoff, 1987), the Shannon's diversity index (Shannon, 1948), generic and species richness (Gaufin, 1957), and percent pollution intolerant Ephemeroptera, Plecoptera, and Trichoptera (EPT). We calculated percent collectors, filterers, gatherers, scrapers, and shredders to describe functional feeding response to increasing levels of watershed imperviousness (Merritt and Cummins, 1996). Although we recognize that there is a difference between functional feeding groups, diversity indices, and other stream quality assessment methods, for purposes of simplicity and following Plafkin *et al.* (1989), we refer to these collectively as stream quality metrics in this paper. Metrics calculated for both habitat types were plotted vs. percent watershed imperviousness to assess the impact of urbanization on the macroinvertebrate communities (Stepenuck *et al.*, 2002) and to compare differences in such impacts between habitat types.

To develop snag-adjusted metric values to assess stream health at sites without riffles, we first confirmed that macroinvertebrate community assemblages differed between the sampled habitats. This was necessary because values of some metrics were dependent upon the taxa that were identified in a sample (i.e., Hilsenhoff BI). To confirm that macroinvertebrate assemblages in snags could be discriminated from those at riffles, we analyzed data using canonical discriminant function analysis (SAS Institute Inc., 1992) in an approach similar to Tonn *et al.* (1983). Final classification success rates were obtained using a split-half validation process with priors set proportional. We only conducted analyses on taxa comprising at least 2% of the entire assemblage to reduce influence of rare species. At sites where two snag samples were collected (i.e., where riffles were not present) we averaged metric values to be used in the discriminant function analysis.

We then assessed metric values to evaluate if the slopes of the regressions of the observed relationships (i.e., between a metric value and percent watershed imperviousness) were uniform between habitat types. When slopes were not uniform, or a significant interaction ( $p \leq 0.10$ ) existed between the covariate (i.e., imperviousness) and the treatment (i.e., habitat type), we did not perform Analysis of Covariance (ANCOVA). In such cases, we could not determine a snag-adjusted value.

For metrics in which no significant interaction existed between the covariate and the treatment, we performed subsequent ANCOVA analyses (SAS Institute Inc., 1992; SPSS Inc., 1998). This allowed us to determine significant differences between snag and riffle habitats in stream quality metrics along a gradient of watershed imperviousness.

Prior to ANCOVA, we transformed metrics to linearize the relationship with the covariate.  $\text{Log}_e(x)$  was used for most metrics, however, we utilized  $\text{log}_e(x + 1)$  to correct for zeroes in the data sets for percent EPT, and percent scrapers, filterers, and shredders. We considered the relationship between a metric and percent watershed imperviousness significant at  $p \leq 0.10$ .

When ANCOVA revealed significant differences between treatments as they covaried with watershed imperviousness, we used the differences between the intercepts of the nonlinear regression equations as snag-adjusted values between the treatments. We reported these snag-adjusted values as the difference in value or percent for each of the metrics.

RESULTS

Discrimination of taxonomic assemblages between snag and riffle habitats was significant (Wilks'  $\lambda = 0.4857$ ,  $p < 0.0001$ ) and resulted in an initial correct classification rate of 85.9% (Figure 3). Nearly unbiased split-half validation of the data resulted in an 80.8% correct classification rate in assignment of habitat type using the discriminant functions defined in the initial classification. Hydropsychid caddisflies [*Hydropsyche betteni* (Ross) and *Cheumatopsyche* spp. (Wellengren)] and elmids beetles [*Optioservus* spp. (Sanderson) and *Stenelmis* spp. (Dufour)] characterized riffle habitat with positively loaded structure coefficients in the discriminant function. Isopods [*Asellus intermedius* (Forbes)] and amphipods [*Hyalella azteca* (Saussure) and *Gammarus pseudolimnaeus* (Bousfield)] represented snag habitat with high negative loadings (Table 1).




FIGURE 3. Canonical Discriminant Axis Discriminating Macroinvertebrate Species at Riffle vs. Snag Habitats.

TABLE 1. Structure Coefficients for the Canonical Discriminant Function.

Taxa	CAN 1
<i>Cheumatopsyche</i> spp.	0.607
<i>Hydropsyche betteni</i>	0.634
<i>Optioservus</i> spp.	0.493
<i>Stenelmis</i> spp.	0.582
<i>Simulium vittatum</i>	0.327
<i>Gammarus pseudolimnaeus</i>	-0.240
<i>Hyalella azteca</i>	-0.398
<i>Asellus intermedius</i>	-0.289
<i>Turbellaria</i> spp.	-0.127
<i>Oligochaeta</i> spp.	0.274
<i>Physa</i> spp.	-0.057

ANCOVA revealed that many macroinvertebrate metrics were significantly ( $p \leq 0.10$ ) related to the degree of imperviousness in the watershed. For example, differences existed between treatments (i.e., snag vs. riffle habitats) as they covaried with the watershed imperviousness for BI, Shannon's diversity index, and percent EPT, shredders, and filterers (Table 2). Species and generic richness were not significantly related to degree of imperviousness in the watershed. The BI score was consistently lower in riffles than in snags by an average of 0.29, indicating better inferred stream quality in riffles across the range of watershed imperviousness (Figure 4, Table 3). Shannon's diversity index was consistently higher in riffles than in snags by a value of 0.58 (Figure 4, Table 3). Riffles had an average of 30.2% more EPT individuals (Figure 4, Table 3) and 44.2% more filterers (Figure 5, Table 3) than snags in our study watersheds. There were 1.5% more shredders in snags across the imperviousness gradient (Figure 5, Table 3). Slopes of the regressions were not uniform

TABLE 2. Nonlinear Regression Parameter Estimates From Analysis of Covariance Results.

Parameter	Coefficient	SE	t	Probability
BI	$(r^2 = 0.315 \ n = 78)$			
Intercept	1.600	0.059	27.04	0.000
Habitat type	-0.105	0.042	-2.52	0.014
Diversity	$(r^2 = 0.204 \ n = 78)$			
Intercept	1.121	0.131	8.580	0.000
Habitat type	0.224	0.092	2.428	0.018
EPT individuals (%)	$(r^2 = 0.295 \ n = 78)$			
Intercept	3.211	0.390	8.232	0.000
Habitat type	1.059	0.276	3.838	0.000
EPT genera (#)	$(r^2 = 0.413 \ n = 78)$			
Intercept	2.166	0.162	13.378	0.000
Habitat type	0.239	0.115	2.085	0.040
Species richness	$(r^2 = 0.120 \ n = 78)$			
Intercept	3.079	0.124	24.771	0.000
Habitat type	-0.117	0.088	-1.328	0.188
Generic richness	$(r^2 = 0.110 \ n = 78)$			
Intercept	3.027	0.123	24.570	0.000
Habitat type	-0.120	0.087	-1.380	0.172
Filterers (%)	$(R^2 = 0.337 \ N = 78)$			
Intercept	2.394	0.359	6.670	0.000
Habitat type	1.476	0.256	5.771	0.000
Shredders (%)	$(R^2 = 0.147 \ N = 78)$			
Intercept	1.473	0.225	6.536	0.000
Habitat type	-0.368	0.161	-2.290	0.025

Note: Nonlinear regression parameter estimates are for log<sub>e</sub> (stream quality metrics) vs. log<sub>e</sub> (% watershed imperviousness) between snag and riffle habitats.

for percent collectors, gatherers, and scrapers, thus a snag-adjusted value could not be determined.

### DISCUSSION

Numerous macroinvertebrate metrics clearly were affected by urbanization that was represented by watershed imperviousness in this study. Watershed urbanization negatively affected BI, Shannon's diversity index, percent EPT, and generic richness. These results concur with the findings of Klein (1979), Lenat and Crawford (1994), Shaver *et al.* (1995), Horner *et al.* (1996), and May *et al.* (1997) and are discussed in detail in Stepenuck *et al.* (2002).

Significant differences between macroinvertebrate communities inhabiting snag and riffle habitats existed in stream quality assessments in our southeastern Wisconsin study streams. These differences were similar to findings by Humphries *et al.* (1998) and Parsons and Norris (1996) who also found differences in macroinvertebrate community assemblages between habitats. Stream quality metrics calculated from samples collected from snag habitats consistently indicated more degraded stream quality than those calculated from samples collected at riffle




FIGURE 4. Fitted Nonlinear Regression Models Relating BI, Shannon's Diversity Index, and Percent EPT to Percent Watershed Imperviousness for Riffle and Snag Habitats.

habitats at the same site. It is likely, therefore, that the physical habitat in these streams directly affects the estimation of water quality attributes based on the macroinvertebrates. Because differences in community assemblage between habitat types at the same stream site may lead to varied stream quality assessments when water quality or stream health is evaluated, metrics derived from different habitats on the same or different streams should not be intermixed or compared without adjusting the scores. Wang *et al.* (2006) suggested using snags as an alternate sampling substrate for streams without riffles. However, this approach does not provide managers with the ability to compare water quality scores between high and low gradient streams. To address this issue, we suggest that snag-adjusted values for metrics shown to differ between snag and riffle habitats may be utilized by managers to evaluate stream health in watersheds without riffles or in which snags were sampled. Using snag-adjusted stream quality metrics could be a valuable management tool in ecoregions that are defined by low gradient, channelized streams, such as those commonly found in

TABLE 3. Nonlinear Regression Equations Relating Snag and Riffle Habitats to Percent Watershed Imperviousness.

Nonlinear Regression Equation	$r^2$	$p$	Habitat Type
BI = 4.94 imperviousness (%) <sup>0.142</sup>	0.28	0.0001	Snag
Shannon's diversity = 3.24 imperviousness (%) <sup>-0.208</sup>	0.11	0.0172	Snag
Percent EPT = 23.0 imperviousness (%) <sup>-0.701</sup>	0.21	0.0008	Snag
Percent filterers = 9.91 imperviousness (%) <sup>-0.402</sup>	0.07	0.061	Snag
Percent shredders = 4.44 imperviousness (%) <sup>-0.287</sup>	0.07	0.058	Snag
BI = 4.65 imperviousness (%) <sup>0.113</sup>	0.25	0.002	Riffle
Shannon's diversity = 3.82 imperviousness (%) <sup>-0.201</sup>	0.31	0.0004	Riffle
Percent EPT = 53.20 imperviousness (%) <sup>-0.557</sup>	0.14	0.023	Riffle
Percent filterers = 54.11 imperviousness (%) <sup>-0.507</sup>	0.14	0.026	Riffle
Percent shredders = 2.96 imperviousness (%) <sup>-0.269</sup>	0.16	0.017	Riffle

Note: Nonlinear regression equations relating snag and riffle habitats to % watershed imperviousness are for BI, Shannon's diversity, % EPT, % filterers, and % shredders.




FIGURE 5. Fitted Nonlinear Regression Models Relating Percent Filterers and Shredders to Percent Watershed for Riffle and Snag Habitats.

the upper Midwestern United States (Omernik and Gallant, 1989).

Resources managers in Wisconsin have reported difficulty locating riffles from which to sample both because of hydrologic modification and because of naturally low gradient streams (S. Galarneau and M. Miller, Wisconsin Department of Natural Resources, February 2007, personal communication). In fact, of 9,650 macroinvertebrate samples collected by Wisconsin Department of Natural Resources staff between 1975 and 2006, 21% were collected from snags, runs, or pools (64% were collected from riffles, which is the prescribed habitat from which to sample whenever possible) (UWSP Aquatic Entomology Lab, 2007). Thus, snag-adjusted metrics could provide managers valuable data to allow for comparisons between sites for a significant portion of their sampling efforts. With limited funding to implement water quality improvement projects, such comparison between sites could help managers prioritize locations at which to implement these projects.

We determined snag-adjusted metric values for BI, Shannon's diversity index, and percent EPT, shredders and filterers. Supplementary studies need to be done, however, to determine if the adjusted values identified in this study are valid for other watersheds along a similar gradient of percent watershed imperviousness. Because of the diversity of habitats present in stream ecosystems, it would be highly useful to derive other nonlinear regression equations for macroinvertebrate metric scores from samples collected from habitats other than snags.

Other research has not addressed adjusting macroinvertebrate community metric scores from snags (or other habitats) to riffles based on regression coefficients across a gradient of watershed imperviousness. Some techniques used to account for differences in habitat types have included sampling at all available habitats and compositing samples to a single sample (Wright *et al.*, 1984), and developing unique sampling methods for high and low gradient streams (Kentucky Division for Environmental Protection, 2002). These techniques tend to be more time intensive, and do not necessarily allow for direct comparison of stream quality across habitat types. However, with increased use and resolution of Geographic Information Systems and the accepted relation between watershed imperviousness and water quality (Schueler, 1994; Arnold and Gibbons, 1996), refined measures of imperviousness will enhance our ability to develop better nonlinear regression models within different ecoregions for future stream quality assessment studies.

Differences in stream quality between snag and riffle habitats were apparent, however, direct comparison of these differences using nonlinear regression

models allowed us to calculate adjusted values for metrics that are more useful when specific habitat types are not necessarily present in a stream. Successful implementation of our regression model can help to simplify some of the common problems associated with calculating macroinvertebrate stream quality metrics from different habitat types. The approach used will allow metric values to be adjusted for samples collected at the most available habitat types, rather than limiting sample collection to sites with riffles.

## ACKNOWLEDGMENTS

We thank Roger Bannerman, Robert Beabal, and John McDougall for preparing watershed land use data; and Tom Aartila, Mary Botzet, Kevin Fassbind, Scott Harpold, and Mike Miller for help with macroinvertebrate sampling. Many thanks to the University of Wisconsin-Stevens Point (UWSP) students who assisted with macroinvertebrate sorting (Sarah Sonsthagen, Sunshine Kapusta, Chris Huff, Pat Short, Kathy Smikrud, Bruce Medhurst, Marty Torpy, and Jenni Stritzel). We are grateful to Jeff Dimick at UWSP for assistance with macroinvertebrate identification and editorial comments and to Dr. Mike Hansen for assistance with nonlinear modeling. Finally, we thank Drs. S.W. Szczytko and C.F. Hartleb who provided valuable editorial comments that helped to improve the document.

## LITERATURE CITED

- Allan, J.D., 1975. The Distributional Ecology and Diversity of Benthic Insects in Cement Creek, Colorado. *Ecology* 56:1040-1053.
- Arnold, C.L. and C.J. Gibbons, 1996. Impervious Surface Coverage: The Emergence of a Key Environmental Indicator. *Journal of the American Planning Association* 62:243-258.
- Baker, S.C. and H.F. Sharp, 1998. Evaluation of the Recovery of a Polluted Urban Stream Using the Ephemeroptera-Plecoptera-Trichoptera Index. *Journal of Freshwater Ecology* 13(2):229-234.
- Booth, D.B. and C.R. Jackson, 1997. Urbanization of Aquatic Systems: Degradation Thresholds, Stormwater Detection, and the Limits of Mitigation. *Journal of the American Water Resources Association* 33(5):1077-1090.
- Brown, A.V. and P.P. Brussock, 1991. Comparison of Benthic Invertebrates Between Riffles and Pools. *Hydrobiologia* 220:99-108.
- Chesters, R.K., 1980. Biological Monitoring Working Party. The 1978 National Testing Exercise. Department of the Environment. Water Data Unit Technical Memorandum 19:1-37.
- Chutter, F.M., 1972. An Empirical Biotic Index of the Quality of Water in South African Streams and Rivers. *Water Research* 6:19-30.
- Clements, W.H., 1987. The Effect of Rock Surface Area on Distribution and Abundance of Stream Insects. *Journal of Freshwater Ecology* 4(1):83-91.
- Elliott, A.G., W.A. Hubert, and S.H. Anderson, 1997. Habitat Associations and Effects of Urbanization on Macroinvertebrates of a Small, High-Plains Stream. *Journal of Freshwater Ecology* 12(1):61-73.
- Erman, D.C. and N.A. Erman, 1984. The Response of Stream Macroinvertebrates to Substrate Size and Heterogeneity. *Hydrobiologia* 108:75-82.
- Gaufin, A.R., 1957. The Use and Value of Aquatic Insects as Indicators of Organic Enrichment. *In: Biological Problems in Water Pollution*, C.M. Tarzwell (Editor). Robert A. Taft Sanitary Engineering Center, Cincinnati, Ohio, pp. 136-143.
- Hilsenhoff, W.L., 1977. Use of Arthropods to Evaluate Water Quality of Streams. Technical Bulletin Number 100. Wisconsin Department of Natural Resources, Madison, Wisconsin, 14 pp.
- Hilsenhoff, W.L., 1987. An Improved Index of Organic Stream Pollution. *The Great Lakes Entomologist* 20(1):31-39.
- Hooper, A.E., 1993. Effects of Season, Habitat, and an Impoundment on Twenty-Five Benthic Community Measures Used to Assess Water Quality. M.S. Thesis. University of Wisconsin-Stevens Point, Stevens Point, Wisconsin.
- Horner, R.R., D.B. Booth, A. Azous, and C.W. May, 1996. Watershed Determinants of Ecosystem Functioning. *In: Effects of Watershed Development and Management on Aquatic Ecosystems*, L.A. Roesner (Editor). Engineering Foundation Conference, Proceedings, Snowbird, Utah, August 4-9, 1996, pp. 251-274.
- Humphries, P., J.E. Gowns, L.G. Serafini, J.H. Hawking, A.J. Chick, and P.S. Lake, 1998. Macroinvertebrate Sampling Methods for Lowland Australian Rivers. *Hydrobiologia* 364:209-218.
- Hynes, H.B.N., 1970a. *The Ecology of Running Waters*. University of Toronto Press, Toronto, Canada.
- Hynes, H.B.N., 1970b. *The Ecology of Stream Insects*. *Annual Review of Entomology* 15:25-42.
- Jones, R.C. and C.C. Clark, 1987. Impact of Watershed Urbanization on Stream Insect Communities. *Water Resources Bulletin* 23(6):1047-1055.
- Kaesler, R.L. and J. Cairns, 1972. Cluster Analysis of Data From Limnological Surveys of the Upper Potomac River. *The American Midland Naturalist* 88(1):56-67.
- Kentucky Division for Environmental Protection, 2002. *Methods for Assessing Biological Integrity of Surface Waters in Kentucky*. Division of Water, Ecological Support Section. Frankfurt, Kentucky.
- King, J.M., 1981. The Distribution of Invertebrate Communities in a Small South African River. *Hydrobiologia* 83:43-65.
- Klein, R.D., 1979. Urbanization and Stream Quality Impairment. *Water Resources Bulletin* 15(4):948-963.
- Lenat, D.R. and J.K. Crawford, 1994. Effects of Land Use on Water Quality and Aquatic Biota of Three North Carolina Piedmont Streams. *Hydrobiologia* 294:185-199.
- Lyons, J., 1992. The Length of Stream to Sample With a Towed Electrofishing Unit When Fish Species Richness is Estimated. *North American Journal of Fisheries Management* 12:198-203.
- Lyons, J., 1996. Patterns in the Species Composition of Fish Assemblages Among Wisconsin Streams. *Environmental Biology of Fishes* 45:329-341.
- Malmqvist, B. and S. Rundle, 2002. Threats to the Running Water Ecosystems of the World. *Environmental Conservation* 29(2):134-153.
- Margalef, R., 1958. Information Theory in Ecology. *General Systems* 3:36-71.
- May, C.W., R.R. Horner, J.R. Karr, B.W. Mar, and E.B. Welch, 1997. Effects of Urbanization on Small Streams in the Puget Sound Lowland Ecoregion. *Watershed Protection Techniques* 2:485-494.
- McCulloch, D.L., 1986. Benthic Macroinvertebrate Distributions in the Riffle-Pool Communities of Two East Texas Streams. *Hydrobiologia* 135:61-70.
- Merritt, R.W. and K.W. Cummins, 1996. *Ecology and Distribution of Aquatic Insects*. *In: An Introduction to the Aquatic Insects of North America*, R.W. Merritt, and K.W. Cummins (Editors). Kendall/Hunt Publishing Company, Dubuque, Iowa, pp. 74-86.
- Minshall, G.W. and J.N. Minshall, 1977. Microdistribution of Benthic Invertebrates in a Rocky Mountain (U.S.A.) Stream. *Hydrobiologia* 55(3):231-249.

- Omernik, J.M. and A.L. Gallant, 1989. Ecoregions of the Upper Midwest States. EPA-600/3-89-060, Environmental Research Laboratory, U.S. Environmental Protection Agency, Corvallis, Oregon.
- Parsons, M. and R.H. Norris, 1996. The Effect of Habitat-Specific Sampling on Biological Assessment of Water Quality Using a Predictive Model. *Freshwater Biology* 36(2):419-434.
- Payne, B.S. and A.C. Miller, 1991. The Structure of Dominant Invertebrate Assemblages in a Small Southeastern Stream. *Journal of Freshwater Ecology* 6(3):257-266.
- Pedersen, E.R. and M.A. Perkins, 1986. The Use of Benthic Macroinvertebrate Data for Evaluating Impacts of Urban Runoff. *Hydrobiologia* 139:13-22.
- Pitt, R. and M. Bozeman, 1983. Sources of Urban Runoff Pollution and its Effects on an Urban Creek. EPA-600/S2-82-090, Center for Environmental Research, U.S. Environmental Protection Agency, Cincinnati, Ohio.
- Plafkin, J.L., M.T. Barbour, K.D. Porter, S.K. Gross, and R.M. Hughes, 1989. Rapid Bioassessment Protocols for Use in Streams and Rivers. Benthic Macroinvertebrates and Fish, EPA-444/4-89-001, Office of Water, U.S. Environmental Protection Agency, Washington, D.C.
- Quinn, J.M. and C.W. Hickey, 1990. Magnitude of Effects of Substrate Particle Size, Recent Flooding, and Catchment Development on Benthic Invertebrates in 88 New Zealand Rivers. *New Zealand Journal of Marine and Freshwater Research* 24:411-427.
- Reice, S.R., 1980. The Role of Substratum in Benthic Macroinvertebrate Microdistribution and Litter Decomposition in a Woodland Stream. *Ecology* 61(3):580-590.
- Richards, C. and G. Host, 1994. Examining Land Use Influences on Stream Habitats and Macroinvertebrates: A GIS Approach. *Water Resources Bulletin* 30(4):729-738.
- Rosenberg, D.M. and V.H. Resh, 1993. Introduction to Freshwater Biomonitoring and Benthic Macroinvertebrates. *In: Freshwater Biomonitoring and Benthic Macroinvertebrates*, D.M. Rosenberg, and V.H. Resh (Editors). Chapman and Hall, New York, New York, pp. 1-9.
- SAS Institute Inc. (Statistical Analysis Systems), 1992. SAS Version 6.12 User's Guide (1992 Edition). SAS Institute Inc., Cary, North Carolina.
- Schueler, T., 1994. The Importance of Imperviousness. *Watershed Protection Techniques* 1(3):100-111.
- Shannon, C., 1948. A Mathematical Theory of Communication. *Bell System Technical Journal* 27:379-423.
- Shaver, E., J. Maxted, G. Curtis, and D. Carter, 1995. Watershed Protection Using an Integrated Approach. *In: Stormwater NPDES Related Monitoring Needs*, H.C. Torno (Editor). American Society of Civil Engineers, New York, New York, pp. 435-459.
- SPSS Inc., 1998. *Systat Version 8.0: Statistics*. SPSS Inc., Chicago, Illinois.
- Stepenuck, K.F., R.L. Crunkilton, and L. Wang, 2002. Impacts of Urban Landuse on Macroinvertebrate Communities in Southeastern Wisconsin Streams. *Journal of the American Water Resources Association* 38(4):1041-1052.
- Tonn, W.M., J.J. Magnuson, and A.M. Forbes, 1983. Community Analysis in Fishery Management: An Application With Northern Wisconsin Lakes. *Transactions of the American Fisheries Society* 112:368-377.
- UWSP Aquatic Entomology Lab, 2007. UWSP Bug Biomonitoring Program Data set 1975-2006. UW-Stevens Point, Stevens Point, Wisconsin. <http://www.uwsp.edu/water/biomonitoring/index3.htm>, accessed January 2007.
- Wang, L., J. Lyons, and R. Bannerman, 2001. Impacts of Urbanization on Stream Habitat and Fish Across Multiple Spatial Scales. *Environmental Management* 28(2):255-266.
- Wang, L., J. Lyons, P. Kanehl, R. Bannerman, and E. Emmons, 2000. Watershed Urbanization and Changes in Fish Communities in Southeastern Wisconsin Streams. *Journal of the American Water Resources Association* 36:1173-1189.
- Wang, L., B. Weigel, P. Kanehl, and K. Lohman, 2006. Influence of Riffle and Snag Habitat Specific Sampling on Stream Macroinvertebrate Assemblage Measures in Bioassessment. *Environmental Monitoring and Assessment* 119:245-273.
- Williams, D.D., 1980. Some Relationships Between Stream Benthos and Substrate Heterogeneity. *Limnology and Oceanography* 25(1):166-172.
- Wright, J.F., D. Moss, P.D. Armitage, and M.T. Furse, 1984. A Preliminary Classification of Running-Water Sites in Great Britain Based on Macro-Invertebrate Species and the Predication of Community Type Using Environmental Data. *Freshwater Biology* 14:221-256.