

Comprehensive Invited Review

PPARs and the Cardiovascular System

Milton Hamblin, Lin Chang, Yanbo Fan, Jifeng Zhang, and Y. Eugene Chen

Reviewing Editors: John Bright, Adnan Erol, M. Faadiel Essop, Shou Wei Han, and Do-Young Yoon

I. Introduction	1416
II. PPAR and the Mechanism of Action	1416
III. PPAR γ Ligands	1417
IV. PPAR γ and Endothelial Cells	1419
A. PPAR γ and the regulation of EC inflammatory response	1419
B. PPAR γ and the regulation of vascular tone	1420
C. PPAR γ and VEGF	1421
D. PPAR γ and EC migration	1421
E. PPAR γ and EC apoptosis	1421
F. PPAR γ and endothelial progenitor cells	1421
V. PPAR γ and Vascular Smooth Muscle Cells	1421
A. PPAR γ and VSMC proliferation	1422
B. PPAR γ and VSMC migration	1423
C. PPAR γ and VSMC apoptosis	1423
D. PPAR γ and the regulation of VSMC inflammatory response	1424
VI. PPAR γ and Monocytes/Macrophages	1424
A. PPAR γ and monocyte/macrophage inflammatory signaling	1424
B. PPAR γ and monocyte/macrophage migration and apoptosis	1424
C. PPAR γ and monocyte/macrophage iNOS expression	1425
D. PPAR γ and monocyte/macrophage CD36 expression	1425
E. PPAR γ and monocyte/macrophage lipid homeostasis	1425
VII. PPAR γ and Atherosclerosis	1425
VIII. PPAR γ and the Heart	1426
IX. PPAR α	1427
A. PPAR α Ligands	1428
B. PPAR α and endothelial cells	1428
C. PPAR α and VSMCs	1428
D. PPAR α and monocytes/macrophages	1429
E. PPAR α and atherosclerosis	1430
F. PPAR α and the heart	1430
XV. PPAR δ	1431
A. PPAR δ ligands	1431
B. PPAR δ and endothelial cells	1432
C. PPAR δ and VSMCs	1432
D. PPAR δ and monocytes/macrophages	1433
E. PPAR δ and atherosclerosis	1433
F. PPAR δ and the heart	1434
XVI. Perspective	1434

Abstract

Peroxisome proliferator-activated receptors (PPARs) belong to the nuclear hormone-receptor superfamily. Originally cloned in 1990, PPARs were found to be mediators of pharmacologic agents that induce hepatocyte peroxisome proliferation. PPARs also are expressed in cells of the cardiovascular system. PPAR γ appears to be highly expressed during atherosclerotic lesion formation, suggesting that increased PPAR γ expression may be a vascular compensatory response. Also, ligand-activated PPAR γ decreases the inflammatory response in cardiovascular cells, particularly in endothelial cells. PPAR α , similar to PPAR γ , also has pleiotropic effects in the cardiovascular system, including antiinflammatory and antiatherosclerotic properties. PPAR α activation inhibits vascular smooth muscle proinflammatory responses, attenuating the development of atherosclerosis. However, PPAR δ overexpression may lead to elevated macrophage inflammation and atherosclerosis. Conversely, PPAR δ ligands are shown to attenuate the pathogenesis of atherosclerosis by improving endothelial cell proliferation and survival while decreasing endothelial cell inflammation and vascular smooth muscle cell proliferation. Furthermore, the administration of PPAR ligands in the form of TZDs and fibrates has been disappointing in terms of markedly reducing cardiovascular events in the clinical setting. Therefore, a better understanding of PPAR-dependent and -independent signaling will provide the foundation for future research on the role of PPARs in human cardiovascular biology. *Antioxid. Redox Signal.* 11, 1415–1452.

I. Introduction

PEROXISOMES are organelles that participate in fatty acid metabolism. Clofibrate analogues, hypolipidemic agents that control plasma cholesterol and triglyceride levels, can induce proliferation of liver cell peroxisomes (300, 301). In addition, two lipid-lowering compounds structurally different from clofibrate, [4-chloro-6-(2,3-xylidino)-2-pyrimidinylthio]acetic acid (Wy-14,643) and 2-chloro-5-(3,5-dimethylpiperidino-sulfonyl)benzoic acid (tibric acid), also were found to stimulate hepatocyte peroxisome proliferation (302). Although hypolipidemic drugs were demonstrated to activate peroxisome proliferation, these studies did not establish a mechanism. Subsequent studies identified a protein whereby peroxisome proliferators bind with affinity (196, 197), and this protein was later identified as a member of the nuclear hormone-receptor superfamily that includes steroid, retinoid, and thyroid hormone receptors (104). The name peroxisome proliferator-activated receptor took origin from the cloning by Issemann *et al.* (172) to identify possible endogenous mediators of peroxisome proliferation-induced gene transcription in rodent livers. The peroxisome proliferator-activated receptors (PPARs) consist of three related transcription factors: PPARalpha (PPAR α), PPARbeta/delta (PPAR β/δ), and PPARgamma (PPAR γ), encoded by the genes *PPARA*, *PPARD*, and *PPARG*, respectively (96). In addition to the role in peroxisome proliferation, these nuclear transcription factors are involved in numerous cellular functions, including insulin sensitivity, glucose homeostasis, fatty acid oxidation, cytokine production, and vasculoprotection.

II. PPAR and the Mechanism of Action


PPARs were initially shown to recognize and bind a DNA sequence upstream of the PPAR target gene. This sequence was termed the peroxisome proliferator response element (PPRE) (251, 362) (Fig. 1). Acyl-CoA oxidase is a peroxisomal enzyme involved in fatty acid oxidation. The promoter of this enzyme was found to contain a DNA sequence that was responsive to stimulation by Wy-14,643, and this stimulatory

response was mediated by PPAR. Of great importance, PPAR was shown to bind to this 5' flanking portion, or peroxisome proliferator response element of the acyl-CoA oxidase gene (362). PPARs, on activation, heterodimerize with the retinoic X receptor (RXR)- α (22, 121, 182, 190), and this is followed by coactivator recruitment, which eventually leads to transcriptional regulation of gene expression (85, 312) (Fig. 1). Besides being involved in transactivation, PPARs also participate in the negative regulation of certain genes by recruiting corepressors (233) (Fig. 1). In addition, other molecular mechanisms are found by which PPARs can inhibit gene expression. First, transrepression can be caused by physical interaction with other transcription factors, including nuclear factor-kappa B (NF- κ B), Smad-3, activator protein-1 (AP-1), and signal transducers and activators of transcription (STAT) proteins (80, 114, 217, 307). Second, PPARs can modulate transrepression through the mitogen-activated protein kinase (MAPK) pathway (157). Coactivators and corepressors, in addition to regulating transcriptional activation, are critical for the repression of certain genes (85, 305, 312). Third, PPARs recruit coactivator proteins and often compete with NF- κ B and AP-1 for binding to these co-regulators (305). Thus, NF- κ B and AP-1 target gene expression is attenuated because of competition with PPARs for coactivator binding.

Finally, PPARs can contribute to transrepression by either inhibiting clearance of co-repressor complexes (123, 287) or releasing co-repressors, which could allow co-repressor binding to NF- κ B, eventually inhibiting NF- κ B target gene expression (305).

The phosphorylation of PPARs is critical to regulating many of the biologic functions of these nuclear receptors. Initially, insulin-induced phosphorylation of PPAR α was shown to increase transcriptional activity (322). Also, stress-activated p38 MAPK has been shown to phosphorylate PPAR α and enhance target gene expression in myocardiocytes (24). Several studies demonstrate that MAPK phosphorylation deactivates PPAR γ and reduces basal and ligand-dependent transcriptional activity (5, 51, 52, 157). However, one study shows that PPAR γ is activated by ERK5 in endothelial cells (ECs), and this particular MAPK does not phosphorylate

FIG. 1. Schematic view of PPAR action. After a ligand binds to PPAR, PPAR heterodimerizes with the retinoid X receptor (RXR) and then binds to the PPRE. Recruiting coactivators and co-repressors leads to activation and repression of PPAR target genes, respectively.



PPAR γ (7). A recent report demonstrates that PPAR γ is under the control of Bcr, a serine/threonine kinase that phosphorylates PPAR γ and prevents transcriptional activity (9). PPAR δ is also considered to be a phosphoprotein because protein kinase A (PKA)-induced phosphorylation of PPAR δ , similar to PPAR α and PPAR γ , has a stimulatory effect on transcription (200). These are just a few of many examples that demonstrate how PPAR signaling may be affected because of phosphorylation by protein kinases.

PPAR γ is most abundantly expressed in adipose tissue, with less expression in the colon and immune system. PPAR γ has been shown to facilitate differentiation of fibroblasts into adipocytes (59). PPAR γ is also involved in the regulation of lipid metabolism, as ligand-dependent activation leads to an increase in genes that regulate fatty acid uptake and storage (320). Furthermore, PPAR γ plays a role in glucose homeostasis and insulin sensitivity (110). Although PPAR γ was initially found to be critical for adipocyte differentiation and function, over time, PPAR γ was discovered to play an important role in the cardiovascular system. As well as in adipocytes and T cells, PPAR γ is also expressed in endothelial cells, vascular smooth muscle cells (VSMCs), and macrophages.

III. PPAR γ Ligands

PPARs possess varying degrees of responsiveness to certain peroxisome proliferating agents (188). Although several compounds were demonstrated to activate PPARs, initially no reports confirmed direct binding to this receptor. However, in 1995, evidence was provided that thiazolidinediones (TZDs), a class of antidiabetic drugs that improve insulin sensitivity, bind to and activate PPAR γ with high affinity (209) (Fig. 2). Furthermore, PPAR γ was shown to be the major target of these insulin-sensitizing agents (110).

Troglitazone (Rezulin), the first FDA-approved TZD used in the clinical setting, was discontinued from the market in 2000 because of reports of liver toxicity (125, 206, 259). Rosiglitazone (Avandia) and pioglitazone (Actos), subsequent TZD agents currently approved for clinical use, are not associated with severe hepatotoxicity (357), although weight gain and edema have been reported as side effects (263). Also,

rosiglitazone has been reported to be associated with increased risks of myocardial infarction and mortality due to cardiovascular complications (265); however, the results are controversial (155, 369). Clinical data from the PROactive study found that pioglitazone reduces the risk of secondary end points, including all-cause mortality, nonfatal myocardial infarction, and stroke in diabetic patients but nonsignificantly decreases the composite primary end-point risk (95). However, a recent meta-analysis that included 19 clinical trials found that pioglitazone reduces primary end-point components, including risk of death, myocardial infarction, and stroke (225).

GW1929 and GW7845 are examples of non-TZD high-affinity ligands for PPAR γ (39, 344) (Fig. 2). In addition, PPAR α/γ dual and PPAR $\alpha/\gamma/\delta$ pan agonists have been developed to promote synergistic antidiabetic and cardiovascular protective effects. Muraglitazar, naveglitazar, tesaglitazar, and netoglitazone are several examples of PPAR α/γ dual agonists (296) (Fig. 2). GW409544 has been shown to be a potent activator of both PPAR α and PPAR γ (390) (Fig. 2). Bezafibrate, a lipid-lowering drug that reduces the risk of myocardial infarction in patients with metabolic syndrome, is a PPAR $\alpha/\gamma/\delta$ pan agonist (353) (Fig. 2).

Several natural PPAR γ ligands have been identified and can be classified into two major groups of compounds, fatty acids and phospholipids. PPAR γ ligands consist of polyunsaturated fatty acids, including linoleic acids (36), linolenic acid (175), arachidonic acid (192), and eicosapentaenoic acid (159) (Fig. 2). Monounsaturated fatty acid compounds that bind PPAR γ include oleic acid (317) (Fig. 2). Oxidatively modified lipids also bind PPAR γ (Fig. 2). 15-Deoxy- δ 12,14-prostaglandin J₂ (15d-PGJ₂) and other J₂ series prostaglandins were identified as natural ligands for PPAR γ (110, 189) (Fig. 2). TZDs were demonstrated to be synthetic analogues of 15d-PGJ₂ (110). Other natural PPAR γ ligands include 12- and 15-hydroxyeicosatetraenoic acid (HETE) (159) and 9- and 13-hydroxyoctadecadienoic acid (HODE) (254) (Fig. 2), oxidized metabolites of arachidonic and linoleic acids, respectively. 1-*O*-hexadecyl-2-azelaoyl-*sn*-glycero-3-phosphocholine (azPC), an oxidized phospholipid, is also a PPAR γ ligand (78) (Fig. 2). In addition, lysophosphatidic acid (LPA) and its naturally




FIG. 2. PPAR γ ligands. Natural and synthetic agonists bind and activate PPAR γ . Natural PPAR γ agonists include 15d-PGJ₂, fatty acids, oxidatively modified lipids, hydroxyeicosatetraenoic acid, hydroxyoctadecadienoic acid, oxidized phospholipids, lysophosphatidic acid, and nitroalkenes. Synthetic PPAR γ agonists include TZDs, GW1929, GW7845, PPAR α/γ dual agonists, and PPAR $\alpha/\gamma/\delta$ pan agonists. Examples of PPAR γ antagonists include BADGE, GW9662, LG100641, PD068235, and SR-202.

occurring analogue, 1-*O*-octadecenyl-2-hydroxy-*sn*-glycero-3-phosphate (AGP) also have affinity for PPAR γ (361, 406) (Fig. 2).

Finally, our research group identified nitroalkenes 9-, 10-, 12-, and 13-nitro-9,12-*cis*-octadecadienoic acid (LNO₂) (319) and 9- and 10-nitro-9-*cis*-octadecenoic acid (OA-NO₂) (19) as natural PPAR γ ligands (Fig. 2). We recently reported the crystal structure of the PPAR γ ligand-binding domain bound to LNO₂ and found that LNO₂ promotes PPAR γ interaction with coactivator motifs of transcriptional coactivators (218). The two charged residues R288 and E343 of PPAR γ that make specific contacts with the NO₂ are not conserved in PPAR α and PPAR δ (218), explaining why LNO₂ preferentially activates PPAR γ rather than the other two PPAR subtypes (319). LNO₂ isomers bind to the two electrostatic regions of the ligand-binding pocket, and these electrostatic clusters allow binding of different ligands at the same time (218, 258). Our studies provide further evidence regarding the interaction between PPAR γ and LNO₂ and serve as a basis for the development of novel PPAR γ ligands that could not only mimic the interactions of LNO₂ on PPAR γ but also extend beyond the current TZD-induced PPAR γ -mediated effects in the cardiovascular system.

PPAR γ ligands can also participate in signaling independent of PPAR γ . Several studies have shown that PPAR γ ligands can directly interact and inhibit transcription factors in a PPAR γ -independent manner. First, although we have shown that nitroalkenes are PPAR γ ligands, nitroalkene-induced inhibition of macrophage proinflammatory cytokine secretion is regulated through nitroalkylation of the p65 subunit, repressing NF- κ B transcriptional activity (76) (Fig. 3).

Second, 15d-PGJ₂ inhibits NF- κ B transcriptional activity by inhibiting I κ B kinase (IKK) (54, 314, 342) and the DNA binding domains of NF- κ B (342). In all likelihood, the effects of

15d-PGJ₂ on IKK activity result in the inhibition of IKK-induced Ser32 and Ser36 phosphorylation of I κ B α (54) (Fig. 3). Compound G, a non-TZD agonist, also inhibits NF- κ B activation by decreasing IKK activity (55). Furthermore, the administration of TZD at higher concentrations attenuates NF- κ B target-gene expression in macrophages lacking PPAR γ (56, 249).

Pioglitazone can bind to mitoNEET, an integral protein located in the outer mitochondrial membrane that regulates oxidative capacity (71) (Fig. 3). MitoNEET received its name because of the Asn-Glu-Glu-Thr (NEET) sequence located in the carboxyl-terminal domain. Isolated mitochondria from the hearts of mitoNEET-null mice display an overall worsening of complex 1-dependent oxygen consumption (384). Because mitoNEET is an iron-sulfur cluster containing protein, and pioglitazone has been shown to increase mitoNEET 2Fe-2S stability (279), it is possible that pioglitazone could regulate the redox potential or function of the mitoNEET iron-binding CDGSH domain [C-X-C-X(2)-(S/T)-X(3)-P-X-C-D-G-(S/A/T)-H] (385).

PPAR γ antagonists are also ligands that can be used as important tools in determining PPAR γ signaling and function in basic science. The safety concerns and adverse side effects of TZDs have spurred an increased effort to study possible therapeutic benefits of administering PPAR γ antagonists in the clinical setting. Bisphenol A diglycidyl ether (BADGE) is often considered to be the first PPAR γ ligand known to inhibit transcriptional activity (386). A potent PPAR γ antagonist is GW9662, a compound that covalently modifies the Cys286 residue of the ligand-binding domain (207). Other examples of PPAR γ antagonists include LG100641 (252), PD068235 (50), and SR-202 (308).

The use of different methods for studying and screening novel PPAR modulators is an important concept of drug




FIG. 3. Schematic view of PPAR γ -dependent and -independent signaling pathways. PPAR γ ligands can exert their effects in cardiovascular cells through PPAR γ -dependent and -independent mechanisms. PPAR γ -mediated increases in IRF-1 and GADD45 result in greater VSMC apoptosis. PPAR γ -dependent decreases in c-fos expression attenuate VSMC proliferation. Ligand-activated PPAR γ inhibits NF- κ B transcriptional activity and inflammation in cardiovascular cells. PPAR γ ligand-independent signaling can decrease I κ B kinase activity, leading to decreased I κ B α phosphorylation, NF- κ B transcriptional activity, and inflammation. Another example of PPAR γ ligand signaling that occurs independent of PPAR γ involves nitroalkylation of the p65 subunit and eventual reduction in NF- κ B activity and inflammation. Pioglitazone can regulate mitochondrial oxidative capacity and normalize lipid oxidation through direct binding to the mitoNEET protein, independent of PPAR γ .

discovery. Several examples are known by which cell-free assays can be used for PPAR-modulator screening. A cell-free competition radioreceptor assay uses recombinant PPAR along with a radioisotope-labeled ligand and competitor ligands (110, 400). The premise of coactivator-dependent receptor ligand assays (CARLAs) includes coactivator recruitment and the use of a pull-down approach to determine the amount of ligand-bound PPAR-coactivator complex. The practice of radioactive labeling is not a requirement in CARLAs, allowing a large, quantitative screening of PPAR compounds (68, 192). The scintillation proximity assays (SPAs) measure receptor–ligand interaction. *Beta emission* from the radioactively labeled ligand is measured, and this is advantageous because of high sensitivity, high reliability, and the lack of a required separation step (100, 262).

The use of radioisotope-free assays is an alternative approach to previous cell-free methods. Surface plasmon resonance (SPR) techniques can be beneficial for detecting ligand–nuclear receptor interactions (401) and ligand-binding effects on nuclear-receptor dimerization (402), as well as screening for ligands from ligand-bound nuclear receptor–coactivator interactions (116). Fluorescence resonance energy transfer (FRET) is a radioisotope-free assay that is used to detect and quantitate PPAR ligand binding. A ligand-induced PPAR conformational change results in coactivator recruitment, allowing the fluorescence donor indirectly linked to PPAR and the fluorescence acceptor indirectly linked to the coactivator to draw into close proximity as the excited fluorescence donor transfers energy to the acceptor (68, 411). A simple ELISA has been developed in which unliganded PPAR weakly binds to the coactivator LXXLL motifs, while ligand-bound PPAR strongly binds to these LXXLL peptides. This radioisotope-free assay uses a specific anti-PPAR antibody to detect PPAR binding (69).

IV. PPAR γ and Endothelial Cells

The first evidence of PPAR γ expression in endothelial cells (34, 179, 235, 387) came from several studies examining the interaction of PPAR γ and plasminogen activator inhibitor type-1 (PAI-1). The expression of PAI-1 in both endothelial cells and adipocytes is involved in limiting fibrinolysis in humans. Elevated PAI-1 has been associated with myocardial ischemia and thrombosis in mice (228). PPAR γ agonists are generally found to increase PAI-I expression in endothelial cells (235, 387), although one study suggests the opposite (179). A later study provided evidence that PPAR γ 1 and not PPAR γ 2 mRNA is present in human umbilical vein endothelial cells (HUVECs) (198).

A. PPAR γ and the regulation of EC inflammatory response

Adhesion molecules can bind to inflammatory cells involved in signaling and regulation on the surface of endothelial cells. These adhesion molecules include vascular cell adhesion molecule-1 (VCAM-1), intercellular adhesion molecule-1 (ICAM-1), platelet–endothelial cell adhesion molecule (PECAM-1), E-selectin, and integrins. Along with monocyte chemoattractant protein-1 (MCP-1) and other chemoattractant molecules, adhesion molecules are responsible for attachment of immune cells to the endothelial layer, followed by eventual immune cell migration across the endothelium (313).

Much evidence demonstrates PPAR γ inhibitory and anti-inflammatory effects in endothelial cells. Several studies have reported that activation of PPAR γ inhibits expression of cellular adhesion molecules, including VCAM-1, ICAM-1, PECAM, and E-selectin, in addition to inflammatory cell migration and adhesion to atherosclerotic plaques (87, 173, 241,




FIG. 4. Schematic view of PPAR γ activation in ECs. Natural or synthetic PPAR γ ligands attenuate VEGF-induced Akt phosphorylation, inhibiting EC proliferation and migration. Ligand-activated PPAR γ exerts its antiinflammatory effects by inhibiting cytokine-induced NF- κ B activation in ECs.

257, 286, 378) (Fig. 4). NF- κ B plays an important role in regulating leukocyte adhesion molecule expression. Cytokines activate NF- κ B in endothelial cells, thereby allowing NF- κ B binding to promoters of adhesion molecule genes. Through NF- κ B binding, cytokine-induced gene expression of ICAM-1, VCAM-1, and E-selectin occur in the endothelium (73). Constitutively active PPAR γ inhibits NF- κ B- and AP-1-regulated gene expression and binding activity in ECs, and PPAR γ activation inhibits adhesion molecule expression by inhibiting NF- κ B and AP-1 signaling, considered the most important transcription factors in endothelial cell signaling (378). Another mechanism that may suppress endothelial cell inflammatory signaling is the inhibition of the diacylglycerol-protein kinase C (PKC) pathway (368). A study examined the effects of PPAR- γ ligands on chemokine expression that is induced by interferon-gamma (IFN- γ) in cultured human endothelial cells. PPAR γ activators decrease IFN-inducible protein of 10 kDa (IP-10), monokine induced by IFN- γ (Mig), and IFN-inducible T-cell α -chemoattractant (I-TAC) expression through the likely inhibition of NF- κ B (237) (Fig. 4). However, expression of MCP-1 is not changed in this study (237), in contrast to a previous report showing that TZDs inhibit tumor necrosis factor-alpha (TNF- α)- and interleukin-1beta (IL-1 β)-induced MCP-1 mRNA expression and secretion (253).

In cultured endothelial cells, TZDs may reduce superoxide production and inflammation (162, 244) by suppressing expression of NAD(P)H oxidase subunits that are critical for superoxide generation (162). Furthermore, a recent study found that mice expressing a dominant negative PPAR γ mutation show elevated oxidative stress and impaired endothelial function in cerebral arteries (32). Next, in cultured endothelial cells, TZDs, along with 15d-PGJ₂, attenuate IFN γ -induced major histocompatibility complex class II (MHC-II), a protein involved in regulating immune responses and T-cell activation (194). Finally, in HUVECs, TZDs promote expres-




FIG. 5. Schematic view of PPAR γ activation in vascular tone regulation. PPAR γ ligands decrease ET-1 and AT-1R expression and increase AT-2R expression. PPAR γ ligands stimulate NO release, and NO can activate endothelial cell PPAR γ through MAPK. TZDs can also decrease oxidative and nitrate stress.

sion of heme-oxygenase 1 (HO-1), a PPAR γ target gene with antiinflammatory properties (193).

B. PPAR γ and the regulation of vascular tone

Endothelin-1 (ET-1) is a vasoconstrictive protein that can also regulate VSMC proliferation. PPAR γ ligands attenuate both ET-1 expression and secretion in endothelial cells by blocking AP-1 signaling (83, 118, 163, 234, 318) (Fig. 5). Angiotensin II (AngII) is also a potent vasoconstrictor that increases angiotensin II type 1 (AT1) receptor expression, leading to narrowing of blood vessels and elevations in oxidative stress. In Sprague-Dawley rats, rosiglitazone and pioglitazone blunt AngII-induced increases in blood pressure by downregulating AT1 receptors and increasing angiotensin II type 2 receptor (AT2) expression (87) (Fig. 5). Both TZDs improve AngII-induced endothelial dysfunction (87). Subsequently, another study reported that in male apoE-knockout (apoE^{-/-}) mice, endothelial dysfunction occurs after AngII treatment in association with decreased PPAR γ gene and protein expression (355). Because human PPAR γ dominant negative mutations are associated with hypertension (27), TZD-induced PPAR γ activation may be one method of treatment for the effects of elevated blood pressure.

Conversely, endothelial cell-derived nitric oxide (NO) is a molecule that is a key participant in vasodilatory activity (280). In 1998, it was found that troglitazone causes vasodilation in humans (117). Subsequent studies showed that PPAR γ ligands increase NO production and release (49, 67, 294) (Fig. 5), although it appears that PPAR γ ligands may stimulate production of endothelial cell NO through different pathways (294). Ligand-activated PPAR γ was found to be critical to heat-shock protein 90/endothelial nitric oxide synthase (eNOS) interaction and eNOS phosphorylation in HUVECs (294). Furthermore, NO was recently reported to activate PPAR γ in endothelial cells through a p38 MAPK signaling pathway (297) (Fig. 5). TZDs possess vasculoprotective effects through the attenuation of oxidative and nitrate stresses (Fig. 5), and elevated NO levels. One study in male hypercholesterolemic rabbits suggests that rosiglitazone protects the endothelium by inhibiting superoxide, peroxynitrite, and excess NO production (351). Similar to adipocytes and VSMCs (94, 165), TZD-induced reduction in

elevated NO levels may be the result of inducible nitric oxide synthase (iNOS) inhibition in endothelial cells (351).

C. PPAR γ and VEGF

PPAR γ activators have been shown to modulate *in vivo* vascular endothelial growth factor (VEGF)-induced angiogenesis and also *in vitro* differentiation of endothelial cells into tubelike structures. In addition, VEGF is known to play a role in endothelial cell proliferation, migration, vascular permeability, and atherosclerosis. Several studies demonstrated that PPAR γ agonist inhibition of VEGF-induced angiogenesis may be PPAR γ dependent (Fig. 4), part of which includes the inhibition of VEGF receptors and urokinase plasminogen activator expression along with increased PAI-1 expression, NO, and apoptosis (185, 387). Rosiglitazone has been shown to decrease VEGF secretion and indirectly to inhibit angiogenesis in tumor endothelial cells (282). However, a recent study found that administration of GW1929, through PPAR γ -mediated signaling, increases *in vitro* endothelial cell tube formation and *in vivo* neovascularization that is associated with elevated VEGF (33).

D. PPAR γ and EC migration

PPAR γ ligands are also involved in antimigratory actions of endothelial cells. VEGF-induced migration of HUVECs is inhibited by troglitazone and ciglitazone, providing evidence of PPAR γ ligand antimigratory effects on endothelial cells. Moreover, the effects of PPAR γ ligands on EC migration include inhibition of Akt phosphorylation (129) (Fig. 4). Leptin, through endothelial *ob* receptor activation, has been shown to promote endothelial cell proliferation, survival, and vascular angiogenesis (38, 331). In addition, leptin can regulate endothelial cell Akt phosphorylation (366) and migration (128). The administration of PPAR γ ligands inhibits leptin-stimulated Akt phosphorylation and EC migration (128). The tumor-suppressor phosphatase and tensin homologue (PTEN), a modulator of the PI3K/Akt signaling pathway, has been reported to attenuate VEGF-induced EC migration through the inhibition of Akt phosphorylation (158), and PTEN levels were found to be elevated after administration of PPAR γ ligands, suggesting the possibility that PTEN plays a role in the inhibitory actions of TZDs on VEGF- and leptin-induced Akt phosphorylation and endothelial cell migration (128) (Fig. 4). Another study, by using scrape-wound and chemotactic assays, found that troglitazone inhibits endothelial cell migration in high-glucose media (146). Troglitazone was shown to accelerate endothelial cell coverage and repair after rat aortic balloon injury. However, the *in vivo* data suggest that endothelial repair may have occurred as a result of troglitazone-induced suppression of endothelial cell apoptosis rather than a reduction in endothelial cell migration (146). A PPAR γ -mediated mechanism for TZD-induced migratory activity is not suggested in this study. Moreover, further evidence suggests that the effects of TZD treatment pertaining to endothelial cell migration might occur through PPAR γ -independent signaling (204).

E. PPAR γ and EC apoptosis

Previous studies suggest that 15d-PGJ₂ and ciglitazone may induce endothelial cell apoptosis through a PPAR γ -

mediated signaling pathway (34, 213). Our laboratory found that administering a PPAR γ antagonist did not block 15d-PGJ₂-induced inhibition of platelet-derived growth factor (PDGF), providing evidence that 15d-PGJ₂ apoptotic and antiproliferative effects may be PPAR γ independent in endothelial cells (409). However, PPAR γ 1 was reported to induce apoptotic genes in HUVECs (169), and a study with PPAR γ gain- and loss-of-function techniques found PPAR γ to be critical to endothelial cell apoptosis (10). Rosiglitazone was shown to inhibit angiogenesis through a PPAR γ -dependent proapoptotic pathway in HUVECs (185). The induction of apoptosis is possibly beneficial, because activated cells may produce cytokines. In cases of severe pulmonary hypertension, lung arterioles consist of phenotypically altered endothelial cells that reduce blood flow and elevate blood pressure. PPAR γ -mediated EC apoptosis could be beneficial in alleviating lumen-obliterating endothelial cell growth (10).

F. PPAR γ and endothelial progenitor cells

Endothelial progenitor cells (EPCs) are circulating vascular progenitor cells that have been shown to stimulate reendothelialization and decrease neointima formation (376). *In vitro* and *in vivo* studies demonstrated that rosiglitazone stimulates angiogenic progenitor cell (APC) differentiation to endothelial cells to promote reendothelialization and vascular protection against injury (377). Rosiglitazone was shown to improve impaired EPC function in diabetic individuals (292). In EPCs isolated from male subjects, rosiglitazone and 15d-PGJ₂ prevented C-reactive protein-induced EPC dysfunction and promoted angiogenesis (367). Rosiglitazone returns migratory activity to baseline in cultured EPCs from diabetic individuals, which may improve impaired EPC function associated with diabetes (291). Pioglitazone has been shown to increase migratory activity of cultured EPCs from patients with coronary artery disease through PPAR γ -dependent signaling (383), as well as to enhance circulating and bone marrow EPC migratory activity (122). Rosiglitazone may also reduce NAD(P)H oxidase and the resultant increase in oxidative stress while enhancing EPC reendothelialization, promoting vessel repair, and improving vascular function (338). Rosiglitazone and pioglitazone, in addition to improving EPC-induced angiogenesis, can attenuate EPC apoptosis (122, 367). A reduction in EPC apoptosis may be of great benefit to individuals with vascular disease (122). PPAR γ inhibition of EPC apoptosis may have significant clinical relevance because previous studies showed that different types of EPCs have different morphology, proliferation rates, survival rates, and gene-expression profiles that contribute to different functions in neovascularogenesis (160, 398). Finally, it has been suggested that many of the beneficial cardiovascular effects from TZD treatment in patients may be due to the positive effects on EPCs (367). The proapoptotic data in ECs and antiapoptotic data in EPCs may be due to different PPAR γ functions in these cells. The role of PPAR γ -independent effects on apoptosis in these cells is a possibility and also should be considered.

V. PPAR γ and Vascular Smooth Muscle Cells

In 1998, three investigative groups reported evidence of PPAR γ expression in rat aortic and human VSMCs (164, 239, 340). Similarly, a later study observed that PPAR γ expression is present in early human vascular lesions and is upregulated

in rat aortic smooth muscle cells after balloon injury (198). Another study reported that both human coronary artery and aortic VSMCs express PPAR γ 1 and PPAR γ 2 isoforms (29). PPAR γ mRNA levels were reported to increase in mesenteric arteries of both young and adult spontaneously hypertensive rats (SHRs), suggesting that PPAR γ expression is differentially regulated in SHRs (88). Similar data regarding mRNA expression in SHRs were reported from our laboratory. However, we found PPAR γ protein expression and function from SHR vascular smooth muscle cells to be lower compared with those in Wistar-Kyoto rats. It is likely that the suppressed PPAR γ function is a result of decreased protein expression, which could explain the increased VSMC proliferative activity in SHRs (388).

A. PPAR γ and VSMC proliferation

TZDs were reported to attenuate VSMC proliferation and regulate vascular tone well before being identified as PPAR γ ligands (98, 337, 407). Troglitazone was initially found to suppress basic fibroblast growth factor (bFGF)-induced vascular smooth muscle cell growth, preventing rat aortic neointima formation after endothelial injury (199) (Fig. 6). Further studies also confirmed the antiproliferative activity of troglitazone on human VSMCs (29, 250). However, these initial studies did not examine whether the vasculoprotective effects of troglitazone were PPAR γ mediated. A later study with a balloon-injury model confirmed that the inhibitory effect of troglitazone on VSMC proliferation occurs through a PPAR γ -mediated pathway (198). TZDs (troglitazone, rosiglitazone, and pioglitazone) inhibit VSMC proliferation in several human vascular cell beds. The particular TZD administered rather than the vascular source is critical for the potential suppression of VSMC proliferation (79).

C-fos is involved in the MAPK pathway, which plays a role in cell proliferation. Troglitazone attenuates bFGF-induced c-fos expression in cultured VSMCs by inhibiting the MAPK signaling pathway (199) (Fig. 6). A later study also found troglitazone to inhibit PDGF-induced c-fos mRNA expression (29) (Fig. 6). Finally, a recent report demonstrated that rosiglitazone and PPAR γ overexpression suppress bFGF-induced c-fos mRNA expression (Fig. 6). Moreover, PPAR γ dominant negative gene transfer attenuates rosiglitazone-induced inhibition of c-fos mRNA expression (223).

Connective tissue growth factor (CTGF) has the ability to regulate many transforming growth factor-beta (TGF- β) responses in VSMCs, including proliferation, migration, and fibrosis. Data from our laboratory demonstrated that PPAR γ interrupts the Smad3 signaling pathway, inhibiting TGF- β -stimulated CTGF expression in human aortic smooth muscle cells (HASMCs) (114) and suggesting crosstalk between PPAR γ and TGF- β pathways (Fig. 6). We found that TGF- β induces early PPAR γ stimulation and late PPAR γ inhibition of gene expression and that growth factor- and cytokine-induced PPAR γ expression is inhibited by TGF- β . Early activation of TGF- β -induced PPAR γ is mediated by early growth response-1 (Egr-1) signaling, whereas inhibition of PPAR γ by TGF- β is mediated by Smad3, AP-1, and Nab2 (112) (Fig. 6).

Studies from our laboratory also provided the first evidence that the PI3-kinase/Akt-dependent pathway is a regulator of PPAR γ 1 gene expression in VSMCs. We reported that PPAR γ 1 is upregulated by PDGF *via* PI3-kinase/Akt signaling (115) (Fig. 6). Dominant negative overexpression of the p85 subunit from PI3-kinase or Akt proteins also suppresses PDGF-induced PPAR γ expression (115). We also found Egr-1 to be the transcriptional regulator of both growth factor- and cytokine-induced VSMC PPAR γ 1 gene expression. Our results demonstrate that PPAR γ is involved in a feedback mechanism that negatively controls VSMC activation (111).

Angiotensin II plays a crucial role in controlling the proliferation and migration of VSMCs. Troglitazone blocks AngII-induced MAPK activation of VSMCs (140) (Fig. 6). One possible mechanism includes the attenuation of PKC nuclear activity and PKC-mediated extracellular signal regulated kinase 1/2 (ERK 1/2) translocation to the nucleus (132). Another mechanism of AngII-induced VSMC proliferation involves the upregulation of AT1 receptors. PPAR γ ligands have been reported to be responsible for the inhibition of AT1 expression in VSMCs (343, 349). Further, it was suggested that ligand-activated PPAR γ inhibits AT1 transcription by blocking Sp1, leading to the suppression of AT1-receptor expression (343). Finally, telmisartan, an AT1-receptor antagonist with partial PPAR γ activator properties, inhibits AT1-receptor expression. Conversely, administration of the PPAR γ antagonist GW9662 attenuates telmisartan-induced inhibition of AT1, confirming a participatory role for PPAR γ in this signaling cascade (167). Both 15d-PGJ₂ and rosiglitazone were shown to decrease




FIG. 6. Schematic view of PPAR γ activation in VSMCs. In VSMCs, TZDs attenuate growth factor-induced (e.g., AngII) cell migration, proliferation, and fibrosis in either a PPAR γ -dependent or -independent manner by interfering with growth factor-stimulated signaling pathways. PPAR γ activation exerts antiinflammatory roles by inhibiting the NF- κ B pathway; PPAR γ activation promotes apoptosis *via* inducing IRF-1 or GADD45 expression.

AngII-stimulated eukaryotic initiation factor 4E-binding protein 1 (4E-BP1) and Src homology (SH) 2-containing inositol phosphatase 2 (SHIP2) phosphorylation, suppressing Ang II-induced VSMC growth (28). Rosiglitazone may directly decrease SHIP2 activity (28). A recent study suggests that pioglitazone and rosiglitazone inhibit AngII-induced Rho kinase, a known modulator of VSMC tonicity and proliferation. This may be accomplished through increased cytosolic Src homology region 2-containing protein tyrosine phosphatase-2 (SHP-2) expression and reduced Vav phosphorylation (372). However, the effects of PPAR γ activators on AngII cell signaling and growth are still unclear.

One of the most important mechanisms in preventing VSMC growth involves suppression of cell-cycle signaling. In PDGF- or insulin-stimulated cultured rat VSMCs, PPAR γ ligands prevent proliferation by inhibiting the G₁/S phase, a rate-determining step in cell-cycle progression (373). Cell-cycle suppression likely occurs through decreased phosphorylation of the retinoblastoma protein (Rb) (373), a mediator of G₁/S progression (327). Moreover, PPAR γ agonists prevent mitogen-induced p27(Kip1) degradation (373), a known inhibitor of cdk and Rb phosphorylation (328). A non-TZD partial PPAR γ agonist can attenuate mitogen-induced downregulation of p27(Kip1) and proliferation in rat aortic vascular smooth muscle cells. Furthermore, functional PPAR γ is necessary to obtain maximal antiproliferative effects in VSMCs (42). PPAR γ ligands also attenuate PDGF-induced p21(Cip1) expression through the likely inhibition of PKC- δ phosphorylation and activity in cultured rat aortic smooth muscle cells (374). p21(Cip1) promotes activation of the cyclin/cdk complex that eventually results in G₁/S phase progression (195, 328). Repression of p21(Cip1) may be another mechanism by which PPAR γ attenuates VSMC proliferation. Minichromosome maintenance proteins (MCMs) 6 and 7 participate in the initial stages of DNA replication (231) and are considered to be E2F target genes (272). On retinoblastoma phosphorylation, E2F dissociates from Rb and is released for transactivation of DNA synthesis target genes (151). PPAR γ ligands attenuate MCM 6 and 7 expression in VSMCs through the prevention of E2F release from Rb transactivation, further demonstrating that PPAR γ agonists inhibit G₁/S cell-cycle progression, in this case by curtailing pRb/E2F/MCM signaling (43).

Telomerase is important for many cellular functions, including VSMC proliferation. PPAR γ ligand administration was shown to downregulate telomerase activity in cultured VSMCs, because of likely inhibition of telomerase reverse transcriptase (TERT) expression, the catalytic subunit of telomerase. Overexpression of TERT abolishes PPAR γ -ligand inhibition of VSMC proliferation. In addition, the Ets-1 transcriptional factor regulates TERT, and PPAR γ agonists inhibit both Ets-1 mRNA expression and binding to the TERT promoter. Thus, it is likely that PPAR γ ligands target TERT for downregulation to counteract the proliferative properties of vascular smooth muscle cells (269).

Another mechanism suggests that PPAR γ ligands inhibit insulin-induced mitogenic signaling by preventing phosphorylation of the Elk-1 transcription factor (130). A recent *in vitro* study showed that troglitazone attenuates LDL-induced VSMC proliferation and production of superoxide, a contributor to proliferation of VSMCs (153). Finally, PPAR γ has also been shown to induce a differentiated phenotype in

proliferative VSMCs. PPAR γ -dependent signaling increases smooth muscle α -actin (SM- α -actin) and smooth muscle myosin heavy chain (SM-MHC), markers of differentiated VSMCs. Moreover, the effects of PPAR γ on VSMC differentiation appear to be mediated by the GATA-6 transcription factor (4).

B. PPAR γ and VSMC migration

Troglitazone has been shown to inhibit PDGF-induced vascular smooth muscle cell migration (29, 199). In addition to troglitazone, 15d-PGJ₂ (198, 239) and rosiglitazone (198) attenuate PDGF-induced VSMC migration. CTGF is known to be involved in VSMC migration, and data from our laboratory provide evidence that PPAR γ inhibits CTGF expression (114). These studies provide strong support for the involvement of activated PPAR γ in the prevention of VSMC migration that leads to subsequent neointima formation.

Angiotensin II is involved in the control of VSMC proliferation and migration. Troglitazone can block AngII-induced MAPK activation of VSMCs, resulting in the inhibition of VSMC migration (140) (Fig. 6). PPAR γ activators can also inhibit PDGF-, thrombin-, and insulin-like growth factor-1 (IGF-1)-induced VSMC migration through MAPK and downstream nuclear signaling (133). Furthermore, PPAR γ ligands were reported to inhibit PDGF-induced Ets-1 nuclear expression in cultured VSMCs (Fig. 6) or from rat aortic balloon injury. Ets-1 is a transcription factor that is part of ERK/MAPK cell migratory signaling. Moreover, Ets-1 is involved in the transcriptional regulation of matrix metalloproteinase-2 (MMP-2) and matrix metalloproteinase-9 (MMP-9), facilitators of VSMC migration (131) (Fig. 6). PPAR γ activators decrease MMP-9 mRNA and protein expression, along with activity, whereas PPAR γ inactivation through phosphorylation reverses agonist-induced inhibition of MMP-9 expression (239).

C. PPAR γ and VSMC apoptosis

In VSMCs, PPAR γ can signal both growth inhibition (405) and apoptosis (44, 148). PPAR γ activation increases GADD45 expression and caspase-mediated apoptosis (Fig. 6). The Oct-1 protein, a transcription factor regulated by PPAR γ , is critical for PPAR γ -induced GADD45 protein expression (44) (Fig. 6). PPAR γ ligand administration and PPAR γ overexpression have been reported to upregulate interferon regulatory factor (IRF-1) expression, mediating PPAR γ -induced apoptosis in VSMCs (Fig. 6). Further evidence of proapoptotic effects is provided by using an anti-sense approach to suppress IRF-1 expression in VSMCs (224). Pioglitazone is shown to increase apoptosis through PPAR γ -dependent TGF- β release in cultured VSMCs, likely facilitating phosphorylated Smad2 nuclear translocation (303) (Fig. 6). TGF- β -induced apoptosis is mediated, in part, by Smad-dependent GADD45 expression, providing further evidence that GADD45 mediates VSMC apoptosis (397) (Fig. 6). Pioglitazone is also reported to induce apoptosis through Smad2 phosphorylation in cultured VSMCs from both nondiabetic and diabetic patients, usually resistant to induced apoptosis (315). Furthermore, troglitazone can induce apoptosis by activating GADD45 and p53 expression independent of PPAR γ activation (275). Rosiglitazone at high concentrations can more potently induce

apoptosis in intimal compared with medial smooth muscle cells (35).

D. PPAR γ and the regulation of VSMC inflammatory response

CCAAT/enhancer-binding proteins (C/EBPs) are involved in transcriptional regulation of inflammatory cytokines and other proteins. PPAR γ ligands attenuate C/EBP δ expression, and C/EBP δ overexpression reverses PPAR γ ligand inhibition of cytokine gene expression (346). Interestingly, elevations in C/EBP δ levels due to inflammation increase PPAR γ expression and strengthen its antiinflammatory effect in VSMCs (347). In addition, PPAR γ ligands suppress C/EBP δ mRNA and protein levels by dephosphorylating STAT-3 (347), suggesting that PPAR γ and C/EBP δ participate in negative autoregulation feedback. Moreover, PPAR γ overexpression decreases C/EBP δ promoter activity, further indicating the presence of receptor-dependent signaling in C/EBP δ expression (347). This mechanism is likely involved in the suppression of inflammatory cytokines during atherosclerosis (347). Other antiinflammatory responses involving PPAR γ activation include the suppression of TNF- α -induced expression of VCAM-1 (Fig. 6), MCP-1, and fractalkine (CX3CL1) in cultured VSMCs through inhibition of NF- κ B (283).

VI. PPAR γ and Monocytes/Macrophages

PPAR γ expression is present in murine macrophages (8, 307), neointimal lesions (198), macrophage-derived foam cells in both early and advanced stages of atherosclerotic lesions (240, 306), and differentiated human monocyte-derived macrophages (64). However, PPAR γ expression, critical for macrophage lipid metabolism, is not a determinant for macrophage differentiation *in vivo* or *in vitro* (56, 249). PPAR γ is also found in other inflammatory cells, including human peripheral blood T cells (395), human CD4⁺ T cells (236), and mature dendritic cells from the spleen (106). PPAR γ expression is also confirmed in mouse T-helper cells (70). The PPAR γ 1 isoform is found in THP-1 and RAW 264.7 cells (306).

A. PPAR γ and monocyte/macrophage inflammatory signaling

Macrophages are often considered to be heterogeneous and respond to various signaling cascades (365). Different cytokines determine the type of stimulatory or inhibitory response on inflammatory signaling by inducing either a "classic" or "alternative" activation pathway in macrophages. Th1 cytokines, including lipopolysaccharide (LPS), IFN- γ , and IL-1 β , tend to be involved in "classic" activation, whereas Th2 cytokines, including IL-4 and IL-13, likely activate the "alternative" pathway. M1 macrophages are involved in proinflammatory cytokine expression and oxidative stress, whereas M2 macrophages play a role in apoptotic cell phagocytosis, sequestering of pathogens, and wound healing (267, 341). Moreover, macrophages demonstrate functional plasticity because they have the ability to switch between M1 and M2 states of activation (295).

PPAR γ was shown to be necessary for monocyte-derived M2 macrophage phenotype expression (37). PPAR γ is also upregulated during M1 switching to an M2 phenotype, which is critical for increased expression of CD36 (31), arginase I


(267), and the mannose receptor (37). PPAR γ has been shown to regulate M1/M2 switching, in part by reducing inflammatory cytokine expression normally associated with an M1 phenotype, such as TNF- α , IL-1 β , and IL-6 (174), and suppressing *in vitro* macrophage activation (307). The suggestion that PPAR γ is an inflammatory regulator is further illustrated by the belief that PPAR γ may reverse suppression of cytotoxic T lymphocytes, normally a function of M2 activation (364). In addition, specific genes from both M1 and M2 macrophages were found to be unaltered when administering TZD (56, 154, 382).

PPAR γ participates in antiinflammatory signaling to protect against atherosclerotic lesion formation, in part, through negative regulation of macrophage transcriptional activity. PPAR γ ligands, in a PPAR γ -dependent manner, attenuate monocyte and macrophage MMP-9 expression and secretion (186, 240, 307, 330), iNOS, and scavenger receptor-A (SR-A) through the likely inhibition of AP-1, STAT, and NF- κ B transcription factor signaling (307). In addition, PPAR γ negatively regulates a specific population of pro-inflammatory genes controlled by these transcription factors (307, 330) (Fig. 7). PPAR γ activation also inhibits macrophage osteopontin (OPN) expression by interfering with nuclear factor binding to the homeobox-like A/T rich region of the OPN promoter, providing another example of PPAR γ inhibition of macrophage gene expression (277, 278). Similarly, PPAR γ ligands were shown to inhibit proinflammatory cytokine (IL-6, IL-1 β , TNF- α) expression in monocytes (174) (Fig. 7). However, PPAR γ may not be required for IFN- α - or LPS-induced proinflammatory cytokine secretion in macrophages (56, 249). Moreover, it is possible that PPAR γ ligands can upregulate antiinflammatory cytokines (Fig. 7), such as the IL-1-receptor antagonist (IL-1Ra), suggesting another way by which PPAR γ can suppress proinflammatory activity (245). PPAR γ also regulates inflammatory signaling in cells other than monocytes and macrophages. PPAR γ activators can suppress IL-2 (70, 236, 395, 396), IFN- γ (236), and TNF- α (236) in human and animal lymphocytes. PPAR γ ligands also decrease CD40-induced IL-12 secretion in dendritic cells (106).

B. PPAR γ and monocyte/macrophage migration and apoptosis

In addition to antiinflammatory properties, PPAR γ ligands inhibit monocyte/macrophage migration. Troglitazone or rosiglitazone administration results in the inhibition of MCP-1-induced monocyte migration (186). Furthermore, oxidized low-density lipoproteins (oxLDLs) may attenuate MCP-1-dependent monocyte migration by inhibiting chemokine receptor 2 (CCR2) expression (145). Both 9-HODE and 13-HODE, components of oxLDL that stimulate monocyte differentiation to macrophages, inhibit macrophage migration and enhance macrophage adhesion to VSMCs by upregulating CX3CR1 and decreasing CCR2 expression through a PPAR γ pathway (26), suggesting a proinflammatory role for macrophage PPAR γ that may lead to the development of atherosclerosis. Moreover, a recent study showed that PPAR γ -dependent signaling increases CXCR2 receptor expression in primary human macrophages, providing further evidence that PPAR γ can also have proinflammatory properties (309). Next, PPAR γ ligands can also induce apoptotic activity by blocking the NF- κ B antiapoptotic signaling

FIG. 7. Schematic view of PPAR γ roles in atherosclerosis. PPAR γ ligands increase CLA, ABCA1, and ABCG1 expression, leading to improved lipid homeostasis. PPAR γ agonists also decrease proinflammatory cytokine and gene expression and increase antiinflammatory cytokine expression. PPAR γ ligands increase SR-B expression, which promotes cholesterol efflux. Conversely, PPAR γ activation upregulates CD36 expression, resulting in increased oxLDL uptake. Increased oxLDL levels further stimulate PPAR γ expression, which leads to increased CD36 expression. Finally, loss of PPAR γ increases CCR2 expression and monocyte recruitment.



cascade in human macrophages (64). Finally, PPAR γ activation during differentiation of human monocytes to macrophages decreases the ability to engulf apoptotic neutrophils (232).

C. PPAR γ and monocyte/macrophage iNOS expression

Studies have shown that the ability of PPAR γ to repress iNOS expression (159, 217, 307) may occur through direct interaction with the CREB-binding protein (CBP) (217). Furthermore, a recent provocative report suggested another mechanism by which PPAR γ represses iNOS and other proinflammatory genes in murine macrophages. SUMO-1 covalently modifies several transcription factors, including PPAR γ (271). SUMOylation of PPAR γ results in binding to the nuclear-receptor co-repressor (N-CoR)-histone deacetylase-3 (HDAC-3) complex, repressing proinflammatory signaling, particularly NF- κ B target genes (270, 287). Furthermore, PPAR γ and the glucocorticoid receptor were found to inhibit iNOS expression through at least two different signaling pathways (270).

D. PPAR γ and monocyte/macrophage CD36 expression

CD36 is a scavenger receptor that promotes uptake of oxLDL (101). Ligand-dependent PPAR γ has been shown to increase CD36 expression through various signaling pathways in both cultured monocytes and macrophages (159, 254, 358). By using embryonic stem cell-derived macrophages, two studies reported that PPAR γ is required for ligand-activated CD36 gene regulation (56, 249). Macrophages from PPAR γ conditional knockout mice are shown to have decreased CD36 expression compared with wild-type macrophages (8). However, although CD36 is a PPAR γ target gene, PPAR γ is not mandatory for oxLDL uptake in differentiated macrophages (56). Moreover, an *in vivo* study showed that TZDs decrease macrophage CD36 protein expression in *ob/ob* mouse models that display characteristics of insulin resistance, diabetes, and obesity, all of which are risk factors for atherosclerosis (219). TGF- β phosphorylation of PPAR γ has

been suggested as an inhibitory mechanism of action regarding PPAR γ -mediated CD36 expression (143).

E. PPAR γ and monocyte/macrophage lipid homeostasis

A role for PPAR γ activation in macrophage cholesterol homeostasis has been established. CLA-1 is a high-density lipoprotein (HDL) receptor involved in cellular cholesterol removal. CLA-1 was shown to be upregulated by PPAR γ ligands in differentiated human macrophages (63) (Fig. 7). PPAR γ ligands also demonstrate a role in reverse-cholesterol transport by upregulating expression of ATP-binding cassette (ABC) transporters ABCA1 (11, 57, 65) and ABCG1 (8, 11) in monocytes and macrophages (Fig. 7), possibly through an LXR- α -mediated transcriptional signaling pathway (57) that may include caveolin-1 (227). This is important, because an atheroprotective role for granulocyte-macrophage colony-stimulating factor (GM-CSF) may involve PPAR γ and ABCA1 signaling (92). Providing further evidence, a PPAR γ conditional knockout mouse model displays a reduction in macrophage cholesterol efflux, although this study found that troglitazone attenuates cholesterol efflux and ABCA1 expression in macrophages from both PPAR γ knockout and wild-type mice, suggesting some PPAR γ -independent effects (8). Finally, although PPAR γ is not required for oxLDL uptake in differentiated macrophages (56), oxLDL uptake is worsened in PPAR γ -deficient macrophages (249). This finding further indicates an important role for PPAR γ in oxLDL lipid trafficking.

VII. PPAR γ and Atherosclerosis

Diabetes has been estimated to increase the risk of developing atherosclerosis by twofold (178). Increasing evidence suggests that failure to maintain normal glycemic control influences the development of atherosclerosis (142, 356). As previously mentioned, PPAR γ is expressed in atherosclerotic lesions (240, 306). Monocytes differentiate into macrophages on migration into the vessel wall. In macrophages, oxLDL uptake occurs through scavenger receptors, promoting the expression of foam cells (127, 254). Initially, PPAR γ was

thought to be proatherosclerotic. PPAR γ ligand administration, combined with an RXR agonist, upregulates oxLDL uptake through increased CD36-receptor expression (Fig. 7). Furthermore, oxLDL exposure increases SR-A and CD36 mRNA expression through a PPAR γ -dependent mechanism, signaling further oxLDL cellular uptake (254) (Fig. 7). Moreover, PPAR γ is highly expressed in foam cells (358). PPAR γ also is found to be highly expressed in cultured CD36⁺ HASMCs, and troglitazone treatment upregulates CD36 expression only in CD36⁺ smooth muscle cells, suggesting that VSMCs may be able to obtain a macrophage-like phenotype and differentiate into foam cells (242). Furthermore, LPA, a PPAR γ ligand synthesized during mild oxidation of LDL (332), and other PPAR γ agonists were also shown to increase neointima formation in rats (406). Collectively, these studies suggest that PPAR γ is involved in the development of atherosclerosis. Another study found that oxLDL uptake was decreased in PPAR γ -deficient macrophages, partly due to loss of CD36 expression. However, troglitazone treatment had no effect on intracellular oxLDL levels (249). A likely explanation is that troglitazone stimulates CD36 while suppressing SR-A expression (249). It is likely that TZD increases neither macrophage intracellular cholesterol levels nor foam cell formation.

However, the majority of studies suggest an atheroprotective role for TZDs and PPAR γ . PPAR γ -ligand treatment increases scavenger receptor B (SR-B) expression in atherosclerotic lesion macrophages of ApoE^{-/-} mice, potentially facilitating cholesterol efflux (63) (Fig. 7). Treatment with rosiglitazone and GW7845 inhibits atherosclerosis in male low-density lipoprotein receptor knockout (LDL-R^{-/-}) mice although CD36 expression is increased. Interestingly, PPAR γ ligand treatment did not reduce atherosclerosis in female mice. Hormonal differences could be an explanation for the dissimilar outcome between genders (215). In male LDL-R^{-/-} mice fed either a high-fructose or high-fat diet, troglitazone can suppress atherosclerotic lesion formation (72). Next, rosiglitazone reduces aortic atherosclerotic lesions in both diabetic and nondiabetic apoE^{-/-} male mice (212). Finally, rosiglitazone treatment is associated with increased ABCA1 gene expression (Fig. 7) and decreased macrophage accumulation in diabetic mice, providing further evidence of an antiatherosclerotic role (48).

LDL-R^{-/-} mice given transplants with bone marrow deficient in PPAR γ demonstrate an overall worsening of atherosclerosis (57). Next, bone marrow generated from macrophage PPAR γ knockout (MacPPAR γ KO) mice was transplanted to LDL-R^{-/-} and wild-type mice. Mice reconstituted with macrophage PPAR γ knockout bone marrow display increased lesion formation in both strains compared with respective controls. In cases of mild or severe hypercholesterolemia, loss of PPAR γ results in increased atherosclerosis, possibly due to increased CCR2 chemokine receptor expression and monocyte recruitment (18) (Fig. 7).

In vitro studies show that functional PPAR γ is more prevalent in intimal VSMCs compared with medial smooth muscle cells. Therefore, intimal vascular smooth muscle cells are a likely target for PPAR γ in regulating antiatherosclerotic effects (35). Another study showed that transfer of the PPAR γ wild-type gene in a rat carotid artery balloon injury model results in decreased neointima formation and that rosiglitazone-induced inhibition of VSMC proliferation and migration


is blunted by PPAR γ -dominant negative gene transfer. However, the effects of rosiglitazone primarily, but not entirely, occur through PPAR γ -mediated signaling (223). In human atherosclerotic plaques, PPAR γ is associated with M2 macrophage marker expression, although PPAR γ activation does not switch M1 macrophages, foam cells, or already differentiated resting macrophages *in vitro* or atherosclerotic plaque macrophages *in vivo* to an M2 phenotype (37).

PPAR γ ligands may also reduce atherosclerotic development by inhibiting IFN- γ -induced increases in MHC-II expression that normally activate T lymphocytes and control immune responses (194). Increased expression of iNOS has been shown in coronary atherosclerotic plaques of patients with unstable angina (84). Troglitazone and 15d-PGJ₂ are found to suppress IL-1 β -induced iNOS production and cytokine-induced NO synthesis in vascular smooth muscle cells. NF- κ B, critical for iNOS transactivation, is down-regulated by both PPAR γ activators in VSMCs (165). Finally, osteoprotegerin (OPG) is involved in the regulation of atherosclerotic lesion calcification. In our laboratory, we demonstrated that PPAR γ ligands or PPAR γ overexpression inhibits OPG expression in human aortic smooth muscle cells (113). The role of PPAR γ in atherosclerosis is controversial, with much of the literature providing the rationale that PPAR γ plays a regulatory role against the development of atherosclerosis. However, several considerations must be taken into account. Pioglitazone binds with less affinity to PPAR γ compared with rosiglitazone, yet has been shown to be more effective at improving patient lipid profiles (135). Many of the beneficial effects of TZD-induced activation of PPAR γ -mediated transcription are still unclear, particularly because the effects of TZDs on PPAR γ -mediated transcriptional activity are tissue specific. Moreover, the biologic effects of PPAR target genes remain largely unestablished, and because PPAR agonists tend to participate in both gene activation and repression, the known biologic effects of PPAR target genes tend to be rather complex. Thus, a need exists for further research regarding the role of PPAR γ and its ligands in atherosclerotic plaque formation, although the literature provides compelling evidence that PPAR γ activation is important for the attenuation of atherosclerosis.

VIII. PPAR γ and the Heart

The role of PPAR γ in the heart is controversial and often paradoxical. First, myocardial PPAR γ expression seems to vary between studies (16, 124, 392, 394). Next, although several reports have demonstrated beneficial effects of PPAR γ agonist administration on the heart (3, 16, 136, 394) (Fig. 8), the effects of TZDs on cardiac function are in question, particularly in humans. A recent study reported that patients who receive rosiglitazone display an increased risk for myocardial infarction and possibly death from cardiovascular events (265) (Fig. 8). *In vivo* administration of TZDs appears to decrease PPAR target gene expression (47, 336). Nonetheless, it is likely PPAR γ agonists exert an indirect action on the heart because PPAR γ has minimal effects on cardiac fatty acid oxidation or PPAR gene expression in cultured myocytes (124). However, a direct role for PPAR γ must be considered because ciglitazone increases insulin-induced glucose transport in cardiomyocytes. Moreover, phosphorylation of Akt residues, Thr308 and Ser473, is required for insulin stimulation of

FIG. 8. Schematic view of PPAR γ roles in the heart. PPAR γ agonists are associated with increased myocardial infarction and cardiovascular events in humans. PPAR γ agonists decrease ischemia/reperfusion injury and cardiac hypertrophy while increasing contractile function in mice. Administration of PPAR γ agonists decreases JNK/AP-1 and NF- κ B signaling pathways and increases carbohydrate oxidation in mice. Mice with cardiac-specific PPAR γ overexpression show a dilated cardiomyopathy phenotype. Moreover, these mice have increased expression of genes involved in glucose transport and fatty acid utilization. Myocardial



PPAR γ -knockout mice display characteristics of cardiac hypertrophy and dilated cardiomyopathy along with increased NF- κ B activity, decreased Akt phosphorylation, and decreased antioxidant gene expression.

glucose transport and is decreased in insulin-resistant cardiomyocytes (248). In particular, because active Akt has been shown to be necessary for glucose transporter 4 (GLUT4) fusion with adipocyte plasma membranes (191), this may support a role for PPAR γ ligand-induced Akt phosphorylation in cardiomyocytes. The possible discrepancy found in endothelial cells (128, 129) and cardiomyocytes may be explained by the use of different stimuli.

Although ciglitazone enhances insulin-stimulated glucose transport, ciglitazone does not improve insulin-stimulated GLUT4 expression in neonatal rat cardiomyocytes (363), adult rat cardiomyocytes (248), or cardiomyoblasts (124). One possible explanation for increased glucose transport is that elevated glucose transporter 1 (GLUT1) expression, not usually seen with insulin-induced glucose uptake, may be a contributing factor, although the mechanisms remain unclear. The cardiomyocyte microtubule network may be important in regulating insulin signaling. Disruption of the microtubule network may prevent the convergence of insulin signaling and GLUT4 vesicle trafficking (248). Conversely, ligand-independent PPAR γ represses GLUT4 gene expression in adipocytes, and rosiglitazone not only alleviates PPAR γ -induced repression of GLUT4, but also facilitates transcription (15). Similarly, PPAR γ 1 and PPAR γ 2 have been shown to repress GLUT4 expression in cardiomyocytes, and this is enhanced by hyperlipidemia, as free fatty acids bind to PPAR γ and further repress GLUT4 transcription (12). Overall, these results suggest that the regulation of glucose transport by insulin may involve PPAR γ -dependent and -independent signaling pathways.

Another proposed mechanism of action involving insulin signaling and PPARs in the cardiovascular system may include the forkhead-box class O (FOXO) family of transcription factors. FOXO1 is highly expressed in adipocytes and may enhance insulin sensitivity (13, 14) through inhibition of PPAR γ 1 and PPAR γ 2 (13). Insulin signaling results in phosphorylation of FOXO1 by Akt (360). FOXO phosphorylation may repress PPAR γ 1 and PPAR γ promoter activity, directly or indirectly leading to increased GLUT4 expression and subsequent improved insulin sensitivity in adipocytes and cardiomyocytes (14).

Transgenic mice overexpressing PPAR γ (MHC-PPAR γ) in the heart were recently generated and characterized (336). However, cardiomyopathy is present at 2 months of age in these mice, with 100% mortality occurring at 5 months. Subsequently, a new transgenic line was generated to circumvent the problem, as these mice display characteristics suggestive of milder cardiomyopathy (Fig. 8). PPAR γ transgenic mice show increases in expression of fatty acid-utilization genes (Fig. 8), similar to MHC-PPAR α mice. Conversely, similar to MHC-PPAR δ mice, glucose-transporter expression is increased in the PPAR γ transgenic model (Fig. 8). Thus, it is possible that combined elevations in cardiac lipid and glucose levels may further potentiate the development of cardiomyopathy (399).

Whole-body PPAR γ deletion is embryo-lethal in murine models (21). To study the function of PPAR γ in the heart, two cardiac-specific PPAR γ -knockout murine models were generated (91, 97); however, these two lines manifest different phenotypes. The first mouse line shows evidence of mild ventricular hypertrophy (Fig. 8) that is further increased by rosiglitazone treatment, suggesting off-target TZD effects on hypertrophy. Systolic function does not seem to be impaired in these cardiac-specific PPAR γ -null mice. NF- κ B activity is increased, and surprisingly, Akt phosphorylation is decreased despite the presence of a hypertrophic phenotype (97) (Fig. 8). The second cardiac-specific murine knockout model demonstrates progressive dilated cardiomyopathy (Fig. 8) in association with mitochondrial oxidative damage and a reduction in the mitochondrial antioxidant, manganese superoxide dismutase (91) (Fig. 8). These models suggest a likely role for PPAR γ in cardiac function as well as in maintaining a proper oxidation/reduction balance.

IX. PPAR α

PPAR α is highly expressed in the liver, with expression in other tissues including heart, kidney, skeletal muscle, small intestine, and brown adipose tissue. Similar to PPAR γ , PPAR α is also expressed in the cardiovascular cells. PPAR α is involved in the expression of genes involved in lipid metabolism, including fatty acid uptake and oxidation. Moreover,

PPAR α , similar to PPAR γ , can play a role in transcriptional repression of certain genes by inhibiting signaling pathways of other transcription factors. The attenuation of proinflammatory signaling is accomplished through this method by downregulating expression of genes involved in promoting the inflammatory response.

In rodent models, PPAR α was shown to be activated by fibrates, hypolipidemic drugs that are involved in peroxisome proliferation and fatty acid oxidation (172). Fibrates include clofibrate, bezafibrate, fenofibrate, and gemfibrozil. Wy-14,643, nafenopin, and clofibrac acid are other hypolipidemic compounds that are PPAR α -activating agents. Warfarin, an anticoagulant, and trichloroacetic acid were also initially described to be stimulators of PPAR α (96). Fatty acids, including linoleic acid and arachidonic acid, were also shown to activate PPAR α and to regulate gene function (138).

A. PPAR α ligands

However, these studies did not demonstrate whether fibrates or fatty acid compounds could directly bind to PPAR α . A ligand-binding assay found that fibrates and certain fatty acids do indeed have binding affinity for PPAR α (109). In addition, GW7647 (40), GW9578 (41), and LY-518674 (393) are known to be PPAR α ligands. PPAR α antagonists are limited in number and include GW6471 (391) and the *N*-acylsulfonamide compounds A and B (325).

B. PPAR α and endothelial cells

PPAR α is expressed in human endothelial cells (83, 170, 240). Moreover, PPAR α activators are involved in several endothelial cell functions. For example, PPAR α agonists can prevent leukocyte recruitment and adhesion to endothelial cells, in part by decreasing VCAM-1 (6, 173, 241, 321), along with ICAM-1 and E-selectin expression (321) (Fig. 9). Downregulation of adhesion molecules by PPAR α activators is likely through inhibition of NF- κ B (241, 321) (Fig. 9). In addition to decreased adhesion molecule expression, PPAR α activators impair leukocyte binding to endothelial cells (6, 173, 241, 321).

PPAR α has been demonstrated to play a role in vascular function. PPAR α ligands inhibit ET-1 synthesis and secretion in endothelial cells through negative regulation of AP-1 (83) (Fig. 9). A possible explanation is that PPAR α activators may suppress, at least in part, PKC activity involved in endothelial cell ET-1 secretion (234). In DOCA-salt rats, fenofibrate prevents increased ET-1 synthesis in mesenteric arteries (163). PPAR α ligands stimulate eNOS expression by PPAR α -mediated signaling (139).

PPAR α has been shown to be involved in endothelial cell inflammatory signaling. One mechanism for endothelial cell PPAR α participation in antiinflammatory pathways may include oxLDL and a phospholipase A2 (PLA2)-dependent pathway, potentially stimulating fatty acid transport protein-1 (FATP-1) expression (81). Another antiinflammatory mechanism suggests that PPAR α ligands may decrease VEGFR2 expression through direct PPAR α /Sp1 interaction in endothelial cells (246). Finally, bezafibrate increases the CuZn superoxide dismutase antioxidant and decreases NAD(P)H oxidase subunit expression in endothelial cells (168). PPAR α ligands have also been shown to attenuate MCP-1 and IL-8 expression in endothelial cells, possibly by PPAR α suppres-




FIG. 9. Schematic view of PPAR α activation in ECs. PPAR α activation attenuates NF- κ B signaling and transcription in ECs, leading to decreased adhesion-molecule expression and inhibition of leukocyte interaction with ECs. PPAR α ligands inhibit ET-1 synthesis by negatively regulating AP-1.


sing NF- κ B (247, 285). Conversely, another study suggests that PPAR α ligands increase MCP-1 and IL-8 expression through a PPAR α -dependent signaling cascade in human aortic endothelial cells (203). Overall, these studies suggest that PPAR α is primarily involved in antiinflammatory signaling, although it is likely that PPAR α may also exert proinflammatory effects.

C. PPAR α and VSMCs

PPAR α is also expressed in human vascular smooth muscle cells (239, 340). As in endothelial cells, PPAR α has an antiinflammatory role in VSMCs. PPAR α activators suppress IL-6 (80, 340), 6-keto-PGF $_{1\alpha}$ (340), along with COX-2 protein and mRNA expression by negatively regulating NF- κ B signaling (340) (Fig. 10). PPAR α agonists may increase VSMC I κ B α , an inhibitory protein that suppresses NF- κ B nuclear translocation (82). HO-1, a PPAR α target gene, is upregulated by PPAR α and contributes to the antiinflammatory effects in VSMCs (193) (Fig. 10). Group IIA secretory phospholipase A2 (sPLA2-II2) is a proinflammatory mediator of atherosclerosis. PPAR α has been shown to repress IL-1 β -induced sPLA2-IIA expression in VSMCs (298).

In vitro studies have shown that PPAR α ligands inhibit VSMC proliferation (264, 404). One possible mechanism may involve PPAR α activation of p16^{INK4a} (Fig. 10), a cdk inhibitor that blocks phosphorylation of the retinoblastoma protein and subsequent G $_1$ /S cell-cycle progression (126). Next, epoxide hydrolase inhibitors activate PPAR α and suppress PDGF-induced VSMC proliferation through negative regulation of cyclin D1 expression (261). Finally, HO-1, in addition to antiinflammatory signaling, also has a role in VSMC anti-proliferation (193) (Fig. 10). PPAR α also has been shown to regulate VSMC migration. Integrins are critical for VSMC migration in atherosclerosis. PPAR α may interact with Smad4 and inhibit TGF- β -induced beta5 integrin expression in VSMCs (187) (Fig. 10). In addition, docosahexaenoic acid may

FIG. 10. Schematic view of PPAR α activation in VSMCs. PPAR α activation in VSMCs inhibits proliferation and migration by interfering with cdk and β 5-integrin signaling pathways. PPAR α activation also exerts anti-inflammatory roles *via* inhibiting NF- κ B mediated-inflammatory factor release.




regulate VSMC apoptosis through PPAR α -dependent p38 MAPK signaling (89).

D. PPAR α and monocytes/macrophages

PPAR α is expressed in differentiated human macrophages (64) and atherosclerotic lesion macrophages (63). This is important because differentiated macrophages play an important role in inflammation and plaque formation. The first evidence for a role of PPAR α in inflammatory control demonstrated that PPAR α -null mice display a prolonged response to inflammatory stimuli. Leukotriene B4 (LTB4) binding to PPAR α results in activation of fatty acid oxidation (FAO) enzymes that degrade fatty acid and disrupt inflammatory signaling (86).

Further evidence that PPAR α plays a protective role against the inflammatory response is shown from experiments using RAW 264.7 mouse macrophages, whereby Wy14,643 reduces nitrate accumulation in association with decreased iNOS and elevated HO-1 (Fig. 11). Interestingly, natural PPAR α ligands, such as LTB4 and 8(S)-HETE, increase nitrate accumulation, an indication of proinflammatory activity. This difference may be due to variable selectivity to PPAR α (74). Other studies in monocytes/macrophages provide evidence of PPAR α -dependent antiinflammatory signaling. Fenofibrate suppresses LPS-induced MMP-9 secretion in monocytes (330) (Fig. 11). PPAR α also has been shown to downregulate the platelet-activating factor (PAF) receptor, possibly regulating monocyte and macrophage inflammatory responses and cellular apoptosis (156).

FIG. 11. Schematic view of PPAR α ligand roles in macrophages and atherosclerosis. PPAR α ligands may prevent atherosclerosis by improving cholesterol homeostasis, decreasing lipid accumulation, and participating in antiinflammatory signaling in macrophages. LDL-R $^{-/-}$ mice transplanted with bone marrow from PPAR α $^{-/-}$ mice have increased atherosclerosis, whereas GW7647 decreases lesion development in LDL-R $^{-/-}$ mice. However, PPAR α $^{-/-}$ /apoE $^{-/-}$ mice are protected against the development of atherosclerosis.



Treatment with Wy-14,643 and bezafibrate inhibits osteopontin expression in human macrophages through AP-1 inhibition (Fig. 11). Moreover, osteopontin expression is not suppressed in macrophages lacking PPAR α expression (255). Another antiinflammatory mechanism points to simvastatin inhibiting PKC-induced phosphorylation of PPAR α that may result in reduced iNOS and IL-6 expression in macrophages (288). PPAR α ligands inhibit IFN γ , TNF- α , and IL-2 proinflammatory cytokine expression in human T cells (236). Conversely, ligand-activated PPAR α can increase ROS production in mouse and human macrophages (352).

ApoB-48R is involved in macrophage lipid accumulation. Wy-14,643 attenuates apoB-48R expression in both monocytes and macrophages (147) (Fig. 11). Lipoprotein lipase (LPL) hydrolyzes the lipids of lipoproteins and is generally considered to be expressed in cells of atherosclerotic plaques, including macrophage-derived foam cells (266). Although several studies demonstrated that PPAR α activators increase LPL mRNA in macrophages (120, 216, 412), conflicting evidence exists regarding LPL secretion. Decreased LPL secretion due to PPAR α activators may reduce glycated LDL uptake witnessed in macrophages (120) (Fig. 11). Conversely, increased LPL secretion could stimulate PPAR α target gene expression in macrophages or provide an antiinflammatory role by reducing VCAM-1 expression in endothelial cells (412). PPAR α ligands are involved in intracellular cholesterol homeostasis and have been shown to reduce cholesteryl ester formation in human macrophages and foam cells, possibly through upregulation of carnitine palmitoyltransferase type 1 (CPT-1), an enzyme involved in fatty acid degradation (66) (Fig. 11). PPAR α ligands can also regulate reverse cholesterol transport or cholesterol efflux. PPAR α ligands increase CLA-1 expression in differentiated human macrophages (63) (Fig. 11). Furthermore, Wy-14,643 was found to elevate ABCA1 expression in macrophages to facilitate apoA1-induced reverse cholesterol transport (65) (Fig. 11). Niemann-Pick type C1 and C2 (NPC1 and NPC2) proteins control intracellular cholesterol mobilization to the plasma membrane for extracellular transport. PPAR α agonists were found to upregulate NPC1 and NPC2 expression in human macrophages (Fig. 11). In addition, NPC1 and NPC2 inhibition has been shown to prevent ABCA1-mediated extracellular cholesterol transport (62) (Fig. 11). Overall, these studies suggest that PPAR α ligands are actively involved in macrophage cholesterol efflux (Fig. 11). Furthermore, PPAR α ligands have been demonstrated to be regulators of cholesterol homeostasis in both normal and atherosclerotic lesion macrophages (Fig. 11).

E. PPAR α and atherosclerosis

A role for PPAR α has been identified in atherosclerotic lesion formation involving several cell types. As mentioned earlier, PPAR α ligands are critical in controlling macrophage cholesterol homeostasis, and PPAR α has been shown to inhibit VSMC proliferation and migration, important steps in the prevention of atherosclerosis. Wy-14,643 induces SR-B1 expression in atherosclerotic lesions (63). PPAR α also may play a role in atherosclerotic thrombosis by inhibiting tissue factor (TF) mRNA and activity in human monocytes and macrophages (238, 260).

Although much evidence suggests that PPAR α ligands protect against atherosclerosis, murine animal models have

yielded conflicting results. The loss of PPAR α is shown to protect against atherosclerosis in apoE $^{-/-}$ mice (359) (Fig. 11). Conversely, fenofibrate attenuates the development of atherosclerotic lesions, with a more pronounced decrease observed in apoE $^{-/-}$ mice that express the human apoA-I transgene (99). Another study showed that GW7647 decreases lesion formation in LDL-R $^{-/-}$ mice (214) (Fig. 11). Furthermore, lesion size is decreased in human apoE2 knockin mice administered fenofibrate (152). Finally, male and female LDL-R $^{-/-}$ mice transplanted with bone marrow from PPAR α $^{-/-}$ mice display increased aortic atherosclerosis (Fig. 11), along with decreased peritoneal macrophage cholesterol efflux (17). Thus, from these studies, the role of PPAR α in atherosclerotic lesion formation is controversial; however, much of the data tends to suggest an atheroprotective effect of PPAR α .

Possible explanations for decreased atherosclerotic development witnessed with the removal of PPAR α in apoE $^{-/-}$ mice may involve systemic or vessel wall effects. Systemic effects may include decreased glucose levels and insulin resistance, lower blood pressure, and the loss of liver PPAR α target genes that lead to atherosclerotic development. Furthermore, the absence of PPAR α may attenuate LPL activity in the subendothelial space of the vessel wall and decrease atherosclerosis. Systemic effects can alter gene expression in vessel walls, making it difficult to confirm the role of vascular wall PPAR α in atherosclerosis (359).

F. PPAR α and the heart

The use of both gain- and loss-of-function techniques has proven useful in evaluating PPAR α and its effects on cardiac energy metabolism. Cultured myocyte treatment with PPAR α ligands or adenoviral overexpression of PPAR α induces several genes involved in fatty acid metabolism (23, 124, 161). However, the effects of PPAR α ligands on myocardial target genes *in vivo* have been disappointing (75). PPAR α ligands decrease cardiac FAO rates in diabetic mice (1, 2). PPAR α , similar to fatty acid metabolism, may also display direct effects on the heart by inhibiting inflammation and collagen deposition resulting from AngII-induced hypertension. Clinically, PPAR α activation may provide a cardioprotective effect against hypertension and hyperlipidemia. Furthermore, fenofibrate activation of PPAR α may decrease hypertension-induced changes in mechanical overload that lead to ventricular hypertrophy (103).

PPAR α ligands are known to have direct effects on mitochondrial function (180, 181). PPAR α activators can differentially inhibit cardiac mitochondrial respiration. The attenuation of cardiac mitochondrial respiratory function is greater with the administration of fenofibrate compared with Wy-14,643 (413). This suggests a possible PPAR α -independent effect because the Wy-14,643 compound has a higher PPAR α affinity than does fenofibrate.

PPAR α is regulated by hypoxia, as shown by the reduction in PPAR α -dependent transcriptional activity of muscle carnitine palmitoyltransferase I, an enzyme involved in mitochondrial FAO. The DNA-binding activity of the PPAR α :RXR heterodimer is reduced in hypoxic cardiomyocytes (161). Furthermore, myocardial hypoxia can decrease PPAR α -dependent gene expression in two *in vivo* rat models (299). The level of PPAR α mRNA, along with its target gene,

medium-chain acyl-CoA dehydrogenase, is decreased after 7 days in a model of hypoxia-induced right ventricular hypertrophy. However, these levels are upregulated at day 14, suggesting a compensatory response by the heart due to increased load. It is likely that part of the transcriptional response to hypoxia-induced right ventricular hypertrophy involves the regulation of PPAR α by hypoxia in the early stages and, in the later stages, by increased load (323).

PPAR α cardiac-specific transgenic mice were developed to discern between PPAR α -induced cardiac effects and ligand-induced systemic effects (107, 108, 149, 284, 316). PPAR α overexpression induces genes involved in cardiac fatty acid metabolism and utilization (108) while suppressing genes known to participate in glucose uptake and utilization (108, 284). Of great interest, these abnormalities are more prominent in mice that are insulin resistant or fed a high-fat diet (107), both of which are capable of elevating circulating lipids. It is likely that increased reliance of fatty acid utilization, along with the concomitant decrease in glucose utilization by the heart, may aggressively promote remodeling, leading to eventual cardiomyopathy. However, the mechanisms whereby altered fatty acid and glucose utilization result in cardiac remodeling are still unclear.

Although a cardiac-specific PPAR α -knockout mouse model has yet to be characterized, murine models with generalized PPAR α -ablated gene expression have been developed and are often used in examining PPAR α function in cardiac energy metabolism and utilization (53, 93, 183, 205, 211, 229, 281, 381). Malonyl-CoA decarboxylase is an important regulator of cardiac fatty acid oxidation (77), and PPAR α knockout mice have decreased malonyl-CoA decarboxylase gene expression (53). PPAR α -null mice show decreased fatty acid oxidation rates (53, 93, 211, 381) along with increased glucose metabolism and oxidation (53, 281). As a result, it is possible that the alterations pertaining to the dependence on each fuel source in PPAR α -deficient mice make it difficult for the heart to adapt to increase workloads (53, 229). Furthermore, increased ventricular afterload is improved in PPAR α -knockout mice with GLUT1 overexpression (229), suggesting that glucose ATP production in PPAR α -null mice may not be sufficient to meet the demands of greater cardiac workload. Moreover, because chronic pressure overload deactivates PPAR α (134, 208), this model may be suitable for studies in cardiac metabolic dysfunction. PPAR α activation can reduce cardiomyocyte hypertrophy, as fenofibrate decreases ET-1-induced neonatal rat cardiomyocyte enlargement (171, 220). A recent investigation with PPAR α -knockout mice demonstrated greater cardiac hypertrophy after pressure overload in association with enhanced inflammatory marker expression (335), and the follow-up study asserts that PPAR α and PPAR δ inhibit inflammation and cardiac hypertrophy by suppressing NF- κ B signaling (334).

XV. PPAR δ

PPAR δ is distributed ubiquitously in almost all tissues, including liver, fat, skeletal muscle, and skin, and differs from the other two PPAR isotypes. Several studies show that PPAR δ has important roles in cell growth, differentiation, placenta growth, colon tumorigenesis, and wound healing (20, 289, 350). Recent studies focused on the effects of PPAR δ regarding lipid metabolism and insulin sensitivity. PPAR δ is

expressed in the vascular system and displays essential regulatory roles in vascular biology.

PPARs, liver X receptors (LXRs), farnesoid X receptor (FXR), and krüpple-like factor (KLF) are transcription factors controlling lipid and glucose metabolism, as well as the inflammatory response. These transcription factors interact with each other and synergistically regulate gene expression. PPAR δ overexpression influences the activity of PPAR α and PPAR γ in 3T3 fibroblasts and nontransformed monkey kidney CV-1 cells (329). PPAR δ inhibits PPAR γ activity by interfering with PPAR γ DNA-binding activity and not PPAR γ gene expression in colon cancer cells, which is identified by PPAR δ knockout and gain-of-function approaches (414). LXR can bind to all three PPAR subtypes, and PPAR ligands can regulate LXR/PPAR interaction, as studied by SPR technology (402). LXR induces fatty acid synthesis, whereas PPAR δ induces fatty acid oxidation. Moreover, the diverging effects of PPAR δ and LXR on metabolic gene regulation are apparent because PPAR δ represses the expression of the LXR target gene *angpt13*, and L-165041 enhances the inhibitory effect. The likely mechanism is that PPAR δ competes with LXR for binding to RXR, and L-165041 increases the affinity between PPAR δ and RXR (243). KLF5, a member of the KLF superfamily, is critical for regulation of adipocyte differentiation and energy metabolism (274). KLF5^{+/-} heterozygous mice are not prone to high-fat diet-induced obesity, insulin resistance, and hypercholesterolemia. Under basal conditions, SUMOylated KLF5, unliganded PPAR δ , and co-repressors form a transcription-repressor complex. Once PPAR δ agonists activate PPAR δ , KLF5 is deSUMOylated and associates with the transcription activation complex composed of liganded PPAR δ and the CREB binding protein (273).

A. PPAR δ ligands

PPAR δ , on activation by ligands, regulates gene expression. For several years, highly selective PPAR δ ligands were not known, and as a result, the progress in PPAR δ research was hampered. Natural ligands, such as unsaturated fatty acids, eicosanoid derivatives, and prostaglandins, have binding affinity for PPAR δ , although natural ligand selectivity tends to be low (389). cPGI activates both PPAR δ and PPAR α (221), whereas retinoid acid activates both RAR and PPAR δ without activating PPAR α and PPAR γ (324). As a result, synthetic ligands were developed to widen this research scope. L-796449, L-165461, and L-783483 have high affinity for PPAR δ , but also to PPAR γ , whereas L-165041 has a high affinity for only PPAR δ (30). Both GW501516 and GW0742 are widely used and are 1,000 times more selective for PPAR δ compared with PPAR α and PPAR γ . The EC₅₀ of PPAR δ transactivation is 1~2 nM (345). PPAR δ and RXR form an obligatory heterodimer and recruit co-repressors such as BCL-6 and SMART to form a transcription complex that binds to the gene-promoter PPRE. Once ligand activated, co-repressors dissociate from the complex, and coactivators such as p300 and SRC-1 bind to the complex, transactivating target gene expression. Recently, Shearer *et al.* (326) identified GSK0660 as a potent antagonist of PPAR δ with a binding assay IC₅₀ of ~160 nM. However, GSK0660 is inactive on PPAR α and PPAR γ , with IC₅₀ levels above ~10 μ M. This antagonist will be useful for elucidating the biologic roles of PPAR δ (326).

B. PPAR δ and endothelial cells

Endothelial dysfunction is characterized by endothelial proinflammatory, procoagulant, and profibrotic states. Impaired endothelial cell permeability, together with the previous clinical entities, is a marker of early-stage atherosclerosis. Endothelial activation is induced by several risk factors, including LDL/oxLDL, hypercholesterolemia, hyperglycemia, and cytokines (TNF- α , IL-1 β), which promote increased adhesion molecule expression and ensuing leukocyte-endothelial adhesion. L-165041 inhibits TNF- α -induced MCP-1 secretion and VCAM-1 expression in the EAhy926 cell line (311). Both GW0742 and GW501516 have potent antiinflammatory effects in endothelial cells (Fig. 12), inhibiting inflammatory cytokine (TNF- α and IL-1 β)-induced adhesion molecule expression and ensuing leukocyte-endothelial adhesion in primary HUVECs. The mechanisms of PPAR δ antiinflammatory effects involve the attenuation of oxidative stress through the upregulation of antioxidant genes catalase, CuZn superoxide dismutase, and thioredoxin, as well as control of BCL-6 co-repressor translocation to proinflammatory genes (105).

In endothelial cells, ligand activation of PPAR δ increases human endothelial cell proliferation and angiogenesis *via* upregulating VEGF expression and release (290). Next, PPAR δ activation by either PGI₂ or L-165041 inhibits H₂O₂-induced EC apoptosis *via* upregulation of 14-3-3 epsilon (226). L-165041 and GW501516 activate the 14-3-3 gene YWHAE promoter, increasing 14-3-3 expression in a C/EBP-dependent manner, and not in a PPRE-dependent fashion. PPAR δ regulates expression of C/EBP and forms a transcriptional complex with C/EBP in ECs (45). PPAR δ activation stimulates proliferation and attenuates apoptosis in EPCs through phosphor-

ylated Akt-dependent signaling. These effects promote enhanced vasculogenesis and may be therapeutically beneficial in the treatment of ischemic cardiovascular disease (144).

C. PPAR δ and VSMCs

PDGF, a neointimal stimulator, induces PPAR δ expression *via* the PI3-kinase/Akt pathway in VSMCs (410). *In vivo* data show that PPAR δ is upregulated during the development of vascular lesion formation (410). Overexpression of PPAR δ in VSMCs increases post-confluent cell proliferation (Fig. 12) by modulating cell-cycle checkpoint genes including cyclin A, cdk2, and p57(Kip2) (410). The suppression of PPAR δ expression may mediate the inhibitory effects of prostacyclin synthase on neointimal formation (166). However, the role of PPAR δ in VSMCs is not yet agreed on. Recently, Lim *et al.* (222) reported that L-165041 suppresses rat VSMC proliferation by inhibiting phosphorylation of the retinoblastoma protein and cell-cycle progression. *In vivo* data show that L-165041 attenuates neointima formation in the carotid artery balloon injury model. GW501516 also dose-dependently suppresses TNF- α -induced VSMC proliferation (184). PPAR δ receptors and agonists may play different roles in VSMC proliferation, accounting for the seemingly inconsistent results. TGF- β 1, known as a potent regulator in the pathogenesis of atherosclerosis and restenosis, is upregulated by PPAR δ in VSMCs as a target gene. GW501516 inhibits IL-1 β -induced MCP-1 expression, which is mediated by TGF- β 1 and its effector, Smad3. The expression of TGF- β 1 is upregulated, and proinflammatory genes are suppressed in the thoracic aorta prepared from GW501516-treated mice. Thus, it is apparent the PPAR δ /TGF- β /MCP-1 pathway stimulates PPAR δ antiinflammatory signaling mechanisms (184).




FIG. 12. Schematic view of PPAR δ roles in atherosclerosis. PPAR δ ligands are beneficial against the development of atherosclerosis by regulating lipid homeostasis in humans. PPAR δ ligands attenuate the development of atherosclerosis in mice by decreasing inflammatory gene expression and macrophage migration while increasing plasma HDL. Inhibition of EC and macrophage inflammatory gene expression by PPAR δ ligands prevents the development of atherosclerosis. However, PPAR δ overexpression stimulates VSMC proliferation and macrophage release of inflammatory factors, which may promote atherosclerotic development.

D. PPAR δ and monocytes/macrophages

Macrophage inflammation and lipid dysfunction are involved in the pathogenesis of atherosclerosis. PPAR δ regulates lipid metabolism in macrophages. VLDL activates expression of genes involved in β -oxidation, thermogenesis, lipid mobilization, and carnitine biosynthesis through PPAR δ -dependent signaling. Knocking out PPAR δ has the same effect as PPAR δ agonists on fatty acid utilization in macrophages, indicating that the endogenous unliganded PPAR δ receptor has an inhibitory effect on lipid oxidation (202). GW501516 increases ABCA1 expression and induces apolipoprotein A1-specific cholesterol efflux in macrophages (276). However, different results are achieved in primary human macrophages and THP-1 human monocytes with compound F. Compound F upregulates genes related to lipid accumulation and downregulates genes involved in lipid efflux and metabolism. Both compound F and PPAR δ overexpression promote lipid accumulation in macrophages (371). Alternatively, activated macrophages are believed to improve the metabolic syndrome although the mechanism of modulating alternative activation of tissue macrophages is still unclear. Adipocyte-derived Th2 cytokines IL-13 and IL-4 induce macrophage PPAR δ expression. Both adipose tissue and liver-resident macrophages are activated to the alternative phenotype by PPAR δ , and this switch is beneficial for fatty acid metabolism and improves insulin sensitivity (177, 268).

The removal of PPAR δ leads to downregulation of MCP-1 and IL-1 β expression, attenuating macrophage proinflammatory responses. Overexpression of PPAR δ enhances the inflammatory response, suggesting that endogenous PPAR δ has a proinflammatory effect in macrophages (Fig. 12). However, similar to endothelial cells, PPAR δ agonists have a potent inhibitory effect on macrophage inflammation (Fig. 12). GW0742 inhibits LPS-induced expression of inflammatory genes *iNOS* and *COX-2* in macrophages (382). Ligand-activated PPAR δ regulates the translocation of nuclear repressor BCL-6 to inflammatory genes and controls the inflammatory switch in a ligand-dependent manner (201). Graham *et al.* (141) reported that GW0742X decreases TNF- α expression in peritoneal macrophages and adipose tissue.

Foam cell and subsequent fatty-streak formation play critical roles in atherogenesis. LDL/oxLDL induces macrophage differentiation into foam cells, in which many genes likely modulate the transformation process. One such example may include the regulation of scavenger-receptor expression by the PPAR family. Both compound F administration and overexpression of PPAR δ stimulate PMA-induced macrophage differentiation (370).

E. PPAR δ and atherosclerosis

A deteriorated plasma lipoprotein profile directly affects vascular function. Elevated plasma levels of low-density lipoproteins (LDLs) increase the risk of atherosclerosis. Conversely, the increase of HDLs has a cardiovascular protective effect. Very low density lipoproteins (VLDLs) and their triglyceride components regulate gene expression *via* activation of PPAR δ in macrophages (58). Accumulating evidence demonstrates that PPAR δ regulates lipid metabolism in metabolically active tissues. Adipose tissue-specific activated

PPAR δ protects against obesity and induces expression of genes required for fatty acid oxidation and energy uncoupling. Adipose-specific PPAR δ transgenic mice also show improved overall lipid profiles and reduced plasma triglyceride levels (379), demonstrating a possible atheroprotective effect. GW501516 increases HDL levels and decreases small dense LDL, triglycerides, and insulin in insulin-resistant middle-aged obese rhesus monkeys (276). In St. Kitts vervet atherosclerotic primate models, GW501516 increases plasma HDL-C, apoA-I, and apoA-II concentrations, demonstrating protective effects of PPAR δ on the cardiovascular system (375). However, considerably less is known about the function of PPAR δ on lipid homeostasis in humans.

In vivo results also were observed in human subjects. A clinical study performed in healthy white normolipidemic male subjects showed that plasma triglyceride and LDL levels significantly decline, whereas HDL-C levels are enhanced after 2 weeks of GW501516 administration (339) (Fig. 12). Consistently, Riserus *et al.* (310) reported that GW501516 treatment significantly reduces plasma triglycerides, apoB, and LDL cholesterol in healthy moderately overweight subjects (Fig. 12). Presently, laboratory and clinical studies indicate that lowering lipid levels can be achieved by administering PPAR δ agonists, resulting in improved lipid homeostasis (Fig. 12). With regard to its genetic basis, the lipid-regulating function of PPAR δ is associated with gene polymorphisms (333). Plasma HDL-C levels are elevated in the PPAR δ exon 4 + 15 C/C and exon 7 + 65 G/G genotypes of healthy white subjects with exposure to endurance training compared with those with other genotypes (150).

The role of PPAR δ in atherosclerosis has been identified in an atherosclerotic animal model. PPAR δ ^{-/-} bone marrow transplanted into γ -irradiated LDL-R^{-/-} mice significantly reduced atherosclerosis lesions, likely as a result of the attenuated inflammatory status of macrophages (201). Li *et al.* (214) reported that PPAR δ agonist GW0742 has no effect on atherosclerotic lesions, whereas PPAR α and PPAR γ agonists strongly inhibit atherosclerosis in hypercholesterolemic diet-fed LDL-R^{-/-} mice. However, PPAR δ agonists inhibit inflammatory gene expression (Fig. 12), including IFN- γ , TNF- α , MCP-1, VCAM-1, and ICAM-1 in atherosclerotic lesions (214). It is likely that the antiinflammatory effect of PPAR δ may not reverse the proatherogenic impact of extreme hypercholesterolemia in this animal model. Treatment with GW0742X reduces atherosclerotic lesions in LDL-R-null mice and decreases MCP-1 and ICAM-1 expression in the aorta (141). In the apoE^{-/-} mouse atherosclerotic model, treatment with GW501516 attenuates atherosclerotic lesion formation through multiple pathways, which may include increases in plasma HDL levels, potent antiinflammatory effects, and suppression of macrophage transmigration (Fig. 12). PPAR δ inhibits the chemokine-receptor signaling pathway by increasing the expression of regulator of G-protein signaling (RGS) genes (25). In the AngII-accelerated atherosclerotic model, GW0742 attenuates AngII-induced atherosclerotic lesion formation. GW0742 increases the expression of BCL-6, RGS4, and RGS5 in the vascular wall, which inhibits inflammatory and atherogenic gene expression (348). In agreement with several *in vitro* studies, these *in vivo* data support an atheroprotective role of PPAR δ agonists (Fig. 12).




FIG. 13. Schematic view of PPAR δ roles in the heart. PPAR δ ligands increase myocardial fatty acid utilization genes. Cardiac-specific PPAR δ -knockout mice have decreased myocardial expression of fatty acid oxidation genes along with increased myocardial lipid accumulation, cardiac hypertrophy, and congestive

heart failure. Myocardial PPAR δ overexpression in mice increases expression of genes involved in glucose utilization, which may prevent further injury after ischemia/reperfusion.

F. PPAR δ and the heart

PPAR δ activation by GW0742 increases palmitate oxidation in neonatal and adult cardiomyocytes, meanwhile upregulating the expression of fatty acid oxidation genes (61) (Fig. 13). Consistently, the expression of key fatty acid oxidation genes (mCPT1, ACOX1, UCP3) and also basal myocardial FAO rates decrease in cardiomyocyte-specific PPAR δ -knockout mice (60) (Fig. 13). Cardiac-specific overexpression of PPAR δ increases the expression of GLUT4 and phosphofruktokinase, a glycolytic gene, promoting myocardial glucose utilization, which may contribute to reduced myocardial injury after ischemia/reperfusion (46) (Fig. 13). GW610742X increases fatty acid oxidation after myocardial infarction in both left and right ventricles, along with the upregulation of PPAR δ metabolic target gene expression, such as CD36, CPT1, and UCP3 (176). These studies suggest PPAR δ increases fatty acid oxidation and related gene expression, providing a physiological benefit for metabolic-related heart disease.

Inflammatory responses are involved in the pathophysiological processes of ischemia/reperfusion, hypertrophy, and fibrosis. Much evidence suggests that PPAR α and PPAR γ suppress myocardial inflammatory responses. PPAR δ attenuates LPS-induced expression of TNF- α through inhibition of NF- κ B in cultured cardiomyocytes (90). PPAR δ interacts with the p65 NF- κ B subunit, inhibiting the LPS-induced NF- κ B signaling pathway and decreasing MCP-1 expression in rat cardiomyocytes (293). Furthermore, GW0742 reduces cardiac expression of IL-6, IL-8, MCP-1, and ICAM-1, which are induced by ischemia/reperfusion (403).

Progressive myocardial lipid accumulation and hypertrophy occur in cardiomyocyte-specific PPAR δ -knockout mice (Fig. 13). The function of the PPAR δ -null heart is impaired, characterized by a decrease in rates of contraction and relaxation, decreased cardiac output, and increased left ventricular end-diastolic pressure (60) (Fig. 13). GW0742X reduces right ventricle hypertrophy and lung congestion (176). Furthermore, PPAR δ activation by L-165041 inhibits phenylephrine-induced protein synthesis and increases carnitine palmitoyl-transferase and pyruvate dehydrogenase kinase 4 expression in cultured rat cardiomyocytes (293).

GW501516 inhibits proliferation of cardiac fibroblasts and myofibroblasts and also suppresses differentiation of fibroblasts into myofibroblasts (354). Collagen accumulation is involved in myocardial fibrosis, and GW501516 attenuates AngII-stimulated collagen synthesis in cardiac fibroblasts (354, 408).

PPAR δ is critical for maintaining normal fatty acid oxidation and energy balance in the heart (60), suggesting that PPAR δ and its ligands may be important for cardiac function, distribution of muscle fiber type, and endurance performance

(60, 150). PPAR δ overexpression or activation may be a contributing factor to increasing endurance and may mimic the effects of exercise on muscle metabolism (102, 119). PPAR δ and its ligands have been shown to improve exercise performance and regulate physical endurance and training in skeletal muscle (380). Conversely, exercise has been shown to promote skeletal muscle PPAR δ accumulation in murine animal models (230). The possibility exists whereby increased exercise may activate PPAR δ by facilitating the internalization of certain fatty acids that act as ligands (380). Another possibility is that exercise increases PPAR γ coactivator-1 α (PGC-1 α) expression (137), and PGC-1 α binding to PPAR δ can potentially activate this transcription factor, irrespective of the presence of ligands (379). Furthermore, plasma HDL-C levels are higher in the PPAR δ exon 4 + 15 C/C and exon 7 + 65 G/G healthy white genotypes with endurance training compared with other genotypes (150), and PPAR δ agonist administration increases plasma HDL-C concentrations in various animal models (210, 276, 375). One explanation may be that increased availability of free fatty acids due to exercise activates PPAR δ and promotes reverse cholesterol transport (150). Finally, a recent study demonstrated that GW501516 and exercise training work synergistically to increase running endurance (256). These studies have important cardiovascular significance because running performance in humans appears to be linked more to cardiovascular performance and not to muscle fiber-type distribution (304).

In summary, although all three PPAR isotypes are involved in the metabolic syndrome and cardiovascular disease, evidence suggests that PPAR δ is different from the other two subtypes. The PPAR δ receptor and agonists can sometimes show distinct modes of action. PPAR δ can repress both PPAR γ and PPAR α target gene activity, and PPAR δ repression is likely PPRE dependent (329). PPAR δ improves the metabolic syndrome and cardiovascular activity through potent antiinflammatory effects and regulation of lipid and glucose metabolism. To date, several studies indicate that PPAR δ is a potential therapeutic target for treatment of the metabolic syndrome and cardiovascular diseases, including atherosclerosis and cardiac hypertrophy. PPAR δ appears to act as a “housekeeper” because of its near-ubiquitous expression. Therefore, it is critical for PPAR δ to be further examined regarding its effects on metabolism and the various tissues related to metabolic function.

XVI. Perspective

PPARs have now been firmly entrenched as key players in the cardiovascular system. During the past decade, considerable evidence has been accumulated regarding the role of peroxisome proliferator-activated receptors in cardiovascular diseases and clinical complications related to cardiovascular

abnormalities. PPARs regulate several cell-signaling mechanisms related to cardiovascular health and disease. A continuing need exists for basic science and clinical investigations to understand fully the role of PPAR in the physiology and pathology of cardiovascular-related diseases. Thus, it is important to gain a better understanding of the regulatory role of PPARs in vascular cells and the heart.

TZDs and fibrates are pharmacologic agents that have pleiotropic effects, many of which are beneficial in alleviating cardiovascular abnormalities in animal models. However, this has not necessarily translated into markedly improved clinical cardiovascular outcomes. This may be because of differences in both uptake and effects on target pathways between various animal species and humans. In addition, increasing evidence shows that several beneficial PPAR agonist effects are not from direct participation of PPAR-signaling pathways. No definitive evidence indicates that activated PPAR γ pathways are critical for the beneficial effects of TZDs in the cardiovascular system. Moreover, greater evidence exists that ligand-activated PPAR signaling may play a role in the witnessed pharmacologic side effects of TZDs.

Hence, dual PPAR agonists were generated to circumvent this problem and simultaneously to activate two PPAR isoforms. However, the administration of dual PPAR agonists in the clinical setting has been somewhat disappointing because of increased risks for cardiovascular events. Selective PPAR modulators (SPPARMs) were developed to find newer, safer, and more effective agonists and have been shown to improve the overall clinical profile. The possibility that cardiovascular diseases in patients may be the result of depleted endogenous PPAR ligand concentrations must also be considered. Furthermore, a need exists to conduct a greater number of studies on the role of PPAR antagonists in the cardiovascular system.

The development of animal model systems specifically for studying PPARs and PPAR agonists has led to greater in-


creases in information regarding the mechanisms of these nuclear transcription factors in the cardiovascular system. Because global deletion of PPAR is embryo-lethal, the use of conditional knockout mice (*e.g.*, ECs, VSMCs, macrophages) has been critical to understanding the development of human cardiovascular diseases. Nonetheless, limitations are found in using the mouse model. Genetically modified mice often do not show characteristics evident of the human phenotype. Thus, we need more suitable animal models that may correct for many, if not all, of these characteristics. The use of genetically modified rabbits, pigs, or monkeys may be more appropriate for studying the effects of PPARs and their agonists in the cardiovascular system and for providing a clearer understanding of the pathophysiology of cardiovascular diseases (Fig. 14).

Finally, although previous studies have successfully targeted PPAR for deletion in cardiovascular cells, the possibility of PPAR cell-cell crosstalk should not be overlooked in the cardiovascular system. For example, does VSMC PPAR γ affect function in PPAR γ -null ECs and *vice versa*? The ability to gain a better understanding of PPARs and agonists in the cardiovascular system will enable us to address the controversy regarding the subsequent administration of pharmacologic agents that not only activate PPAR pathways, but may also have PPAR-independent effects.

Acknowledgments

Dr. Chen's laboratory is funded by National Institutes of Health (HL68878, HL89544, HL75397, and HL92421). M.H. is supported by a postdoctoral fellowship from the National Institutes of Health (T32 HL007853). L.C. and J.Z. are supported by American Heart Association Midwest Affiliate Fellowship (0625705Z) and National Career Development Grant (0835237N), respectively. Y.E.C. is an established investigator of American Heart Association.

FIG. 14. Perspective view of PPARs and PPAR ligands in the cardiovascular system. The use of mouse models has shown that PPAR ligands have many beneficial effects in the cardiovascular system. However, PPAR ligand administration (*e.g.*, rosiglitazone) in the clinical setting has not necessarily translated into markedly improved cardiovascular outcomes. Furthermore, some question exists as to whether the beneficial effects of PPAR agonists involve PPAR-dependent signaling. Although the development of animal model systems specifically for studying PPAR agonists and PPAR gain- and loss-of-function has elucidated important findings regarding the molecular mechanisms of cardiovascular disease, certain limitations pertain to the use of mouse models. In many cases, genetically altered murine models do not display characteristics similar to those of humans. Hence, a need exists for using genetically modified animals, such as rabbits, pigs, or monkeys, that have a closer phenotypic resemblance to humans and therefore may be more appropriate for studying PPARs and PPAR agonists in the cardiovascular system.



Abbreviations

4E-BP1, 4E-binding protein 1; 15d-PGJ₂, 15-deoxy- δ 12,14-prostaglandin J₂; ABC, ATP-binding cassette; AGP, 1-O-octadecenyl-2-hydroxy-*sn*-glycero-3-phosphate; AngII, angiotensin II; AP-1, activator protein-1; APC, angiogenic progenitor cell; apoE^{-/-}, apo E knockout; AT1, angiotensin II type 1 receptor; AT2, angiotensin II type 2 receptor; azPC, 1-O-hexadecyl-2-azelaoyl-*sn*-glycero-3-phosphocholine; BADGE, bisphenol A diglycidyl ether; bFGF, basic fibroblast growth factor; CARLA, coactivator-dependent receptor ligand assay; CBP, CREB-binding protein; CCR2, chemokine receptor 2; C/EBP, CCAAT/enhancer-binding protein; CPT-1, carnitine palmitoyltransferase type 1; CTGF, connective tissue growth factor; ECs, endothelial cells; Egr-1, early growth response-1; eNOS, endothelial nitric oxide synthase; EPC, endothelial progenitor cell; ERK 1/2, extracellular signal regulated kinase 1/2; ET-1, endothelin-1; FAO, fatty acid oxidation; FATP-1, fatty acid transport protein-1; FRET, fluorescence resonance energy transfer; FXR, farnesoid X receptor; FOXO, forkhead-box class O; GLUT1, glucose transporter 1; GLUT4, glucose transporter 4; GM-CSF, granulocyte-macrophage colony-stimulating factor; HASMCs, human aortic smooth muscle cells; HDAC-3, histone deacetylase-3; HDL, high-density lipoprotein; HETE, hydroxyeicosatetraenoic acid; HO-1, heme-oxygenase 1; HODE, hydroxyoctadecadienoic acid; HUVECs, human umbilical vein endothelial cells; ICAM-1, intercellular adhesion molecule-1; IFN, interferon; IFN- γ , interferon-gamma; IGF, insulin-like growth factor; I κ B α , I κ appaB-alpha; IKK, I κ appaB kinase; IL, interleukin; IL-1 β , interleukin-1beta; IL-1Ra, IL-1 receptor antagonist; iNOS, inducible nitric oxide synthase; IP-10, IFN-inducible protein of 10 kDa; IRF-1, interferon regulatory factor; I-TAC, IFN-inducible T-cell α -chemoattractant; KLF, kruppel-like factor; LDL, low-density lipoprotein; LDL-R^{-/-}, low-density lipoprotein receptor knockout; LNO₂, nitro-9,12-*cis*-octadecadienoic acid; LPA, lysophosphatidic acid; LPL, lipoprotein lipase; LPS, lipopolysaccharide; LTB₄, leukotriene B₄; LXR, liver X receptor; MAPK, mitogen-activated protein kinase; MCM, minichromosome maintenance protein; MCP-1, monocyte chemoattractant protein-1; MHC-II, major histocompatibility complex class II; Mig, monokine induced by IFN- γ ; MMP-2, matrix metalloproteinase-2; MMP-9, matrix metalloproteinase-9; N-CoR, nuclear receptor co-repressor; NF- κ B, nuclear factor-kappa B; NO, nitric oxide; NPC, Niemann-Pick, type C; OA-NO₂, nitro-9-*cis*-octadecenoic acid; OPG, osteoprotegerin; OPN, osteopontin; oxLDL, oxidized LDL; PAF, platelet-activating factor; PAI-1, plasminogen activator inhibitor type-1; PDGF, platelet-derived growth factor; PECAM-1, platelet-endothelial cell adhesion molecule; PGC-1 α , PPAR γ coactivator-1 α ; PKC, protein kinase C; PLA, phospholipase A2; PPAR, peroxisome proliferator-activated receptor; PPAR α , PPARalpha; PPAR β/δ , PPARbeta/delta; PPAR γ , PPARgamma; PPAR γ E null, endothelial cell PPARgamma knockout; PPRE, peroxisome proliferator response element; PTEN, phosphatase and tensin homologue; Rb, retinoblastoma protein; RGS, regulator of G-protein signaling; RXR, retinoic X receptor; SHIP2, Src homology (SH) 2-containing inositol phosphatase 2; SHP-2, Src homology region 2-containing protein tyrosine phosphatase-2; SHRs, spontaneously hypertensive rats; SM- α -actin, smooth muscle alpha-actin; SM-MHC, smooth muscle myosin heavy chain;

SPA, scintillation proximity assay; sPLA2-II2, secretory phospholipase A2; SPPARMs, selective PPAR modulators; SPR, surface plasmon resonance; SR-A, scavenger receptor A; SR-B, scavenger receptor B; STAT, signal transduction and activator of transcription; TERT, telomerase reverse transcriptase; TF, tissue factor; TGF- β , transforming growth factor-beta; TNF- α , tumor necrosis factor-alpha; TZD, thiazolidinedione; VCAM-1, vascular cell adhesion molecule-1; VEGF, vascular endothelial growth factor; VLDLs, very low density lipoproteins; VSMCs, vascular smooth muscle cells.

References

1. Aasum E, Belke DD, Severson DL, Riemersma RA, Cooper M, Andreassen M, and Larsen TS. Cardiac function and metabolism in type 2 diabetic mice after treatment with BM 17.0744, a novel PPAR-alpha activator. *Am J Physiol Heart Circ Physiol* 283: H949-H957, 2002.
2. Aasum E, Cooper M, Severson DL, and Larsen TS. Effect of BM 17.0744, a PPARalpha ligand, on the metabolism of perfused hearts from control and diabetic mice. *Can J Physiol Pharmacol* 83: 183-190, 2005.
3. Abdelrahman M, Sivarajah A, and Thiemermann C. Beneficial effects of PPAR-gamma ligands in ischemia-reperfusion injury, inflammation and shock. *Cardiovasc Res* 65: 772-781, 2005.
4. Abe M, Hasegawa K, Wada H, Morimoto T, Yanazume T, Kawamura T, Hirai M, Furukawa Y, and Kita T. GATA-6 is involved in PPARgamma-mediated activation of differentiated phenotype in human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* 23: 404-410, 2003.
5. Adams M, Reginato MJ, Shao D, Lazar MA, and Chatterjee VK. Transcriptional activation by peroxisome proliferator-activated receptor gamma is inhibited by phosphorylation at a consensus mitogen-activated protein kinase site. *J Biol Chem* 272: 5128-5132, 1997.
6. Ahmed W, Orasanu G, Nehra V, Asatryan L, Rader DJ, Ziouzenkova O, and Plutzky J. High-density lipoprotein hydrolysis by endothelial lipase activates PPARalpha: a candidate mechanism for high-density lipoprotein-mediated repression of leukocyte adhesion. *Circ Res* 98: 490-498, 2006.
7. Akaike M, Che W, Marmarosh NL, Ohta S, Osawa M, Ding B, Berk BC, Yan C, and Abe J. The hinge-helix 1 region of peroxisome proliferator-activated receptor gamma1 (PPARgamma1) mediates interaction with extracellular signal-regulated kinase 5 and PPARgamma1 transcriptional activation: involvement in flow-induced PPARgamma activation in endothelial cells. *Mol Cell Biol* 24: 8691-8704, 2004.
8. Akiyama TE, Sakai S, Lambert G, Nicol CJ, Matsusue K, Pimprale S, Lee YH, Ricote M, Glass CK, Brewer HB, Jr., and Gonzalez FJ. Conditional disruption of the peroxisome proliferator-activated receptor gamma gene in mice results in lowered expression of ABCA1, ABCG1, and apoE in macrophages and reduced cholesterol efflux. *Mol Cell Biol* 22: 2607-2619, 2002.
9. Alexis JD, Wang N, Che W, Lerner-Marmarosh N, Sahni A, Korshunov VA, Zou Y, Ding B, Yan C, Berk BC, and Abe JI. Bcr kinase activation by angiotensin II inhibits peroxisome proliferator-activated receptor {gamma} transcriptional activity in vascular smooth muscle cells. *Circ Res* 1: 69-78, 2009.

10. Ameshima S, Golpon H, Cool CD, Chan D, Vandivier RW, Gardai SJ, Wick M, Nemenoff RA, Geraci MW, and Voelkel NF. Peroxisome proliferator-activated receptor gamma (PPARgamma) expression is decreased in pulmonary hypertension and affects endothelial cell growth. *Circ Res* 92: 1162–1169, 2003.
11. Argmann CA, Sawyez CG, McNeil CJ, Hegele RA, and Huff MW. Activation of peroxisome proliferator-activated receptor gamma and retinoid X receptor results in net depletion of cellular cholesteryl esters in macrophages exposed to oxidized lipoproteins. *Arterioscler Thromb Vasc Biol* 23: 475–482, 2003.
12. Armoni M, Harel C, Bar-Yoseph F, Milo S, and Karnieli E. Free fatty acids repress the GLUT4 gene expression in cardiac muscle via novel response elements. *J Biol Chem* 280: 34786–34795, 2005.
13. Armoni M, Harel C, Karni S, Chen H, Bar-Yoseph F, Ver MR, Quon MJ, and Karnieli E. FOXO1 represses peroxisome proliferator-activated receptor-gamma1 and -gamma2 gene promoters in primary adipocytes: a novel paradigm to increase insulin sensitivity. *J Biol Chem* 281: 19881–19891, 2006.
14. Armoni M, Harel C, and Karnieli E. Transcriptional regulation of the GLUT4 gene: from PPAR-gamma and FOXO1 to FFA and inflammation. *Trends Endocrinol Metab: TEM* 18: 100–107, 2007.
15. Armoni M, Kritiz N, Harel C, Bar-Yoseph F, Chen H, Quon MJ, and Karnieli E. Peroxisome proliferator-activated receptor-gamma represses GLUT4 promoter activity in primary adipocytes, and rosiglitazone alleviates this effect. *J Biol Chem* 278: 30614–30623, 2003.
16. Asakawa M, Takano H, Nagai T, Uozumi H, Hasegawa H, Kubota N, Saito T, Masuda Y, Kadowaki T, and Komuro I. Peroxisome proliferator-activated receptor gamma plays a critical role in inhibition of cardiac hypertrophy *in vitro* and *in vivo*. *Circulation* 105: 1240–1246, 2002.
17. Babaev VR, Ishiguro H, Ding L, Yancey PG, Dove DE, Kovacs WJ, Semenkovich CF, Fazio S, and Linton MF. Macrophage expression of peroxisome proliferator-activated receptor-alpha reduces atherosclerosis in low-density lipoprotein receptor-deficient mice. *Circulation* 116: 1404–1412, 2007.
18. Babaev VR, Yancey PG, Ryzhov SV, Kon V, Breyer MD, Magnuson MA, Fazio S, and Linton MF. Conditional knockout of macrophage PPARgamma increases atherosclerosis in C57BL/6 and low-density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 25: 1647–1653, 2005.
19. Baker PR, Lin Y, Schopfer FJ, Woodcock SR, Groeger AL, Bathyany C, Sweeney S, Long MH, Iles KE, Baker LM, Branchaud BP, Chen YE, and Freeman BA. Fatty acid transduction of nitric oxide signaling: multiple nitrated unsaturated fatty acid derivatives exist in human blood and urine and serve as endogenous peroxisome proliferator-activated receptor ligands. *J Biol Chem* 280: 42464–42475, 2005.
20. Barak Y, Liao D, He W, Ong ES, Nelson MC, Olefsky JM, Boland R, and Evans RM. Effects of peroxisome proliferator-activated receptor delta on placentation, adiposity, and colorectal cancer. *Proc Natl Acad Sci U S A* 99: 303–308, 2002.
21. Barak Y, Nelson MC, Ong ES, Jones YZ, Ruiz-Lozano P, Chien KR, Koder A, and Evans RM. PPAR gamma is required for placental, cardiac, and adipose tissue development. *Mol Cell* 4: 585–595, 1999.
22. Bardot O, Aldridge TC, Latruffe N, and Green S. PPAR-RXR heterodimer activates a peroxisome proliferator response element upstream of the bifunctional enzyme gene. *Biochem Biophys Res Commun* 192: 37–45, 1993.
23. Barger PM, Brandt JM, Leone TC, Weinheimer CJ, and Kelly DP. Deactivation of peroxisome proliferator-activated receptor-alpha during cardiac hypertrophic growth. *J Clin Invest* 105: 1723–1730, 2000.
24. Barger PM, Browning AC, Garner AN, and Kelly DP. p38 mitogen-activated protein kinase activates peroxisome proliferator-activated receptor alpha: a potential role in the cardiac metabolic stress response. *J Biol Chem* 276: 44495–44501, 2001.
25. Barish GD, Atkins AR, Downes M, Olson P, Chong LW, Nelson M, Zou Y, Hwang H, Kang H, Curtiss L, Evans RM, and Lee CH. PPARdelta regulates multiple proinflammatory pathways to suppress atherosclerosis. *Proc Natl Acad Sci U S A* 105: 4271–4276, 2008.
26. Barlic J, Zhang Y, Foley JF, and Murphy PM. Oxidized lipid-driven chemokine receptor switch, CCR2 to CX3CR1, mediates adhesion of human macrophages to coronary artery smooth muscle cells through a peroxisome proliferator-activated receptor gamma-dependent pathway. *Circulation* 114: 807–819, 2006.
27. Barroso I, Gurnell M, Crowley VE, Agostini M, Schwabe JW, Soos MA, Maslen GL, Williams TD, Lewis H, Schafer AJ, Chatterjee VK, and O'Rahilly S. Dominant negative mutations in human PPARgamma associated with severe insulin resistance, diabetes mellitus and hypertension. *Nature* 402: 880–883, 1999.
28. Benkirane K, Amiri F, Diep QN, El Mabrouk M, and Schiffrin EL. PPAR-gamma inhibits ANG II-induced cell growth via SHIP2 and 4E-BP1. *Am J Physiol Heart Circ Physiol* 290: H390–H397, 2006.
29. Benson S, Wu J, Padmanabhan S, Kurtz TW, and Pershadsingh HA. Peroxisome proliferator-activated receptor (PPAR)-gamma expression in human vascular smooth muscle cells: inhibition of growth, migration, and c-fos expression by the peroxisome proliferator-activated receptor (PPAR)-gamma activator troglitazone. *Am J Hypertens* 13: 74–82, 2000.
30. Berger J, Leibowitz MD, Doebber TW, Elbrecht A, Zhang B, Zhou G, Biswas C, Cullinan CA, Hayes NS, Li Y, Tanen M, Ventre J, Wu MS, Berger GD, Mosley R, Marquis R, Santini C, Sahoo SP, Tolman RL, Smith RG, and Moller DE. Novel peroxisome proliferator-activated receptor (PPAR) gamma and PPARdelta ligands produce distinct biological effects. *J Biol Chem* 274: 6718–6725, 1999.
31. Berry A, Balard P, Coste A, Olagner D, Lagane C, Authier H, Benoit-Vical F, Lepert JC, Seguela JP, Magnaval JF, Chambon P, Metzger D, Desvergne B, Wahli W, Auwerx J, and Pipy B. IL-13 induces expression of CD36 in human monocytes through PPARgamma activation. *Eur J Immunol* 37: 1642–1652, 2007.
32. Beyer AM, Baumbach GL, Halabi CM, Modrick ML, Lynch CM, Gerhold TD, Ghoneim SM, de Lange WJ, Keen HL, Tsai YS, Maeda N, Sigmund CD, and Faraci FM. Interference with PPARgamma signaling causes cerebral vascular dysfunction, hypertrophy, and remodeling. *Hypertension* 51: 867–871, 2008.

33. Biscetti F, Gaetani E, Flex A, Aprahamian T, Hopkins T, Straface G, Pecorini G, Stigliano E, Smith RC, Angelini F, Castellot JJ Jr, and Pola R. Selective activation of peroxisome proliferator-activated receptor (PPAR)alpha and PPAR gamma induces neoangiogenesis through a vascular endothelial growth factor-dependent mechanism. *Diabetes* 57: 1394–1404, 2008.
34. Bishop-Bailey D and Hla T. Endothelial cell apoptosis induced by the peroxisome proliferator-activated receptor (PPAR) ligand 15-deoxy-delta12, 14-prostaglandin J2. *J Biol Chem* 274: 17042–17048, 1999.
35. Bishop-Bailey D, Hla T, and Warner TD. Intimal smooth muscle cells as a target for peroxisome proliferator-activated receptor-gamma ligand therapy. *Circulation research* 91: 210–217, 2002.
36. Bocos C, Gottlicher M, Gearing K, Banner C, Enmark E, Teboul M, Crickmore A, and Gustafsson JA. Fatty acid activation of peroxisome proliferator-activated receptor (PPAR). *J Steroid Biochem Mol Biol* 53: 467–473, 1995.
37. Bouhrel MA, Derudas B, Rigamonti E, Dievart R, Brozek J, Haulon S, Zawadzki C, Jude B, Torpier G, Marx N, Staels B, and Chinetti-Gbaguidi G. PPARgamma activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab* 6: 137–143, 2007.
38. Bouloumie A, Drexler HC, Lafontan M, and Busse R. Leptin, the product of Ob gene, promotes angiogenesis. *Circ Res* 83: 1059–1066, 1998.
39. Brown KK, Henke BR, Blanchard SG, Cobb JE, Mook R, Kaldor I, Kliewer SA, Lehmann JM, Lenhard JM, Harrington WW, Novak PJ, Faison W, Binz JG, Hashim MA, Oliver WO, Brown HR, Parks DJ, Plunket KD, Tong WQ, Menius JA, Adkison K, Noble SA, and Willson TM. A novel N-aryl tyrosine activator of peroxisome proliferator-activated receptor-gamma reverses the diabetic phenotype of the Zucker diabetic fatty rat. *Diabetes* 48: 1415–1424, 1999.
40. Brown PJ, Stuart LW, Hurley KP, Lewis MC, Winegar DA, Wilson JG, Wilkison WO, Ittoop OR, and Willson TM. Identification of a subtype selective human PPARalpha agonist through parallel-array synthesis. *Bioorg Med Chem Lett* 11: 1225–1227, 2001.
41. Brown PJ, Winegar DA, Plunket KD, Moore LB, Lewis MC, Wilson JG, Sundseth SS, Koble CS, Wu Z, Chapman JM, Lehmann JM, Kliewer SA, and Willson TM. A ureido-thioisobutyric acid (GW9578) is a subtype-selective PPARalpha agonist with potent lipid-lowering activity. *J Med Chem* 42: 3785–3788, 1999.
42. Bruemmer D, Berger JP, Liu J, Kintscher U, Wakino S, Fleck E, Moller DE, and Law RE. A non-thiazolidinedione partial peroxisome proliferator-activated receptor gamma ligand inhibits vascular smooth muscle cell growth. *Eur J Pharmacol* 466: 225–234, 2003.
43. Bruemmer D, Yin F, Liu J, Berger JP, Kiyono T, Chen J, Fleck E, Van Herle AJ, Forman BM, and Law RE. Peroxisome proliferator-activated receptor gamma inhibits expression of minichromosome maintenance proteins in vascular smooth muscle cells. *Mol Endocrinol (Baltimore)* 17: 1005–1018, 2003.
44. Bruemmer D, Yin F, Liu J, Berger JP, Sakai T, Blaschke F, Fleck E, Van Herle AJ, Forman BM, and Law RE. Regulation of the growth arrest and DNA damage-inducible gene 45 (GADD45) by peroxisome proliferator-activated receptor gamma in vascular smooth muscle cells. *Circ Res* 93: e38–e47, 2003.
45. Brunelli L, Cieslik KA, Alcorn JL, Vatta M, and Baldini A. Peroxisome proliferator-activated receptor-delta upregulates 14-3-3 epsilon in human endothelial cells via CCAAT/enhancer binding protein-beta. *Circ Res* 100: e59–e71, 2007.
46. Burkart EM, Sambandam N, Han X, Gross RW, Courtois M, Gierasch CM, Shoghi K, Welch MJ, and Kelly DP. Nuclear receptors PPARbeta/delta and PPARalpha direct distinct metabolic regulatory programs in the mouse heart. *J Clin Invest* 117: 3930–3939, 2007.
47. Cabrero A, Jove M, Planavila A, Merlos M, Laguna JC, and Vazquez-Carrera M. Down-regulation of acyl-CoA oxidase gene expression in heart of troglitazone-treated mice through a mechanism involving chicken ovalbumin upstream promoter transcription factor II. *Mol Pharmacol* 64: 764–772, 2003.
48. Calkin AC, Forbes JM, Smith CM, Lassila M, Cooper ME, Jandeleit-Dahm KA, and Allen TJ. Rosiglitazone attenuates atherosclerosis in a model of insulin insufficiency independent of its metabolic effects. *Arterioscler, Thromb Vasc Biol* 25: 1903–1909, 2005.
49. Calnek DS, Mazzella L, Roser S, Roman J, and Hart CM. Peroxisome proliferator-activated receptor gamma ligands increase release of nitric oxide from endothelial cells. *Arterioscler Thromb Vasc Biol* 23: 52–57, 2003.
50. Camp HS, Chaudhry A, and Leff T. A novel potent antagonist of peroxisome proliferator-activated receptor gamma blocks adipocyte differentiation but does not revert the phenotype of terminally differentiated adipocytes. *Endocrinology* 142: 3207–3213, 2001.
51. Camp HS and Tafuri SR. Regulation of peroxisome proliferator-activated receptor gamma activity by mitogen-activated protein kinase. *J Biol Chem* 272: 10811–10816, 1997.
52. Camp HS, Tafuri SR, and Leff T. c-Jun N-terminal kinase phosphorylates peroxisome proliferator-activated receptor-gamma1 and negatively regulates its transcriptional activity. *Endocrinology* 140: 392–397, 1999.
53. Campbell FM, Kozak R, Wagner A, Altarejos JY, Dyck JR, Belke DD, Severson DL, Kelly DP, and Lopaschuk GD. A role for peroxisome proliferator-activated receptor alpha (PPARalpha) in the control of cardiac malonyl-CoA levels: reduced fatty acid oxidation rates and increased glucose oxidation rates in the hearts of mice lacking PPARalpha are associated with higher concentrations of malonyl-CoA and reduced expression of malonyl-CoA decarboxylase. *J Biol Chem* 277: 4098–4103, 2002.
54. Castrillo A, Diaz-Guerra MJ, Hortelano S, Martin-Sanz P, and Bosca L. Inhibition of IkappaB kinase and IkappaB phosphorylation by 15-deoxy-delta(12,14)-prostaglandin J(2) in activated murine macrophages. *Mol Cell Biol* 20: 1692–1698, 2000.
55. Castrillo A, Mojena M, Hortelano S, and Bosca L. Peroxisome proliferator-activated receptor-gamma-independent inhibition of macrophage activation by the non-thiazolidinedione agonist L-796,449: comparison with the effects of 15-deoxy-delta(12,14)-prostaglandin J(2). *J Biol Chem* 276: 34082–34088, 2001.
56. Chawla A, Barak Y, Nagy L, Liao D, Tontonoz P, and Evans RM. PPAR-gamma dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation. *Nat Med* 7: 48–52, 2001.
57. Chawla A, Boisvert WA, Lee CH, Laffitte BA, Barak Y, Joseph SB, Liao D, Nagy L, Edwards PA, Curtiss LK,

- Evans RM, and Tontonoz P. A PPAR gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. *Mol Cell* 7: 161–171, 2001.
58. Chawla A, Lee CH, Barak Y, He W, Rosenfeld J, Liao D, Han J, Kang H, and Evans RM. PPARdelta is a very low-density lipoprotein sensor in macrophages. *Proc Natl Acad Sci U S A* 100: 1268–1273, 2003.
59. Chawla A, Schwarz EJ, Dimaculangan DD, and Lazar MA. Peroxisome proliferator-activated receptor (PPAR) gamma: adipose-predominant expression and induction early in adipocyte differentiation. *Endocrinology* 135: 798–800, 1994.
60. Cheng L, Ding G, Qin Q, Huang Y, Lewis W, He N, Evans RM, Schneider MD, Brako FA, Xiao Y, Chen YE, and Yang Q. Cardiomyocyte-restricted peroxisome proliferator-activated receptor-delta deletion perturbs myocardial fatty acid oxidation and leads to cardiomyopathy. *Nat Med* 10: 1245–1250, 2004.
61. Cheng L, Ding G, Qin Q, Xiao Y, Woods D, Chen YE, and Yang Q. Peroxisome proliferator-activated receptor delta activates fatty acid oxidation in cultured neonatal and adult cardiomyocytes. *Biochem Biophys Res Commun* 313: 277–286, 2004.
62. Chinetti-Gbaguidi G, Rigamonti E, Helin L, Mutka AL, Lepore M, Fruchart JC, Clavey V, Ikonen E, Lestavel S, and Staels B. Peroxisome proliferator-activated receptor alpha controls cellular cholesterol trafficking in macrophages. *J Lipid Res* 46: 2717–2725, 2005.
63. Chinetti G, Gbaguidi FG, Griglio S, Mallat Z, Antonucci M, Poulain P, Chapman J, Fruchart JC, Tedgui A, Najib-Fruchart J, and Staels B. CLA-1/SR-BI is expressed in atherosclerotic lesion macrophages and regulated by activators of peroxisome proliferator-activated receptors. *Circulation* 101: 2411–2417, 2000.
64. Chinetti G, Griglio S, Antonucci M, Torra IP, Delerive P, Majd Z, Fruchart JC, Chapman J, Najib J, and Staels B. Activation of proliferator-activated receptors alpha and gamma induces apoptosis of human monocyte-derived macrophages. *J Biol Chem* 273: 25573–25580, 1998.
65. Chinetti G, Lestavel S, Bocher V, Remaley AT, Neve B, Torra IP, Teissier E, Minnich A, Jaye M, Duverger N, Brewer HB, Fruchart JC, Clavey V, and Staels B. PPAR-alpha and PPAR-gamma activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway. *Nat Med* 7: 53–58, 2001.
66. Chinetti G, Lestavel S, Fruchart JC, Clavey V, and Staels B. Peroxisome proliferator-activated receptor alpha reduces cholesterol esterification in macrophages. *Circ Res* 92: 212–217, 2003.
67. Cho DH, Choi YJ, Jo SA, and Jo I. Nitric oxide production and regulation of endothelial nitric-oxide synthase phosphorylation by prolonged treatment with troglitazone: evidence for involvement of peroxisome proliferator-activated receptor (PPAR) gamma-dependent and PPAR-gamma-independent signaling pathways. *J Biol Chem* 279: 2499–2506, 2004.
68. Cho MC, Lee K, Paik SG, and Yoon DY. Peroxisome proliferator-activated receptor (PPAR) modulators and metabolic disorders. *PPAR Res* 2008: 679137, 2008.
69. Cho MC, Yoon HE, Kang JW, Park SW, Yang Y, Hong JT, Song EY, Paik SG, Kim SH, and Yoon DY. A simple method to screen ligands of peroxisome proliferator-activated receptor delta. *Eur J Pharm Sci* 29: 355–360, 2006.
70. Clark RB, Bishop-Bailey D, Estrada-Hernandez T, Hla T, Puddington L, and Padula SJ. The nuclear receptor PPAR gamma and immunoregulation: PPAR gamma mediates inhibition of helper T cell responses. *J Immunol* 164: 1364–1371, 2000.
71. Colca JR, McDonald WG, Waldon DJ, Leone JW, Lull JM, Bannow CA, Lund ET, and Mathews WR. Identification of a novel mitochondrial protein ("mitoNEET") cross-linked specifically by a thiazolidinedione photoprobe. *Am J Physiol* 286: E252–E260, 2004.
72. Collins AR, Meehan WP, Kintscher U, Jackson S, Wakino S, Noh G, Palinski W, Hsueh WA, and Law RE. Troglitazone inhibits formation of early atherosclerotic lesions in diabetic and nondiabetic low density lipoprotein receptor-deficient mice. *Arterioscler Thromb Vasc Biol* 21: 365–371, 2001.
73. Collins T, Read MA, Neish AS, Whitley MZ, Thanos D, and Maniatis T. Transcriptional regulation of endothelial cell adhesion molecules: NF-kappa B and cytokine-inducible enhancers. *FASEB J* 9: 899–909, 1995.
74. Colville-Nash PR, Qureshi SS, Willis D, and Willoughby DA. Inhibition of inducible nitric oxide synthase by peroxisome proliferator-activated receptor agonists: correlation with induction of heme oxygenase 1. *J Immunol* 161: 978–984, 1998.
75. Cook WS, Yeldandi AV, Rao MS, Hashimoto T, and Reddy JK. Less extrahepatic induction of fatty acid beta-oxidation enzymes by PPAR alpha. *Biochem Biophys Res Commun* 278: 250–257, 2000.
76. Cui T, Schopfer FJ, Zhang J, Chen K, Ichikawa T, Baker PR, Batthyany C, Chacko BK, Feng X, Patel RP, Agarwal A, Freeman BA, and Chen YE. Nitrated fatty acids: endogenous anti-inflammatory signaling mediators. *J Biol Chem* 281: 35686–35698, 2006.
77. Cuthbert KD and Dyck JR. Malonyl-CoA decarboxylase is a major regulator of myocardial fatty acid oxidation. *Curr Hypertens Rep* 7: 407–411, 2005.
78. Davies SS, Pontsler AV, Marathe GK, Harrison KA, Murphy RC, Hinshaw JC, Prestwich GD, Hilaire AS, Prescott SM, Zimmerman GA, and McIntyre TM. Oxidized alkyl phospholipids are specific, high affinity peroxisome proliferator-activated receptor gamma ligands and agonists. *J Biol Chem* 276: 16015–16023, 2001.
79. de Dios ST, Bruemmer D, Dille RJ, Ivey ME, Jennings GL, Law RE, and Little PJ. Inhibitory activity of clinical thiazolidinedione peroxisome proliferator activating receptor-gamma ligands toward internal mammary artery, radial artery, and saphenous vein smooth muscle cell proliferation. *Circulation* 107: 2548–2550, 2003.
80. Delerive P, De Bosscher K, Besnard S, Vanden Berghe W, Peters JM, Gonzalez FJ, Fruchart JC, Tedgui A, Haegeman G, and Staels B. Peroxisome proliferator-activated receptor alpha negatively regulates the vascular inflammatory gene response by negative cross-talk with transcription factors NF-kappaB and AP-1. *J Biol Chem* 274: 32048–32054, 1999.
81. Delerive P, Furman C, Teissier E, Fruchart J, Duriez P, and Staels B. Oxidized phospholipids activate PPARalpha in a phospholipase A2-dependent manner. *FEBS Lett* 471: 34–38, 2000.
82. Delerive P, Gervois P, Fruchart JC, and Staels B. Induction of IkappaBalpha expression as a mechanism contributing to the anti-inflammatory activities of peroxisome proliferator-activated receptor-alpha activators. *J Biol Chem* 275: 36703–36707, 2000.

83. Delerive P, Martin-Nizard F, Chinetti G, Trottein F, Fruchart JC, Najib J, Duriez P, and Staels B. Peroxisome proliferator-activated receptor activators inhibit thrombin-induced endothelin-1 production in human vascular endothelial cells by inhibiting the activator protein-1 signaling pathway. *Circ Res* 85: 394–402, 1999.
84. Depre C, Havaux X, Renkin J, Vanoverschelde JL, and Wijns W. Expression of inducible nitric oxide synthase in human coronary atherosclerotic plaque. *Cardiovasc Res* 41: 465–472, 1999.
85. Desvergne B and Wahli W. Peroxisome proliferator-activated receptors: nuclear control of metabolism. *Endocr Rev* 20: 649–688, 1999.
86. Devchand PR, Keller H, Peters JM, Vazquez M, Gonzalez FJ, and Wahli W. The PPARalpha-leukotriene B4 pathway to inflammation control. *Nature* 384: 39–43, 1996.
87. Diep QN, El Mabrouk M, Cohn JS, Endemann D, Amiri F, Viridis A, Neves MF, and Schiffrin EL. Structure, endothelial function, cell growth, and inflammation in blood vessels of angiotensin II-infused rats: role of peroxisome proliferator-activated receptor-gamma. *Circulation* 105: 2296–2302, 2002.
88. Diep QN and Schiffrin EL. Increased expression of peroxisome proliferator-activated receptor-alpha and -gamma in blood vessels of spontaneously hypertensive rats. *Hypertension* 38: 249–254, 2001.
89. Diep QN, Touyz RM, and Schiffrin EL. Docosahexaenoic acid, a peroxisome proliferator-activated receptor-alpha ligand, induces apoptosis in vascular smooth muscle cells by stimulation of p38 mitogen-activated protein kinase. *Hypertension* 36: 851–855, 2000.
90. Ding G, Cheng L, Qin Q, Frontin S, and Yang Q. PPAR-delta modulates lipopolysaccharide-induced TNFalpha inflammation signaling in cultured cardiomyocytes. *J Mol Cell Cardiol* 40: 821–828, 2006.
91. Ding G, Fu M, Qin Q, Lewis W, Kim HW, Fukai T, Bacanamwo M, Chen YE, Schneider MD, Mangelsdorf DJ, Evans RM, and Yang Q. Cardiac peroxisome proliferator-activated receptor gamma is essential in protecting cardiomyocytes from oxidative damage. *Cardiovasc Res* 76: 269–279, 2007.
92. Ditiatkovski M, Toh BH, and Bobik A. GM-CSF deficiency reduces macrophage PPAR-gamma expression and aggravates atherosclerosis in ApoE-deficient mice. *Arterioscler Thromb Vasc Biol* 26: 2337–2344, 2006.
93. Djouadi F, Weinheimer CJ, Saffitz JE, Pitchford C, Bastin J, Gonzalez FJ, and Kelly DP. A gender-related defect in lipid metabolism and glucose homeostasis in peroxisome proliferator-activated receptor alpha-deficient mice. *J Clin Invest* 102: 1083–1091, 1998.
94. Dobashi K, Asayama K, Nakane T, Kodera K, Hayashibe H, and Nakazawa S. Troglitazone inhibits the expression of inducible nitric oxide synthase in adipocytes *in vitro* and *in vivo* study in 3T3-L1 cells and Otsuka Long-Evans Tokushima fatty rats. *Life Sci* 67: 2093–2101, 2000.
95. Dormandy JA, Charbonnel B, Eckland DJ, Erdmann E, Massi-Benedetti M, Moules IK, Skene AM, Tan MH, Lefebvre PJ, Murray GD, Standl E, Wilcox RG, Wilhelmsen L, Betteridge J, Birkeland K, Golay A, Heine RJ, Koranyi L, Laakso M, Mokan M, Norkus A, Pirags V, Podar T, Scheen A, Scherbaum W, Schernthaner G, Schmitz O, Skrha J, Smith U, and Taton J. Secondary prevention of macrovascular events in patients with type 2 diabetes in the PROactive study (PROspective pioglitazone clinical trial in macrovascular events): a randomised controlled trial. *Lancet* 366: 1279–1289, 2005.
96. Dreyer C, Krey G, Keller H, Givel F, Helftenbein G, and Wahli W. Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell* 68: 879–887, 1992.
97. Duan SZ, Ivashchenko CY, Russell MW, Milstone DS, and Mortensen RM. Cardiomyocyte-specific knockout and agonist of peroxisome proliferator-activated receptor-gamma both induce cardiac hypertrophy in mice. *Circ Res* 97: 372–379, 2005.
98. Dube RK, Zhang HY, Reddy SR, Boegehold MA, and Kotchen TA. Pioglitazone attenuates hypertension and inhibits growth of renal arteriolar smooth muscle in rats. *Am J Physiol* 265: R726–R732, 1993.
99. Duez H, Chao YS, Hernandez M, Torpier G, Poulain P, Mundt S, Mallat Z, Teissier E, Burton CA, Tedgui A, Fruchart JC, Fievet C, Wright SD, and Staels B. Reduction of atherosclerosis by the peroxisome proliferator-activated receptor alpha agonist fenofibrate in mice. *J Biol Chem* 277: 48051–48057, 2002.
100. Elbrecht A, Chen Y, Adams A, Berger J, Griffin P, Klatt T, Zhang B, Menke J, Zhou G, Smith RG, and Moller DE. L-764406 is a partial agonist of human peroxisome proliferator-activated receptor gamma: the role of Cys313 in ligand binding. *J Biol Chem* 274: 7913–7922, 1999.
101. Endemann G, Stanton LW, Madden KS, Bryant CM, White RT, and Protter AA. CD36 is a receptor for oxidized low density lipoprotein. *J Biol Chem* 268: 11811–11816, 1993.
102. Erol A. The functions of PPARs in aging and longevity. *PPAR Res* 2007: 39654, 2007.
103. Erol A. PPARalpha activators may play role for the regression of ventricular hypertrophy in hypertensive and hyperlipidemic patients. *Med Hypoth* 66: 1044–1045, 2006.
104. Evans RM. The steroid and thyroid hormone receptor superfamily. *Science (New York)* 240: 889–895, 1988.
105. Fan Y, Wang Y, Tang Z, Zhang H, Qin X, Zhu Y, Guan Y, Wang X, Staels B, Chien S, and Wang N. Suppression of pro-inflammatory adhesion molecules by PPAR-delta in human vascular endothelial cells. *Arterioscler, Thromb Vasc Biol* 28: 315–321, 2008.
106. Faveeuw C, Fougeray S, Angeli V, Fontaine J, Chinetti G, Gosset P, Delerive P, Maliszewski C, Capron M, Staels B, Moser M, and Trottein F. Peroxisome proliferator-activated receptor gamma activators inhibit interleukin-12 production in murine dendritic cells. *FEBS Lett* 486: 261–266, 2000.
107. Finck BN, Han X, Courtois M, Aimond F, Nerbonne JM, Kovacs A, Gross RW, and Kelly DP. A critical role for PPARalpha-mediated lipotoxicity in the pathogenesis of diabetic cardiomyopathy: modulation by dietary fat content. *Proc Natl Acad Sci U S A* 100: 1226–1231, 2003.
108. Finck BN, Lehman JJ, Leone TC, Welch MJ, Bennett MJ, Kovacs A, Han X, Gross RW, Kozak R, Lopaschuk GD, and Kelly DP. The cardiac phenotype induced by PPARalpha overexpression mimics that caused by diabetes mellitus. *J Clin Invest* 109: 121–130, 2002.
109. Forman BM, Chen J, and Evans RM. Hypolipidemic drugs, polyunsaturated fatty acids, and eicosanoids are ligands for peroxisome proliferator-activated receptors alpha and delta. *Proc Natl Acad Sci U S A* 94: 4312–4317, 1997.
110. Forman BM, Tontonoz P, Chen J, Brun RP, Spiegelman BM, and Evans RM. 15-Deoxy-delta 12, 14-prostaglandin J2 is a ligand for the adipocyte determination factor PPAR gamma. *Cell* 83: 803–812, 1995.

111. Fu M, Zhang J, Lin Y, Zhu X, Ehrenguber MU, and Chen YE. Early growth response factor-1 is a critical transcriptional mediator of peroxisome proliferator-activated receptor-gamma 1 gene expression in human aortic smooth muscle cells. *J Biol Chem* 277: 26808–26814, 2002.
112. Fu M, Zhang J, Lin Y, Zhu X, Zhao L, Ahmad M, Ehrenguber MU, and Chen YE. Early stimulation and late inhibition of peroxisome proliferator-activated receptor gamma (PPAR gamma) gene expression by transforming growth factor beta in human aortic smooth muscle cells: role of early growth-response factor-1 (Egr-1), activator protein 1 (AP1) and Smads. *Biochem J* 370: 1019–1025, 2003.
113. Fu M, Zhang J, Lin Yg Y, Zhu X, Willson TM, and Chen YE. Activation of peroxisome proliferator-activated receptor gamma inhibits osteoprotegerin gene expression in human aortic smooth muscle cells. *Biochem Biophys Res Commun* 294: 597–601, 2002.
114. Fu M, Zhang J, Zhu X, Myles DE, Willson TM, Liu X, and Chen YE. Peroxisome proliferator-activated receptor gamma inhibits transforming growth factor beta-induced connective tissue growth factor expression in human aortic smooth muscle cells by interfering with Smad3. *J Biol Chem* 276: 45888–45894, 2001.
115. Fu M, Zhu X, Wang Q, Zhang J, Song Q, Zheng H, Ogawa W, Du J, and Chen YE. Platelet-derived growth factor promotes the expression of peroxisome proliferator-activated receptor gamma in vascular smooth muscle cells by a phosphatidylinositol 3-kinase/Akt signaling pathway. *Circ Res* 89: 1058–1064, 2001.
116. Fujino T, Sato Y, Une M, Kanayasu-Toyoda T, Yamaguchi T, Shudo K, Inoue K, and Nishimaki-Mogami T. *In vitro* farnesoid X receptor ligand sensor assay using surface plasmon resonance and based on ligand-induced coactivator association. *J Steroid Biochem Mol Biol* 87: 247–252, 2003.
117. Fujishima S, Ohya Y, Nakamura Y, Onaka U, Abe I, and Fujishima M. Troglitazone, an insulin sensitizer, increases forearm blood flow in humans. *Am J Hypertens* 11: 1134–1137, 1998.
118. Fukunaga Y, Itoh H, Doi K, Tanaka T, Yamashita J, Chun TH, Inoue M, Masatsugu K, Sawada N, Saito T, Hosoda K, Kook H, Ueda M, and Nakao K. Thiazolidinediones, peroxisome proliferator-activated receptor gamma agonists, regulate endothelial cell growth and secretion of vasoactive peptides. *Atherosclerosis* 158: 113–119, 2001.
119. Gaudel C and Grimaldi PA. Metabolic functions of peroxisome proliferator-activated receptor beta/delta in skeletal muscle. *PPAR Res* 2007: 86394, 2007.
120. Gbaguidi FG, Chinetti G, Milosavljevic D, Teissier E, Chapman J, Olivecrona G, Fruchart JC, Griglio S, Fruchart-Najib J, and Staels B. Peroxisome proliferator-activated receptor (PPAR) agonists decrease lipoprotein lipase secretion and glycated LDL uptake by human macrophages. *FEBS Lett* 512: 85–90, 2002.
121. Gearing KL, Gottlicher M, Teboul M, Widmark E, and Gustafsson JA. Interaction of the peroxisome-proliferator-activated receptor and retinoid X receptor. *Proc Natl Acad Sci U S A* 90: 1440–1444, 1993.
122. Gensch C, Clever YP, Werner C, Hanhoun M, Bohm M, and Laufs U. The PPAR-gamma agonist pioglitazone increases neoangiogenesis and prevents apoptosis of endothelial progenitor cells. *Atherosclerosis* 192: 67–74, 2007.
123. Ghisletti S, Huang W, Ogawa S, Pascual G, Lin ME, Willson TM, Rosenfeld MG, and Glass CK. Parallel SUMOylation-dependent pathways mediate gene- and signal-specific transrepression by LXRs and PPARgamma. *Mol Cell* 25: 57–70, 2007.
124. Gilde AJ, van der Lee KA, Willemsen PH, Chinetti G, van der Leij FR, van der Vusse GJ, Staels B, and van Bilsen M. Peroxisome proliferator-activated receptor (PPAR) alpha and PPARbeta/delta, but not PPARgamma, modulate the expression of genes involved in cardiac lipid metabolism. *Circ Res* 92: 518–524, 2003.
125. Gitlin N, Julie NL, Spurr CL, Lim KN, and Juarbe HM. Two cases of severe clinical and histologic hepatotoxicity associated with troglitazone. *Ann Intern Med* 129: 36–38, 1998.
126. Gizard F, Amant C, Barbier O, Bellosta S, Robillard R, Percevault F, Sevestre H, Krimpenfort P, Corsini A, Rochette J, Glineur C, Fruchart JC, Torpier G, and Staels B. PPAR alpha inhibits vascular smooth muscle cell proliferation underlying intimal hyperplasia by inducing the tumor suppressor p16INK4a. *J Clin Invest* 115: 3228–3238, 2005.
127. Glass CK and Witztum JL. Atherosclerosis: the road ahead. *Cell* 104: 503–516, 2001.
128. Goetze S, Bungenstock A, Czupalla C, Eilers F, Stawowy P, Kintscher U, Spencer-Hansch C, Graf K, Nurnberg B, Law RE, Fleck E, and Grafe M. Leptin induces endothelial cell migration through Akt, which is inhibited by PPARgamma-ligands. *Hypertension* 40: 748–754, 2002.
129. Goetze S, Eilers F, Bungenstock A, Kintscher U, Stawowy P, Blaschke F, Graf K, Law RE, Fleck E, and Grafe M. PPAR activators inhibit endothelial cell migration by targeting Akt. *Biochem Biophys Res Commun* 293: 1431–1437, 2002.
130. Goetze S, Kim S, Xi XP, Graf K, Yang DC, Fleck E, Meehan WP, Hsueh WA, and Law RE. Troglitazone inhibits mitogenic signaling by insulin in vascular smooth muscle cells. *J Cardiovasc Pharmacol* 35: 749–757, 2000.
131. Goetze S, Kintscher U, Kim S, Meehan WP, Kaneshiro K, Collins AR, Fleck E, Hsueh WA, and Law RE. Peroxisome proliferator-activated receptor-gamma ligands inhibit nuclear but not cytosolic extracellular signal-regulated kinase/mitogen-activated protein kinase-regulated steps in vascular smooth muscle cell migration. *J Cardiovasc Pharmacol* 38: 909–921, 2001.
132. Goetze S, Xi XP, Graf K, Fleck E, Hsueh WA, and Law RE. Troglitazone inhibits angiotensin II-induced extracellular signal-regulated kinase 1/2 nuclear translocation and activation in vascular smooth muscle cells. *FEBS Lett* 452: 277–282, 1999.
133. Goetze S, Xi XP, Kawano H, Gotlibowski T, Fleck E, Hsueh WA, and Law RE. PPAR gamma-ligands inhibit migration mediated by multiple chemoattractants in vascular smooth muscle cells. *J Cardiovasc Pharmacol* 33: 798–806, 1999.
134. Goikoetxea MJ, Beaumont J, and Diez J. Peroxisome proliferator-activated receptor alpha and hypertensive heart disease. *Drugs* 64(suppl 2): 9–18, 2004.
135. Goldberg RB, Kendall DM, Deeg MA, Buse JB, Zagar AJ, Pinaire JA, Tan MH, Khan MA, Perez AT, and Jacober SJ. A comparison of lipid and glycemic effects of pioglitazone and rosiglitazone in patients with type 2 diabetes and dyslipidemia. *Diabetes Care* 28: 1547–1554, 2005.
136. Golfman LS, Wilson CR, Sharma S, Burgmaier M, Young ME, Guthrie PH, Van Arsdall M, Adrogue JV, Brown KK, and Taegtmeier H. Activation of PPARgamma enhances myocardial glucose oxidation and improves contractile function in isolated working hearts of ZDF rats. *Am J Physiol* 289: E328–E336, 2005.

137. Goto M, Terada S, Kato M, Katoh M, Yokozeki T, Tabata I, and Shimokawa T. cDNA Cloning and mRNA analysis of PGC-1 in epitrochlearis muscle in swimming-exercised rats. *Biochem Biophys Res Commun* 274: 350–354, 2000.
138. Gottlicher M, Widmark E, Li Q, and Gustafsson JA. Fatty acids activate a chimera of the clofibrilic acid-activated receptor and the glucocorticoid receptor. *Proc Natl Acad Sci U S A* 89: 4653–4657, 1992.
139. Goya K, Sumitani S, Xu X, Kitamura T, Yamamoto H, Kurebayashi S, Saito H, Kouhara H, Kasayama S, and Kawase I. Peroxisome proliferator-activated receptor alpha agonists increase nitric oxide synthase expression in vascular endothelial cells. *Arterioscler Thromb Vasc Biol* 24: 658–663, 2004.
140. Graf K, Xi XP, Hsueh WA, and Law RE. Troglitazone inhibits angiotensin II-induced DNA synthesis and migration in vascular smooth muscle cells. *FEBS Lett* 400: 119–121, 1997.
141. Graham TL, Mookherjee C, Suckling KE, Palmer CN, and Patel L. The PPARdelta agonist GW0742X reduces atherosclerosis in LDLR(-/-) mice. *Atherosclerosis* 181: 29–37, 2005.
142. Greenland P, Knoll MD, Stamler J, Neaton JD, Dyer AR, Garside DB, and Wilson PW. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA* 290: 891–897, 2003.
143. Han J, Hajjar DP, Tauras JM, Feng J, Gotto AM Jr, and Nicholson AC. Transforming growth factor-beta1 (TGF-beta1) and TGF-beta2 decrease expression of CD36, the type B scavenger receptor, through mitogen-activated protein kinase phosphorylation of peroxisome proliferator-activated receptor-gamma. *J Biol Chem* 275: 1241–1246, 2000.
144. Han JK, Lee HS, Yang HM, Hur J, Jun SI, Kim JY, Cho CH, Koh GY, Peters JM, Park KW, Cho HJ, Lee HY, Kang HJ, Oh BH, Park YB, and Kim HS. Peroxisome proliferator-activated receptor-delta agonist enhances vasculogenesis by regulating endothelial progenitor cells through genomic and nongenomic activations of the phosphatidylinositol 3-kinase/Akt pathway. *Circulation* 118: 1021–1033, 2008.
145. Han KH, Chang MK, Boullier A, Green SR, Li A, Glass CK, and Quehenberger O. Oxidized LDL reduces monocyte CCR2 expression through pathways involving peroxisome proliferator-activated receptor gamma. *J Clin Invest* 106: 793–802, 2000.
146. Hannan KM, Dilley RJ, de Dios ST, and Little PJ. Troglitazone stimulates repair of the endothelium and inhibits neointimal formation in denuded rat aorta. *Arterioscler Thromb Vasc Biol* 23: 762–768, 2003.
147. Haraguchi G, Kobayashi Y, Brown ML, Tanaka A, Isobe M, Gianturco SH, and Bradley WA. PPAR(alpha) and PPAR(gamma) activators suppress the monocyte-macrophage apoB-48 receptor. *J Lipid Res* 44: 1224–1231, 2003.
148. Harkin DP, Bean JM, Miklos D, Song YH, Truong VB, Englert C, Christians FC, Ellisen LW, Maheswaran S, Oliner JD, and Haber DA. Induction of GADD45 and JNK/SAPK-dependent apoptosis following inducible expression of BRCA1. *Cell* 97: 575–586, 1999.
149. Harris IS, Treskov I, Rowley MW, Heximer S, Kaltenbronn K, Finck BN, Gross RW, Kelly DP, Blumer KJ, and Muslin AJ. G-protein signaling participates in the development of diabetic cardiomyopathy. *Diabetes* 53: 3082–3090, 2004.
150. Hautala AJ, Leon AS, Skinner JS, Rao DC, Bouchard C, and Rankinen T. Peroxisome proliferator-activated receptor-delta polymorphisms are associated with physical performance and plasma lipids: the HERITAGE Family Study. *Am J Physiol Heart Circ Physiol* 292: H2498–H2505, 2007.
151. Helin K. Regulation of cell proliferation by the E2F transcription factors. *Curr Opin Genet Dev* 8: 28–35, 1998.
152. Hennuyer N, Tailleux A, Torpier G, Mezdour H, Fruchart JC, Staels B, and Fievet C. PPARalpha, but not PPAR-gamma, activators decrease macrophage-laden atherosclerotic lesions in a nondiabetic mouse model of mixed dyslipidemia. *Arterioscler Thromb Vasc Biol* 25: 1897–1902, 2005.
153. Heo KS, Kim DU, Ryoo S, Nam M, Baek ST, Kim L, Park SK, Myung CS, and Hoe KL. PPARgamma activation abolishes LDL-induced proliferation of human aortic smooth muscle cells via SOD-mediated down-regulation of superoxide. *Biochem Biophys Res Commun* 359: 1017–1023, 2007.
154. Hevener AL, Olefsky JM, Reichart D, Nguyen MT, Bandyopadhyay G, Leung HY, Watt MJ, Benner C, Febbraio MA, Nguyen AK, Folian B, Subramaniam S, Gonzalez FJ, Glass CK, and Ricote M. Macrophage PPAR gamma is required for normal skeletal muscle and hepatic insulin sensitivity and full antidiabetic effects of thiazolidinediones. *J Clin Invest* 117: 1658–1669, 2007.
155. Home PD, Pocock SJ, Beck-Nielsen H, Gomis R, Hanefeld M, Jones NP, Komajda M, and McMurray JJ. Rosiglitazone evaluated for cardiovascular outcomes: an interim analysis. *N Engl J Med* 357: 28–38, 2007.
156. Hourton D, Delerive P, Stankova J, Staels B, Chapman MJ, and Ninio E. Oxidized low-density lipoprotein and peroxisome-proliferator-activated receptor alpha down-regulate platelet-activating-factor receptor expression in human macrophages. *Biochem J* 354: 225–232, 2001.
157. Hu E, Kim JB, Sarraf P, and Spiegelman BM. Inhibition of adipogenesis through MAP kinase-mediated phosphorylation of PPARgamma. *Science (New York)* 274: 2100–2103, 1996.
158. Huang J and Kontos CD. PTEN modulates vascular endothelial growth factor-mediated signaling and angiogenic effects. *J Biol Chem* 277: 10760–10766, 2002.
159. Huang JT, Welch JS, Ricote M, Binder CJ, Willson TM, Kelly C, Witztum JL, Funk CD, Conrad D, and Glass CK. Interleukin-4-dependent production of PPAR-gamma ligands in macrophages by 12/15-lipoxygenase. *Nature* 400: 378–382, 1999.
160. Hur J, Yoon CH, Kim HS, Choi JH, Kang HJ, Hwang KK, Oh BH, Lee MM, and Park YB. Characterization of two types of endothelial progenitor cells and their different contributions to neovasclogenesis. *Arterioscler Thromb Vasc Biol* 24: 288–293, 2004.
161. Huss JM, Levy FH, and Kelly DP. Hypoxia inhibits the peroxisome proliferator-activated receptor alpha/retinoid X receptor gene regulatory pathway in cardiac myocytes: a mechanism for O₂-dependent modulation of mitochondrial fatty acid oxidation. *J Biol Chem* 276: 27605–27612, 2001.
162. Hwang J, Kleinhenz DJ, Lassegue B, Griendling KK, Dikalov S, and Hart CM. Peroxisome proliferator-activated receptor-gamma ligands regulate endothelial membrane superoxide production. *Am J Physiol Cell Physiol* 288: C899–C905, 2005.
163. Iglarz M, Touyz RM, Amiri F, Lavoie MF, Diep QN, and Schiffrin EL. Effect of peroxisome proliferator-activated receptor-alpha and -gamma activators on vascular re-

- modeling in endothelin-dependent hypertension. *Arterioscler Thromb Vasc Biol* 23: 45–51, 2003.
164. Iijima K, Yoshizumi M, Ako J, Eto M, Kim S, Hashimoto M, Sugimoto N, Liang YQ, Sudoh N, Toba K, and Ouchi Y. Expression of peroxisome proliferator-activated receptor gamma (PPARgamma) in rat aortic smooth muscle cells. *Biochem Biophys Res Commun* 247: 353–356, 1998.
 165. Ikeda U, Shimpo M, Murakami Y, and Shimada K. Peroxisome proliferator-activated receptor-gamma ligands inhibit nitric oxide synthesis in vascular smooth muscle cells. *Hypertension* 35: 1232–1236, 2000.
 166. Imai H, Numaguchi Y, Ishii M, Kubota R, Yokouchi K, Ogawa Y, Kondo T, Okumura K, and Murohara T. Prostacyclin synthase gene transfer inhibits neointimal formation by suppressing PPAR delta expression. *Atherosclerosis* 195: 322–332, 2007.
 167. Imayama I, Ichiki T, Inanaga K, Ohtsubo H, Fukuyama K, Ono H, Hashiguchi Y, and Sunagawa K. Telmisartan downregulates angiotensin II type 1 receptor through activation of peroxisome proliferator-activated receptor gamma. *Cardiovasc Res* 72: 184–190, 2006.
 168. Inoue I, Goto S, Matsunaga T, Nakajima T, Awata T, Hokari S, Komoda T, and Katayama S. The ligands/activators for peroxisome proliferator-activated receptor alpha (PPARalpha) and PPARgamma increase Cu²⁺/Zn²⁺-superoxide dismutase and decrease p22phox message expressions in primary endothelial cells. *Metabolism* 50: 3–11, 2001.
 169. Inoue I, Hayashi K, Yagasaki F, Nakamura K, Matsunaga T, Xu H, Inukai K, Awata T, Komoda T, and Katayama S. Apoptosis of endothelial cells may be mediated by genes of peroxisome proliferator-activated receptor gamma 1 (PPARgamma 1) and PPARalpha genes. *J Atheroscler Thromb* 10: 99–108, 2003.
 170. Inoue I, Shino K, Noji S, Awata T, and Katayama S. Expression of peroxisome proliferator-activated receptor alpha (PPAR alpha) in primary cultures of human vascular endothelial cells. *Biochem Biophys Res Commun* 246: 370–374, 1998.
 171. Irukayama-Tomobe Y, Miyauchi T, Sakai S, Kasuya Y, Ogata T, Takanashi M, Iemitsu M, Sudo T, Goto K, and Yamaguchi I. Endothelin-1-induced cardiac hypertrophy is inhibited by activation of peroxisome proliferator-activated receptor-alpha partly via blockade of c-Jun NH2-terminal kinase pathway. *Circulation* 109: 904–910, 2004.
 172. Issemann I and Green S. Activation of a member of the steroid hormone receptor superfamily by peroxisome proliferators. *Nature* 347: 645–650, 1990.
 173. Jackson SM, Parhami F, Xi XP, Berliner JA, Hsueh WA, Law RE, and Demer LL. Peroxisome proliferator-activated receptor activators target human endothelial cells to inhibit leukocyte-endothelial cell interaction. *Arterioscler Thromb Vasc Biol* 19: 2094–2104, 1999.
 174. Jiang C, Ting AT, and Seed B. PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 391: 82–86, 1998.
 175. Jiang WG, Redfern A, Bryce RP, and Mansel RE. Peroxisome proliferator activated receptor-gamma (PPAR-gamma) mediates the action of gamma linolenic acid in breast cancer cells. *Prostaglandins Leukot Essent Fatty Acids* 62: 119–127, 2000.
 176. Jucker BM, Doe CP, Schnackenberg CG, Olzinski AR, Maniscalco K, Williams C, Hu TC, Lenhard SC, Costell M, Bernard R, Sarov-Blat L, Stepleski K, and Willette RN. PPARdelta activation normalizes cardiac substrate metabolism and reduces right ventricular hypertrophy in congestive heart failure. *J Cardiovasc Pharmacol* 50: 25–34, 2007.
 177. Kang K, Reilly SM, Karabacak V, Gangl MR, Fitzgerald K, Hatano B, and Lee CH. Adipocyte-derived Th2 cytokines and myeloid PPARdelta regulate macrophage polarization and insulin sensitivity. *Cell Metab* 7: 485–495, 2008.
 178. Kannel WB and McGee DL. Diabetes and cardiovascular disease: the Framingham study. *JAMA* 241: 2035–2038, 1979.
 179. Kato K, Satoh H, Endo Y, Yamada D, Midorikawa S, Sato W, Mizuno K, Fujita T, Tsukamoto K, and Watanabe T. Thiazolidinediones down-regulate plasminogen activator inhibitor type 1 expression in human vascular endothelial cells: a possible role for PPARgamma in endothelial function. *Biochem Biophys Res Commun* 258: 431–435, 1999.
 180. Keller BJ, Bradford BU, Marsman DS, Cattley RC, Popp JA, Bojes HK, and Thurman RG. The nongenotoxic hepatocarcinogen Wy-14,643 is an uncoupler of oxidative phosphorylation *in vivo*. *Toxicol Appl Pharmacol* 119: 52–58, 1993.
 181. Keller BJ, Marsman DS, Popp JA, and Thurman RG. Several nongenotoxic carcinogens uncouple mitochondrial oxidative phosphorylation. *Biochim Biophys Acta* 1102: 237–244, 1992.
 182. Keller H, Dreyer C, Medin J, Mahfoudi A, Ozato K, and Wahli W. Fatty acids and retinoids control lipid metabolism through activation of peroxisome proliferator-activated receptor-retinoid X receptor heterodimers. *Proc Natl Acad Sci U S A* 90: 2160–2164, 1993.
 183. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, and Wahli W. Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting. *J Clin Invest* 103: 1489–1498, 1999.
 184. Kim HJ, Ham SA, Kim SU, Hwang JY, Kim JH, Chang KC, Yabe-Nishimura C, Kim JH, and Seo HG. Transforming growth factor-beta1 is a molecular target for the peroxisome proliferator-activated receptor delta. *Circ Res* 102: 193–200, 2008.
 185. Kim KY and Cheon HG. Antiangiogenic effect of rosiglitazone is mediated via peroxisome proliferator-activated receptor gamma-activated maxi-K channel opening in human umbilical vein endothelial cells. *J Biol Chem* 281: 13503–13512, 2006.
 186. Kintscher U, Goetze S, Wakino S, Kim S, Nagpal S, Chandraratna RA, Graf K, Fleck E, Hsueh WA, and Law RE. Peroxisome proliferator-activated receptor and retinoid X receptor ligands inhibit monocyte chemotactic protein-1-directed migration of monocytes. *Eur J Pharmacol* 401: 259–270, 2000.
 187. Kintscher U, Lyon C, Wakino S, Bruemmer D, Feng X, Goetze S, Graf K, Moustakas A, Staels B, Fleck E, Hsueh WA, and Law RE. PPARalpha inhibits TGF-beta-induced beta5 integrin transcription in vascular smooth muscle cells by interacting with Smad4. *Circ Res* 91: e35–e44, 2002.
 188. Kliewer SA, Forman BM, Blumberg B, Ong ES, Borgmeyer U, Mangelsdorf DJ, Umesono K, and Evans RM. Differential expression and activation of a family of murine peroxisome proliferator-activated receptors. *Proc Natl Acad Sci U S A* 91: 7355–7359, 1994.
 189. Kliewer SA, Lenhard JM, Willson TM, Patel I, Morris DC, and Lehmann JM. A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 83: 813–819, 1995.

190. Kliewer SA, Umesonon K, Noonan DJ, Heyman RA, and Evans RM. Convergence of 9-cis retinoic acid and peroxisome proliferator signalling pathways through heterodimer formation of their receptors. *Nature* 358: 771-774, 1992.
191. Koumanov F, Jin B, Yang J, and Holman GD. Insulin signaling meets vesicle traffic of GLUT4 at a plasma-membrane-activated fusion step. *Cell Metab* 2: 179-189, 2005.
192. Krey G, Braissant O, L'Horsset F, Kalkhoven E, Perroud M, Parker MG, and Wahli W. Fatty acids, eicosanoids, and hypolipidemic agents identified as ligands of peroxisome proliferator-activated receptors by coactivator-dependent receptor ligand assay. *Mol Endocrinol (Baltimore)* 11: 779-791, 1997.
193. Kronke G, Kadl A, Ikonomu E, Bluml S, Furnkranz A, Sarembock JJ, Bochkov VN, Exner M, Binder BR, and Leitinger N. Expression of heme oxygenase-1 in human vascular cells is regulated by peroxisome proliferator-activated receptors. *Arterioscler Thromb Vasc Biol* 27: 1276-1282, 2007.
194. Kwak BR, Myit S, Mulhaupt F, Veillard N, Rufer N, Roosnek E, and Mach F. PPARgamma but not PPARalpha ligands are potent repressors of major histocompatibility complex class II induction in atheroma-associated cells. *Circ Res* 90: 356-362, 2002.
195. LaBaer J, Garrett MD, Stevenson LF, Slingerland JM, Sandhu C, Chou HS, Fattaey A, and Harlow E. New functional activities for the p21 family of CDK inhibitors. *Genes Dev* 11: 847-862, 1997.
196. Lalwani ND, Alvares K, Reddy MK, Reddy MN, Parikh I, and Reddy JK. Peroxisome proliferator-binding protein: identification and partial characterization of nafenopin-, clofibrac acid-, and ciprofibrate-binding proteins from rat liver. *Proc Natl Acad Sci U S A* 84: 5242-5246, 1987.
197. Lalwani ND, Fahl WE, and Reddy JK. Detection of a nafenopin-binding protein in rat liver cytosol associated with the induction of peroxisome proliferation by hypolipidemic compounds. *Biochem Biophys Res Commun* 116: 388-393, 1983.
198. Law RE, Goetze S, Xi XP, Jackson S, Kawano Y, Demer L, Fishbein MC, Meehan WP, and Hsueh WA. Expression and function of PPARgamma in rat and human vascular smooth muscle cells. *Circulation* 101: 1311-1318, 2000.
199. Law RE, Meehan WP, Xi XP, Graf K, Wuthrich DA, Coats W, Faxon D, and Hsueh WA. Troglitazone inhibits vascular smooth muscle cell growth and intimal hyperplasia. *J Clin Invest* 98: 1897-1905, 1996.
200. Lazennec G, Canaple L, Saugy D, and Wahli W. Activation of peroxisome proliferator-activated receptors (PPARs) by their ligands and protein kinase A activators. *Mol Endocr (Baltimore)* 14: 1962-1975, 2000.
201. Lee CH, Chawla A, Urbiztondo N, Liao D, Boisvert WA, Evans RM, and Curtiss LK. Transcriptional repression of atherogenic inflammation: modulation by PPARdelta. *Science (New York)* 302: 453-457, 2003.
202. Lee CH, Olson P, Hevener A, Mehl I, Chong LW, Olefsky JM, Gonzalez FJ, Ham J, Kang H, Peters JM, and Evans RM. PPARdelta regulates glucose metabolism and insulin sensitivity. *Proc Natl Acad Sci U S A* 103: 3444-3449, 2006.
203. Lee H, Shi W, Tontonoz P, Wang S, Subbanagounder G, Hedrick CC, Hama S, Borromeo C, Evans RM, Berliner JA, and Nagy L. Role for peroxisome proliferator-activated receptor alpha in oxidized phospholipid-induced synthesis of monocyte chemoattractant protein-1 and interleukin-8 by endothelial cells. *Circ Res* 87: 516-521, 2000.
204. Lee KS, Park JH, Lee S, Lim HJ, Jang Y, and Park HY. Troglitazone inhibits endothelial cell proliferation through suppression of casein kinase 2 activity. *Biochem Biophys Res Commun* 346: 83-88, 2006.
205. Lee SS, Pineau T, Drago J, Lee EJ, Owens JW, Kroetz DL, Fernandez-Salguero PM, Westphal H, and Gonzalez FJ. Targeted disruption of the alpha isoform of the peroxisome proliferator-activated receptor gene in mice results in abolishment of the pleiotropic effects of peroxisome proliferators. *Mol Cell Biol* 15: 3012-3022, 1995.
206. Lee WM. Drug-induced hepatotoxicity. *N Engl J Med* 349: 474-485, 2003.
207. Leesnitzer LM, Parks DJ, Bledsoe RK, Cobb JE, Collins JL, Consler TG, Davis RG, Hull-Ryde EA, Lenhard JM, Patel L, Plunket KD, Shenk JL, Stimmel JB, Therapontos C, Willson TM, and Blanchard SG. Functional consequences of cysteine modification in the ligand binding sites of peroxisome proliferator activated receptors by GW9662. *Biochemistry* 41: 6640-6650, 2002.
208. Lehman JJ and Kelly DP. Gene regulatory mechanisms governing energy metabolism during cardiac hypertrophic growth. *Heart Fail Rev* 7: 175-185, 2002.
209. Lehmann JM, Moore LB, Smith-Oliver TA, Wilkison WO, Willson TM, and Kliewer SA. An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J Biol Chem* 270: 12953-12956, 1995.
210. Leibowitz MD, Fievet C, Hennuyer N, Peinado-Onsurbe J, Duez H, Bergera J, Cullinan CA, Sparrow CP, Baffic J, Berger GD, Santini C, Marquis RW, Tolman RL, Smith RG, Moller DE, and Auwerx J. Activation of PPARdelta alters lipid metabolism in db/db mice. *FEBS Lett* 473: 333-336, 2000.
211. Leone TC, Weinheimer CJ, and Kelly DP. A critical role for the peroxisome proliferator-activated receptor alpha (PPARalpha) in the cellular fasting response: the PPAR-alpha-null mouse as a model of fatty acid oxidation disorders. *Proc Natl Acad Sci U S A* 96: 7473-7478, 1999.
212. Levi Z, Shaish A, Yacov N, Levkovitz H, Trestman S, Gerber Y, Cohen H, Dvir A, Rhachmani R, Ravid M, and Harats D. Rosiglitazone (PPARgamma-agonist) attenuates atherogenesis with no effect on hyperglycaemia in a combined diabetes-atherosclerosis mouse model. *Diabetes Obes Metab* 5: 45-50, 2003.
213. Levonen AL, Dickinson DA, Moellering DR, Mulcahy RT, Forman HJ, and Darley-Usmar VM. Biphasic effects of 15-deoxy-delta(12,14)-prostaglandin J(2) on glutathione induction and apoptosis in human endothelial cells. *Arterioscler Thromb Vasc Biol* 21: 1846-1851, 2001.
214. Li AC, Binder CJ, Gutierrez A, Brown KK, Plotkin CR, Pattison JW, Valledor AF, Davis RA, Willson TM, Witztum JL, Palinski W, and Glass CK. Differential inhibition of macrophage foam-cell formation and atherosclerosis in mice by PPARalpha, beta/delta, and gamma. *J Clin Invest* 114: 1564-1576, 2004.
215. Li AC, Brown KK, Silvestre MJ, Willson TM, Palinski W, and Glass CK. Peroxisome proliferator-activated receptor gamma ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. *J Clin Invest* 106: 523-531, 2000.
216. Li L, Beauchamp MC, and Renier G. Peroxisome proliferator-activated receptor alpha and gamma agonists upregulate human macrophage lipoprotein lipase expression. *Atherosclerosis* 165: 101-110, 2002.

217. Li M, Pascual G, and Glass CK. Peroxisome proliferator-activated receptor gamma-dependent repression of the inducible nitric oxide synthase gene. *Mol Cell Biol* 20: 4699–4707, 2000.
218. Li Y, Zhang J, Schopfer FJ, Martynowski D, Garcia-Barrio MT, Kovach A, Suino-Powell K, Baker PR, Freeman BA, Chen YE, and Xu HE. Molecular recognition of nitrated fatty acids by PPAR gamma. *Nat Struct Mol Biol* 15: 865–867, 2008.
219. Liang CP, Han S, Okamoto H, Carnemolla R, Tabas I, Accili D, and Tall AR. Increased CD36 protein as a response to defective insulin signaling in macrophages. *J Clin Invest* 113: 764–773, 2004.
220. Liang F, Wang F, Zhang S, and Gardner DG. Peroxisome proliferator activated receptor (PPAR)alpha agonists inhibit hypertrophy of neonatal rat cardiac myocytes. *Endocrinology* 144: 4187–4194, 2003.
221. Lim H, Gupta RA, Ma WG, Paria BC, Moller DE, Morrow JD, DuBois RN, Trzaskos JM, and Dey SK. Cyclooxygenase-2-derived prostacyclin mediates embryo implantation in the mouse via PPARdelta. *Genes Dev* 13: 1561–1574, 1999.
222. Lim HJ, Lee S, Park JH, Lee KS, Choi HE, Chung KS, Lee HH, and Park HY. PPARdelta agonist L-165041 inhibits rat vascular smooth muscle cell proliferation and migration via inhibition of cell cycle. *Atherosclerosis* 2009 (in press).
223. Lim S, Jin CJ, Kim M, Chung SS, Park HS, Lee IK, Lee CT, Cho YM, Lee HK, and Park KS. PPARgamma gene transfer sustains apoptosis, inhibits vascular smooth muscle cell proliferation, and reduces neointima formation after balloon injury in rats. *Arterioscler Thromb Vasc Biol* 26: 808–813, 2006.
224. Lin Y, Zhu X, McLntee FL, Xiao H, Zhang J, Fu M, and Chen YE. Interferon regulatory factor-1 mediates PPARgamma-induced apoptosis in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* 24: 257–263, 2004.
225. Lincoff AM, Wolski K, Nicholls SJ, and Nissen SE. Pioglitazone and risk of cardiovascular events in patients with type 2 diabetes mellitus: a meta-analysis of randomized trials. *JAMA* 298: 1180–1188, 2007.
226. Liou JY, Lee S, Ghelani D, Matijevic-Aleksic N, and Wu KK. Protection of endothelial survival by peroxisome proliferator-activated receptor-delta mediated 14-3-3 upregulation. *Arterioscler Thromb Vasc Biol* 26: 1481–1487, 2006.
227. Llaverias G, Vazquez-Carrera M, Sanchez RM, Noe V, Ciudad CJ, Laguna JC, and Alegret M. Rosiglitazone upregulates caveolin-1 expression in THP-1 cells through a PPAR-dependent mechanism. *J Lipid Res* 45: 2015–2024, 2004.
228. Loskutoff DJ and Samad F. The adipocyte and hemostatic balance in obesity: studies of PAI-1. *Arterioscler Thromb Vasc Biol* 18: 1–6, 1998.
229. Luptak I, Balschi JA, Xing Y, Leone TC, Kelly DP, and Tian R. Decreased contractile and metabolic reserve in peroxisome proliferator-activated receptor-alpha-null hearts can be rescued by increasing glucose transport and utilization. *Circulation* 112: 2339–2346, 2005.
230. Luquet S, Lopez-Soriano J, Holst D, Fredenrich A, Melki J, Rassoulzadegan M, and Grimaldi PA. Peroxisome proliferator-activated receptor delta controls muscle development and oxidative capability. *FASEB J* 17: 2299–2301, 2003.
231. Maiorano D, Lutzmann M, and Mechali M. MCM proteins and DNA replication. *Curr Opin Cell Biol* 18: 130–136, 2006.
232. Majai G, Sarang Z, Csomos K, Zahuczky G, and Fesus L. PPARgamma-dependent regulation of human macrophages in phagocytosis of apoptotic cells. *Eur J Immunol* 37: 1343–1354, 2007.
233. Marcus SL, Miyata KS, Zhang B, Subramani S, Rachubinski RA, and Capone JP. Diverse peroxisome proliferator-activated receptors bind to the peroxisome proliferator-responsive elements of the rat hydratase/dehydrogenase and fatty acyl-CoA oxidase genes but differentially induce expression. *Proc Natl Acad Sci U S A* 90: 5723–5727, 1993.
234. Martin-Nizard F, Furman C, Delerive P, Kandoussi A, Fruchart JC, Staels B, and Duriez P. Peroxisome proliferator-activated receptor activators inhibit oxidized low-density lipoprotein-induced endothelin-1 secretion in endothelial cells. *J Cardiovasc Pharmacol* 40: 822–831, 2002.
235. Marx N, Bourcier T, Sukhova GK, Libby P, and Plutzky J. PPARgamma activation in human endothelial cells increases plasminogen activator inhibitor type-1 expression: PPARgamma as a potential mediator in vascular disease. *Arterioscler Thromb Vasc Biol* 19: 546–551, 1999.
236. Marx N, Kehrle B, Kohlhammer K, Grub M, Koenig W, Hombach V, Libby P, and Plutzky J. PPAR activators as antiinflammatory mediators in human T lymphocytes: implications for atherosclerosis and transplantation-associated arteriosclerosis. *Circ Res* 90: 703–710, 2002.
237. Marx N, Mach F, Sauty A, Leung JH, Sarafi MN, Ransohoff RM, Libby P, Plutzky J, and Luster AD. Peroxisome proliferator-activated receptor-gamma activators inhibit IFN-gamma-induced expression of the T cell-active CXC chemokines IP-10, Mig, and I-TAC in human endothelial cells. *J Immunol* 164: 6503–6508, 2000.
238. Marx N, Mackman N, Schonbeck U, Yilmaz N, Hombach V, Libby P, and Plutzky J. PPARalpha activators inhibit tissue factor expression and activity in human monocytes. *Circulation* 103: 213–219, 2001.
239. Marx N, Schonbeck U, Lazar MA, Libby P, and Plutzky J. Peroxisome proliferator-activated receptor gamma activators inhibit gene expression and migration in human vascular smooth muscle cells. *Circ Res* 83: 1097–1103, 1998.
240. Marx N, Sukhova G, Murphy C, Libby P, and Plutzky J. Macrophages in human atheroma contain PPARgamma: differentiation-dependent peroxisomal proliferator-activated receptor gamma(PPARgamma) expression and reduction of MMP-9 activity through PPARgamma activation in mononuclear phagocytes *in vitro*. *Am J Pathol* 153: 17–23, 1998.
241. Marx N, Sukhova GK, Collins T, Libby P, and Plutzky J. PPARalpha activators inhibit cytokine-induced vascular cell adhesion molecule-1 expression in human endothelial cells. *Circulation* 99: 3125–3131, 1999.
242. Matsumoto K, Hirano K, Nozaki S, Takamoto A, Nishida M, Nakagawa-Toyama Y, Janabi MY, Ohya T, Yamashita S, and Matsuzawa Y. Expression of macrophage (Mphi) scavenger receptor, CD36, in cultured human aortic smooth muscle cells in association with expression of peroxisome proliferator activated receptor-gamma, which regulates gain of Mphi-like phenotype *in vitro*, and its implication in atherogenesis. *Arterioscler Thromb Vasc Biol* 20: 1027–1032, 2000.
243. Matsusue K, Miyoshi A, Yamano S, and Gonzalez FJ. Ligand-activated PPARbeta efficiently represses the induction of LXR-dependent promoter activity through competition with RXR. *Mol Cell Endocrinol* 256: 23–33, 2006.

244. Mehta JL, Hu B, Chen J, and Li D. Pioglitazone inhibits LOX-1 expression in human coronary artery endothelial cells by reducing intracellular superoxide radical generation. *Arterioscler Thromb Vasc Biol* 23: 2203–2208, 2003.
245. Meier CA, Chicheportiche R, Juge-Aubry CE, Dreyer MG, and Dayer JM. Regulation of the interleukin-1 receptor antagonist in THP-1 cells by ligands of the peroxisome proliferator-activated receptor gamma. *Cytokine* 18: 320–328, 2002.
246. Meissner M, Stein M, Urbich C, Reisinger K, Suske G, Staels B, Kaufmann R, and Gille J. PPARalpha activators inhibit vascular endothelial growth factor receptor-2 expression by repressing Sp1-dependent DNA binding and transactivation. *Circ Res* 94: 324–332, 2004.
247. Mishra A, Chaudhary A, and Sethi S. Oxidized omega-3 fatty acids inhibit NF-kappaB activation via a PPARalpha-dependent pathway. *Arterioscler Thromb Vasc Biol* 24: 1621–1627, 2004.
248. Montessuit C, Papageorgiou I, and Lerch R. Nuclear receptor agonists improve insulin responsiveness in cultured cardiomyocytes through enhanced signaling and preserved cytoskeletal architecture. *Endocrinology* 149: 1064–1074, 2008.
249. Moore KJ, Rosen ED, Fitzgerald ML, Randow F, Andersson LP, Altshuler D, Milstone DS, Mortensen RM, Spiegelman BM, and Freeman MW. The role of PPAR-gamma in macrophage differentiation and cholesterol uptake. *Nat Med* 7: 41–47, 2001.
250. Morikang E, Benson SC, Kurtz TW, and Pershadsingh HA. Effects of thiazolidinediones on growth and differentiation of human aorta and coronary myocytes. *Am J Hypertens* 10: 440–446, 1997.
251. Muerhoff AS, Griffin KJ, and Johnson EF. The peroxisome proliferator-activated receptor mediates the induction of CYP4A6, a cytochrome P450 fatty acid omega-hydroxylase, by clofibrac acid. *J Biol Chem* 267: 19051–19053, 1992.
252. Mukherjee R, Hoener PA, Jow L, Bilakovics J, Klausner K, Mais DE, Faulkner A, Croston GE, and Paterniti JR Jr. A selective peroxisome proliferator-activated receptor-gamma (PPARgamma) modulator blocks adipocyte differentiation but stimulates glucose uptake in 3T3-L1 adipocytes. *Mol Endocr (Baltimore)* 14: 1425–1433, 2000.
253. Murao K, Imachi H, Momoi A, Sayo Y, Hosokawa H, Sato M, Ishida T, and Takahara J. Thiazolidinedione inhibits the production of monocyte chemoattractant protein-1 in cytokine-treated human vascular endothelial cells. *FEBS Lett* 454: 27–30, 1999.
254. Nagy L, Tontonoz P, Alvarez JG, Chen H, and Evans RM. Oxidized LDL regulates macrophage gene expression through ligand activation of PPARgamma. *Cell* 93: 229–240, 1998.
255. Nakamachi T, Nomiya T, Gizard F, Heywood EB, Jones KL, Zhao Y, Fuentes L, Takebayashi K, Aso Y, Staels B, Inukai T, and Bruemmer D. PPARalpha agonists suppress osteopontin expression in macrophages and decrease plasma levels in patients with type 2 diabetes. *Diabetes* 56: 1662–1670, 2007.
256. Narkar VA, Downes M, Yu RT, Emblar E, Wang YX, Banayo E, Mihaylova MM, Nelson MC, Zou Y, Juguilon H, Kang H, Shaw RJ, and Evans RM. AMPK and PPARdelta agonists are exercise mimetics. *Cell* 134: 405–415, 2008.
257. Nawa T, Nawa MT, Cai Y, Zhang C, Uchimura I, Narumi S, Numano F, and Kitajima S. Repression of TNF-alpha-induced E-selectin expression by PPAR activators: involvement of transcriptional repressor LRF-1/ATF3. *Biochem Biophys Res Commun* 275: 406–411, 2000.
258. Nettles KW. Insights into PPARgamma from structures with endogenous and covalently bound ligands. *Nat Struct Mol Biol* 15: 893–895, 2008.
259. Neuschwander-Tetri BA, Isley WL, Oki JC, Ramrakhiani S, Quiason SG, Phillips NJ, and Brunt EM. Troglitazone-induced hepatic failure leading to liver transplantation: a case report. *Ann Intern Med* 129: 38–41, 1998.
260. Neve BP, Corseaux D, Chinetti G, Zawadzki C, Fruchart JC, Duriez P, Staels B, and Jude B. PPARalpha agonists inhibit tissue factor expression in human monocytes and macrophages. *Circulation* 103: 207–212, 2001.
261. Ng VY, Morisseau C, Falck JR, Hammock BD, and Kroetz DL. Inhibition of smooth muscle proliferation by urea-based alkanolic acids via peroxisome proliferator-activated receptor alpha-dependent repression of cyclin D1. *Arterioscler Thromb Vasc Biol* 26: 2462–2468, 2006.
262. Nichols JS, Parks DJ, Consler TG, and Blanchard SG. Development of a scintillation proximity assay for peroxisome proliferator-activated receptor gamma ligand binding domain. *Anal Biochem* 257: 112–119, 1998.
263. Niemeyer NV and Janney LM. Thiazolidinedione-induced edema. *Pharmacotherapy* 22: 924–929, 2002.
264. Nigro J, Dilley RJ, and Little PJ. Differential effects of gemfibrozil on migration, proliferation and proteoglycan production in human vascular smooth muscle cells. *Atherosclerosis* 162: 119–129, 2002.
265. Nissen SE and Wolski K. Effect of rosiglitazone on the risk of myocardial infarction and death from cardiovascular causes. *N Engl J Med* 356: 2457–2471, 2007.
266. O'Brien KD, Gordon D, Deeb S, Ferguson M, and Chait A. Lipoprotein lipase is synthesized by macrophage-derived foam cells in human coronary atherosclerotic plaques. *J Clin Invest* 89: 1544–1550, 1992.
267. Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, Morel CR, Subramanian V, Mukundan L, Eagle AR, Vats D, Brombacher F, Ferrante AW, and Chawla A. Macrophage-specific PPARgamma controls alternative activation and improves insulin resistance. *Nature* 447: 1116–1120, 2007.
268. Odegaard JI, Ricardo-Gonzalez RR, Red Eagle A, Vats D, Morel CR, Goforth MH, Subramanian V, Mukundan L, Ferrante AW, and Chawla A. Alternative M2 activation of Kupffer cells by PPARdelta ameliorates obesity-induced insulin resistance. *Cell Metabol* 7: 496–507, 2008.
269. Ogawa D, Nomiya T, Nakamachi T, Heywood EB, Stone JF, Berger JP, Law RE, and Bruemmer D. Activation of peroxisome proliferator-activated receptor gamma suppresses telomerase activity in vascular smooth muscle cells. *Circ Res* 98: e50–e59, 2006.
270. Ogawa S, Lozach J, Benner C, Pascual G, Tangirala RK, Westin S, Hoffmann A, Subramaniam S, David M, Rosenfeld MG, and Glass CK. Molecular determinants of crosstalk between nuclear receptors and toll-like receptors. *Cell* 122: 707–721, 2005.
271. Ohshima T, Koga H, and Shimotohno K. Transcriptional activity of peroxisome proliferator-activated receptor gamma is modulated by SUMO-1 modification. *J Biol Chem* 279: 29551–29557, 2004.
272. Ohtani K, Iwanaga R, Nakamura M, Ikeda M, Yabuta N, Tsuruga H, and Nojima H. Cell growth-regulated expression of mammalian MCM5 and MCM6 genes mediated by the transcription factor E2F. *Oncogene* 18: 2299–2309, 1999.

273. Oishi Y, Manabe I, Tobe K, Ohsugi M, Kubota T, Fujiu K, Maemura K, Kubota N, Kadowaki T, and Nagai R. SUMOylation of Kruppel-like transcription factor 5 acts as a molecular switch in transcriptional programs of lipid metabolism involving PPAR-delta. *Nat Med* 14: 656–666, 2008.
274. Oishi Y, Manabe I, Tobe K, Tsushima K, Shindo T, Fujiu K, Nishimura G, Maemura K, Yamauchi T, Kubota N, Suzuki R, Kitamura T, Akira S, Kadowaki T, and Nagai R. Kruppel-like transcription factor KLF5 is a key regulator of adipocyte differentiation. *Cell Metab* 1: 27–39, 2005.
275. Okura T, Nakamura M, Takata Y, Watanabe S, Kitami Y, and Hiwada K. Troglitazone induces apoptosis via the p53 and Gadd45 pathway in vascular smooth muscle cells. *Eur J Pharmacol* 407: 227–235, 2000.
276. Oliver WR Jr., Shenk JL, Snaith MR, Russell CS, Plunket KD, Bodkin NL, Lewis MC, Winegar DA, Sznaidman ML, Lambert MH, Xu HE, Sternbach DD, Kliewer SA, Hansen BC, and Willson TM. A selective peroxisome proliferator-activated receptor delta agonist promotes reverse cholesterol transport. *Proc Natl Acad Sci U S A* 98: 5306–5311, 2001.
277. Oyama Y, Akuzawa N, Nagai R, and Kurabayashi M. PPARgamma ligand inhibits osteopontin gene expression through interference with binding of nuclear factors to A/T-rich sequence in THP-1 cells. *Circ Res* 90: 348–355, 2002.
278. Oyama Y, Kurabayashi M, Akuzawa N, and Nagai R. Troglitazone, a PPARgamma ligand, inhibits osteopontin gene expression in human monocytes/macrophage THP-1 cells. *J Atheroscler Thromb* 7: 77–82, 2000.
279. Paddock ML, Wiley SE, Axelrod HL, Cohen AE, Roy M, Abresch EC, Capraro D, Murphy AN, Nechushtai R, Dixon JE, and Jennings PA. MitoNEET is a uniquely folded 2Fe 2S outer mitochondrial membrane protein stabilized by pioglitazone. *Proc Natl Acad Sci U S A* 104: 14342–14347, 2007.
280. Palmer RM, Ferrige AG, and Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature* 327: 524–526, 1987.
281. Panagia M, Gibbons GF, Radda GK, and Clarke K. PPAR-alpha activation required for decreased glucose uptake and increased susceptibility to injury during ischemia. *Am J Physiol Heart Circ Physiol* 288: H2677–H2683, 2005.
282. Panigrahy D, Singer S, Shen LQ, Butterfield CE, Freedman DA, Chen EJ, Moses MA, Kilroy S, Duensing S, Fletcher C, Fletcher JA, Hlatky L, Hahnfeldt P, Folkman J, and Kaipainen A. PPARgamma ligands inhibit primary tumor growth and metastasis by inhibiting angiogenesis. *J Clin Invest* 110: 923–932, 2002.
283. Park KG, Lee KM, Chang YC, Magae J, Ando K, Kim KB, Kim YN, Kim HS, Park JY, Lee KU, and Lee IK. The ascochlorin derivative, AS-6, inhibits TNF-alpha-induced adhesion molecule and chemokine expression in rat vascular smooth muscle cells. *Life Sci* 80: 120–126, 2006.
284. Park SY, Cho YR, Finck BN, Kim HJ, Higashimori T, Hong EG, Lee MK, Danton C, Deshmukh S, Cline GW, Wu JJ, Bennett AM, Rothermel B, Kalinowski A, Russell KS, Kim YB, Kelly DP, and Kim JK. Cardiac-specific overexpression of peroxisome proliferator-activated receptor-alpha causes insulin resistance in heart and liver. *Diabetes* 54: 2514–2524, 2005.
285. Pasceri V, Cheng JS, Willerson JT, and Yeh ET. Modulation of C-reactive protein-mediated monocyte chemoattractant protein-1 induction in human endothelial cells by anti-atherosclerosis drugs. *Circulation* 103: 2531–2534, 2001.
286. Pasceri V, Wu HD, Willerson JT, and Yeh ET. Modulation of vascular inflammation *in vitro* and *in vivo* by peroxisome proliferator-activated receptor-gamma activators. *Circulation* 101: 235–238, 2000.
287. Pascual G, Fong AL, Ogawa S, Gamliel A, Li AC, Perissi V, Rose DW, Willson TM, Rosenfeld MG, and Glass CK. A SUMOylation-dependent pathway mediates transrepression of inflammatory response genes by PPAR-gamma. *Nature* 437: 759–763, 2005.
288. Paumelle R, Blanquart C, Briand O, Barbier O, Duhem C, Woerly G, Percevault F, Fruchart JC, Dombrowicz D, Glineur C, and Staels B. Acute antiinflammatory properties of statins involve peroxisome proliferator-activated receptor-alpha via inhibition of the protein kinase C signaling pathway. *Circ Res* 98: 361–369, 2006.
289. Peters JM, Lee SS, Li W, Ward JM, Gavrilova O, Everett C, Reitman ML, Hudson LD, and Gonzalez FJ. Growth, adipose, brain, and skin alterations resulting from targeted disruption of the mouse peroxisome proliferator-activated receptor beta(delta). *Mol Cell Biol* 20: 5119–5128, 2000.
290. Piqueras L, Reynolds AR, Hodivala-Dilke KM, Alfranca A, Redondo JM, Hatae T, Tanabe T, Warner TD, and Bishop-Bailey D. Activation of PPARbeta/delta induces endothelial cell proliferation and angiogenesis. *Arterioscler Thromb Vasc Biol* 27: 63–69, 2007.
291. Pistrosch F, Herbrig K, Oelschlaegel U, Richter S, Passauer J, Fischer S, and Gross P. PPARgamma-agonist rosiglitazone increases number and migratory activity of cultured endothelial progenitor cells. *Atherosclerosis* 183: 163–167, 2005.
292. Pistrosch F, Passauer J, Fischer S, Fuecker K, Hanefeld M, and Gross P. In type 2 diabetes, rosiglitazone therapy for insulin resistance ameliorates endothelial dysfunction independent of glucose control. *Diabetes Care* 27: 484–490, 2004.
293. Planavila A, Rodriguez-Calvo R, Jove M, Michalik L, Wahli W, Laguna JC, and Vazquez-Carrera M. Peroxisome proliferator-activated receptor beta/delta activation inhibits hypertrophy in neonatal rat cardiomyocytes. *Cardiovasc Res* 65: 832–841, 2005.
294. Polikandriotis JA, Mazzella LJ, Rupnow HL, and Hart CM. Peroxisome proliferator-activated receptor gamma ligands stimulate endothelial nitric oxide production through distinct peroxisome proliferator-activated receptor gamma-dependent mechanisms. *Arterioscler Thromb Vasc Biol* 25: 1810–1816, 2005.
295. Porcheray F, Viaud S, Rimaniol AC, Leone C, Samah B, Dereuddre-Bosquet N, Dormont D, and Gras G. Macrophage activation switching: an asset for the resolution of inflammation. *Clin Exp Immunol* 142: 481–489, 2005.
296. Pourcet B, Fruchart JC, Staels B, and Glineur C. Selective PPAR modulators, dual and pan PPAR agonists: multimodal drugs for the treatment of type 2 diabetes and atherosclerosis. *Expert Opin Emerg Drugs* 11: 379–401, 2006.
297. Ptasinska A, Wang S, Zhang J, Wesley RA, and Danner RL. Nitric oxide activation of peroxisome proliferator-activated receptor gamma through a p38 MAPK signaling pathway. *FASEB J* 21: 950–961, 2007.
298. Ravau L, Denoyelle C, Monne C, Limon I, Raymondjean M, and El Hadri K. Inhibition of interleukin-1beta-induced group IIA secretory phospholipase A2 expression by peroxisome proliferator-activated receptors (PPARs) in rat vascular smooth muscle cells: cooperation between

- PPARbeta and the proto-oncogene BCL-6. *Mol Cell Biol* 27: 8374–8387, 2007.
299. Razeghi P, Young ME, Abbasi S, and Taegtmeier H. Hypoxia *in vivo* decreases peroxisome proliferator-activated receptor alpha-regulated gene expression in rat heart. *Biochem Biophys Res Commun* 287: 5–10, 2001.
 300. Reddy J, Svoboda D, and Azarnoff D. Microbody proliferation in liver induced by nafenopin, a new hypolipidemic drug: comparison with CPIB. *Biochem Biophys Res Commun* 52: 537–543, 1973.
 301. Reddy JK. Hepatic microbody proliferation and catalase synthesis induced by methyl clofenapate, a hypolipidemic analog of CPIB. *Am J Pathol* 75: 103–118, 1974.
 302. Reddy JK and Krishnakantha TP. Hepatic peroxisome proliferation: induction by two novel compounds structurally unrelated to clofibrate. *Science (New York)* 190: 787–789, 1975.
 303. Redondo S, Ruiz E, Santos-Gallego CG, Padilla E, and Tejerina T. Pioglitazone induces vascular smooth muscle cell apoptosis through a peroxisome proliferator-activated receptor-gamma, transforming growth factor-beta1, and a Smad2-dependent mechanism. *Diabetes* 54: 811–817, 2005.
 304. Richter EA, Kiens B, and Wojtaszewski JF. Can exercise mimetics substitute for exercise? *Cell Metab* 8: 96–98, 2008.
 305. Ricote M and Glass CK. PPARs and molecular mechanisms of transrepression. *Biochim Biophys Acta* 1771: 926–935, 2007.
 306. Ricote M, Huang J, Fajas L, Li A, Welch J, Najib J, Witztum JL, Auwerx J, Palinski W, and Glass CK. Expression of the peroxisome proliferator-activated receptor gamma (PPAR-gamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc Natl Acad Sci U S A* 95: 7614–7619, 1998.
 307. Ricote M, Li AC, Willson TM, Kelly CJ, and Glass CK. The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature* 391: 79–82, 1998.
 308. Rieusset J, Touri F, Michalik L, Escher P, Desvergne B, Niesor E, and Wahli W. A new selective peroxisome proliferator-activated receptor gamma antagonist with anti-obesity and antidiabetic activity. *Mol Endocr (Baltimore)* 16: 2628–2644, 2002.
 309. Rigamonti E, Fontaine C, Lefebvre B, Duhem C, Lefebvre P, Marx N, Staels B, and Chinetti-Gbaguidi G. Induction of CXCR2 receptor by peroxisome proliferator-activated receptor gamma in human macrophages. *Arterioscler Thromb Vasc Biol* 28: 932–939, 2008.
 310. Riserus U, Sprecher D, Johnson T, Olson E, Hirschberg S, Liu A, Fang Z, Hegde P, Richards D, Sarov-Blat L, Strum JC, Basu S, Cheeseman J, Fielding BA, Humphreys SM, Danoff T, Moore NR, Murgatroyd P, O'Rahilly S, Sutton P, Willson T, Hassall D, Frayn KN, and Karpe F. Activation of peroxisome proliferator-activated receptor (PPAR)delta promotes reversal of multiple metabolic abnormalities, reduces oxidative stress, and increases fatty acid oxidation in moderately obese men. *Diabetes* 57: 332–339, 2008.
 311. Rival Y, Beneteau N, Taillandier T, Pezet M, Dupont-Passelaigue E, Patoiseau JF, Junquero D, Colpaert FC, and Delhon A. PPARalpha and PPARdelta activators inhibit cytokine-induced nuclear translocation of NF-kappaB and expression of VCAM-1 in EAhy926 endothelial cells. *Eur J Pharmacol* 435: 143–151, 2002.
 312. Robyr D, Wolffe AP, and Wahli W. Nuclear hormone receptor coregulators in action: diversity for shared tasks. *Mol Endocr (Baltimore)* 14: 329–347, 2000.
 313. Ross R. Atherosclerosis: an inflammatory disease. *N Engl J Med* 340: 115–126, 1999.
 314. Rossi A, Kapahi P, Natoli G, Takahashi T, Chen Y, Karin M, and Santoro MG. Anti-inflammatory cyclopentenone prostaglandins are direct inhibitors of IkappaB kinase. *Nature* 403: 103–108, 2000.
 315. Ruiz E, Redondo S, Gordillo-Moscoso A, and Tejerina T. Pioglitazone induces apoptosis in human vascular smooth muscle cells from diabetic patients involving the transforming growth factor-beta/activin receptor-like kinase-4/5/7/Smad2 signaling pathway. *J Pharmacol Exp Ther* 321: 431–438, 2007.
 316. Sambandam N, Morabito D, Wagg C, Finck BN, Kelly DP, and Lopaschuk GD. Chronic activation of PPARalpha is detrimental to cardiac recovery after ischemia. *Am J Physiol Heart Circ Physiol* 290: H87–H95, 2006.
 317. Santini E, Fallahi P, Ferrari SM, Masoni A, Antonelli A, and Ferrannini E. Effect of PPAR-gamma activation and inhibition on glucose-stimulated insulin release in INS-1e cells. *Diabetes* 53(suppl 3): S79–S83, 2004.
 318. Satoh H, Tsukamoto K, Hashimoto Y, Hashimoto N, Togo M, Hara M, Maekawa H, Isoo N, Kimura S, and Watanabe T. Thiazolidinediones suppress endothelin-1 secretion from bovine vascular endothelial cells: a new possible role of PPARgamma on vascular endothelial function. *Biochem Biophys Res Commun* 254: 757–763, 1999.
 319. Schopfer FJ, Lin Y, Baker PR, Cui T, Garcia-Barrio M, Zhang J, Chen K, Chen YE, and Freeman BA. Nitrolinoleic acid: an endogenous peroxisome proliferator-activated receptor gamma ligand. *Proc Natl Acad Sci U S A* 102: 2340–2345, 2005.
 320. Semple RK, Meirhaeghe A, Vidal-Puig AJ, Schwabe JW, Wiggins D, Gibbons GF, Gurnell M, Chatterjee VK, and O'Rahilly S. A dominant negative human peroxisome proliferator-activated receptor (PPAR){alpha} is a constitutive transcriptional corepressor and inhibits signaling through all PPAR isoforms. *Endocrinology* 146: 1871–1882, 2005.
 321. Sethi S, Ziouzenkova O, Ni H, Wagner DD, Plutzky J, and Mayadas TN. Oxidized omega-3 fatty acids in fish oil inhibit leukocyte-endothelial interactions through activation of PPAR alpha. *Blood* 100: 1340–1346, 2002.
 322. Shalev A, Siegrist-Kaiser CA, Yen PM, Wahli W, Burger AG, Chin WW, and Meier CA. The peroxisome proliferator-activated receptor alpha is a phosphoprotein: regulation by insulin. *Endocrinology* 137: 4499–4502, 1996.
 323. Sharma S, Taegtmeier H, Adroque J, Razeghi P, Sen S, Ngumbela K, and Essop MF. Dynamic changes of gene expression in hypoxia-induced right ventricular hypertrophy. *Am J Physiol Heart Circ Physiol* 286: H1185–H1192, 2004.
 324. Shaw N, Elholm M, and Noy N. Retinoic acid is a high affinity selective ligand for the peroxisome proliferator-activated receptor beta/delta. *J Biol Chem* 278: 41589–41592, 2003.
 325. Shearer BG and Billin AN. The next generation of PPAR drugs: do we have the tools to find them? *Biochim Biophys Acta* 1771: 1082–1093, 2007.
 326. Shearer BG, Steger DJ, Way JM, Stanley TB, Lobe DC, Grillot DA, Iannone MA, Lazar MA, Willson TM, and Billin AN. Identification and characterization of a selective per-

- oxisome proliferator-activated receptor beta/delta (NR1C2) antagonist. *Mol Endocr (Baltimore)* 22: 523–529, 2008.
327. Sherr CJ. G1 phase progression: cycling on cue. *Cell* 79: 551–555, 1994.
 328. Sherr CJ and Roberts JM. CDK inhibitors: positive and negative regulators of G1-phase progression. *Genes Dev* 13: 1501–1512, 1999.
 329. Shi Y, Hon M, and Evans RM. The peroxisome proliferator-activated receptor delta, an integrator of transcriptional repression and nuclear receptor signaling. *Proc Natl Acad Sci U S A* 99: 2613–2618, 2002.
 330. Shu H, Wong B, Zhou G, Li Y, Berger J, Woods JW, Wright SD, and Cai TQ. Activation of PPARalpha or gamma reduces secretion of matrix metalloproteinase 9 but not interleukin 8 from human monocytic THP-1 cells. *Biochem Biophys Res Commun* 267: 345–349, 2000.
 331. Sierra-Honigsmann MR, Nath AK, Murakami C, Garcia-Cardena G, Papapetropoulos A, Sessa WC, Madge LA, Schechner JS, Schwabb MB, Polverini PJ, and Flores-Riveros JR. Biological action of leptin as an angiogenic factor. *Science (New York)* 281: 1683–1686, 1998.
 332. Siess W, Zangl KJ, Essler M, Bauer M, Brandl R, Corrinth C, Bittman R, Tigyi G, and Aepfelbacher M. Lysophosphatidic acid mediates the rapid activation of platelets and endothelial cells by mildly oxidized low density lipoprotein and accumulates in human atherosclerotic lesions. *Proc Natl Acad Sci U S A* 96: 6931–6936, 1999.
 333. Skogsberg J, Kannisto K, Cassel TN, Hamsten A, Eriksson P, and Ehrenborg E. Evidence that peroxisome proliferator-activated receptor delta influences cholesterol metabolism in men. *Arterioscler Thromb Vasc Biol* 23: 637–643, 2003.
 334. Smeets PJ, Teunissen BE, Planavila A, de Vogel-van den Bosch H, Willemsen PH, van der Vusse GJ, and van Bilsen M. Inflammatory pathways are activated during cardiomyocyte hypertrophy and attenuated by peroxisome proliferator-activated receptors PPAR[alpha] and PPAR [delta]. *J Biol Chem* 283: 29109–29118, 2008.
 335. Smeets PJ, Teunissen BE, Willemsen PH, van Nieuwenhoven FA, Brouns AE, Janssen BJ, Cleutjens JP, Staels B, van der Vusse GJ, and van Bilsen M. Cardiac hypertrophy is enhanced in PPAR alpha^{-/-} mice in response to chronic pressure overload. *Cardiovasc Res* 78: 79–89, 2008.
 336. Son NH, Park TS, Yamashita H, Yokoyama M, Huggins LA, Okajima K, Homma S, Szabolcs MJ, Huang LS, and Goldberg JJ. Cardiomyocyte expression of PPARgamma leads to cardiac dysfunction in mice. *J Clin Invest* 117: 2791–2801, 2007.
 337. Song J, Walsh MF, Igwe R, Ram JL, Barazi M, Dominguez LJ, and Sowers JR. Troglitazone reduces contraction by inhibition of vascular smooth muscle cell Ca²⁺ currents and not endothelial nitric oxide production. *Diabetes* 46: 659–664, 1997.
 338. Sorrentino SA, Bahlmann FH, Besler C, Muller M, Schulz S, Kirchhoff N, Doerries C, Horvath T, Limbourg A, Limbourg F, Fliser D, Haller H, Drexler H, and Landmesser U. Oxidant stress impairs *in vivo* reendothelialization capacity of endothelial progenitor cells from patients with type 2 diabetes mellitus: restoration by the peroxisome proliferator-activated receptor-gamma agonist rosiglitazone. *Circulation* 116: 163–173, 2007.
 339. Sprecher DL, Massien C, Pearce G, Billin AN, Perlstein I, Willson TM, Hassall DG, Ancellin N, Patterson SD, Lobe DC, and Johnson TG. Triglyceride:high-density lipoprotein cholesterol effects in healthy subjects administered a peroxisome proliferator activated receptor delta agonist. *Arterioscler Thromb Vasc Biol* 27: 359–365, 2007.
 340. Staels B, Koenig W, Habib A, Merval R, Lebret M, Torra IP, Delerive P, Fadel A, Chinetti G, Fruchart JC, Najib J, Maclouf J, and Tedgui A. Activation of human aortic smooth-muscle cells is inhibited by PPARalpha but not by PPARgamma activators. *Nature* 393: 790–793, 1998.
 341. Stout RD and Suttles J. Functional plasticity of macrophages: reversible adaptation to changing microenvironments. *J Leukoc Biol* 76: 509–513, 2004.
 342. Straus DS, Pascual G, Li M, Welch JS, Ricote M, Hsiang CH, Sengchanthalangsy LL, Ghosh G, and Glass CK. 15-Deoxy-delta 12,14-prostaglandin J2 inhibits multiple steps in the NF-kappa B signaling pathway. *Proc Natl Acad Sci U S A* 97: 4844–4849, 2000.
 343. Sugawara A, Takeuchi K, Uruno A, Ikeda Y, Arima S, Kudo M, Sato K, Taniyama Y, and Ito S. Transcriptional suppression of type 1 angiotensin II receptor gene expression by peroxisome proliferator-activated receptor-gamma in vascular smooth muscle cells. *Endocrinology* 142: 3125–3134, 2001.
 344. Suh N, Wang Y, Williams CR, Risingsong R, Gilmer T, Willson TM, and Sporn MB. A new ligand for the peroxisome proliferator-activated receptor-gamma (PPAR-gamma), GW7845, inhibits rat mammary carcinogenesis. *Cancer Res* 59: 5671–5673, 1999.
 345. Sznajdman ML, Haffner CD, Maloney PR, Fivush A, Chao E, Goreham D, Sierra ML, LeGrumelec C, Xu HE, Montana VG, Lambert MH, Willson TM, Oliver WR Jr, and Sternbach DD. Novel selective small molecule agonists for peroxisome proliferator-activated receptor delta (PPARdelta): synthesis and biological activity. *Bioorg Med Chem Lett* 13: 1517–1521, 2003.
 346. Takata Y, Kitami Y, Okura T, and Hiwada K. Peroxisome proliferator-activated receptor-gamma activation inhibits interleukin-1beta-mediated platelet-derived growth factor-alpha receptor gene expression via CCAAT/enhancer-binding protein-delta in vascular smooth muscle cells. *J Biol Chem* 276: 12893–12897, 2001.
 347. Takata Y, Kitami Y, Yang ZH, Nakamura M, Okura T, and Hiwada K. Vascular inflammation is negatively autoregulated by interaction between CCAAT/enhancer-binding protein-delta and peroxisome proliferator-activated receptor-gamma. *Circ Res* 91: 427–433, 2002.
 348. Takata Y, Liu J, Yin F, Collins AR, Lyon CJ, Lee CH, Atkins AR, Downes M, Barish GD, Evans RM, Hsueh WA, and Tangirala RK. PPARdelta-mediated antiinflammatory mechanisms inhibit angiotensin II-accelerated atherosclerosis. *Proc Natl Acad Sci U S A* 105: 4277–4282, 2008.
 349. Takeda K, Ichiki T, Tokunou T, Funakoshi Y, Iino N, Hirano K, Kanaide H, and Takeshita A. Peroxisome proliferator-activated receptor gamma activators down-regulate angiotensin II type 1 receptor in vascular smooth muscle cells. *Circulation* 102: 1834–1839, 2000.
 350. Tan NS, Michalik L, Noy N, Yasmin R, Pacot C, Heim M, Fluhmann B, Desvergne B, and Wahli W. Critical roles of PPAR beta/delta in keratinocyte response to inflammation. *Genes Dev* 15: 3263–3277, 2001.
 351. Tao L, Liu HR, Gao E, Teng ZP, Lopez BL, Christopher TA, Ma XL, Batinic-Haberle I, Willette RN, Ohlstein EH, and Yue TL. Antioxidative, antinitrative, and vasculoprotective effects of a peroxisome proliferator-activated receptor-gamma agonist in hypercholesterolemia. *Circulation* 108: 2805–2811, 2003.

352. Teissier E, Nohara A, Chinetti G, Paumelle R, Cariou B, Fruchart JC, Brandes RP, Shah A, and Staels B. Peroxisome proliferator-activated receptor alpha induces NADPH oxidase activity in macrophages, leading to the generation of LDL with PPAR-alpha activation properties. *Circ Res* 95: 1174–1182, 2004.
353. Tenenbaum A, Motro M, Fisman EZ, Tanne D, Boyko V, and Behar S. Bezafibrate for the secondary prevention of myocardial infarction in patients with metabolic syndrome. *Arch Intern Med* 165: 1154–1160, 2005.
354. Teunissen BE, Smeets PJ, Willemsen PH, De Windt LJ, Van der Vusse GJ, and Van Bilsen M. Activation of PPARdelta inhibits cardiac fibroblast proliferation and the transdifferentiation into myofibroblasts. *Cardiovasc Res* 75: 519–529, 2007.
355. Tham DM, Martin-McNulty B, Wang YX, Wilson DW, Vergona R, Sullivan ME, Dole W, and Rutledge JC. Angiotensin II is associated with activation of NF-kappaB-mediated genes and downregulation of PPARs. *Physiol Genom* 11: 21–30, 2002.
356. Thom T, Haase N, Rosamond W, Howard VJ, Rumsfeld J, Manolio T, Zheng ZJ, Flegal K, O'Donnell C, Kittner S, Lloyd-Jones D, Goff DC Jr, Hong Y, Adams R, Friday G, Furie K, Gorelick P, Kissela B, Marler J, Meigs J, Roger V, Sidney S, Sorlie P, Steinberger J, Wasserthiel-Smoller S, Wilson M, and Wolf P. Heart disease and stroke statistics: 2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 113: e85–e151, 2006.
357. Tolman KG and Chandramouli J. Hepatotoxicity of the thiazolidinediones. *Clin Liver Dis* 7: 369–379, vi, 2003.
358. Tontonoz P, Nagy L, Alvarez JG, Thomazy VA, and Evans RM. PPARgamma promotes monocyte/macrophage differentiation and uptake of oxidized LDL. *Cell* 93: 241–252, 1998.
359. Tordjman K, Bernal-Mizrachi C, Zemany L, Weng S, Feng C, Zhang F, Leone TC, Coleman T, Kelly DP, and Semenkovich CF. PPARalpha deficiency reduces insulin resistance and atherosclerosis in apoE-null mice. *J Clin Invest* 107: 1025–1034, 2001.
360. Tran H, Brunet A, Griffith EC, and Greenberg ME. The many forks in FOXO's road. *Sci STKE* 2003: RE5, 2003.
361. Tsukahara T, Tsukahara R, Yasuda S, Makarova N, Valentine WJ, Allison P, Yuan H, Baker DL, Li Z, Bittman R, Parrill A, and Tigyi G. Different residues mediate recognition of 1-O-oleyllysophosphatidic acid and rosiglitazone in the ligand binding domain of peroxisome proliferator-activated receptor gamma. *J Biol Chem* 281: 3398–3407, 2006.
362. Tugwood JD, Issemann I, Anderson RG, Bundell KR, McPheat WL, and Green S. The mouse peroxisome proliferator activated receptor recognizes a response element in the 5' flanking sequence of the rat acyl CoA oxidase gene. *EMBO J* 11: 433–439, 1992.
363. van der Lee KA, Vork MM, De Vries JE, Willemsen PH, Glatz JF, Reneman RS, Van der Vusse GJ, and Van Bilsen M. Long-chain fatty acid-induced changes in gene expression in neonatal cardiac myocytes. *J Lipid Res* 41: 41–47, 2000.
364. Van Ginderachter JA, Meerschaut S, Liu Y, Brys L, De Groeve K, Hassanzadeh Ghassabeh G, Raes G, and De Baetselier P. Peroxisome proliferator-activated receptor gamma (PPARgamma) ligands reverse CTL suppression by alternatively activated (M2) macrophages in cancer. *Blood* 108: 525–535, 2006.
365. Van Ginderachter JA, Movahedi K, Hassanzadeh Ghassabeh G, Meerschaut S, Beschin A, Raes G, and De Baetselier P. Classical and alternative activation of mononuclear phagocytes: picking the best of both worlds for tumor promotion. *Immunobiology* 211: 487–501, 2006.
366. Vecchione C, Maffei A, Colella S, Aretini A, Poulet R, Frati G, Gentile MT, Fratta L, Trimarco V, Trimarco B, and Lembo G. Leptin effect on endothelial nitric oxide is mediated through Akt-endothelial nitric oxide synthase phosphorylation pathway. *Diabetes* 51: 168–173, 2002.
367. Verma S, Kuliszewski MA, Li SH, Szmítko PE, Zucco L, Wang CH, Badiwala MV, Mickle DA, Weisel RD, Fedak PW, Stewart DJ, and Kutryk MJ. C-reactive protein attenuates endothelial progenitor cell survival, differentiation, and function: further evidence of a mechanistic link between C-reactive protein and cardiovascular disease. *Circulation* 109: 2058–2067, 2004.
368. Verrier E, Wang L, Wadham C, Albanese N, Hahn C, Gamble JR, Chatterjee VK, Vadas MA, and Xia P. PPARgamma agonists ameliorate endothelial cell activation via inhibition of diacylglycerol-protein kinase C signaling pathway: role of diacylglycerol kinase. *Circ Res* 94: 1515–1522, 2004.
369. Villacorta L, Schopfer FJ, Zhang J, Freeman BA, and Chen YE. PPARgamma and its ligands: therapeutic implications in cardiovascular disease. *Clin Sci (Lond)* 116: 205–218, 2009.
370. Vosper H, Khoudoli GA, and Palmer CN. The peroxisome proliferator activated receptor delta is required for the differentiation of THP-1 monocytic cells by phorbol ester. *Nucl Recept* 1: 9, 2003.
371. Vosper H, Patel L, Graham TL, Khoudoli GA, Hill A, Macphee CH, Pinto I, Smith SA, Suckling KE, Wolf CR, and Palmer CN. The peroxisome proliferator-activated receptor delta promotes lipid accumulation in human macrophages. *J Biol Chem* 276: 44258–44265, 2001.
372. Wakino S, Hayashi K, Kanda T, Tatematsu S, Homma K, Yoshioka K, Takamatsu I, and Saruta T. Peroxisome proliferator-activated receptor gamma ligands inhibit Rho/Rho kinase pathway by inducing protein tyrosine phosphatase SHP-2. *Circ Res* 95: e45–e55, 2004.
373. Wakino S, Kintscher U, Kim S, Yin F, Hsueh WA, and Law RE. Peroxisome proliferator-activated receptor gamma ligands inhibit retinoblastoma phosphorylation and G1->S transition in vascular smooth muscle cells. *J Biol Chem* 275: 22435–22441, 2000.
374. Wakino S, Kintscher U, Liu Z, Kim S, Yin F, Ohba M, Kuroki T, Schonthal AH, Hsueh WA, and Law RE. Peroxisome proliferator-activated receptor gamma ligands inhibit mitogenic induction of p21(Cip1) by modulating the protein kinase Cdelta pathway in vascular smooth muscle cells. *J Biol Chem* 276: 47650–47657, 2001.
375. Wallace JM, Schwarz M, Coward P, Houze J, Sawyer JK, Kelley KL, Chai A, and Rudel LL. Effects of peroxisome proliferator-activated receptor alpha/delta agonists on HDL-cholesterol in vervet monkeys. *J Lipid Res* 46: 1009–1016, 2005.
376. Walter DH, Rittig K, Bahlmann FH, Kirchmair R, Silver M, Murayama T, Nishimura H, Losordo DW, Asahara T, and Isner JM. Statin therapy accelerates reendothelialization: a novel effect involving mobilization and incorporation of bone marrow-derived endothelial progenitor cells. *Circulation* 105: 3017–3024, 2002.
377. Wang CH, Ciliberti N, Li SH, Szmítko PE, Weisel RD, Fedak PW, Al-Omran M, Cherng WJ, Li RK, Stanford WL,

- and Verma S. Rosiglitazone facilitates angiogenic progenitor cell differentiation toward endothelial lineage: a new paradigm in glitazone pleiotropy. *Circulation* 109: 1392–1400, 2004.
378. Wang N, Verna L, Chen NG, Chen J, Li H, Forman BM, and Stemberman MB. Constitutive activation of peroxisome proliferator-activated receptor-gamma suppresses proinflammatory adhesion molecules in human vascular endothelial cells. *J Biol Chem* 277: 34176–34181, 2002.
 379. Wang YX, Lee CH, Tiep S, Yu RT, Ham J, Kang H, and Evans RM. Peroxisome-proliferator-activated receptor delta activates fat metabolism to prevent obesity. *Cell* 113: 159–170, 2003.
 380. Wang YX, Zhang CL, Yu RT, Cho HK, Nelson MC, Bayuga-Ocampo CR, Ham J, Kang H, and Evans RM. Regulation of muscle fiber type and running endurance by PPARdelta. *PLoS Biol* 2: e294, 2004.
 381. Watanabe K, Fujii H, Takahashi T, Kodama M, Aizawa Y, Ohta Y, Ono T, Hasegawa G, Naito M, Nakajima T, Kamijo Y, Gonzalez FJ, and Aoyama T. Constitutive regulation of cardiac fatty acid metabolism through peroxisome proliferator-activated receptor alpha associated with age-dependent cardiac toxicity. *J Biol Chem* 275: 22293–22299, 2000.
 382. Welch JS, Ricote M, Akiyama TE, Gonzalez FJ, and Glass CK. PPARgamma and PPARdelta negatively regulate specific subsets of lipopolysaccharide and IFN-gamma target genes in macrophages. *Proc Natl Acad Sci U S A* 100: 6712–6717, 2003.
 383. Werner C, Kamani CH, Gensch C, Bohm M, and Laufs U. The peroxisome proliferator-activated receptor-gamma agonist pioglitazone increases number and function of endothelial progenitor cells in patients with coronary artery disease and normal glucose tolerance. *Diabetes* 56: 2609–2615, 2007.
 384. Wiley SE, Murphy AN, Ross SA, van der Geer P, and Dixon JE. MitoNEET is an iron-containing outer mitochondrial membrane protein that regulates oxidative capacity. *Proc Natl Acad Sci U S A* 104: 5318–5323, 2007.
 385. Wiley SE, Paddock ML, Abresch EC, Gross L, van der Geer P, Nechushtai R, Murphy AN, Jennings PA, and Dixon JE. The outer mitochondrial membrane protein mitoNEET contains a novel redox-active 2Fe-2S cluster. *J Biol Chem* 282: 23745–23749, 2007.
 386. Wright HM, Clish CB, Mikami T, Hauser S, Yanagi K, Hiramatsu R, Serhan CN, and Spiegelman BM. A synthetic antagonist for the peroxisome proliferator-activated receptor gamma inhibits adipocyte differentiation. *J Biol Chem* 275: 1873–1877, 2000.
 387. Xin X, Yang S, Kowalski J, and Gerritsen ME. Peroxisome proliferator-activated receptor gamma ligands are potent inhibitors of angiogenesis *in vitro* and *in vivo*. *J Biol Chem* 274: 9116–9121, 1999.
 388. Xiong C, Mou Y, Zhang J, Fu M, Chen YE, Akinbami MA, and Cui T. Impaired expression of PPAR gamma protein contributes to the exaggerated growth of vascular smooth muscle cells in spontaneously hypertensive rats. *Life Sci* 77: 3037–3048, 2005.
 389. Xu HE, Lambert MH, Montana VG, Parks DJ, Blanchard SG, Brown PJ, Sternbach DD, Lehmann JM, Wisely GB, Willson TM, Kliewer SA, and Milburn MV. Molecular recognition of fatty acids by peroxisome proliferator-activated receptors. *Mol Cell* 3: 397–403, 1999.
 390. Xu HE, Lambert MH, Montana VG, Plunket KD, Moore LB, Collins JL, Oplinger JA, Kliewer SA, Gampe RT Jr, McKee DD, Moore JT, and Willson TM. Structural determinants of ligand binding selectivity between the peroxisome proliferator-activated receptors. *Proc Natl Acad Sci U S A* 98: 13919–13924, 2001.
 391. Xu HE, Stanley TB, Montana VG, Lambert MH, Shearer BG, Cobb JE, McKee DD, Galardi CM, Plunket KD, Nolte RT, Parks DJ, Moore JT, Kliewer SA, Willson TM, and Stimmel JB. Structural basis for antagonist-mediated recruitment of nuclear co-repressors by PPARalpha. *Nature* 415: 813–817, 2002.
 392. Xu Y, Gen M, Lu L, Fox J, Weiss SO, Brown RD, Perlov D, Ahmad H, Zhu P, Greyson C, Long CS, and Schwartz GG. PPAR-gamma activation fails to provide myocardial protection in ischemia and reperfusion in pigs. *Am J Physiol Heart Circ Physiol* 288: H1314–H1323, 2005.
 393. Xu Y, Mayhugh D, Saeed A, Wang X, Thompson RC, Dominianni SJ, Kauffman RF, Singh J, Bean JS, Bensch WR, Barr RJ, Osborne J, Montrose-Rafizadeh C, Zink RW, Yumibe NP, Huang N, Luffer-Atlas D, Rungta D, Maise DE, and Mantlo NB. Design and synthesis of a potent and selective triazolone-based peroxisome proliferator-activated receptor alpha agonist. *J Med Chem* 46: 5121–5124, 2003.
 394. Yamamoto K, Ohki R, Lee RT, Ikeda U, and Shimada K. Peroxisome proliferator-activated receptor gamma activators inhibit cardiac hypertrophy in cardiac myocytes. *Circulation* 104: 1670–1675, 2001.
 395. Yang XY, Wang LH, Chen T, Hodge DR, Resau JH, DaSilva L, and Farrar WL. Activation of human T lymphocytes is inhibited by peroxisome proliferator-activated receptor gamma (PPARgamma) agonists: PPARgamma co-association with transcription factor NFAT. *J Biol Chem* 275: 4541–4544, 2000.
 396. Yang XY, Wang LH, Mihalic K, Xiao W, Chen T, Li P, Wahl LM, and Farrar WL. Interleukin (IL)-4 indirectly suppresses IL-2 production by human T lymphocytes via peroxisome proliferator-activated receptor gamma activated by macrophage-derived 12/15-lipoxygenase ligands. *J Biol Chem* 277: 3973–3978, 2002.
 397. Yoo J, Ghiassi M, Jirmanova L, Balliet AG, Hoffman B, Fornace AJ Jr, Liebermann DA, Bottinger EP, and Roberts AB. Transforming growth factor-beta-induced apoptosis is mediated by Smad-dependent expression of GADD45b through p38 activation. *J Biol Chem* 278: 43001–43007, 2003.
 398. Yoon CH, Hur J, Park KW, Kim JH, Lee CS, Oh IY, Kim TY, Cho HJ, Kang HJ, Chae IH, Yang HK, Oh BH, Park YB, and Kim HS. Synergistic neovascularization by mixed transplantation of early endothelial progenitor cells and late outgrowth endothelial cells: the role of angiogenic cytokines and matrix metalloproteinases. *Circulation* 112: 1618–1627, 2005.
 399. Young ME, McNulty P, and Taegtmeier H. Adaptation and maladaptation of the heart in diabetes, Part II: potential mechanisms. *Circulation* 105: 1861–1870, 2002.
 400. Young PW, Buckle DR, Cantello BC, Chapman H, Clapham JC, Coyle PJ, Haigh D, Hindley RM, Holder JC, Kallender H, Latter AJ, Lawrie KW, Mossakowska D, Murphy GJ, Roxbee Cox L, and Smith SA. Identification of high-affinity binding sites for the insulin sensitizer rosiglitazone (BRL-49653) in rodent and human adipocytes using a radioiodinated ligand for peroxisomal proliferator-activated receptor gamma. *J Pharmacol Exp Ther* 284: 751–759, 1998.

401. Yu C, Chen L, Luo H, Chen J, Cheng F, Gui C, Zhang R, Shen J, Chen K, Jiang H, and Shen X. Binding analyses between human PPAR γ -LBD and ligands. *Eur J Biochem FEBS* 271: 386–397, 2004.
402. Yue L, Ye F, Gui C, Luo H, Cai J, Shen J, Chen K, Shen X, and Jiang H. Ligand-binding regulation of LXR/RXR and LXR/PPAR heterodimerizations: SPR technology-based kinetic analysis correlated with molecular dynamics simulation. *Protein Sci* 14: 812–822, 2005.
403. Yue TL, Nerurkar SS, Bao W, Jucker BM, Sarov-Blat L, Steplewski K, Ohlstein EH, and Willette RN. *In vivo* activation of peroxisome proliferator-activated receptor- δ protects the heart from ischemia/reperfusion injury in Zucker fatty rats. *J Pharmacol Exp Ther* 325: 466–474, 2008.
404. Zahradka P, Yurkova N, Litchie B, Moon MC, Del Rizzo DF, and Taylor CG. Activation of peroxisome proliferator-activated receptors α and γ 1 inhibits human smooth muscle cell proliferation. *Mol Cell Biochem* 246: 105–110, 2003.
405. Zhan Q, Lord KA, Alamo I, Jr., Hollander MC, Carrier F, Ron D, Kohn KW, Hoffman B, Liebermann DA, and Fornace AJ Jr. The gadd and MyD genes define a novel set of mammalian genes encoding acidic proteins that synergistically suppress cell growth. *Mol Cell B* 14: 2361–2371, 1994.
406. Zhang C, Baker DL, Yasuda S, Makarova N, Balazs L, Johnson LR, Marathe GK, McIntyre TM, Xu Y, Prestwich GD, Byun HS, Bittman R, and Tigyi G. Lysophosphatidic acid induces neointima formation through PPAR γ activation. *J Exp Med* 199: 763–774, 2004.
407. Zhang F, Sowers JR, Ram JL, Standley PR, and Peuler JD. Effects of pioglitazone on calcium channels in vascular smooth muscle. *Hypertension* 24: 170–175, 1994.
408. Zhang H, Pi R, Li R, Wang P, Tang F, Zhou S, Gao J, Jiang J, Chen S, and Liu P. PPAR β / δ activation inhibits angiotensin II-induced collagen type I expression in rat cardiac fibroblasts. *Arch Biochem Biophys* 460: 25–32, 2007.
409. Zhang J, Fu M, Zhao L, and Chen YE. 15-Deoxy-prostaglandin J(2) inhibits PDGF-A and -B chain expression in human vascular endothelial cells independent of PPAR γ . *Biochem Biophys Res Commun* 298: 128–132, 2002.
410. Zhang J, Fu M, Zhu X, Xiao Y, Mou Y, Zheng H, Akinbami MA, Wang Q, and Chen YE. Peroxisome proliferator-activated receptor δ is up-regulated during vascular lesion formation and promotes post-confluent cell proliferation in vascular smooth muscle cells. *J B Chem* 277: 11505–11512, 2002.
411. Zhou G, Cummings R, Li Y, Mitra S, Wilkinson HA, Elbrecht A, Hermes JD, Schaeffer JM, Smith RG, and Moller DE. Nuclear receptors have distinct affinities for coactivators: characterization by fluorescence resonance energy transfer. *Mol Endocr (Baltimore)* 12: 1594–1604, 1998.
412. Ziouzenkova O, Perrey S, Asatryan L, Hwang J, MacNaull KL, Moller DE, Rader DJ, Sevanian A, Zechner R, Hoefler G, and Plutzky J. Lipolysis of triglyceride-rich lipoproteins generates PPAR ligands: evidence for an antiinflammatory role for lipoprotein lipase. *Proc Natl Acad Sci U S A* 100: 2730–2735, 2003.
413. Zungu M, Felix R, and Essop MF. Wy-14,643 and fenofibrate inhibit mitochondrial respiration in isolated rat cardiac mitochondria. *Mitochondrion* 6: 315–322, 2006.
414. Zuo X, Wu Y, Morris JS, Stimmel JB, Leesnitzer LM, Fischer SM, Lippman SM, and Shureiqi I. Oxidative metabolism of linoleic acid modulates PPAR- β / δ suppression of PPAR- γ activity. *Oncogene* 25: 1225–1241, 2006.

Address reprint requests to:
Y. Eugene Chen, M.D., Ph.D.

Cardiovascular Center
Department of Internal Medicine
University of Michigan Medical Center
1150 W. Medical Center Drive
Ann Arbor, MI 48109

E-mail: echenum@umich.edu

Date of first submission to ARS Central, September 4, 2008;
date of final revised submission, December 6, 2008; date of
acceptance, December 7, 2008.

This article has been cited by:

1. Wen-Hsuan Fong, Hsin-Da Tsai, Yu-Chang Chen, Jui-Sheng Wu, Teng-Nan Lin. 2010. Anti-apoptotic Actions of PPAR- γ Against Ischemic Stroke. *Molecular Neurobiology* **41**:2-3, 180-186. [[CrossRef](#)]