

Genetic analysis of petrobactin transport in *Bacillus anthracis*

Paul E. Carlson Jr,¹ Shandee D. Dixon,¹
Brian K. Janes,¹ Katherine A. Carr,¹
Tyler D. Nusca,^{1,2} Erica C. Anderson,¹
Sarra E. Keene,¹ David H. Sherman^{1,2} and
Philip C. Hanna^{1*}

¹Department of Microbiology and Immunology, and ²Life Sciences Institute and Departments of Medicinal Chemistry and Chemistry, University of Michigan Medical School, Ann Arbor, MI 48104, USA.

Summary

Iron acquisition mechanisms play an important role in the pathogenesis of many infectious microbes. In *Bacillus anthracis*, the siderophore petrobactin is required for both growth in iron-depleted conditions and for full virulence of the bacterium. Here we demonstrate the roles of two putative petrobactin binding proteins FatB and FpuA (encoded by GBAA5330 and GBAA4766 respectively) in *B. anthracis* iron acquisition and pathogenesis. Markerless deletion mutants were created using allelic exchange. The $\Delta fatB$ strain was capable of wild-type levels of growth in iron-depleted conditions, indicating that FatB does not play an essential role in petrobactin uptake. In contrast, $\Delta fpuA$ bacteria exhibited a significant decrease in growth under low-iron conditions when compared with wild-type bacteria. This mutant could not be rescued by the addition of exogenous purified petrobactin. Further examination of this strain demonstrated increased levels of petrobactin accumulation in the culture supernatants, suggesting no defect in siderophore synthesis or export but, instead, an inability of $\Delta fpuA$ to import this siderophore. $\Delta fpuA$ spores were also significantly attenuated in a murine model of inhalational anthrax. These results provide the first genetic evidence demonstrating the role of FpuA in petrobactin uptake.

Introduction

Bacillus anthracis, the causative agent of anthrax, is a Gram-positive, spore-forming bacterium that readily infects a variety of mammals. In the environment, *B. anthracis* exists primarily as dormant spores, which are the infectious form of this microorganism (Dixon *et al.*, 1999). Spores can enter a mammalian host through three distinct routes: cutaneous, gastrointestinal or inhalational, each leading to distinct symptoms and disease course (Dixon *et al.*, 1999; Inglesby *et al.*, 1999; 2002; Doganay and Welsby, 2006). Infection through the inhalational route leads to the most severe and potentially fatal form of the disease (Dixon *et al.*, 1999; Inglesby *et al.*, 1999; 2002; Doganay and Welsby, 2006). Upon entry into the host, spores are phagocytosed by resident macrophages and transported to the regional lymph nodes (Dutz and Kohout, 1971; Dutz and Kohout-Dutz, 1981; Guidi-rontani *et al.*, 1999; Dixon *et al.*, 2000). While in the intracellular environment, *B. anthracis* spores germinate, transforming into rapidly dividing vegetative cells (Dixon *et al.*, 2000). It is thought that these bacilli begin to multiply within macrophages prior to the induction of cell death and escape from these immune cells (Dixon *et al.*, 2000). Once out of the macrophages, *B. anthracis* vegetative cells continue to grow rapidly in the bloodstream. Although symptoms during these early stages are relatively mild, the bacteria soon replicate to very high levels, approaching 10^8 organisms per milliliter of blood. Production of two binary toxins by these bacteria can lead to serious complications, including hypotension, shock and eventually death (Dixon *et al.*, 1999).

Iron acquisition mechanisms are vital for the survival and pathogenesis of many infectious microbes, including *B. anthracis* (Ratledge and Dover, 2000; Faraldo-Gomez and Sansom, 2003; Wandersman and Delepelaire, 2004). In the mammalian host the concentration of free iron is limiting, with most of the iron sequestered into complexes such as transferrin and haemoglobin (Ratledge and Dover, 2000; Glanfield *et al.*, 2007). *B. anthracis*, and other pathogenic microbes, succeed in such environments by utilizing various iron acquisition methods including the production of siderophores and a variety of iron transport systems (Ratledge and Dover, 2000; Faraldo-Gomez and Sansom, 2003; Glanfield *et al.*, 2007). Siderophores are high-affinity




Fig. 1. Construction of *B. anthracis* mutants. Gene deletion mutants were created by allelic exchange removing either (A) GBAA4766 (*fpuA*) or (B) GBAA5330 (*fatB*). For each deletion, the initial 30 and final 30 nucleotides of the gene were fused creating a markerless deletion. Deleted genetic material was replaced by three stop codons, represented by 'X'.

iron binding molecules that are secreted by bacteria to sequester insoluble ferric iron from host complexes (Ratledge and Dover, 2000). *B. anthracis* produces two different siderophore molecules, bacillibactin and petrobactin (Cendrowski *et al.*, 2004; Koppisch *et al.*, 2005; Wilson *et al.*, 2006). Although the genes encoding bacillibactin biosynthesis are induced during iron starvation, this molecule is not required for growth of *B. anthracis* under iron-limiting conditions (Cendrowski *et al.*, 2004). *B. anthracis* strains deficient in bacillibactin biosynthesis are fully virulent in a murine model of anthrax infection (Cendrowski *et al.*, 2004). In contrast, petrobactin biosynthesis has been clearly associated with anthrax pathogenesis. Strains unable to synthesize this siderophore exhibit attenuated growth under iron-depleted conditions and are highly attenuated for murine virulence (Cendrowski *et al.*, 2004; Abergel *et al.*, 2006; Lee *et al.*, 2007; Pflieger *et al.*, 2007).

Once secreted siderophores bind extracellular iron they must be reacquired for the organism to take advantage of this system. Uptake of iron-bound siderophores typically occurs through specific bacterial membrane receptor complexes (Faraldo-Gomez and Sansom, 2003). The specific transport system involved in petrobactin uptake has yet to be identified. Recently, Zawadzka *et al.* reported the identification of two *B. cereus* proteins, FatB and FpuA, capable of binding multiple forms of petrobactin *in vitro* (Zawadzka *et al.*, 2009a). The *B. anthracis* homologues of these proteins are encoded by genes designated GBAA5330 and GBAA4766, and have high homology to the *B. cereus* proteins (95% and 99% amino acid identity respectively). Expression of these genes was also identified as significantly upregulated by *B. anthracis* during iron starvation (Carlson *et al.*, 2009). In the current study, we sought to elucidate the role of these two petrobactin binding proteins in the pathogenesis of *B. anthracis*.

Results

Bacillus anthracis $\Delta fpuA$ exhibits a growth defect in iron-depleted media

We sought to determine the role of two genes encoding homologues of the *B. cereus* petrobactin binding proteins FatB and FpuA (encoded by GBAA5330 and GBAA4766 respectively) in *B. anthracis* iron acquisition. *B. cereus* homologues of these proteins are able to bind petrobactin *in vitro* (Zawadzka *et al.*, 2009a); however, the function of these proteins in bacterial iron acquisition and pathogenesis remains to be elucidated. To accomplish this mutant *B. anthracis*, strains lacking either GBAA4766 ($\Delta fpuA$), GBAA5330 ($\Delta fatB$), or both genes ($\Delta fpuA \Delta fatB$) were created by markerless gene replacement (Fig. 1 and Table 1). PCR was used to screen all mutant strains for the expected gene deletions, as well as the presence of the virulence plasmid pXO1, prior to use in experiments (data not shown).

We hypothesized that if these proteins were involved in the uptake of petrobactin, mutants lacking the respective genes would exhibit significant growth defects in iron-depleted media (IDM), similar to a mutant that cannot synthesize petrobactin. To test this, vegetative bacilli of wild-type *B. anthracis*, $\Delta fpuA$, $\Delta fatB$ and the $\Delta fpuA \Delta fatB$ double mutant were inoculated into IDM at an initial OD₆₀₀

Table 1. *Bacillus anthracis* strains used in this study.

Strain	Relevant characteristics	Reference
Sterne 34F ₂	Wild type (pXO1 ⁺ , pXO2 ⁻)	Sterne (1939)
BA850	34F ₂ , $\Delta asbABCDEF$	Lee <i>et al.</i> (2007)
PC101	34F ₂ , $\Delta fatB$	This work
PC102	34F ₂ , $\Delta fpuA$	This work
PC103	34F ₂ , $\Delta fpuA \Delta fatB$	This work


Fig. 2. Growth of *B. anthracis* mutant strains in IDM.

A and B. Wild-type (solid diamonds), $\Delta fatB$ (solid squares), $\Delta fpuA$ (open triangles), $\Delta fpuA \Delta fatB$ (open squares) and $\Delta asbABCDEF$ (open circles) were grown in either IDM (A) or BHI (B). C. Wild-type (solid diamonds), $\Delta fpuA$ (open triangles), $\Delta fpuA/pfpuA+$ were grown in IDM at 26°C. All cultures were inoculated with vegetative bacilli at an initial $OD_{600} = 0.05$ and growth was monitored by measuring change in OD_{600} over time. Data presented are representative of at least three individual experiments.

of 0.05 and growth was monitored hourly by measurement of optical density at 600 nm. A mutant strain deficient for petrobactin biosynthesis ($\Delta asbABCDEF$) was also included as a control (Cendrowski *et al.*, 2004; Lee *et al.*, 2007). Cultures of wild-type *B. anthracis* are capable of rapid growth under these conditions, reaching high density within four hours (Fig. 2A, solid diamonds). As previously reported, the petrobactin-deficient mutant exhibited a growth defect in this medium, emphasizing the importance of this siderophore in the ability of *B. anthracis* to replicate in low-iron conditions (Fig. 2A, open circles) (Cendrowski *et al.*, 2004). The $\Delta fatB$ mutant grew as well as wild-type *B. anthracis* in IDM, suggesting that this gene is not required for growth in low-iron conditions (Fig. 2A, solid squares). In sharp contrast, both mutant strains lacking *fpuA* ($\Delta fpuA$ and $\Delta fpuA \Delta fatB$) exhibited severe growth defects in IDM, with each growing to levels nearly identical to the petrobactin-deficient strain (Fig. 2A). Introduction of the $\Delta fpuA$ mutant allele into the petrobactin biosynthesis-deficient strain ($\Delta asbABCDEF$) did not result in a stronger phenotype, further suggesting a role for FpuA in petrobactin-dependent iron acquisition (data not shown). It should also be noted that none of the gene deletions studied here resulted in a general growth defect and all mutant strains grew equally well in brain heart infusion (BHI) broth (Fig. 2B). Importantly, plasmid-based expression of *fpuA* driven by its native promoter resulted in complementation of $\Delta fpuA$ as demonstrated by increased growth in IDM (Fig. 2C). The complementation studies were performed at room temperature as the pBKJ236 vector cannot replicate at 37°C. These data indicate that FpuA, but not FatB, is necessary for *B. anthracis* growth in iron-depleted conditions and $\Delta fpuA$ mutants are unable to efficiently utilize iron at the low levels available in this medium.

ΔfpuA mutants are able to produce and secrete petrobactin

It was possible that the growth defect observed for the $\Delta fpuA$ mutant was due to an unexpected loss of the ability to produce or secrete petrobactin, rather than an inability to import the siderophore. As strains deficient in petrobactin production can be rescued by the addition of






Fig. 3. Addition of supplemental petrobactin does not rescue binding protein mutants. Wild type (solid diamonds), $\Delta fpuA$ (open triangles) and $\Delta asbABCDEF$ (open circles) were grown in either (A) IDM or (B) IDM supplemented with 2.5 μM petrobactin. Cultures were inoculated with spores and grown in microtitre plates. Data presented are representative of three individual experiments.

exogenous petrobactin to the media (Lee *et al.*, 2007), we sought to determine if the $\Delta fpuA$ mutants could also be complemented in the same manner. Wild-type, $\Delta fpuA$ and $\Delta asbABCDEF$ spores were inoculated into IDM with or without supplemental petrobactin (2.5 μM) and growth was monitored by increase in optical density (600 nm) over a 10 h time-course (Fig. 3). These experiments were performed in triplicate in 96-well plates due to the limited availability of purified petrobactin. All strains used in this study germinated with equal efficiencies (data not shown). As expected, wild-type spores exhibited significant growth in IDM with or without supplemental petrobactin. As pre-

viously observed, the $\Delta asbABCDEF$ strain, while unable to grow in IDM, grew at a rate similar to wild type in cultures supplemented with petrobactin. Conversely, the $\Delta fpuA$ mutant was not rescued by the addition of supplemental petrobactin. This finding is consistent with the hypothesis that this protein is required for the uptake of iron-bound petrobactin molecules.

If FpuA is indeed the petrobactin receptor, we hypothesized that the $\Delta fpuA$ mutant would exhibit increased levels of petrobactin in culture supernatants due to their inability to import this molecule. The Arnow assay was used to measure the concentration of total catechols in culture supernatants (Arnow, 1937). This colorimetric assay measures the presence of both petrobactin and its immediate biosynthetic precursor, 3,4-dihydroxybenzoic acid (DHB) (Cendrowski *et al.*, 2004; Garner *et al.*, 2004). Vegetative bacilli from wild-type, $\Delta fpuA$ and $\Delta asbABCDEF$ were inoculated into IDM, as described previously. Supernatants were isolated from each of the cultures hourly for 6 h post inoculation and analysed for catechol levels (Fig. 4A). No signal could be detected above background from supernatants isolated during the first 2 h of growth. Importantly, supernatants from $\Delta asbABCDEF$ cultures, which should not contain any petrobactin, did not exhibit a reaction in this assay at any of the time points tested. The lack of detectable signal in supernatants from these mutants suggests that, at least at these time points, petrobactin and 3,4-DHB are the only catechols in the media. Wild-type *B. anthracis* cultures did show an increase in catechol levels in supernatants over the time-course studied. The $\Delta fpuA$ mutant exhibited significantly higher levels of extracellular catechols, presumably petrobactin, at all time points when compared with wild-type bacteria (Fig. 4A).

As the Arnow assay is able to detect multiple molecules, we next sought to measure petrobactin levels directly from culture supernatants. Supernatants obtained from both wild-type and $\Delta fpuA$ mutant cultures grown for 6 h were normalized to equivalent OD₆₀₀ and examined using high-performance liquid chromatography (HPLC). Peaks corresponding in retention time (37.5 min), UV absorbance spectra (peaks at 260 and 290 nm), and mass spectra (m/z : $[M + 2H]^{2+} = 360$ and $[M + 2H - H_2O]^{2+} = 351$) to an authentic petrobactin standard were observed in both wild-type and $\Delta fpuA$ traces. HPLC peak integration revealed an approximate 65% increase of petrobactin accumulation per OD₆₀₀ in $\Delta fpuA$ supernatants over that observed in wild-type samples (Fig. 4B). Importantly, this peak was not observed in the $\Delta asbABCDEF$ strain which is unable to produce petrobactin (data not shown).

In order to examine the possibility that the altered petrobactin levels were due to increased production, we performed quantitative real-time PCR to determine levels of *asbA* expression after 6 h of growth in IDM. Loss of *fpuA* expression in either single or double mutant strains did




Fig. 4. Increased petrobactin levels observed in $\Delta fpuA$ strains.

A. Extracellular catechol levels were measured over time in culture supernatants using the Arnow assay. Data were normalized to the OD₆₀₀ of cultures at each time point and are presented as per cent of wild-type petrobactin levels at 3 h. No signal could be detected at time points prior to 3 h. Error bars represent standard deviation of triplicate samples within one experiment. Data are representative of four individual experiments.

B. Comparison of petrobactin accumulation by HPLC. Supernatants from wild-type and $\Delta fpuA$ cultures that were grown for 6 h and normalized to equivalent OD₆₀₀ were observed by HPLC. A peak corresponding to an authentic petrobactin standard is observed in both wild-type and $\Delta fpuA$ traces (arrow).

not lead to a significant increase in the transcription of this gene (\log_2 [fold change] = 0.6 ± 0.15). While it is more likely that the increase in petrobactin accumulation detected in $\Delta fpuA$ culture supernatants was due to the inability of this strain to import iron-bound petrobactin, we cannot exclude the possibility that this slight increase in transcription is responsible for this observation.

Gallium-associated growth inhibition requires FpuA

One test that has been used to show siderophore transport in other systems is gallium sensitivity. Gallium has been shown to bind to various siderophores, including petrobactin (Banin *et al.*, 2008; Zawadzka *et al.*, 2009b). Gallium is toxic to most bacterial cells and transport of siderophore–gallium complexes across the membrane often leads to cell death (Olanmi *et al.*, 2000; Banin *et al.*, 2008). It is therefore possible to correlate defects in siderophore transport with increased resistance to gallium. This system was used here to more definitively show a decrease in petrobactin in mutant strains. Wild-type, $\Delta asbABCDEF$ and $\Delta fpuA$ strains were grown in IDM with or without 20 μ M gallium sulphate. Growth was monitored by change in optical density (600 nm) over time. The addition of gallium to cultures of wild-type *B. anthracis* led to a significant decrease in growth over time, with a peak inhibition of 40% observed at 2 h (Table 2). As would be expected, growth of the petrobactin-deficient strain $\Delta as-$

bABCDEF was not significantly impaired by the addition of gallium to the culture media (Table 2). Finally, deletion of the putative petrobactin binding protein, FpuA, protects *B. anthracis* from gallium-induced growth inhibition, suggesting that this strain is defective in transport of gallium across the membrane. The fact that strains lacking the putative receptor have the same phenotype as those deficient in petrobactin biosynthesis supports a decreased transport of petrobactin in these mutants. These data provide further evidence that FpuA is required for transport of petrobactin into *B. anthracis*.

Role of FpuA in *B. anthracis* virulence

Because it is known that mutants unable to synthesize petrobactin exhibit significant attenuation in murine virulence (Cendrowski *et al.*, 2004; Pflieger *et al.*, 2008), we

Table 2. Resistance of $\Delta fpuA$ to gallium treatment.

	Hours			
	1	2	3	4
Wild type	25.0 ^{a,b}	40.2	33.4	27.5
$\Delta asbABCDEF$	4.6	6.0	8.7	5.0
$\Delta fpuA$	4.6	4.6	4.6	8.0

a. % growth inhibition.

b. Results are representative of four independent experiments.




Fig. 5. Attenuation of virulence of the $\Delta fpuA$ strain in a murine model of infection. DBA/2J mice were infected by intratracheal infection with wild-type (filled diamonds) or $\Delta fpuA$ (open triangles) spores at 1×10^5 (A) or 1×10^6 (B) spores per mouse. Mice were monitored for 14 days. Survival curves for $\Delta fpuA$ were significantly different from wild type at all doses tested by the log-rank test (P -values indicated above).

hypothesized that strains unable to import this siderophore would exhibit similar levels of attenuation. To test this hypothesis, we employed a murine model of inhalational anthrax. DBA/2J mice were used for these studies as these mice are known to be susceptible to *B. anthracis* Sterne strain (Welkos *et al.*, 1986). DBA/2J mice ($n = 8$) were inoculated intratracheally with wild-type, $\Delta fpuA$ or $\Delta asbABCDEF$ spores and monitored over the course of 14 days. When inoculated with 1.5×10^5 spores, mice receiving *B. anthracis* $\Delta fpuA$ exhibited increased survival compared with those infected with wild-type spores (Fig. 5). At this dose, all mice infected with wild-type spores succumbed to infection within 5 days post infection, while 87.5% of mice

infected with $\Delta fpuA$ ($P = 0.0006$) survived the full 2 week experiment, with the only death occurring on the first day post infection (Fig. 5A). In fact, even at a 10-fold higher dose (1.5×10^6 spores/mouse), the $\Delta fpuA$ mutant showed attenuated murine virulence compared with wild-type spores (Fig. 5B). At this dose, 75% of mice infected with $\Delta fpuA$ ($P = 0.01$) showed no sign of disease after 14 days, while all of the mice infected with wild-type spores succumbed to infection by day four (Fig. 5B). These results are similar to what was observed for the mutant strain defective in petrobactin biosynthesis (data not shown and Cendrowski *et al.*, 2004). In order to calculate an accurate LD_{50} for both the $\Delta fpuA$ and $\Delta asbABCDEF$ strains, mice were infected with 1.5×10^7 spores of each of the mutants. Even at this dose four of eight mice infected with $\Delta fpuA$ failed to show any sign of disease during the course of the infection. The LD_{50} of $\Delta fpuA$ was determined to be 2.2×10^7 spores/mouse, approximately three logs higher than that of wild-type spores (Table 3). Interestingly, when mice were infected with a similar dose of the petrobactin biosynthesis-deficient strain, $\Delta asbABCDEF$, 100% of mice succumbed to disease by day four post infection, indicating that $\Delta fpuA$ is slightly more attenuated than this strain ($P = 0.0002$). In fact, the $\Delta asbABCDEF$ mutant spores exhibited an LD_{50} nearly 10-fold lower than that of the $\Delta fpuA$ mutants (Table 3).

Discussion

The ability of a microorganism to survive in low-iron environments, such as a mammalian host, typically requires specialized systems of iron acquisition and uptake. Although *B. anthracis* encodes the biosynthesis machinery for two siderophores, only one of these, petrobactin, is required for efficient bacterial growth in low-iron conditions and for full virulence in a murine model of anthrax infection (Cendrowski *et al.*, 2004; Lee *et al.*, 2007). As this siderophore plays such an important role in *B. anthracis* pathogenesis, we sought to elucidate the role of two proteins recently implicated in petrobactin binding. We created deletion mutants lacking two putative siderophore binding proteins, FatB and FpuA, encoded by GBAA5330 and GBAA4766 respectively. *B. cereus* homologues of these proteins were recently shown to have the ability to bind to petrobactin; however, their role in the pathogenesis of the bacterium remained untested (Zawadzka *et al.*,

Table 3. Attenuated virulence of $\Delta fpuA$ strain in DBA/2J mice.

Strain	LD_{50}
Wild type	7.5×10^3
$\Delta fpuA$	2.2×10^7
$\Delta asbABCDEF$	2.7×10^6

2009a). Here we have demonstrated the role of these genes during growth in low-iron conditions as well as their effect on murine virulence.

Despite the ability of both FpuA and FatB to bind petrobactin *in vitro*, only FpuA appears important for growth of *B. anthracis* under iron-limiting conditions. Mutants lacking *fpuA* exhibit a severe growth defect when either vegetative bacteria or spores were inoculated into IDM (Figs 2 and 3A). These cultures also exhibit growth kinetics nearly identical to those of *B. anthracis* strains unable to synthesize petrobactin (Fig. 2, 3A and Cendrowski *et al.*, 2004; Lee *et al.*, 2007). Importantly, although vegetative bacilli of these strains ($\Delta asbABCDEF$ and $\Delta fpuA$) are capable of some growth in IDM (Fig. 2A), these strains are not capable of outgrowth from spores in this medium (Fig. 3). Although the growth kinetics were slightly slower than wild-type bacteria, replacement of this gene *in trans* did restore the ability of the bacteria to grow exponentially in IDM (Fig. 2C). It should be noted, a *B. anthracis* strain with deletions in both putative petrobactin binding proteins ($\Delta fpuA \Delta fatB$) grew to the same levels as $\Delta fpuA$ single mutants (Fig. 1 and data not shown). In contrast, mutants lacking FatB are not significantly impaired in their ability to grow in IDM (Fig. 2). In fact, growth curves for the $\Delta fatB$ mutant in IDM were nearly identical to those of wild-type bacilli (Fig. 2, filled squares). These data indicate that FatB does not play a role in uptake of petrobactin by *B. anthracis*. Curiously, the FatB homologue in *Bacillus subtilis* has recently been shown to be required for uptake of petrobactin (Zawadzka *et al.*, 2009b). It should be noted that *B. subtilis* does not make petrobactin, nor does it have an orthologue of *fpuA*. A role for this protein in iron metabolism in *B. cereus* remains to be tested.

Importantly, we have shown that $\Delta fpuA$ bacteria are not deficient in petrobactin synthesis or secretion, but rather the growth defect of this strain appears to be caused by an inability to use this siderophore. The addition of supplemental petrobactin had no effect on the growth of the $\Delta fpuA$ strain (Fig. 3B). Addition of excess siderophore is able to restore the growth of *B. anthracis* strains deficient in petrobactin biosynthesis, indicating that the $\Delta fpuA$ is phenotypically distinct from these strains. Petrobactin production from $\Delta fpuA$ bacteria was conclusively shown using both the Arnow assay and HPLC (Fig. 4). At the time points tested in the Arnow assay, no signal could be detected in the supernatants of strains unable to synthesize petrobactin, raising the possibility that bacillibactin was not being secreted at these time points. This finding is not surprising given the fact that most bacillibactin secretion has been seen following 24 h cultures and recently it was shown that bacillibactin secretion was undetectable until the bacteria had grown for 10 or more hours (Wilson *et al.*, 2006; 2009; Pflieger *et al.*, 2008). Although no catechols were detected using the Arnow assay on $\Delta asbABCDEF$ supernatants,

these molecules were observed from both wild-type bacteria and the $\Delta fpuA$ mutant. Also, petrobactin was specifically detected in both wild-type and $\Delta fpuA$ supernatants using HPLC. Interestingly, petrobactin levels were significantly higher in supernatants of $\Delta fpuA$ cultures than in those of wild-type *B. anthracis*. It is likely that the $\Delta fpuA$ mutant strain is synthesizing petrobactin at levels comparable to wild-type bacteria, as only a minimal increase was observed in the transcription of the petrobactin biosynthetic operon during growth of this strain in IDM. These results suggest that the increased accumulation of petrobactin in the supernatants of $\Delta fpuA$ cultures is due to the inability of these strains to import petrobactin, thus trapping the iron-bound siderophore in the extracellular environment and depriving the bacterium of the necessary iron. Strains lacking the putative petrobactin receptor also exhibit a resistance to gallium-induced growth inhibition. As gallium can be bound by petrobactin, these results further support the idea that the $\Delta fpuA$ mutant is deficient in petrobactin uptake. Taken together, the inability of supplemental petrobactin to restore the growth of $\Delta fpuA$ mutants, the increased accumulation of petrobactin in the $\Delta fpuA$ culture supernatants, the increased resistance of $\Delta fpuA$ to gallium, and the fact that the homologous protein from *B. cereus* binds petrobactin, clearly implicate FpuA in the uptake of iron-bound petrobactin.

Although it was clear that $\Delta fpuA$ strains were deficient for growth under iron-limiting conditions *in vitro*, it was also important to determine the role of this protein during infection. We have shown that the loss of a single putative petrobactin receptor in the $\Delta fpuA$ strain leads to severe attenuation in this infection model (Fig. 5). The LD₅₀ of the $\Delta fpuA$ mutant was found to be nearly 3000 times that of wild-type *B. anthracis* Sterne (Table 3). Similar to what was observed in the *in vitro* experiments, no significant difference was seen in the virulence of the double mutant ($\Delta fpuA \Delta fatB$) when compared with the mutant lacking only *fpuA* (data not shown). The level of attenuation observed for these mutants further emphasizes the importance of iron acquisition through petrobactin and its receptor complex in *B. anthracis* pathogenesis.

The observed attenuation of strains unable to import petrobactin is consistent with published reports on the attenuation of petrobactin biosynthesis mutants (Cendrowski *et al.*, 2004; Pflieger *et al.*, 2008). Although the growth phenotypes of $\Delta fpuA$ and $\Delta asbABCDEF$ strains appear very similar in IDM, slight differences in murine virulence were observed between these two strains. The virulence of these two mutant strains appears similar at lower doses, with neither strain killing a significant number of the mice tested. A difference in virulence was observed at higher doses however, and the LD₅₀ of $\Delta asbABCDEF$ mutants was found to be approximately eightfold lower than that of the $\Delta fpuA$ strain (Table 3). These results indi-

cate that $\Delta fpuA$ mutants are slightly less virulent than even the petrobactin biosynthesis mutant, leading to the hypothesis that this receptor protein plays a secondary function during mammalian infection. It is possible that this receptor is able to scavenge siderophore molecules produced by other bacterial species, providing the bacterium an increased survival advantage within its host. Uptake of exogenous siderophores has been shown in other bacterial species including *B. cereus* and *Pseudomonas aeruginosa* (Ollinger *et al.*, 2006; Greenwald *et al.*, 2009); however, the exact mechanism involved in *B. anthracis* remains to be elucidated.

The data presented here, combined with the fact that FpuA is able to bind petrobactin *in vitro* (Zawadzka *et al.*, 2009a), implicate this protein as a part of the petrobactin uptake system. FpuA has homology to substrate binding proteins from ABC family transporters (Miethke and Marahiel, 2007). In Gram-positive bacteria, these are membrane-bound proteins which interact with a transmembrane permease to transport molecules into the bacterium (Miethke and Marahiel, 2007). The identity of the remaining components of this ABC transport system remains to be elucidated. Although some candidate proteins have been identified (Zawadzka *et al.*, 2009a), further study is required to show a role for these proteins in petrobactin uptake.

Iron uptake systems are thought to be strong candidate targets for vaccine and therapeutic treatments, because they are often implicated in pathogen virulence and their surface localization leaves them exposed to extracellular molecules and the immune system (Glanfield *et al.*, 2007). Studies in *E. coli*, *Streptococcus pneumoniae* and *Strep-tococcus aureus* have shown that vaccines targeting iron uptake systems can improve immune responses and disease survival in animal infection models (Jomaa *et al.*, 2006; Kuklin *et al.*, 2006; Alteri *et al.*, 2009). It has been postulated that iron uptake proteins from *B. anthracis*, including the two examined here, could serve as vaccine targets (Gat *et al.*, 2006). Antibodies to these proteins were present in the serum of infected guinea pigs, indicating that they are expressed by the bacterium during infection (Gat *et al.*, 2006). As disruption of a single siderophore binding protein/iron uptake system in the $\Delta fpuA$ mutant leads to such dramatic phenotypes in *B. anthracis* growth and virulence it is likely that this protein could be a prime target for the development of new therapeutic measures. The data presented here indicate that a therapeutic blocking this receptor could greatly reduce the bacterium's ability to replicate within the low-iron environment of the human host. Such a treatment could, hypothetically, allow the host immune system to eliminate the pathogen before bacterial replication and toxin production reach lethal levels.

Given that recombinant forms of *B. cereus* FpuA have been shown to bind petrobactin directly, that these pro-

teins are nearly identical to their *B. anthracis* homologues, and the genetic evidence provided in this study, it is highly likely that FpuA serves as the receptor for petrobactin. In contrast, our data indicate that FatB does not play a role in petrobactin uptake in *B. anthracis*. Finally, the severe attenuation of $\Delta fpuA$ strains in a murine model of inhalation anthrax raises the possibility that this protein might be a useful target for the development of new vaccines and therapeutics.

Experimental procedures

Bacterial strains and growth conditions

All of the work described in this manuscript was performed using the Sterne 34F₂ (pXO1⁺, pXO2⁻) strain of *B. anthracis*. Initial spore stocks were prepared as follows. Strains were grown in modified G medium (Kim and Goepfert, 1974) for 3 days at 37°C with shaking and spores were prepared as previously described (Passalacqua *et al.*, 2006). Spores were stored at room temperature in sterile water and titered by haemocytometer (spores ml⁻¹) for animal studies. For all iron depletion studies, IDM was used (Cendrowski *et al.*, 2004). Experimental design for growth curves, Arnow assays and RNA isolation experiments were done as follows. Spores were germinated in BHI broth and incubated overnight with shaking, at room temperature. The following day, vegetative bacilli were diluted 1:100 in fresh BHI and grown 1–2 h at 37°C. Actively growing bacilli were then washed 3× with phosphate-buffered saline (Invitrogen) and 3× with IDM to ensure removal of nutrients and potential iron sources carried over from the BHI. Washed bacteria were then used to inoculate cultures of IDM or iron-replete media (IDM + 20 µM ferrous sulphate) at an OD₆₀₀ = 0.05. Experiments performed with petrobactin were started with 1 × 10⁵ spores per well in 96-well plates and performed in a Molecular Devices M2 plate reader. Growth was monitored by change in OD₆₀₀ over 10 h. For gallium-resistance experiments, strains were grown in IDM with or without 20 µM gallium sulphate and growth was monitored over time. Per cent growth inhibition was calculated as $\{[(OD_{600} \text{ IDM} + \text{gallium}) / (OD_{600} \text{ IDM})] * 100\}$.

Mutant construction

Mutant *B. anthracis* strains were constructed using allelic exchange as previously described (Janes and Stibitz, 2006). The $\Delta fpuA$ and $\Delta fatB$ single mutants were created in the wild-type Sterne 34F₂ background by deletion of either GBAA4766 or GBAA5330 respectively. The double mutant was created by deleting GBAA4766 in the $\Delta fatB$ single deletion strain. The $\Delta asbABCDEF$ *B. anthracis* strain used in these studies has been reported previously (Lee *et al.*, 2007). The sequence of all oligonucleotides used in creating and screening these mutant strains is available on request.

Complementation of gene deletions *fpuA* along with the upstream promoter region (120 bp) was cloned into the temperature-sensitive plasmid, pBKJ401 [a derivative of pBKJ236 (Janes and Stibitz, 2006)]. The resulting vector, *pFpuA+*, was then introduced to $\Delta fpuA$ by conjugal transfer.

Conjugates were selected by screening for erythromycin resistance. Because this vector cannot replicate at 37°C, all growth of the plasmid containing strain was performed at 25°C.

Real-time RT-PCR

cDNA synthesis was performed using Superscript III (Invitrogen) and 1 µg of total RNA. Real-time reactions were performed with a 1:500 final dilution of template cDNA. Primer sets were designed for genes indicated using Primer 3 (Rozen and Skaletsky, 2000) and reactions were carried out on a Bio-Rad ICycler real-time machine using SYBR Green (Bio-Rad). The *acoB* gene (GBAA2775) was used as the internal reference, as its expression was previously determined to remain unchanged during iron starvation (data not shown).

Measurement of petrobactin

Extracellular levels of catechols were measured using the Arnow assay (Arnow, 1937). Bacteria were grown in IDM and samples of cultures were removed at the indicated time points. Cultures were centrifuged and filtered using 0.2 µm filters to remove bacteria. Samples were mixed with equal volumes of 0.5 M HCl, nitrate-molybdate reagent (10% sodium nitrate and 10% sodium molybdate), and 1 N NaOH. Positive reactions produce a red colour and absorbance was determined at 515 nm. Samples were normalized to OD₆₀₀ of the original culture and data are presented as per cent of wild-type petrobactin levels at the 3 h time point.

For direct measurement of petrobactin, supernatants were obtained from three cultures of both wild-type Sterne and *ΔfpuA* following 6 h of growth in IDM. All cultures were diluted with IDM to equal OD₆₀₀ and sterile-filtered. Samples of individual culture supernatants were mixed with 9 vols of LCMS grade methanol and incubated at -20°C for 20 min. Methanol crashes were centrifuged at 14 000 *g* and 1600 µl of each supernatant was transferred to disposable 1.7 ml tubes for evaporation by vacuum centrifugation. Dried pellets were re-dissolved in 80 µl of ddH₂O and directly injected onto a Beckman Coulter System Gold for HPLC analysis. Peaks were separated using a C₁₈ analytical reverse-phase column (Waters XBridge C₁₈, 5 µm, 6 × 250 mm) at a rate of 1 ml min⁻¹ with the following solvent system: 5% methanol (MeOH) in water with 0.1% formic acid (FA) for 25 min then increased to 50% MeOH plus 0.1% FA over the next 28.5 min followed by a cleanup step with 100% MeOH plus 0.1% FA for the remaining 10 min. Mass spectrometric detection was performed by direct injection of a collected peak onto a Finnigan LTQ Linear Ion Trap (ThermoElectron) scanning in positive mode. An authentic standard of petrobactin was used for comparisons. The A₂₉₀ peaks corresponding to petrobactin were integrated using the area annotation feature included in the 32 Karat Software package (Beckman Coulter). Peak areas corresponding to petrobactin were normalized with the mean of wild-type sample values equalling 100% petrobactin production.

Murine infections

Intratracheal infections of DBA/2J mice (Jackson Laboratories) were performed as previously described (Heffernan

et al., 2007). Groups of eight mice were infected with either wild-type or mutant spores at a variety of doses ranging from 1 × 10⁵ to 1 × 10⁸ spores per mouse. Mice were monitored over a period of 14 days. LD₅₀ values were calculated using the Moving Average Interpolation program available at (<http://falkow.stanford.edu/whatwedo/software/software.html>) (Kim *et al.*, 2003).

Acknowledgements

The authors would like to thank Yousong Ding for assistance with mass spectrometry analysis. This work was sponsored by the NIH/NIAID Regional Center of Excellence for Bio-defense and Emerging Infectious Diseases Research (RCE) Program. The authors wish to acknowledge membership within and support from the Region V 'Great Lakes' RCE (NIH award 1-U54-AI-057053). This research was also supported by HHS contract N266200400059C-N01-AI-40059. P.E.C. received funding from training Grant T32-AI07528, 'Molecular Mechanisms of Microbial Pathogenesis'.

References

- Abergel, R.J., Wilson, M.K., Arceneaux, J.E., Hoette, T.M., Strong, R.K., Byers, B.R., *et al.* (2006) Anthrax pathogen evades the mammalian immune system through stealth siderophore production. *Proc Natl Acad Sci USA* **103**: 18499–18503.
- Alteri, C.J., Hagan, E.C., Sivick, K.E., Smith, S.N., and Mobley, H.L. (2009) Mucosal immunization with iron receptor antigens protects against urinary tract infection. *PLoS Pathog* **5**: e1000586.
- Arnow, L.E. (1937) Colorimetric determination of the components of 3, 4-dihydroxyphenylalanine-tyrosine mixtures. *J Biol Chem* **118**: 531–537.
- Banin, E., Lozinski, A., Brady, K.M., Berenshtein, E., Butterfield, P.W., Moshe, M., *et al.* (2008) The potential of desferrioxamine-gallium as an anti-Pseudomonas therapeutic agent. *Proc Natl Acad Sci USA* **105**: 16761–16766.
- Carlson, P.E., Jr, Carr, K.A., Janes, B.K., Anderson, E.C., and Hanna, P.C. (2009) Transcriptional profiling of *Bacillus anthracis* Sterne (34F₂) during iron starvation. *PLoS One* **4**: e6988.
- Cendrowski, S., MacArthur, W., and Hanna, P. (2004) *Bacillus anthracis* requires siderophore biosynthesis for growth in macrophages and mouse virulence. *Mol Microbiol* **51**: 407–417.
- Dixon, T.C., Meselson, M., Guillemin, J., and Hanna, P.C. (1999) Anthrax. *N Engl J Med* **341**: 815–826.
- Dixon, T.C., Fadl, A.A., Koehler, T.M., Swanson, J.A., and Hanna, P.C. (2000) Early *Bacillus anthracis*-macrophage interactions: intracellular survival and escape. *Cell Microbiol* **2**: 453–463.
- Doganay, L., and Welsby, P.D. (2006) Anthrax: a disease in waiting? *Postgrad Med J* **82**: 754–756.
- Dutz, W., and Kohout, E. (1971) Anthrax. *Pathol Annu* **6**: 209–248.
- Dutz, W., and Kohout-Dutz, E. (1981) Anthrax. *Int J Dermatol* **20**: 203–206.
- Faraldo-Gomez, J.D., and Sansom, M.S. (2003) Acquisition

- of siderophores in gram-negative bacteria. *Nat Rev Mol Cell Biol* **4**: 105–116.
- Garner, B.L., Arceneaux, J.E., and Byers, B.R. (2004) Temperature control of a 3,4-dihydroxybenzoate (protocatechuate)-based siderophore in *Bacillus anthracis*. *Curr Microbiol* **49**: 89–94.
- Gat, O., Grosfeld, H., Ariel, N., Inbar, I., Zaide, G., Broder, Y., *et al.* (2006) Search for *Bacillus anthracis* potential vaccine candidates by a functional genomic-serologic screen. *Infect Immun* **74**: 3987–4001.
- Glanfield, A., McManus, D.P., Anderson, G.J., and Jones, M.K. (2007) Pumping iron: a potential target for novel therapeutics against schistosomes. *Trends Parasitol* **23**: 583–588.
- Greenwald, J., Nader, M., Celia, H., Gruffaz, C., Geoffroy, V., Meyer, J.M., *et al.* (2009) FpvA bound to non-cognate pyoverdines: molecular basis of siderophore recognition by an iron transporter. *Mol Microbiol* **72**: 1246–1259.
- Guidi-Rontani, C., Weber-Levy, M., Labruyere, E., and Mock, M. (1999) Germination of *Bacillus anthracis* spores within alveolar macrophages. *Mol Microbiol* **31**: 9–17.
- Heffernan, B.J., Thomason, B., Herring-Palmer, A., and Hanna, P. (2007) *Bacillus anthracis* anthrolysin O and three phospholipases C are functionally redundant in a murine model of inhalation anthrax. *FEMS Microbiol Lett* **271**: 98–105.
- Inglesby, T.V., Henderson, D.A., Bartlett, J.G., Ascher, M.S., Eitzen, E., Friedlander, A.M., *et al.* (1999) Anthrax as a biological weapon: medical and public health management. Working Group on Civilian Biodefense. *JAMA* **281**: 1735–1745.
- Inglesby, T.V., O'Toole, T., Henderson, D.A., Bartlett, J.G., Ascher, M.S., Eitzen, E., *et al.* (2002) Anthrax as a biological weapon, 2002: updated recommendations for management. *JAMA* **287**: 2236–2252.
- Janes, B.K., and Stibitz, S. (2006) Routine markerless gene replacement in *Bacillus anthracis*. *Infect Immun* **74**: 1949–1953.
- Jomaa, M., Terry, S., Hale, C., Jones, C., Dougan, G., and Brown, J. (2006) Immunization with the iron uptake ABC transporter proteins PiaA and PiuA prevents respiratory infection with *Streptococcus pneumoniae*. *Vaccine* **24**: 5133–5139.
- Kim, H.U., and Goepfert, J.M. (1974) A sporulation medium for *Bacillus anthracis*. *J Appl Bacteriol* **37**: 265–267.
- Kim, C.C., Monack, D., and Falkow, S. (2003) Modulation of virulence by two acidified nitrite-responsive loci of *Salmonella enterica* serovar Typhimurium. *Infect Immun* **71**: 3196–3205.
- Koppisch, A.T., Browder, C.C., Moe, A.L., Shelley, J.T., Kinkel, B.A., Hersman, L.E., *et al.* (2005) Petrobactin is the primary siderophore synthesized by *Bacillus anthracis* str. Sterne under conditions of iron starvation. *Biometals* **18**: 577–585.
- Kuklin, N.A., Clark, D.J., Secore, S., Cook, J., Cope, L.D., McNeely, T., *et al.* (2006) A novel *Staphylococcus aureus* vaccine: iron surface determinant B induces rapid antibody responses in rhesus macaques and specific increased survival in a murine *S. aureus* sepsis model. *Infect Immun* **74**: 2215–2223.
- Lee, J.Y., Janes, B.K., Passalacqua, K.D., Pflieger, B.F., Bergman, N.H., Liu, H., *et al.* (2007) Biosynthetic analysis of the petrobactin siderophore pathway from *Bacillus anthracis*. *J Bacteriol* **189**: 1698–1710.
- Miethke, M., and Marahiel, M.A. (2007) Siderophore-based iron acquisition and pathogen control. *Microbiol Mol Biol Rev* **71**: 413–451.
- Olakanmi, O., Britigan, B.E., and Schlesinger, L.S. (2000) Gallium disrupts iron metabolism of mycobacteria residing within human macrophages. *Infect Immun* **68**: 5619–5627.
- Ollinger, J., Song, K.B., Antelmann, H., Hecker, M., and Helmann, J.D. (2006) Role of the Fur regulon in iron transport in *Bacillus subtilis*. *J Bacteriol* **188**: 3664–3673.
- Passalacqua, K.D., Bergman, N.H., Herring-Palmer, A., and Hanna, P. (2006) The superoxide dismutases of *Bacillus anthracis* do not cooperatively protect against endogenous superoxide stress. *J Bacteriol* **188**: 3837–3848.
- Pflieger, B.F., Lee, J.Y., Somu, R.V., Aldrich, C.C., Hanna, P.C., and Sherman, D.H. (2007) Characterization and analysis of early enzymes for petrobactin biosynthesis in *Bacillus anthracis*. *Biochemistry* **46**: 4147–4157.
- Pflieger, B.F., Kim, Y., Nusca, T.D., Maltseva, N., Lee, J.Y., Rath, C.M., *et al.* (2008) Structural and functional analysis of AsbF: origin of the stealth 3,4-dihydroxybenzoic acid subunit for petrobactin biosynthesis. *Proc Natl Acad Sci USA* **105**: 17133–17138.
- Ratledge, C., and Dover, L.G. (2000) Iron metabolism in pathogenic bacteria. *Annu Rev Microbiol* **54**: 881–941.
- Rozen, S., and Skaletsky, H.J. (2000) Primer3 on the WWW for general users and for biologist programmers. In *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Krawetz, S., and Misener, S. (eds). Totowa, NJ: Humana Press, pp. 365–386.
- Sterne, M. (1939) The immunization of laboratory animals against anthrax. Onderstepoort. *J Vet Sci Anim Ind* **13**: 313–317.
- Wandersman, C., and Delepelaire, P. (2004) Bacterial iron sources: from siderophores to hemophores. *Annu Rev Microbiol* **58**: 611–647.
- Welkos, S.L., Keener, T.J., and Gibbs, P.H. (1986) Differences in susceptibility of inbred mice to *Bacillus anthracis*. *Infect Immun* **51**: 795–800.
- Wilson, M.K., Abergel, R.J., Raymond, K.N., Arceneaux, J.E., and Byers, B.R. (2006) Siderophores of *Bacillus anthracis*, *Bacillus cereus*, and *Bacillus thuringiensis*. *Biochem Biophys Res Commun* **348**: 320–325.
- Wilson, M.K., Abergel, R.J., Arceneaux, J.E., Raymond, K.N., and Byers, B.R. (2009) Temporal production of the two *Bacillus anthracis* siderophores, petrobactin and bacillibactin. *Biometals* **23**: 129–134.
- Zawadzka, A.M., Abergel, R.J., Nichiporuk, R., Andersen, U.N., and Raymond, K.N. (2009a) Siderophore-mediated iron acquisition systems in *Bacillus cereus*: identification of receptors for anthrax virulence-associated petrobactin. *Biochemistry* **48**: 3645–3657.
- Zawadzka, A.M., Kim, Y., Maltseva, N., Nichiporuk, R., Fan, Y., Joachimiak, A., *et al.* (2009b) Characterization of a *Bacillus subtilis* transporter for petrobactin, an anthrax stealth siderophore. *Proc Natl Acad Sci USA* **106**: 21854–21859.