

Role of *Escherichia coli* YbeY, a highly conserved protein, in rRNA processing

Bryan W. Davies,¹ Caroline Köhrer,¹ Asha I. Jacob,^{1†}
Lyle A. Simmons,^{1,2†} Jianyu Zhu,³
Lourdes M. Aleman,¹ Uttam L. RajBhandary¹ and
Graham C. Walker^{1*}

¹Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139, USA.

²Department of Molecular, Cellular, and Developmental Biology, University of Michigan, Ann Arbor, MI, USA.

³Department of Molecular, Cell and Developmental Biology and Center for Molecular Biology of RNA, University of California at Santa Cruz, Santa Cruz, CA, USA.

Summary

The UPF0054 protein family is highly conserved with homologues present in nearly every sequenced bacterium. In some bacteria, the respective gene is essential, while in others its loss results in a highly pleiotropic phenotype. Despite detailed structural studies, a cellular role for this protein family has remained unknown. We report here that deletion of the *Escherichia coli* homologue, YbeY, causes striking defects that affect ribosome activity, translational fidelity and ribosome assembly. Mapping of 16S, 23S and 5S rRNA termini reveals that YbeY influences the maturation of all three rRNAs, with a particularly strong effect on maturation at both the 5'- and 3'-ends of 16S rRNA as well as maturation of the 5'-termini of 23S and 5S rRNAs. Furthermore, we demonstrate strong genetic interactions between *ybeY* and *rnc* (encoding RNase III), *ybeY* and *rnr* (encoding RNase R), and *ybeY* and *pnp* (encoding PNPase), further suggesting a role for YbeY in rRNA maturation. Mutation of highly conserved amino acids in YbeY, allowed the identification of two residues (H114, R59) that were found to have a significant effect *in vivo*. We discuss the implications of these findings for rRNA maturation and ribosome assembly in bacteria.

Introduction

Ribosome maturation and assembly occur in a cooperative and ordered fashion (Noller and Nomura, 1987; Kaczanowska and Ryden-Aulin, 2007). In bacteria, a 50S ribosomal subunit associates with a 30S ribosomal subunit to form an active 70S ribosome. The 50S ribosomal subunit is comprised of 23S and 5S rRNAs and 33 ribosomal proteins, while the 30S ribosomal subunit is comprised of 16S rRNA and 21 ribosomal proteins (Wilson and Nierhaus, 2007). 16S, 23S and 5S rRNAs are cotranscribed as part of a large rRNA precursor. Before transcription is complete, ribosomal proteins associate with rRNA to form ribonucleoprotein complexes that are acted on by RNase III (Robertson *et al.*, 1967). RNase III cleaves the initial transcript into 17S, 25S and 9S rRNA precursors that undergo further processing at their 5'- and 3'-termini to yield mature 16S, 23S and 5S rRNAs respectively (Gegenheimer *et al.*, 1977; Nierhaus, 1991).

Our understanding of the steps by which rRNA precursors are processed to their mature forms is still incomplete. In *Escherichia coli*, RNase G and RNase E are required for the maturation of the 5'-terminus of 16S rRNA (Li *et al.*, 1999a). RNase T is required for maturation of the 3'-terminus of 23S rRNA (Li *et al.*, 1999b). RNase E is required for partial maturation of the 5'-terminus of 5S rRNA (Misra and Apirion, 1979), and RNase E and RNase T are required for the maturation of the 3'-terminus of 5S rRNA (Misra and Apirion, 1979; Li and Deutscher, 1995). RNases responsible for the final step(s) in maturation of the 3'-terminus of 16S rRNA, 5'-terminus of 23S rRNA and 5'-terminus of 5S rRNA remain unknown (Deutscher, 2009) (Fig. 1).

Reconstitution of active 30S and 50S ribosomal subunits has been performed *in vitro* using mature rRNAs and ribosomal proteins (Nomura and Erdmann, 1970; Wireman and Sypherd, 1974; Nierhaus, 1980; Green and Noller, 1999). However, the conditions required for *in vitro* reconstitution are far from physiological and it is well recognized that many additional proteins, which include processing, modification and assembly factors, are required for rRNA maturation and eventual 50S, 30S and 70S ribosome formation *in vivo*. Several of these factors, such as Era and ObgE, are essential for cell viability (Wilson

Accepted 13 August, 2010. *For correspondence. E-mail gwalker@mit.edu; Tel. (+ 617) 253 6711; Fax (+ 617) 253 2643. †These authors contributed equally to this work.




Fig. 1. Steps in the processing of rRNA in *E. coli*. All enzymes indicated are endonucleases with the exception of RNase T that generates mature 3'-termini by trimming the rRNA precursors.

and Nierhaus, 2007). While these accessory factors are clearly important, in many cases we lack a good understanding of their role in ribosome biogenesis (Bunner *et al.*, 2010).

The UPF0054 protein family is highly conserved among prokaryotes. A gene encoding a member of this protein family is found in almost all sequenced bacterial genomes and it is one of the 206 genes that comprise the predicted minimal bacterial genome set (Gil *et al.*, 2004). In some bacteria including *Bacillus subtilis* (Kobayashi *et al.*, 2003), *Haemophilus influenzae* (Akerley *et al.*, 2002) and *Vibrio cholerae* (B.W. Davies *et al.*, in preparation), the respective homologue is essential. In the plant symbiont, *Sinorhizobium meliloti*, the respective homologue is not essential, but loss of its activity results in extreme pleiotropy and an inability of *S. meliloti* to form a symbiosis with plant hosts (Davies and Walker, 2008).

Here, we show that loss of the *E. coli* homologue, YbeY, results in viable cells with a pleiotropic phenotype and striking defects in ribosome function including decreased ribosome activity, reduced translational fidelity and altered translation initiation factor binding. Furthermore, we demonstrate that deletion of *ybeY* strongly affects the maturation of all three rRNAs causing a particularly strong defect in maturation of the 16S rRNA 5'- and 3'-termini, as well as maturation of the 5'-termini of 23S and 5S rRNAs. Strong genetic interactions between *ybeY* and *rnc* (encoding RNase III), *ybeY* and *rnr* (encoding RNase R), and *ybeY* and *pnp* (encoding PNPase) additionally suggest a requirement for YbeY activity in rRNA maturation. We discuss the implications

of these findings for rRNA maturation and ribosome assembly in bacteria.

Results

UPF0054 protein family members in bacteria are functionally equivalent

Disruption of the gene encoding the *S. meliloti* UPF0054 protein family member, *SMc01113*, results in extreme pleiotropy (Davies and Walker, 2008). Deletion of the *E. coli* UPF0054 family member gene, *ybeY*, from the reference strain MC4100 similarly results in an extremely pleiotropic phenotype that includes a modest decrease in growth rate in rich medium (Fig. 2A), but a significant sensitivity to numerous physiologically diverse stresses including β -lactam antibiotics, temperature, detergents and oxidative stress (Fig. 2B and C; Fig. S1). Deletion of *ybeY* from a second reference *E. coli* strain, MG1655, resulted in the same phenotypes (data not shown), indicating that the effects are not strain-specific. All $\Delta ybeY$ mutant phenotypes were rescued by ectopic expression of *ybeY* (Fig. 2A–C; Fig. S1). Expression of either the *S. meliloti* homologue *SMc01113* or the *B. subtilis* homologue *yqfG* rescues the $\Delta ybeY$ mutant phenotypes as effectively as *ybeY* itself (Fig. 2A–C; Fig. S1). Conversely, expression of *ybeY* in the *S. meliloti* *SMc01113* mutant background rescues all free-living and symbiotic phenotypes of the *SMc01113* mutant (Fig. S2A–C and data not shown) (Jones *et al.*, 2007; Davies and Walker, 2008). Taken together, these results demonstrate a universally conserved function for UPF0054 members in bacteria.




Fig. 2. Phenotypic analysis of the *E. coli* *ΔybeY* mutant.

A–C. Growth curves of MC4100 (WT) and the *ΔybeY* mutant complemented strains in LB at 37°C. Doubling times: 40 ± 2 min (*ΔybeY* mutant) vs. 28 ± 3 min (MC4100). Sensitivity of the *ΔybeY* mutant to stresses (B) cefotaxime and (C) temperature. The *ΔybeY* mutant with empty vector only (*ΔybeY*+vector) is shown on each plot for clarity. UPF0054 homologues: *ybeY* (*E. coli*), *yqfG* (*B. subtilis*) and *SMc01113* (*S. meliloti*). WT+vector (■), *ΔybeY*+vector (●), *ΔybeY*+*pybeY* (▲), *ΔybeY*+*pyqfG* (▼) and *ΔybeY*+*pSMc01113* (◄). 'p' indicates that the gene indicated is expressed from a plasmid. In (C), the pairs of samples show a 10-fold dilution each.

D. Polysome profiles for MC4100 and the *ΔybeY* mutant. The positions of polysomes, 70S ribosomes and 50S and 30S ribosomal subunits are indicated.

E–G. *In vitro* translation assay under saturating substrate conditions. MC4100 S100 fractions were mixed with a polyU template and equal amounts of MC4100 or *ΔybeY* mutant 70S ribosomes in an *in vitro* translation reaction as described in the *Experimental procedures*. Translational activity is normalized to MC4100 70S ribosome reactions. MC4100 and the *ΔybeY* mutant were transformed with plasmids expressing *lacZ* containing (F) frameshift mutations (+1 or –1) or (G) nonsense codons. *LacZ* activity was assayed as described in the *Experimental procedures*. The per cent *LacZ* activity (in Miller units) of each mutant *lacZ* allele relative to the wild-type *lacZ* allele for MC4100 or the *ΔybeY* mutant is reported. The value of *LacZ* activity from the wild-type allele (18921 Miller units in the *ΔybeY* mutant and 12247 Miller units in MC4100, respectively) was set as 100% activity. For clarity, wild-type *LacZ* activity has been omitted from the plots. Each assay was performed in triplicate.

H. Immunoblots identifying IF2 and IF3 in whole cell lysates, 30S, 50S and 70S ribosome fractions from MC4100 and the *ΔybeY* mutant. Immunoblotting for OmpA is used as a loading control. Equal A₂₆₀ amounts were loaded for the 30S, 50S and 70S ribosome fractions. The experiment was repeated three times and a representative result is shown.

Deletion of ybeY causes defects in polysome formation, ribosome activity, translational fidelity and translation initiation factor binding

Compared with the polysome profile of the parental strain, the $\Delta ybeY$ mutant has a decrease in polysomes and a large increase in both free 50S and free 30S ribosomal subunits relative to 70S ribosomes (Fig. 2D). Ectopic expression of *ybeY* in the $\Delta ybeY$ mutant restored the normal polysome profile (Fig. S3A–C). After we had completed much of this work, a study investigating the role of YbeY in the *E. coli* heat shock response also reported that deletion of *ybeY* caused loss of polysomes (Rasouly *et al.*, 2009); however, there was no mention of a potential function for YbeY in ribosome maturation.

The decreased polysome population led us to examine the functional state of the intact ribosomes in the $\Delta ybeY$ mutant. We reconstituted an *in vitro* translation system with an S100 extract from the wild-type strain MC4100 and 70S ribosomes from either the $\Delta ybeY$ mutant or its parent MC4100. Under saturating substrate conditions, ribosomes from the $\Delta ybeY$ mutant had reduced translational activity compared with 70S ribosomes from MC4100 (Fig. 2E). Furthermore, both frameshifting and readthrough of nonsense codons were increased *in vivo* in the $\Delta ybeY$ mutant, with the largest effect being for +1 frameshifts (Fig. 2F and G). The accumulation of defects in ribosomal function and polysome assembly, could account for the pleiotropic nature of the $\Delta ybeY$ mutant as many stress response programmes rely on the upregulation of defence proteins to ward off stress or to repair damage (Imlay and Linn, 1987; Friedberg *et al.*, 2005).

The increased pools of free 30S and 50S ribosomal subunits found in the $\Delta ybeY$ mutant suggest that a portion of these ribosomes may be defective in assembly of the 70S ribosome. In bacteria, translation initiation requires three protein initiation factors; IF1, IF2 and IF3. IF2 correctly positions the initiator tRNA in the ribosomal P site and promotes association of 30S and 50S ribosomal subunits. While IF1 is also known to stabilize the binding of the initiator tRNA, IF3 acts as a proofreading and anti-association factor (Kaczanowska and Ryden-Aulin, 2007). Analysis of total cell lysates showed only a slight but reproducible decrease in IF2 and an increase in IF3 in the $\Delta ybeY$ mutant (Fig. 2H). However, we found a substantial decrease in IF2 present in the fraction containing 30S subunits isolated from the $\Delta ybeY$ mutant, suggesting that many of the mutant 30S ribosomal subunits are not available for 70S ribosome assembly. We also observed a slight increase in IF3 associated with $\Delta ybeY$ mutant 30S subunits and 70S ribosomes, but the change was modest compared with IF2. This suggests that the 30S ribosomal

subunits in a $\Delta ybeY$ mutant have a defect that prevents ribosome assembly and/or promotes disassembly of 70S ribosomes when assembly is attempted.

The $\Delta ybeY$ mutant is defective in rRNA maturation

The ribosome is primarily composed of rRNA (Wilson and Nierhaus, 2007) and incomplete processing of rRNA can lead to defects in ribosome function (Wireman and Sypherd, 1974). Comparison of total rRNA profiles by agarose gel electrophoresis from MC4100 and the $\Delta ybeY$ mutant showed the presence of a substantial amount of 17S rRNA in the $\Delta ybeY$ mutant, as well as a faster migrating species, which we annotate as 16S*, indicating a defect in 16S rRNA maturation (Fig. 3A). 17S rRNA is the precursor of 16S rRNA (Fig. 1) and is generated from the initial rRNA transcript by RNase III cleavage (Gegenheimer *et al.*, 1977). The aberrant 16S* rRNA species may be a product of misprocessing and/or degradation of the 17S or 16S rRNA. Ectopic expression of *ybeY* in the $\Delta ybeY$ mutant restored normal 16S rRNA maturation (Fig. S3D). We were unable to assess the state of 5S and 23S rRNA processing in these total rRNA profile experiments (see below, however) because the difference between mature and immature 5S and 23S rRNAs is only a few nucleotides, respectively, and thus does not cause a detectable shift on agarose gels.

In strain MC4100, 30S ribosomal subunits contained mostly 16S rRNA along with some 17S rRNA, whereas 70S ribosomes contained only 16S rRNA (Fig. 3A). In contrast, 30S ribosomal subunits from the $\Delta ybeY$ mutant contained much more 17S rRNA than 16S rRNA as well as a substantial amount of 16S* rRNA (Fig. 3A). Furthermore, there was also a substantial amount of 17S rRNA in 70S ribosomes isolated from the $\Delta ybeY$ mutant (Fig. 3A). The accumulation of 17S rRNA was not due to the slower growth rate of the $\Delta ybeY$ mutant. When the strains were grown in minimal medium where MC4100 and the $\Delta ybeY$ mutant show similar doubling times, we still observed similar levels of 17S rRNA accumulation (data not shown). 17S rRNA is thought not to be competent for translation *in vitro* (Wireman and Sypherd, 1974). The increased abundance of 17S and 16S* rRNA in the 30S ribosomal subunits could account for altered translation initiation factor binding and the other translational defects described above for the $\Delta ybeY$ mutant. Similarly, the increased abundance of 17S rRNA in 70S ribosomes could explain, at least in part, the reduced translation activity and decreased translational accuracy of the $\Delta ybeY$ 70S ribosomes. Interestingly, 16S* rRNA was not found in $\Delta ybeY$ 70S ribosomes suggesting that it is not competent for assembly into the 70S ribosome and may thus be a non-functional product of rRNA processing in the $\Delta ybeY$ mutant.




Fig. 3. Analysis of rRNA from *E. coli* MC4100 and the $\Delta ybeY$ mutant.

A. Total RNA isolated from whole cells, 30S ribosomal subunits and 70S ribosomes from MC4100 and the $\Delta ybeY$ mutant. The positions of 23S, 17S, 16S and 16S* rRNAs are indicated based on their mobility.

B. Northern blot analysis using probes directed against the 5'- and 3'-termini of 17S rRNA. Equal amounts of total RNA from MC4100 and the $\Delta ybeY$ mutant were used. The locations of the probes are shown in the diagram below the blots.

C and D. Primer extension and site-specific RNase H cleavage assays to map the 5'- and 3'-termini of 16S, 23S and 5S rRNAs from MC4100 and the $\Delta ybeY$ mutant. 'P' and 'M' indicate the location of bands corresponding to precursor and mature forms of the rRNA respectively. Annotation of the positions of the 5' and 3'-termini of 16S, 23S and 5S rRNAs, mature and precursor species, were based on previous observations (Li *et al.*, 1999a). Total RNA was prepared from MC4100 and the $\Delta ybeY$ mutant strains as described in the *Experimental procedures*.

In agreement with the observed increase in 17S rRNA in the $\Delta ybeY$ mutant, Northern blot analysis showed that both 5'- and 3'-termini of 17S rRNA were present at much higher levels in total RNA extracted from the $\Delta ybeY$ mutant than in total RNA from the parental strain MC4100 (Fig. 3B). Use of primer extension to map the 5'-terminus (Fig. 3C) and site-specific RNase H cleavage followed by Northern hybridization (Li *et al.*, 1999b) to map the 3'-terminus of 16S rRNA (Fig. 3D) showed that the termini of the mature and immature 16S rRNA in total RNA extracted from the $\Delta ybeY$ mutant and the parental strain MC4100 were identical. However, consistent with the above results, both immature 5'- and 3'-termini of 16S rRNA were present at higher levels in total RNA extracted from the $\Delta ybeY$ mutant (Fig. 3C and D). We did not observe either 5'- or 3'-termini shorter than that of mature 16S rRNA, although, if the 5'- or 3'-terminus of the 16S* species were extensively degraded, the assays used would not have detected them.

Primer extension and site-specific RNase H cleavage were also used to determine the maturation state of 23S and 5S rRNAs. Strikingly, along with 16S rRNA, the maturation of the 5'- and 3'-termini of both 23S and 5S rRNA were also affected, with all four termini showing increased amounts of the immature form in the $\Delta ybeY$ mutant (Fig. 3C and D), indicating that *ybeY* function is required for normal 5' and 3' processing of 16S, 23S and 5S rRNAs. Nevertheless, the most significant defects in the

$\Delta ybeY$ mutant appeared to be in the complete maturation of the 5'- and 3'-termini of 16S rRNA.

Rasouly *et al.* demonstrated a defect in the polysome profile of a $\Delta ybeY$ mutant after heat shock (Rasouly *et al.*, 2009). After shifting the cultures to 45°C for 1 h, we observed little mature 16S rRNA in the $\Delta ybeY$ mutant by agarose gel electrophoresis; instead we saw two bands that migrate near the 17S and 16S* rRNA positions (Fig. 3E). In contrast, there were no significant changes in the rRNA profile of MC4100 samples shifted to 45°C compared with those at 37°C. As 17S rRNA is not competent for translation (Wireman and Sypherd, 1974), and 16S* rRNA appears to be excluded from 70S ribosomes (Fig. 3A), the increase in 17S and 16S* rRNA-like species and the decrease in 16S rRNA in the $\Delta ybeY$ mutant following heat shock could explain the defect in polysome formation at elevated temperatures (Rasouly *et al.*, 2009) and the overall temperature sensitivity of the $\Delta ybeY$ mutant described above (Fig. 2C).

ybeY shows strong genetic interactions with *rnc* (RNase III), *rnr* (RNase R) and *pnp* (PNPase)

The serious defects in translation and rRNA maturation in the $\Delta ybeY$ mutant suggested that YbeY might be involved in rRNA metabolism. To explore the cellular role of YbeY, we compared the rRNA profile of the $\Delta ybeY$ mutant with that of several well-characterized *E. coli* RNase mutants




Fig. 4. Analysis of rRNA from the $\Delta ybeY$ mutant and seven well-characterized *E. coli* RNase mutant strains. The relevant genotype from which the rRNA was extracted is indicated under each lane. The parental strain MC4100 rRNA is shown in each case as a control. A and B. Agarose gel electrophoresis of total rRNA from single and double RNase mutant strains. The positions of 23S, 18S*, 17S, 16S and 16S* rRNAs are indicated. C–H. Primer extension and site-specific RNase H cleavage assays to map the 5'- and 3'-termini of 16S, 23S and 5S rRNAs from single and double RNase mutant strains. 'P' and 'M' indicate the position of bands corresponding to the precursor and mature form for each rRNA. Annotation of the positions of the 5'- and 3'-termini of 16S, 23S and 5S rRNAs, mature and precursor species, were based on previous observations (Li *et al.*, 1999a).

disrupted in the following genes: *cafA* (RNase G), *rnc* (RNase III), *rnr* (RNase R), *pnp* (PNPase), *rnd* (RNase D), *rnt* (RNase T) and *rph* (RNase PH) (Fig. 4A). We also transduced each of these RNase mutations into the $\Delta ybeY$ mutant background and examined the rRNA profiles of these double mutants (Fig. 4B). We then mapped the 5'- and 3'-termini of 16S, 23S and 5S rRNAs for each

strain using primer extension and site-specific RNase H cleavage assays (Fig. 4C–H) as described above.

Profiles of total rRNA showed that, of all the mutants tested, the $\Delta ybeY$ mutant had the greatest defect in maturation of 17S rRNA to 16S rRNA (Fig. 4A). Primer extension and site-specific RNase H cleavage assays confirmed this result showing high levels of both immature

5' and 3' 16S rRNA termini in the $\Delta ybeY$ mutant compared with the other RNase mutants (Fig. 4C and D). As previously described (Li *et al.*, 1999a), the $\Delta cafA$ mutant also showed severe defects in maturation of the 16S rRNA 5'-terminus. However, in contrast to the $\Delta ybeY$ mutant, which accumulates the full 17S rRNA 5' precursor (+ 115 nucleotides), the $\Delta cafA$ mutant accumulates the shorter + 66 nucleotides 5' precursor (Fig. 4C). Also, of all the mutants tested, only the $\Delta ybeY$ mutant showed accumulation of high levels of immature 16S rRNA 3'-terminus (Fig. 4D).

The introduction of either the *mnc* or *rnr* mutation into the $\Delta ybeY$ mutant resulted in major alterations in the gross rRNA profile of the $\Delta ybeY$ mutant (Fig. 4B). Strikingly, 17S rRNA was no longer present in the $\Delta ybeY \Delta rnc$ mutant, instead 16S* rRNA and a new rRNA species migrating more slowly than 17S rRNA accumulated. Primer extension and site-specific RNase H cleavage assays confirmed this new precursor, which we designate 18S* rRNA, as a 16S rRNA derivative with additional nucleotides on both the 5'- and 3'-termini of 17S rRNA (Fig. 4C and D). Gegenheimer *et al.* reported a transient 18S rRNA species in a Δrnc mutant (Gegenheimer *et al.*, 1977) that is matured to 16S rRNA and has a similar mobility as the new precursor species we observe. We have not mapped the termini of 18S* rRNA or the 18S species found by Gegenheimer *et al.* to the nucleotide level, but based on their relative mobility, we suggest that the 18S* rRNA observed in the $\Delta ybeY \Delta rnc$ mutant may be 18S rRNA. If 18S* and 18S rRNA are identical, our data suggest that the absence of YbeY influences the processing of 18S rRNA to 16S rRNA in a Δrnc mutant. Further characterization of the 18S* rRNA species in the $\Delta ybeY \Delta rnc$ mutant could be of much interest.

The $\Delta ybeY \Delta rnr$ mutant showed a substantial decrease in 16S rRNA and reciprocal increase in both 17S and 16S* rRNA (Fig. 4B). This result was confirmed by mapping of the 5'- and 3'-termini (Fig. 4C and D). Furthermore, primer extension of 16S rRNA 5'-termini showed an increase in a set of new rRNA species of minor intensity that migrated between 17S and 16S rRNAs. These new species may represent partially mature or non-functional 16S-like rRNA species. RNase R has been shown to function as a scavenging RNase that removes non-functional rRNA from the cell (Cheng and Deutscher, 2003). Loss of this function is consistent with the appearance of 16S rRNA intermediates observed by primer extension in the $\Delta ybeY \Delta rnr$ mutant. It also suggests that, in the absence of *ybeY*, the cell misprocesses large amounts of 17S rRNA generating abnormal species that RNase R normally helps to degrade.

Strikingly, we observed very little of the mature 16S rRNA 3'-terminus in either the $\Delta ybeY \Delta rnr$ mutant or the $\Delta ybeY \Delta pnp$ mutant rRNA (Fig. 4D). Deletion of either

rnr or *pnp* alone does not significantly affect the maturation of the 16S rRNA 3'-terminus (Fig. 4D). The RNase(s) required for maturation of the 16S rRNA 3'-terminus has not been identified (Deutscher, 2009). Our results suggest that PNPase and RNase R may act in maturation of the 16S rRNA 3'-terminus and their role may be modulated by YbeY. Interestingly, while both the $\Delta ybeY \Delta rnr$ mutant and the $\Delta ybeY \Delta pnp$ mutant affect maturation of the 16S rRNA 3'-terminus, only the $\Delta ybeY \Delta rnr$ mutant exhibits additional defects in 16S rRNA metabolism as shown by the appearance of extra aberrant rRNA species (Fig. 4B and C). This suggests that RNase R and YbeY interact in aspects of rRNA metabolism beyond 16S rRNA 3'-terminus maturation, such as quality control of rRNA.

Mapping of the 5'- and 3'-termini showed that deletion of *ybeY* also affected 5' maturation of 23S and 5S rRNAs (Fig. 4E and H). The $\Delta ybeY$ mutant had a greater effect in 23S and 5S rRNA maturation than nearly all of the RNase mutants tested. As expected, deletion of *mnc* abolishes formation of mature 23S rRNA 5'- and 3'-termini (King *et al.*, 1984). Also as expected, deletion of *rnt* greatly inhibits maturation of the 23S and 5S rRNA 3'-termini (Li *et al.*, 1999b). Interestingly, we have also found that in the $\Delta ybeY \Delta rnt$ mutant the 17S rRNA precursor was extended by one or two nucleotides at the 3'-terminus (Fig. 4D), suggesting the possible involvement of RNase T in trimming the 17S precursor at the 3'-terminus prior to complete 3' maturation of 16S rRNA. Deletion of *pnp* and *rph* also caused accumulation of 5'-immature 5S rRNA precursor implicating a role of these two RNases in this step of rRNA maturation (Fig. 4G).

RNase E is known to function in the maturation of the 5'-terminus of 16S rRNA (Li *et al.*, 1999a), and the 5'- and 3'-termini of 5S rRNA (Misra and Apirion, 1979; Li and Deutscher, 1995). RNase E is essential in *E. coli* and temperature-sensitive mutants have been used to explore its potential roles in rRNA maturation (Li *et al.*, 1999a). Using the *rne-1* temperature-sensitive RNase E allele, we constructed an *rne-1 \Delta ybeY::cat^R* double mutant and attempted to explore their genetic interaction in rRNA processing at the non-permissive temperature (45°C). The rRNA profile is shown (Fig. S3F); mapping of the 16S rRNA termini showed that RNase E contributes to the maturation of the 5'-terminus of 16S rRNA as expected (Li *et al.*, 1999a), but does not contribute significantly to the maturation of the 3'-terminus under the conditions used (data not shown). Unfortunately, it is difficult to interpret the results from the *rne-1 \Delta ybeY::cat^R* double mutant grown at the non-permissive temperature because of the strong temperature sensitivity of the $\Delta ybeY$ mutant by itself (Rasouly *et al.*, 2009; see also Fig. 2C and Fig. S3E). Because of this, effects of the *rne-1* mutation at 45°C are overshadowed by effects of the $\Delta ybeY$ deletion.




Fig. 5. Growth of MC4100, $\Delta ybeY$ mutant and several $\Delta ybeY$ double mutants in rich medium at 37°C. Most $\Delta ybeY$ double mutants did not show a growth defect (data not shown); the $\Delta ybeY \Delta pnp$ double mutant is shown as an example. In contrast, the $\Delta ybeY \Delta rnc$ and $\Delta ybeY \Delta rrn$ mutants showed a significant decrease in growth rate.

A temperature-independent approach for conditional *rnc* inactivation will be necessary for such experiments.

Growth rate analysis of the $\Delta ybeY$ double mutants

We next examined the growth behaviour in rich medium of single and double mutants described above. Further strengthening a connection between YbeY and RNase R, we observed the greatest effect in the $\Delta ybeY \Delta rrn$ mutant (Fig. 5). This mutant shows reduced growth rate in addition to an extended lag phase. Interestingly, while both the $\Delta ybeY \Delta rrn$ and the $\Delta ybeY \Delta pnp$ mutants affect 16S rRNA 3'-terminus maturation strongly, the $\Delta ybeY \Delta pnp$ grows identically to the $\Delta ybeY$ mutant, showing only a modest growth rate reduction. The change in growth behaviour of the $\Delta ybeY \Delta rrn$ mutant compared with the $\Delta ybeY$ mutant could be due to defects other than the incomplete maturation of the 16S rRNA, e.g. quality control of rRNA. Also, the significant accumulation of 16S* rRNA, a species that can be assembled in 30S ribosomal subunits, but not in functional 70S ribosomes (Fig. 3A) may be detrimental to the cell and be responsible for the growth behaviour of the $\Delta ybeY \Delta rrn$ mutant.

We also observed a decrease in growth rate in the $\Delta ybeY \Delta rnc$ mutant in rich medium (Fig. 5) compared with the respective single mutants, although this was less pronounced than for the $\Delta ybeY \Delta rrn$ mutant. Thus, the strong genetic interaction observed in the $\Delta ybeY \Delta rrn$ and $\Delta ybeY \Delta rnc$ mutants is manifested at both the rRNA processing and physiological levels.

Highly conserved amino acid residues required for YbeY activity in vivo

All UPF0054 family members contain a conserved H3XH5XH motif (Fig. 6A). The presence of this motif has led to the categorization of UPF0054 family members as putative metal-dependent hydrolases and, more specifically, by some as putative metal-dependent proteases due to the presence of a similar motif found in certain mammalian proteases (E.V. Koonin, pers. comm.; Tatusov *et al.*, 2001; Verma *et al.*, 2002). The recently determined structures of *E. coli* YbeY and two additional UPF0054 family members (Oganessian *et al.*, 2003; Yeh *et al.*, 2005; Zhan *et al.*, 2005) support a putative metal-dependent hydrolytic function; however, no substrate has been identified for this protein family despite extensive screening (Oganessian *et al.*, 2003).

To study the structure–function relationship of the conserved UPF0054 motif and other highly conserved amino acids in this protein family, we mutated several conserved amino acid residues in *ybeY* on our complementation plasmid to Ala (Fig. 6A). We then determined the ability of these mutant *ybeY* alleles to complement the growth and rRNA processing defects of the $\Delta ybeY$ mutant (Fig. 6B and C). The first His residue (H114) in the conserved H3XH5XH motif was required for partial recovery of growth of the $\Delta ybeY$ mutant at 45°C and to rescue rRNA processing defects of the $\Delta ybeY$ mutant, supporting the importance of this motif for YbeY function (Fig. 6B and C). Interestingly, changing the second and third His residues from this motif (H118, H124) to Ala did not have much of an effect on complementation. A highly conserved Arg residue (R59) was also found to be necessary for growth complementation. The loss of complementation by H114A and R59A alleles was not due to insufficient protein expression as plasmid-driven expression of all *ybeY* mutant alleles exceeded *ybeY* expression from the endogenous locus (Fig. S3G). It should be noted that some of the *ybeY* mutant alleles that did not show an effect on complementation when expressed from a multi-copy plasmid, may do so if they are expressed at levels of YbeY normally present in wild-type cells.

Discussion

We have shown that the functionally conserved UPF0054 protein family member, YbeY, influences rRNA maturation and ribosome formation in *E. coli*. Deletion of *ybeY* from *E. coli* results in defects in ribosome assembly and activity as well as in attenuation of 16S, 23S and 5S rRNA processing.

rRNA processing, ribosome maturation and translation are intertwined. Several reports have implicated partial ribosome assembly as a requirement for full rRNA




Fig. 6. A. Sequence alignment of UPF0054 homologues from bacteria and eukaryotes. Alignments were performed using T-coffee (Poiret *et al.*, 2003). The red bar underlines the conserved H3XH5XH motif that is used to classify members of this family. Red asterisks indicate amino acid residues that were analysed in this study by Ala mutagenesis.

B. Sensitivity of the $\Delta ybeY$ mutant strain expressing mutant *ybeY* alleles to high temperature. Strains were serially diluted (1:10), plated on LB plates and incubated at 45°C.

C. rRNA profiles of the $\Delta ybeY$ mutant expressing the mutant *ybeY* alleles H114A and R59A. The positions of 23S, 17S, 16S and 16S* rRNAs are indicated.

maturation (Mangiarotti *et al.*, 1974; 1975; Srivastava and Schlessinger, 1988) and, while deletion of specific RNases can lead to specific defects in rRNA maturation (Li and Deutscher, 1995; Li *et al.*, 1999a,b), loss of non-RNase ribosome maturation factors can also lead to rRNA maturation defects. For example, both Era and ObgE are GTPases required for ribosome assembly. Era and ObgE are essential for viability in *E. coli* and loss of either activity causes accumulation of 17S rRNA (Inoue *et al.*, 2003; Sato *et al.*, 2005). Similarly, deletion of ribosome maturation factors RimB and RimP results in slower growth rates, reduced number of polysomes and accumu-

lation of 16S rRNA precursor (Lövgren *et al.*, 2004). Furthermore, loss of RsgA, a GTPase activated by the 30S ribosomal subunit, results in the accumulation of both 17S rRNA and another species shorter than 16S rRNA (Himeno *et al.*, 2004). Our results could also mean that loss of YbeY affects a critical step in ribosome biogenesis and thereby slows rRNA maturation. It will be interesting to explore physical interactions of YbeY with the ribosome and rRNA directly as well as possible effects of YbeY on ribosomal proteins, e.g. protein modifications. Supporting a potential association of YbeY with the ribosome, recent work has identified interactions between YbeY and both

30S and 50S ribosomal proteins in a large-scale pull-down experiment (Arifuzzaman *et al.*, 2006).

Active translation has also been suggested to be required for certain steps in rRNA maturation (Hayes and Vasseur, 1976; Sirdeshmukh and Schlessinger, 1985; Srivastava and Schlessinger, 1988). The dramatic decrease in IF2 binding to the 30S ribosomal subunit in the $\Delta ybeY$ mutant (Fig. 2H) could suggest that YbeY is needed, directly or indirectly, to form the translation initiation complex. A decrease in efficiency of initiation complex formation could slow translation to a point that negatively impacts maturation of rRNA.

The strongest evidence of a role for YbeY in rRNA and/or ribosome maturation comes from the analysis of rRNA in strains carrying the $\Delta ybeY$ mutation alone or in combination with other mutations in known RNases (Figs 3 and 4). RNases required for the processing of the 16S rRNA 3'-terminus, 23S rRNA 5'-terminus and 5S rRNA 5'-terminus have not been identified. Interestingly, we observe significant defects in processing of all of these termini in the $\Delta ybeY$ mutant (Fig. 3C and D). The $\Delta ybeY$ mutant shows a much stronger overall effect on rRNA maturation than any of the RNases tested, even those with a known role in rRNA processing (Fig. 4C–H). Only RNase III, which is required for the initial cleavages of the rRNA precursor, also affects the maturation of multiple rRNA termini. Deletion of both *ybeY* and *mrc* stabilizes the accumulation of an 18S-like rRNA precursor with similar mobility to that of a transient 18S rRNA species previously observed in an *mrc* mutant (Gegenheimer *et al.*, 1977). Loss of YbeY activity may therefore exacerbate a specific rRNA maturation defect that is only observed in the absence of the RNase III required for the initial cleavage events in rRNA maturation.

Deletion of *ybeY* along with either *mr* or *pnp* prevents, almost completely, the maturation of the 16S rRNA 3'-terminus (Fig. 4D), while maturation of all other rRNA termini proceeds to some degree in both of these mutants. Furthermore, large amounts of 16S rRNA are improperly processed in the $\Delta ybeY \Delta mnr$ double mutant, as shown by the accumulation of additional aberrant 16S rRNA maturation intermediates. These results suggest that YbeY may function along with RNase R and PNPase in the maturation of the 16S rRNA 3'-terminus and potentially in other steps of rRNA metabolism such as rRNA quality control.

Site-directed mutagenesis of YbeY supports the requirement for one of the His residues (H114) in the highly conserved putative hydrolase domain and a highly conserved Arg residue (R59) for YbeY function *in vivo* (Fig. 6). For example, ectopic expression from a plasmid of *ybeY* mutants H114A and R59A in the $\Delta ybeY$ mutant failed to complement the growth defect of the mutant strain at elevated temperatures. The H3XH5XH motif is

postulated to co-ordinate a Zn²⁺ ion that could act catalytically in a hydrolysis reaction (Zhan *et al.*, 2005). The crystal structure of YbeY also shows the highly conserved R59 pointing inward towards the H3XH5XH motif (Zhan *et al.*, 2005). The location of R59 along with its positive charge could imply that R59 helps position a substrate near the H3XH5XH motif. Given the strong genetic interactions of *ybeY* with *mrc*, *mr* and *pnp* (Fig. 4C–H) established here, along with the requirement of the putative hydrolase domain for *ybeY* activity, YbeY may also have RNase activity in the cell.

It is currently unclear why only one (H114) out of three His residues within the highly conserved H3XH5XH motif appears to be required for *ybeY* activity. It is possible that contributions of the other two His residues (H118, H124) towards YbeY activity may also be seen, when YbeY mutants are expressed at wild type-like levels. It is interesting to note the H114A and R59A mutants show similar sensitivity to growth at elevated temperature, but the rRNA processing defects are markedly different (Fig. 6B and C). While the rRNA profile of the H114A mutant resembles that of the *ybeY* mutant, the rRNA profile of the R59A mutant more closely resembles that of the MC4100, suggesting that the different mutations affect different functions of YbeY or affect the same rRNA processing function to different degrees.

Finally, although deletion of *ybeY* results in severe rRNA maturation defects suggesting a role in rRNA and/or ribosome maturation it remains possible that YbeY actually has an alternative function in the cell that impacts these processes. Although this remains a formal possibility, the simplest explanation for the striking effects of deletion of *ybeY* on rRNA processing and ribosome assembly is that YbeY functions directly in rRNA and/or ribosome maturation and assembly.

Experimental procedures

Strains, plasmids, growth conditions and DNA manipulations

Strains and plasmids are shown in Table S1. Strains were grown aerobically in Luria–Bertani (LB) medium at 37°C except for heat shock experiments where strains were grown at 45°C. Antibiotics for strain selection were used at the following concentrations, ampicillin 100 µg ml⁻¹, chloramphenicol 20 µg ml⁻¹ and kanamycin 25 µg ml⁻¹. Deletion of *ybeY* and the addition of a C-terminal FLAG tag to the genomic copy of *ybeY* (strain BWD55) was performed using the methods of Wanner (Datsenko and Wanner, 2000). For deletion, *ybeY* was first replaced with a *cat^R* cassette creating the $\Delta ybeY::cat^R$ mutant. The *cat^R* cassette was then excised to create a clean $\Delta ybeY$ deletion mutant in MC4100. Deletion of *ybeY* maintains the ATG start of *ybeY* and the overlapping stop codon of the upstream gene *ybeZ*. Translation from the ATG start of *ybeY* terminates eight amino acids into the flip

recombinase site of the λ -red gene targeting recombinase system (Datsenko and Wanner, 2000). DNA primers used for deletion of *ybeY* and addition of the C-terminal FLAG tag are listed in Table S2. Allele transfers were done by P1 transduction. DNA manipulations were performed according to the methods of Sambrook (Sambrook and Russell, 2001). Site-directed mutagenesis was performed using the Quikchange II kit from Stratagene. DNA primers used for site-directed mutagenesis are listed in Table S2. For complementation studies, *ybeY*, *SMc01113* and *yqfG* were cloned downstream of the tetracycline promoter in pBR322 or the tryptophan promoter in pMS03. The DNA primers used for complementation cloning are shown in Table S2.

Phenotypic analysis

Stress and plant assays were performed as previously described (Davies and Walker, 2008).

Polysome and rRNA analysis

rRNA and polysomes were extracted from cells growing exponentially at 37°C, unless otherwise stated. Cells were pelleted, resuspended in buffer A (20 mM HEPES pH 7.5, 5 mM β -mercaptoethanol, 10 mM MgCl₂, 50 mM NH₄Cl and 0.1 mM PMSF) and lysed by repeated freeze/thaw cycles. After clarification by centrifugation, lysates were loaded on a 5–20% sucrose gradient. Polysomes and ribosomes were separated by ultracentrifugation followed by fractionation using a peristaltic pump. The A₂₆₀ value was determined for each fraction.

rRNA was extracted from logarithmically growing cultures or separated ribosome fractions using Qiagen RNeasy Mini Kit. For gross analysis of 16S and 23S rRNA, 800 ng of rRNA was separated by synergel/agarose gel electrophoresis as described (Wachi *et al.*, 1999) using 0.9% synergel and 0.7% agarose.

Primer extension assays were performed using Superscript II reverse transcriptase (Invitrogen) as per manufacturer's instructions with primers annealing to the 5'-terminal region of mature 16S, 23S and 5S rRNA respectively. The site-specific RNase H cleavage assay was performed as described (Li and Deutscher, 1995; Li *et al.*, 1999a,b) followed by Northern hybridization using probes specific for the mature 3'-termini of 16S, 23S and 5S rRNAs respectively. One hundred nanograms of total RNA was used for each assay. Reaction products were separated on denaturing polyacrylamide gels; 10% polyacrylamide/7 M urea for primer extensions and 6% polyacrylamide/7 M urea for site-specific RNase H cleavage assays. DNA oligonucleotides for primer extension and RNase H mapping are shown in Table S2.

For the Northern blot in Fig. 2B, 250 ng of total MC4100 and $\Delta ybeY$ RNA was separated on a 1% agarose gel containing 2% formamide. Probes used to identify 17S rRNA 5'- and 3'-termini are shown in Table S2. The probes were detected with the AlkPhos Direct Labeling Kit (Amersham).

For temperature shift assays, cultures were grown to exponential phase at 37°C. Samples were taken for rRNA analysis and the cultures were then shifted to 45°C. After 1 h samples were taken for rRNA analysis.

In vitro translation and lacZ assays

In vitro translation assays were performed essentially as described (Matthaei and Nirenberg, 1961). Exponentially growing cells were pelleted, resuspended in buffer A and lysed by French press. Lysates were centrifuged twice at 30 000 *g* and once at 100 000 *g*. The resulting supernatant was kept as the S100 fraction, and the pelleted ribosomes were washed and resuspended in buffer A. Ribosomes were further purified by pelleting at 200 000 *g* through a 20% sucrose cushion. Pelleted ribosomes were resuspended in buffer A and quantified by UV absorbance at 260 nm, and stored at –80°C.

Mix I (25 μ g of polyU mRNA and 10 pmoles of ribosomes) was pre-incubated at 37°C for 15 min in polyU-mix (20 mM HEPES pH 7.5, 2 mM DTT, 10 mM MgCl₂, 50 mM NH₄Cl, 1 mM spermidine, 2 mM ATP, 0.5 mM GTP, 8 mM phosphoenolpyruvate and 2 μ M pyruvate kinase). Mix II (50 μ g of bulk tRNA, 100 μ g S100 extract and 200 pmoles of C¹⁴-Phe) was incubated at 37°C for 15 min in polyU-mix. Mix I and II were mixed and incubated at 37°C for 15 min. ¹⁴C-Phe incorporation into protein was determined by TCA precipitation and scintillation counting. All reactions were performed in triplicate.

For LacZ assays, strains carrying wild-type or mutant *lacZ* alleles on a plasmid were harvested in exponential phase. The assay for LacZ activity was then performed as described (Miller, 1972). The per cent LacZ activity (in Miller units) of each mutant *lacZ* allele relative to the wild-type *lacZ* allele for MC4100 or the $\Delta ybeY$ mutant is reported.

Acknowledgements

We thank Dr Harry Noller for helpful discussion, Dr Murray Deutscher for his generous gifts of *mbb*, *rnd*, *rph*, *mt*, *rne-1* and *mnr* *E. coli* mutant strains and Dr Albert Dahlberg for insightful discussion and for his gift of plasmids pSG25, pSG163, pSG853, pSG3/4, *plac7* and *plac10*. This work was supported by National Institutes of Health Grant GM31030 to G.C.W. and Grant GM17151 to U.L.R., MIT Center for Environmental Health Sciences Grant P30ES002109, a National Sciences and Engineering Research Council of Canada graduate scholarship to B.W.D. and an NCI postdoctoral fellowship to L.A.S. G.C.W. is an American Cancer Society Research Professor.

References

- Akerley, B.J., Rubin, E.J., Novick, V.L., Amaya, K., Judson, N., and Mekalanos, J.J. (2002) A genome-scale analysis for identification of genes required for growth or survival of *Haemophilus influenzae*. *Proc Natl Acad Sci USA* **99**: 966–971.
- Arifuzzaman, M., Maeda, M., Itoh, A., Nishikata, K., Takita, C., Saito, R., *et al.* (2006) Large-scale identification of protein-protein interaction of *Escherichia coli* K-12. *Genome Res* **16**: 686–691.
- Bunner, A.E., Nord, S., Wikström, P.M., and Williamson, J.R. (2010) The effect of ribosome assembly cofactors on *in vitro* 30S subunit reconstitution. *J Mol Biol* **398**: 1–7.

- Cheng, Z.F., and Deutscher, M.P. (2003) Quality control of ribosomal RNA mediated by polynucleotide phosphorylase and RNase R. *Proc Natl Acad Sci USA* **100**: 6388–6393.
- Datsenko, K.A., and Wanner, B.L. (2000) One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci USA* **97**: 6640–6645.
- Davies, B.W., and Walker, G.C. (2008) A highly conserved protein of unknown function is required by *Sinorhizobium meliloti* for symbiosis and environmental stress protection. *J Bacteriol* **190**: 1118–1123.
- Deutscher, M.P. (2009) Maturation and degradation of ribosomal RNA in bacteria. *Prog Mol Biol Transl Sci* **85**: 369–391.
- Friedberg, E.C., Walker, G.C., Siede, W., Wood, R.D., Schultz, R.A., and Ellenberger, T. (2005) *DNA Repair and Mutagenesis*. Washington, DC: ASM Press.
- Gegenheimer, P., Watson, N., and Apirion, D. (1977) Multiple pathways for primary processing of ribosomal RNA in *Escherichia coli*. *J Biol Chem* **252**: 3064–3073.
- Gil, R., Silva, F.J., Pereto, J., and Moya, A. (2004) Determination of the core of a minimal bacterial gene set. *Microbiol Mol Biol Rev* **68**: 518–537.
- Green, R., and Noller, H.F. (1999) Reconstitution of functional 50S ribosomes from in vitro transcripts of *Bacillus stearothermophilus* 23S rRNA. *Biochemistry* **38**: 1772–1779.
- Hayes, F., and Vasseur, M. (1976) Processing of the 17-S *Escherichia coli* precursor RNA in the 27-S pre-ribosomal particle. *Eur J Biochem* **61**: 433–442.
- Himeno, H., Hanawa-Suetsugu, K., Kimura, T., Takagi, K., Sugiyama, W., Shirata, S., et al. (2004) A novel GTPase activated by the small subunit of ribosome. *Nucl Acids Res* **32**: 5303–5309.
- Imlay, J.A., and Linn, S. (1987) Mutagenesis and stress responses induced in *Escherichia coli* by hydrogen peroxide. *J Bacteriol* **169**: 2967–2976.
- Inoue, K., Alsina, J., Chen, J., and Inouye, M. (2003) Suppression of defective ribosome assembly in a rbfA deletion mutant by overexpression of Era, an essential GTPase in *Escherichia coli*. *Mol Microbiol* **48**: 1005–1016.
- Jones, K.M., Kobayashi, H., Davies, B.W., Taga, M.E., and Walker, G.C. (2007) How rhizobial symbionts invade plants: the *Sinorhizobium-Medicago* model. *Nat Rev Microbiol* **5**: 619–633.
- Kaczanowska, M., and Ryden-Aulin, M. (2007) Ribosome biogenesis and the translation process in *Escherichia coli*. *Microbiol Mol Biol Rev* **71**: 477–494.
- King, T.C., Sirdeshmukh, R., and Schlessinger, D. (1984) RNase III cleavage is obligate for maturation but not for function of *Escherichia coli* pre-23S rRNA. *Proc Natl Acad Sci USA* **81**: 185–188.
- Kobayashi, K., Ehrlich, S.D., Albertini, A., Amati, G., Andersen, K.K., Arnaud, M., et al. (2003) Essential *Bacillus subtilis* genes. *Proc Natl Acad Sci USA* **100**: 4678–4683.
- Li, Z., and Deutscher, M.P. (1995) The tRNA processing enzyme RNase T is essential for maturation of 5S RNA. *Proc Natl Acad Sci USA* **92**: 6883–6886.
- Li, Z., Pandit, S., and Deutscher, M.P. (1999a) RNase G (CafA protein) and RNase E are both required for the 5' maturation of 16S ribosomal RNA. *EMBO J* **18**: 2878–2885.
- Li, Z., Pandit, S., and Deutscher, M.P. (1999b) Maturation of 23S ribosomal RNA requires the exoribonuclease RNase T. *RNA* **5**: 139–146.
- Lövgren, J.M., Bylund, G.O., Srivastava, M.K., Lundberg, L.A., Persson, O.P., Wingsle, G., and Wikström, P.M. (2004) The PRC-barrel domain of the ribosome maturation protein RimM mediates binding to ribosomal protein S19 in the 30S ribosomal subunits. *RNA* **10**: 1798–1812.
- Mangiarotti, G., Turco, E., Ponsetto, A., and Altruda, F. (1974) Precursor 16S RNA in active 30S ribosomes. *Nature* **247**: 147–148.
- Mangiarotti, G., Turco, E., Perlo, C., and Altruda, F. (1975) Role of precursor 16S RNA in assembly of *E. coli* 30S ribosomes. *Nature* **253**: 569–571.
- Matthaei, H., and Nirenberg, M.W. (1961) The dependence of cell-free protein synthesis in *E. coli* upon RNA prepared from ribosomes. *Biochem Biophys Res Commun* **4**: 404–408.
- Miller, J.H. (1972) *Experiments in Molecular Cloning*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory.
- Misra, T.K., and Apirion, D. (1979) RNase E, an RNA processing enzyme from *Escherichia coli*. *J Biol Chem* **254**: 11154–11159.
- Nierhaus, K.H. (1980) The assembly of the prokaryotic ribosome. *Biosystems* **12**: 273–282.
- Nierhaus, K.H. (1991) The assembly of prokaryotic ribosomes. *Biochimie* **73**: 739–755.
- Noller, H.F., and Nomura, M. (1987) *Ribosomes*. *Escherichia coli and Salmonella typhimurium; Cellular and Molecular Biology*. Washington DC: American Society for Microbiology, pp. 104–125.
- Nomura, M., and Erdmann, V.A. (1970) Reconstitution of 50S ribosomal subunits from dissociated molecular components. *Nature* **228**: 744–748.
- Oganessian, V., Busso, D., Brandsen, J., Chen, S., Jancarik, J., Kim, R., and Kim, S.H. (2003) Structure of the hypothetical protein AQ_1354 from *Aquifex aeolicus*. *Acta Crystallogr D Biol Crystallogr* **59**: 1219–1223.
- Poirot, O., O'Toole, E., and Notredame, C. (2003) Tcoffee@igs: a web server for computing, evaluating and combining multiple sequence alignments. *Nucl Acids Res* **31**: 3503–3506.
- Rasouly, A., Schonbrun, M., Shenhar, Y., and Ron, E.Z. (2009) YbeY – a heat shock protein involved in translation in *Escherichia coli*. *J Bacteriol* **191**: 2649–2655.
- Robertson, H.D., Webster, R.E., and Zinder, N.D. (1967) A nuclease specific for double-stranded RNA. *Virology* **32**: 718–719.
- Sambrook, J., and Russell, D.W. (2001) *Molecular Cloning. A Laboratory Manual*. Cold Spring Harbour, NY: Cold Spring Harbor Laboratory.
- Sato, A., Kobayashi, G., Hayashi, H., Yoshida, H., Wada, A., Maeda, M., et al. (2005) The GTP binding protein Obg homolog ObgE is involved in ribosome maturation. *Genes Cells* **10**: 393–408.
- Sirdeshmukh, R., and Schlessinger, D. (1985) Why is processing of 23 S ribosomal RNA in *Escherichia coli* not obligate for its function? *J Mol Biol* **186**: 669–672.
- Srivastava, A.K., and Schlessinger, D. (1988) Coregulation of processing and translation: mature 5' termini of *Escherichia coli* 23S ribosomal RNA form in polysomes. *Proc Natl Acad Sci USA* **85**: 7144–7148.

- Tatusov, R.L., Natale, D.A., Garkavtsev, I.V., Tatusova, T.A., Shankavaram, U.T., Rao, B.S., *et al.* (2001) The COG database: new developments in phylogenetic classification of proteins from complete genomes. *Nucl Acids Res* **29**: 22–28.
- Verma, R., Aravind, L., Oania, R., McDonald, W.H., Yates, J.R., 3rd, Koonin, E.V., and Deshaies, R.J. (2002) Role of Rpn11 metalloprotease in deubiquitination and degradation by the 26S proteasome. *Science* **298**: 611–615.
- Wachi, M., Umitsuki, G., Shimizu, M., Takada, A., and Nagai, K. (1999) *Escherichia coli* cafA gene encodes a novel RNase, designated as RNase G, involved in processing of the 5' end of 16S rRNA. *Biochem Biophys Res Commun* **259**: 483–488.
- Wilson, D.N., and Nierhaus, K.H. (2007) The weird and wonderful world of bacterial ribosome regulation. *Crit Rev Biochem Mol Biol* **42**: 187–219.
- Wireman, J.W., and Sypherd, P.S. (1974) In vitro assembly of 30S ribosomal particles from precursor 16S RNA of *Escherichia coli*. *Nature* **247**: 552–554.
- Yeh, D.C., Parsons, L.M., Parsons, J.F., Liu, F., Eisenstein, E., and Orban, J. (2005) NMR structure of HI0004, a putative essential gene product from *Haemophilus influenzae*, and comparison with the X-ray structure of an *Aquifex aeolicus* homolog. *Protein Sci* **14**: 424–430.
- Zhan, C., Fedorov, E.V., Shi, W., Ramagopal, U.A., Thirumuruhan, R., Manjasetty, B.A., *et al.* (2005) The ybeY protein from *Escherichia coli* is a metalloprotein. *Acta Crystallogr Sect F Struct Biol Cryst Commun* **61**: 959–963.

Supporting information

Additional supporting information may be found in the online version of this article.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.