

# Two Iranian Families With a Novel Mutation in *GJB2* Causing Autosomal Dominant Nonsyndromic Hearing Loss

Niloofar Bazazzadegan,<sup>1</sup> Abraham M. Sheffield,<sup>2</sup> Masoomeh Sobhani,<sup>3</sup> Kimia Kahrizi,<sup>1</sup> Nicole C. Meyer,<sup>2</sup> Guy Van Camp,<sup>4</sup> Nele Hilgert,<sup>4</sup> Seyedeh Sedigheh Abedini,<sup>1</sup> Farkhondeh Habibi,<sup>5</sup> Ahmad Daneshi,<sup>6</sup> Carla Nishimura,<sup>2</sup> Matthew R. Avenarius,<sup>7</sup> Mohammad Farhadi,<sup>6</sup> Richard J.H. Smith,<sup>2</sup> and Hossein Najmabadi<sup>1\*</sup>

<sup>1</sup>Genetics Research Center, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran

<sup>2</sup>Molecular Otolaryngology Research Laboratories, Department of Otolaryngology Head and Neck Surgery, University of Iowa, IA

<sup>3</sup>Prevention Department, Talesh Welfare Center, Gilan Province, Iran

<sup>4</sup>Department of Medical Genetics, University of Antwerp (UA), Antwerp, Belgium

<sup>5</sup>Prevention Department, Rasht Welfare Center, Gilan Province, Iran

<sup>6</sup>Research Center of Ear, Nose, Throat, Head and Neck Surgery, Iran University of Medical Sciences, Tehran, Iran

<sup>7</sup>Department of Human Genetics, University of Michigan School of Medicine, Ann Arbor, Michigan

Received 6 May 2008; Accepted 17 October 2009

Mutations in *GJB2*, encoding connexin 26 (Cx26), cause both autosomal dominant and autosomal recessive nonsyndromic hearing loss (ARNSHL) at the *DFNA3* and *DFNB1* loci, respectively. Most of the over 100 described *GJB2* mutations cause ARNSHL. Only a minority has been associated with autosomal dominant hearing loss. In this study, we present two families with autosomal dominant nonsyndromic hearing loss caused by a novel mutation in *GJB2* (p.Asp46Asn). Both families were ascertained from the same village in northern Iran consistent with a founder effect. This finding implicates the D46N missense mutation in Cx26 as a common cause of deafness in this part of Iran mandating mutation screening of *GJB2* for D46N in all persons with hearing loss who originate from this geographic region. © 2011 Wiley-Liss, Inc.

**Key words:** connexin 26; D46N; autosomal dominant nonsyndromic hearing loss; *DFNA3*; Iran

## INTRODUCTION

Approximately one in 1,000 newborns has severe-to-profound hearing loss and in more than half of this group of babies the etiology is genetic [Smith et al., 2005]. The pattern of inheritance is most frequently autosomal recessive, while autosomal dominant, X-linked and mitochondrial inheritances occur less frequently. Mutations in one gene, *GJB2*, have been found to cause up to half of autosomal recessive nonsyndromic hearing loss (ARNSHL) cases in developed countries [Estivill et al., 1998; Kelley et al., 1998; Green et al., 1999].

### How to Cite this Article:

Bazazzadegan N, Sheffield AM, Sobhani M, Kahrizi K, Meyer NC, Van Camp G, Hilgert N, Abedini SS, Habibi F, Daneshi A, Nishimura C, Avenarius MR, Farhadi M, Smith RJH, Najmabadi H. 2011. Two Iranian families with a novel mutation in *GJB2* causing autosomal dominant nonsyndromic hearing loss.

Am J Med Genet Part A 155:1202–1211.

*GJB2* encodes connexin 26 (Cx26), a member of a highly conserved protein family found throughout the animal kingdom [van Steensel et al., 2004a]. Six connexins oligomerize to form pore-like plasma membrane hemichannels called connexons. Connexons

Grant sponsor: Iranian National Science Foundation; Grant numbers: 85073/23, 85033/10; Grant sponsor: National Institutes of Health (NIH)-NIDCD; Grant numbers: RO1 DCOO2842, RO1 DCO3544.

\*Correspondence to:

Hossein Najmabadi, Ph.D., Professor of Medical & Molecular Genetics, Head & Director of the Genetics Research Center, Director of National Prenatal Reference Laboratory, University of Social Welfare and Rehabilitation Sciences, Koodakyar St, Daneshjoo Blvd, Evin, Tehran 19834, Iran. E-mail: hnajm12@yahoo.com

Published online 11 April 2011 in Wiley Online Library (wileyonlinelibrary.com).

DOI 10.1002/ajmg.a.33209

in adjacent cells then align to form gap junction channels, which facilitate electrical and biochemical coupling between cells. Although the function of gap junction channels in the inner ear has not been definitively determined, a role in maintaining cochlear

homeostasis by recycling  $K^+$  ions following hair cell depolarization has been postulated [Kikuchi et al., 1995].

The majority of the over 100 deafness-causing mutations described in *GJB2* causes ARNSHL, however 29 *GJB2* mutations have

TABLE I. AD Nonsyndromic and Syndromic *GJB2* Mutations

Mutation		Protein domain	Number of affecteds	Phenotype	Refs.
cDNA level	Protein level				
c.131G > C	W44S	E1	—	Nonsyndromic	<a href="http://davinci.crg.es/deafness/index.php?seccion=mut_db&amp;db=nonsynd&amp;nonsynd=cx26mut">http://davinci.crg.es/deafness/index.php?seccion=mut_db&amp;db=nonsynd&amp;nonsynd=cx26mut</a>
c.132G > C	W44C	E1	2 French families	Nonsyndromic	Denoyelle et al. [1998]
c.223C > T	R75W	M2	—	Nonsyndromic	Janecke et al. [2001]; Mani et al. [2008]
c.224G > A	R75Q	M2	4	Nonsyndromic	<a href="http://davinci.crg.es/deafness/index.php?seccion=mut_db&amp;db=nonsynd&amp;nonsynd=cx26mut">http://davinci.crg.es/deafness/index.php?seccion=mut_db&amp;db=nonsynd&amp;nonsynd=cx26mut</a>
c.428G > A	R143Q	M3	1 Austrian family	Nonsyndromic	Löffler et al. [2001]
c.487A > C	M163L	E2	—	Nonsyndromic	Matos et al. [2008]
c.535G > A	D179N	E2	1 Italian family	Nonsyndromic	<a href="http://davinci.crg.es/deafness/index.php?seccion=mut_db&amp;db=nonsynd&amp;nonsynd=cx26mut">http://davinci.crg.es/deafness/index.php?seccion=mut_db&amp;db=nonsynd&amp;nonsynd=cx26mut</a>
c.551G > A	R184Q	E2	—	Nonsyndromic	Hamelmann et al. [2001]
c.605G > T	C202F	M4	A large French family	Nonsyndromic	Morlé et al. [2000]
c.34G > C	G12R	NT	1	Syndromic (keratitis-ichthyosis-deafness)	Richard et al. [2002]
c.42C > G	N14K	NT	1	Hypotrichosis-deafness	van Steensel et al. [2004b]
c.40A > C	N14Y	NT	1	Syndromic (keratitis-ichthyosis-deafness)	Arita et al. [2006]
c.50C > T	S17F	NT	1	Syndromic (keratitis-ichthyosis-deafness)	Richard et al. [2002]
c.119C > T	A40V	M1	1	Syndromic (keratitis-ichthyosis-deafness and the follicular occlusion triad)	Montgomery et al. [2004]
125_127delAGG	E42del	M1	1 family	Syndromic (palmoplantar keratoderma with deafness)	Rouan et al. [2001]
c.134G > A	G45E	E1	—	Syndromic (Keratitis-Ichthyosis-Deafness)	Janecke et al. [2005]; Jonard et al. [2008]
c.148G > A	D50N	E1	?	Syndromic (keratitis-ichthyosis-deafness; hystrix-like ichthyosis-deafness)	van Steensel et al. [2002]
c.148G > T	D50Y	E1	1	Syndromic (keratitis-ichthyosis-deafness)	Yotsumoto et al. [2003]
c.160A > C	N54H	E1	1	Syndromic (Bart-Pumphrey)	Akiyama et al. [2007]
not described	N54K	E1	1 family	Syndromic (Bart-Pumphrey)	Richard et al. [2000]
c.175G > C	G59R	E1	1	Syndromic (palmoplantar keratoderma with deafness)	Leonard et al. [2005]
c.175G > A	G59S	E1	1	Syndromic (palmoplantar keratoderma with deafness)	Alexandrino et al. [2005]
c.176C > G	G59A	E1	1 family	Syndromic (palmoplantar keratoderma with deafness)	Heathcote et al. [2000]
c.196G > C	D66H	E1	10	Syndromic (Vohwinkel)	Maestrini et al. [1999]
c.219A > G	H73R	M2	1 family	Syndromic (palmoplantar keratoderma with deafness)	de Zwart-Storm et al. [2008b]
c.223C > T	R75W	M2	—	Syndromic (palmoplantar keratoderma with deafness)	Richard et al. [1998]; Yuan et al. [2009]
c.224G > A	R75Q	E1	1 Turkish family	Syndromic (palmoplantar keratoderma with deafness)	Uyguner et al. [2002]
c.389G > T	G130V	CL	1 family	Syndromic (Vohwinkel)	Snoeckx et al. [2005]
c.424T > C	F142L	M3	1	Syndromic (mucositis-deafness)	Brown et al. [2003]
c.548C > T	S183F	E2	1 family	Syndromic (palmoplantar keratoderma with deafness)	de Zwart-Storm et al. [2008a]


been associated with autosomal dominant hearing loss (ADHL) (Table I). About two-thirds of these mutations cause a syndrome in which hearing loss is associated with distinct skin manifestations. Examples of these syndromes include Keratitis-Ichthyosis-Deafness (KID) syndrome [MIM 148210], Vohwinkel syndrome [MIM 124500], Bart-Pumphrey syndrome [MIM 149200], and palmoplantar keratoderma (PPK) with deafness [MIM 148350]. The remaining one-third of dominant *GJB2* mutations causes autosomal dominant nonsyndromic hearing loss (ADNSHL). In

this study, we present a novel *GJB2* mutation responsible for ADNSHL in two families from a village in northern Iran.

## MATERIALS AND METHODS

### Subjects

This study was approved by the Institutional Review Boards of the University of Social Welfare and Rehabilitation Sciences in Tehran, Iran and the University of Iowa, Iowa City, IA, United States.



**FIG. 1.** A: Pedigrees of two families, L-1301 [upper pedigree] and L-3057 [lower pedigree] with hearing loss caused by a novel mutation in *GJB2*, D46N. The proband of each family is indicated by an arrow. All individuals who did not have audiograms are indicated with an asterisk. Studied branches are shown with capital letters. B: All genotypes are indicated on pedigrees beneath the studied individuals.




FIG. 1. (Continued)

Informed consent was obtained from all family members wishing to participate in this research. For each consenting person, clinical assessment and genetic counseling were performed. Audiometric testing was performed by measuring air and bone conduction thresholds.

### Haplotyping

Genomic DNA was isolated from peripheral blood using established techniques [Miller et al., 1988] and used to complete haplotype reconstruction for microsatellite markers that map to chromosome 13q12, the genomic region that includes *GJB2*. Markers were PCR amplified and products were resolved on the ABI Prism 3130 Genetic Analyzer. Alleles were assigned using GeneMapper 3.0 software (Applied Biosystems, Inc., Foster city, CA), with haplotypes reconstructed using custom-made software.

### GJB2 Screening

Mutation screening was completed on an Applied Biosystems (ABI) model 3700 automated sequencer. Sequence data were compared to published sequence for *GJB2* using the Sequencher 4.1.2 software program package (Gene Codes, MI, Ann Arbor, MI). The putative

structural and functional significance of observed mutations was assessed using the ConSeq web server (<http://conseq.tau.ac.il/>). Conseq scores obtained using this alignment method range from 1 (variable) to 9 (conserved).

## RESULTS

### Subjects

Family members from two families, L-1301 and L-3057, segregating ADNSHL were recruited for this study (Fig. 1). Although most persons consented to participate, some individuals (shown with asterisk in Fig. 1) declined to have audiograms taken. For those persons with hearing loss, audiometry confirmed the loss to be sensorineural. Figure 2 shows hearing thresholds for the better hearing ear in six persons from family L-1301 and in four persons from family L-3057. In most persons, the hearing loss was postlingual with an onset in the first decade, however VI:13 and VI:9 had prelingual hearing loss that quickly progressed to become severe to profound in VI:13 (Fig. 2: L-1301:1 and L-1301:2). Intrafamilial variation was observed (Fig. 2). Pure tone audiograms by age group are shown in Figure 3. No features of syndromic hearing loss were observed in any person with hearing loss in these families (Fig. 4).




FIG. 1. (Continued)

### Haplotype Analysis and Mutation Screening

Haplotype reconstruction was consistent with linkage to the DFNA3 locus on chromosome 13q12. Mutation analysis of *GJB2* identified a novel mutation, 351G > A, in affected subjects (Fig. 1). This missense mutation replaces the acidic aspartate residue at position 46 with the neutral asparagine residue (D46N). The mutation was not present in nonaffected family members and was similarly absent in 200 Iranian control samples. D46 is located in the

first extracellular loop (E1) of Cx26, a region known to be highly conserved among connexin family members and to play an important role in voltage gating and in facilitating docking between opposing connexons [Rubin et al., 1992; White et al., 1995; Martin and Evans, 2004; Mese et al., 2007]. The ConSeq alignment confirmed conservation of this amino acid in multiple species (ConSeq score = 9; Fig. 5). Based on these data, D46N is believed to be the disease-causing mutation in both families segregating ADNSHL.




FIG. 1. (Continued)





FIG. 3. Audioprofiles showing average audiograms by age group for both families.

FIG. 2. Audiograms associated with *GJB2* D46N hearing loss [better hearing ear is shown]. Top: Sequential audiograms from two persons in family L-1301 showing progression to severe-to-profound hearing loss. Middle: Audiograms from five persons in family L-1301 at different ages showing relatively uniform audioprofiles. Bottom: Audiograms from four persons in family L-3057 at different ages showing variably in hearing thresholds by age.





**FIG. 4.** Photos of one affected individual showing absence of skin symptoms.

## DISCUSSION

In this report we present two families with ADNSHL caused by a novel amino acid change—D46N—in *GJB2*. To date, 9 mutations in *GJB2* have been linked to ADNSHL whilst 21 mutations cause syndromic hearing loss associated with a variety of skin diseases. Two mutations, R75W and R75Q, cause both syndromic and nonsyndromic hearing loss.

Most of the known dominant mutations, whether syndromic and nonsyndromic, occur in the highly conserved first extracellular loop (E1) of Cx26, which is crucial for voltage gating and connexon–connexon docking (Fig. 6) [Richard et al., 2002; Martin and Evans, 2004]. Prominent among these mutations are G45E, D50N, and D50Y, which cause KID syndrome [Richard et al., 2002; Sonoda

et al., 2004; Janecke et al., 2005; OMIM 148210], and W44C and W44S, which cause ADNSHL [Denoyelle et al., 1998; Tekin et al., 2001; Marziano et al., 2003; MIM 601544]. The D46N mutation affects the same region of E1 (Fig. 6). The recently resolved crystal structure of a Cx26 gap junction channel has shown that this segment of E1 forms a  $3_{10}$  helix structure which lines the channel pore [Maeda et al., 2009].

Many mutations in *GJB2* that cause ARNSHL cause a simple loss-of-channel activity, however it is less clear how dominant mutations affect gap junction function [Bruzzone et al., 2001]. The broad phenotypic heterogeneity seen with autosomal dominant mutations suggests variability in the molecular defects caused by individual mutations. For example, some dominant *GJB2* mutations, like R75W, are known to inhibit formation of functional channels. Expression studies in *Xenopus* oocytes have shown that R75W, which occurs in a highly conserved region involved in voltage gating, inhibits normal gap junction function of co-expressed wild-type Cx26 in a dominant-negative manner [Richard et al., 1998]. Dye transfer studies using the gap junction permeant tracer Cascade Blue have also demonstrated a disruption of intercellular coupling with W44S, R75W, G59A, and D66H. For G59A and D66H mutants, this change correlates with impaired intracellular trafficking and targeting to the plasma membrane [Marziano et al., 2003]. Functional studies also indicate that the mutation T55N produces a protein that is deeply impaired in its intracellular trafficking and fails to reach the plasma membrane [Melchionda et al., 2005].

Although functional studies of D46N are needed to determine the effect of this mutation on channel activity, useful information can be gleaned by studying the structure of wild-type Cx26 gap junction channels. Based on the structural analysis by Maeda and coworkers, D46 is one of the key residues involved in interactions between adjacent connexins. Additionally, D46 and D50 line the gap junction pore at the level of the cell membrane and face the interior of the channel, creating a negatively charged path between adjacent cells [Maeda et al., 2009]. This region is hypothesized to contribute to charge selectivity and size restriction of the gap junction. While the size of the side chain would not be significantly altered, the D46N mutation would reduce the negative charge, potentially altering the permselectivity of the channel.


The strong evolutionary conservation of D46 between species and among connexins, its location lining the pore of the Cx26 gap junction, the predicted importance of its negative charge, and its absence in 200 Iranian control subjects are convincing evidence for the disease-causing effect of this mutation. The identification of this mutation in two families from the same small village in northern Iran is consistent with a founder effect, which we confirmed by genotyping additional microsatellite markers in the DFNB1 region. On this basis, we recommend screening for this mutation in deaf persons from the Northern provinces in Iran.

## ACKNOWLEDGMENTS

We would like to thank our patients and their families for their collaboration in this research. This project was sponsored by the Iranian National Science Foundation grant numbers 85073/23 and 85033/10. Richard J. H. Smith is the Sterba Hearing Research







Professor, University of Iowa College of Medicine, who supported the project with National Institutes of Health (NIH)-NIDCD grants RO1 DCOO2842 and RO1 DCO3544.

## REFERENCES

- Akiyama M, Sakai K, Arita K, Nomura Y, Ito K, Kodama K, McMillan JR, Kobayashi K, Sawamura D, Shimizu H. 2007. A novel GJB2 mutation p. Asn54His in a patient with palmoplantar keratoderma, sensorineural hearing loss and knuckle pads. *J Invest Dermatol* 127:1540–1543.
- Alexandrino F, Sartorato EL, Marques-de-Faria AP, Steiner CE. 2005. G59S mutation in the GJB2 (connexin 26) gene in a patient with Bart-Pumphrey syndrome. *Am J Med Genet Part A* 136A:282–284.
- Arita K, Aiyama M, Aizawa T, Umetsu Y, Segawa I, Goto M, Sawamura D, Demura M, Kawano K, Shimizu H. 2006. A novel N14Y mutation in Connexin26 in keratitis-ichthyosis-deafness syndrome: Analyses of altered gap junctional communication and molecular structure of N terminus of mutated Connexin26. *Am J Pathol* 169:416–423.
- Brown CW, Levy ML, Flaitz CM, Reid BS, Manolidis S, Hebert AA, Bender MM, Heilstedt HA, Plunkett KS, Fang P, Roa BB, Chung P, Tang HY, Richard G, Alford RL. 2003. A novel GJB2 connexin 26 mutation, F142L, in a patient with unusual mucocutaneous findings and deafness. *J Invest Dermatol* 121:1221–1223.
- Bruzzone R, Gomes D, Denoyelle E, Duval N, Perea J, Veronesi V, Weil D, Petit C, Gabellec MM, D'Andrea P, White TW. 2001. Functional analysis of a dominant mutation of human connexin26 associated with non-syndromic deafness. *Cell Commun Adhes* 8:425–431.
- de Zwart-Storm EA, van Geel M, van Neer PA, Steijlen PM, Martin PE, van Steensel MA. 2008a. A novel missense mutation in the second extracellular domain of GJB2, p.Ser183Phe, causes a syndrome of focal palmoplantar keratoderma with deafness. *Am J Pathol* 173:1113–1119.
- de Zwart-Storm EA, Hamm H, Stoevesandt J, Steijlen PM, Martin PE, van Geel M, van Steensel MA. 2008b. A novel missense mutation in GJB2 disturbs gap junction protein transport and causes focal palmoplantar keratoderma with deafness. *J Med Genet* 45:161–166.
- Denoyelle F, Lina-Granade G, Plauchu H, Bruzzone R, Chaïb H, Lévi-Acobas F, Weil D, Petit C. 1998. Connexin 26 gene linked to a dominant deafness. *Nature* 393:319–320.
- Estivill X, Fortina P, Surrey S, Rabionet R, Melchionda S, D'Agruma I. 1998. Connexin-26 mutations in sporadic and inherited sensorineural deafness. *Lancet* 351:394–398.
- Green GE, Scott DA, McDonald JM, Woodworth GG, Sheffield VC, Smith RJH. 1999. Carrier rates in the midwestern United States for GJB2 mutations causing inherited deafness. *JAMA* 281:2211–2216.
- Hamelmann C, Amedofu GK, Albrecht K, Muntau B, Gelhaus A, Brobby GW, Horstmann RD. 2001. Pattern of connexin 26 (GJB2) mutations causing sensorineural hearing impairment in Ghana. *Hum Mutat* 18:84–85.
- Heathcote K, Syrris P, Carter ND, Patton MA. 2000. A connexin 26 mutation causes a syndrome of sensorineural hearing loss and palmoplantar hyperkeratosis. *J Med Genet* 37:50–51.
- Janecke AR, Nekahm D, Löffler J, Hirst-Stadlmann A, Müller T, Utermann G. 2001. De novo mutation of the connexin 26 gene associated with dominant non-syndromic sensorineural hearing loss. *Hum Genet* 108:269–270.
- Janecke AR, Hennies HC, Günther B, Gansl G, Smolle J, Messmer EM, Utermann G, Rittinger O. 2005. GJB2 mutations in keratitis-ichthyosis-deafness syndrome including its fatal form. *Am J Med Genet Part A* 133A:128–131.
- Jonard L, Feldmann D, Parsy C, Freitag S, Sinico M, Koval C, Grati M, Couderc R, Denoyelle F, Bodemer C, Marlin S, Hadj-Rabia S. 2008. A familial case of Keratitis-Ichthyosis-Deafness (KID) syndrome with the GJB2 mutation G45E. *Eur J Med Genet* 51:35–43.
- Kelley PM, Harris DJ, Comer BC, Askew JW, Fowler T, Smith SD, Kimberling WJ. 1998. Novel mutations in the connexin 26 gene (GJB2) that cause autosomal recessive (DFNB1) hearing loss. *Am J Hum Genet* 62:792–799.
- Kikuchi T, Kimura RS, Paul DL, Adams JC. 1995. Gap junctions in the rat cochlea: Immunohistochemical and ultrastructural analysis. *Anal Embryol* 191:101–118.
- Leonard NJ, Krol AL, Bleoo S, Somerville MJ. 2005. Sensorineural hearing loss, striate palmoplantar hyperkeratosis, and knuckle pads in a patient with a novel connexin 26 (GJB2) mutation. *J Med Genet* 42:e2.
- Löffler J, Nekahm D, Hirst-Stadlmann A, Günther B, Menzel HJ, Utermann G, Janecke AR. 2001. Sensorineural hearing loss and the incidence of Cx26 mutations in Austria. *Eur J Hum Genet* 9:226–230.
- Maeda S, Nakagawa S, Suga M, Yamashita E, Oshima A, Fujiyoshi Y, Tsukihara T. 2009. Structure of the connexin 26 gap junction channel at 3.5 Å resolution. *Nature* 458:597–602.
- Maestrini E, Korge BP, Ocaña-Sierra J, Calzolari E, Cambiaghi S, Scudder PM, Hovnanian A, Monaco AP, Munro CS. 1999. A missense mutation in connexin26, D66H, causes mutilating keratoderma with sensorineural deafness (Vohwinkel's syndrome) in three unrelated families. *Hum Mol Genet* 8:1237–1243.
- Mani RS, Ganapathy A, Jalvi R, Srikumari Srisailapathy CR, Malhotra V, Chadha S, Agarwal A, Ramesh A, Rangasayee RR, Anand A. 2009. Functional consequences of novel connexin 26 mutations associated with hereditary hearing loss. *Eur J Hum Genet* 17:502–509.
- Martin PEM, Evans HW. 2004. Incorporation of connexins into plasma membranes and gap junctions. *Cardiovasc Res* 62:378–387.
- Marziano NK, Casalotti SO, Portelli AE, Becker DL, Forge A. 2003. Mutations in the gene for connexin 26 (GJB2) that cause hearing loss have a dominant negative effect on connexin 30. *Hum Mol Genet* 12:805–812.
- Matos T, Caria H, Galhardo I, Dias O, Andrea M, Correia C, Fialho G. 2008. A novel M163L mutation in GJB2 gene associated with autosomal dominant isolated hearing loss. *Hear Res* 240:87–92.

- Melchionda S, Bicego M, Marciano E, Franze A, Morgutti M, Bortone G, Zelante L, Carella M, D' Andrea P. 2005. Functional characterization of novel CX26 (T55N) mutation associated to non-syndromic hearing loss. *Biochem Biophys Res Commun* 337:799–805.
- Mese G, Richard G, White TW. 2007. Gap junctions: Basic structure and function. *J Invest Dermatol* 127:2516–2524.
- Miller SA, Dykes DD, Polesky HF. 1988. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16:1215.
- Montgomery JR, White TW, Martin BL, Turner ML, Holland SM. 2004. A novel connexin 26 mutation associated with features of the keratitis-ichthyosis-deafness syndrome and the follicular occlusion triad. *J Am Dermatol* 51:377–382.
- Morlé L, Bozon M, Alloisio N, Latour P, Vandenberghe A, Plauchu H, Collet L, Edery P, Godet J, Lina-Granade G. 2000. A novel C202F mutation in the connexin26 gene (GJB2) associated with autosomal dominant isolated hearing loss. *J Med Genet* 37:368–370.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: 124500: March 13, 2001. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: 148210: September 2, 2009. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: 148350: November 13, 2008. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: 149200: October 4, 2005. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.
- Online Mendelian Inheritance in Man, OMIM (TM). Johns Hopkins University, Baltimore, MD. MIM Number: 601544: December 22, 2010. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>.
- Richard G, White TW, Smith LE, Bailey RA, Compton JG, Paul DL, Bale SJ. 1998. Functional defects of Cx26 resulting from a heterozygous missense mutation in a family with dominant deaf-mutism and palmoplantar keratoderma. *Hum Genet* 103:393–399.
- Richard G, Brown N, Smith LE, Terrinoni A, Melino G, Mackie RM, Bale SJ, Uitto J. 2000. The spectrum of mutations in erythrokeratodermias—Novel and de novo mutations in GJB3. *Hum Genet* 106:321–329.
- Richard G, Rouan F, Willoughby CE, Brown N, Chung P, Ryyänen M, Jabs EW, Bale SJ, DiGiovanna JJ, Uitto J, Russell L. 2002. Missense mutations in GJB2 encoding connexin-26 cause the ectodermal dysplasia keratitis-ichthyosis-deafness syndrome. *Am J Hum Genet* 70:1341–1348.
- Rouan F, White TW, Brown N, Taylor AM, Lucke TW, Paul DL, Munro CS, Uitto J, Hodgins MB, Richard G. 2001. Trans-dominant inhibition of connexin-43 by mutant connexin-26: Implications for dominant connexin disorders affecting epidermal differentiation. *J Cell Sci* 114: 2105–2113.
- Rubin JB, Verselis VK, Bennett MV, Bargiello TA. 1992. Molecular analysis of voltage dependence of heterotypic gap junctions formed by connexins 26 and 32. *Biophys J* 62:183–193;discussion 193–195.
- Smith RJ, Bale JF Jr, White KR. 2005. Sensorineural hearing loss in children. *Lancet* 365:879–890.
- Snoeckx RL, Hassan DM, Kamal NM, Van Den Bogaert K, Van Camp G. 2005. Mutation analysis of the GJB2 (connexin 26) gene in Egypt. *Hum Mutat* 26:60–61.
- Sonoda S, Uchino E, Sonoda KH, Yotsumoto S, Uchio E, Isashiki Y, Sakamoto T. 2004. Two patients with severe corneal disease in KID syndrome. *Am J Ophthalmol* 137:181–183.
- Tekin M, Arnos KS, Xia XJ, Oelrich MK, Liu XZ, Nance WE, Pandya A. 2001. W44C mutation in the connexin 26 gene associated with dominant non-syndromic deafness. *Clin Genet* 59:269–273.
- Uyguner O, Tukul T, Baykal C, Eris H, Emiroglu M, Hafiz G, Ghanbari A, Baserer N, Yuksel-Apak M, Wollnik B. 2002. The novel R75Q mutation in the GJB2 gene causes autosomal dominant hearing loss and palmoplantar keratoderma in a Turkish family. *Clin Genet* 62:306–309.
- van Steensel MA. 2004a. Gap junction diseases of the skin. *Am J Med Genet Part C* 131C:12–19.
- van Steensel MA, van Geel M, Nahuys M, Smitt JH, Steijlen PM. 2002. A novel connexin 26 mutation in a patient diagnosed with keratitis-ichthyosis-deafness syndrome. *J Invest Derm* 118:724–727.
- van Steensel MA, Steijlen PM, Bladergroen RS, Hoefslot EH, van Ravenswaaij-Arts CM, van Geel M. 2004b. A phenotype resembling the Clouston syndrome with deafness is associated with a novel missense GJB2 mutation. *J Invest Dermatol* 123:291–293.
- White TW, Paul DL, Goodenough DA, Bruzzone R. 1995. Functional analysis of selective interactions among rodent connexins. *Mol Biol Cell* 6:459–470.
- Yotsumoto S, Hashiguchi T, Chen X, Ohtake N, Tomitaka A, Akamatsu H, Matsunaga K, Shiraishi S, Miura H, Adachi J, Kanzaki T. 2003. Novel mutations in GJB2 encoding connexin-26 in Japanese patients with keratitis-ichthyosis-deafness syndrome. *Br J Dermatol* 148:649–653.
- Yuan Y, Huang D, Yu F, Zhu X, Kang D, Yuan H, Han D, Dai P. 2009. A de novo GJB2 (connexin 26) mutation, R75W, in a Chinese pedigree with hearing loss and palmoplantar keratoderma. *Am J Med Genet A* 149A:689–692.