

ORIGINAL  
ARTICLE

## Neuronal Nogo-A regulates glutamate receptor subunit expression in hippocampal neurons

Xiangmin Peng, Jeeyong Kim, Zhigang Zhou, David J. Fink and Marina Mata

*Department of Neurology, University of Michigan and VA Ann Arbor Healthcare System, Ann Arbor, Michigan, USA***Abstract**

Nogo-A and its cognate receptor NgR1 (NgR1) are both expressed in neurons. To explore the function of these proteins in neurons of the CNS, we carried out a series of studies using postnatal hippocampal neurons in culture. Interfering with the binding of Nogo-A to NgR1 either by adding truncated soluble fragment of NgR1 (NgSR) or by reducing NgR1 protein with a specific siRNA, resulted in a marked reduction in Nogo-A expression. Inhibition of Rho-ROCK or MEK-MAPK signaling resulted in a similar reduction in neuronal Nogo-A mRNA and protein. Reducing Nogo-A protein levels by siRNA resulted in an increase in the post-synaptic scaffolding protein PSD95, as well as increases in GluA1/GluA2 AMPA receptor and GluN1/GluN2A/GluN2B NMDA glutamate receptor su-

bunits. siRNA treatment to reduce Nogo-A resulted in phosphorylation of mTOR; addition of rapamycin to block mTOR signaling prevented the up-regulation in glutamate receptor subunits. siRNA reduction of NgR1 resulted in increased expression of the same glutamate receptor subunits. Taken together the results suggest that transcription and translation of Nogo-A in hippocampal neurons is regulated by a signaling through NgR1, and that interactions between neuronal Nogo-A and NgR1 regulate glutamatergic transmission by altering NMDA and AMPA receptor levels through an rapamycin-sensitive mTOR-dependent translation mechanism.

**Keywords:** Nogo-A, Nogo receptor, synaptic plasticity. *J. Neurochem.* (2011) **119**, 1183–1193.

Nogo-A was originally identified as a growth inhibitor. Myelin-derived Nogo-A binds preferentially to neuronal NgR1, triggering growth cone collapse and inhibition of neurite growth (Chen *et al.* 2000; Fournier *et al.* 2001). In the setting of CNS injury, Nogo-A expressed in oligodendrocytes plays important roles in blocking regeneration of central axons (Liu *et al.* 2006; Cheatwood *et al.* 2008a; Gonzenbach and Schwab 2008; Nash *et al.* 2009). Subsequent studies revealed that Nogo-A is expressed also by neurons (Huber *et al.* 2002), and neuronal Nogo-A expression is increased after injury and in some disease states; NogoA in hippocampus is increased after deafferentation or kainate-induced seizures, in human brains with Alzheimer disease or temporal lobe epilepsy, in cortical neurons after focal ischemia and in dorsal root ganglia after axonal injury (Meier *et al.* 2003; Bandtlow *et al.* 2004; Gil *et al.* 2006; Cheatwood *et al.* 2008b; Peng *et al.* 2010). We have previously reported that exposure to myelin inhibitors *in vitro* results in an increase in Nogo-A expression in dorsal root ganglion neurons; an effect mediated by NgR1 (Peng *et al.* 2010). However, reducing neuronal Nogo-A enhances growth cone motility and neurite growth, while reducing

axon branching (Craveiro *et al.* 2008; Montani *et al.* 2009; Peng *et al.* 2010; Petrinovic *et al.* 2010).

NgR1 is expressed at both pre-synaptic and post-synaptic sites (Wang *et al.* 2002; Barrette *et al.* 2007), and expression levels correlate with synaptic activity (Josephson *et al.* 2003). After spinal cord injury, enhanced synaptic plasticity in sensory cortex correlates with down-regulation of NgR1 (Endo *et al.* 2007). In visual cortex, NgR1 signaling modulates experience-dependent plasticity and the period of ocular dominance plasticity is prolonged in NgR1 null

Received May 6, 2011; revised manuscript received September 20, 2011; accepted October 4, 2011.

Address correspondence and reprint requests to Dr Marina Mata, Department of Neurology, University of Michigan and VA Ann Arbor Healthcare System, 3027 BSRB, 109 Zina Pitcher Pl, Ann Arbor, MI 48109, USA. E-mail: mmata@umich.edu

**Abbreviations used:** AMPA, 2-amino-3-(5-methyl-3-oxo-1,2-oxazol-4-yl) propionic acid; ERK, extracellular signal-related kinase; HSV, herpes simplex virus; LTD, long term depression; LTP, long term potentiation; MEK1/2, mitogen activated protein kinase kinase 1/2; mTOR, mammalian target of rapamycin; PSD95, post synaptic density protein 95; SRE, serum response element.

mice (McGee *et al.* 2005). NgR1 modulates activity dependent synaptic strength and spine morphology and Nogo66 peptide applied to CA1 dendritic field suppresses long term potentiation (LTP) in hippocampal slices (Raiker *et al.* 2010). Long term depression (LTD) is reduced and the LTP response enhanced in Schaffer collateral-CA1 of NgR1 null mice, while mice over-expressing NgR1 have impairment of long term memory (Lee *et al.* 2008; Karlen *et al.* 2009). Conversely, over-expression of Nogo-A (with Nogo-B) in cerebellar Purkinje cells results in synaptic destabilization of GABAergic terminals (Aloy *et al.* 2006).

These observations suggest that neuronal Nogo-A may play a role in regulating glutamatergic synapses. To investigate the role of neuronal Nogo-A interactions with NgR1 in the development of synapses we studied postnatal hippocampal neurons in culture. We found that Nogo-A transcription in neurons is under control of NgR1 signaling through Rho-ROCK and MAPK pathways, and reducing neuronal Nogo-A with siRNA promoted increases in NMDA and 2-amino-3-(5-methyl-3-oxo-1,2-oxazol-4-yl) propionic acid (AMPA) receptor subunit expression and dendritic post-synaptic density protein 95 (PSD95) through a mammalian target of rapamycin (mTOR) mediated and rapamycin-sensitive pathway.

## Materials and methods

### Tissue culture and *in vitro* experiments

The study was reviewed and approved by our institutional animal studies committee. Hippocampal neurons were isolated from P2 rats and cortical neurons were obtained from E17 rat pups of both sexes obtained from litters produced by timed pregnant Sprague-Dawley female rats (Charles River). The cells were cultured in defined Neurobasal medium (Gibco) containing B27, Glutamax I, Albumax I, and penicillin/streptomycin. A solution of mitotic inhibitors fluoro-2-deoxyuridine (2.5 µg/mL) and uridine (2.7 µg/mL) (Sigma, St Louis, MO, USA) was added to the cultures twice weekly. Hippocampal neurons were maintained *in vitro* for 15–19 days (DIV15–19) for these studies. ROCK inhibitor Y-27632 (Calbiochem, San Diego, CA, USA) or the highly selective and potent mitogen activated protein kinase kinase 1/2 (MEK1/2) inhibitor UO126 (Promega, Madison, WI, USA) were added for 24 h. UO126 was chosen because of its pharmacokinetic properties and minimal, if any, effect on other kinase pathways (Favata *et al.* 1998). Similarly Y-27632 has been shown to be a specific inhibitor of ROCKI/II with  $K_i$  more than 100-fold lower than those for protein kinase A (PKA), protein kinase C (PKC), myosin light chain kinase (MLCK), p21 activated kinase (PAK) and does not affect extracellular signal-related kinase (ERK) or c-Jun N-terminal kinase (JNK) activity at the concentrations employed in these studies (Uehata *et al.* 1997; Davies *et al.* 2000; Ishizaki *et al.* 2000; Narumiya *et al.* 2000). Cortical neurons DIV7 were infected for 2 h with herpes simplex virus (HSV)-based a vector we termed QHNgSR expressing the soluble fragment of NgR1 (aa 1-310; NgSR), or a vector we named QHGFP expressing green fluorescent protein (GFP) at a multiplicity of infection of 1. Media from

transfected cortical neurons, containing NgSR released from QHNgSR or from QHGFP control vector was applied to the hippocampal neurons for 24 h (Peng *et al.* 2010).

### siRNA preparation and transfection

ON-TARGET plus SMARTpool siRNA directed against Nogo-A and NgR1 (Dharmacon, Chicago, IL, USA). The siRNA sequences used for Nogo-A were as follows: sequence 1, 5'-CCAAAUCACU-ACGAAAGA-3'; sequence 2, 5'-UCUAGAAGUAUCCGACAAA-3'; sequence 3, 5'-GAAUGAAGCCACAGGUACA-3'; sequence 4, 5'-GAAUAAAGGACUCGGGGAA-3'; The siRNA sequences used for NgR1 were as follows: sequence 1, 5'-GCCACGGCACAUC-CAAUGA-3'; sequence 2, 5'-AGAAAGAACCGCACCCGUA-3'; sequence 3, 5'-CUGCAGAAGUCCGAGGUU-3'; sequence 4, 5'-GGAAGUGGGAGCAGUGGAA-3'; ON-TARGET plus siCONTROL non-targeting pool siRNA (Dharmacon) was used as control. For siRNA transfection, 2.5 µL of siRNA in 47.5 µL of antibiotic-free cultured medium and 2 µL of DharmaFECT siRNA transfection reagent 3 (Dharmacon) in 48 µL of cultured medium were incubated for 20 min at 21°C. Hippocampal neurons (DIV15) were treated with siRNA transfection solution for 72 h.

### RT-PCR

RNA was isolated from hippocampal neurons using TRIzol (Invitrogen, Carlsbad, CA, USA). cDNA was amplified using the following primer sets: β-actin forward, 5'-CAG TTC GCC ATG GAT GAC GAT ATC-3', and β-actin reverse, 5'-CAC GCT CGG TCA GGA TCT TCA TG-3'; Nogo-A forward, 5'-GAG ACC CTT TTT GCT CTT CCT G-3', and Nogo-A reverse, 5'-AAT GAT GGG CAA AGC TGT GCT G-3'. All reactions involved initial denaturation at 94°C for 5 min followed by 24 cycles (94°C for 30 s, 68°C for 2 min, and 1 cycle 68°C for 8 min using a GeneAmp PCR 2700 (Applied Biosystems, Carlsbad, CA, USA). No changes in β-actin mRNA levels were observed across different experimental groups and β-actin was used as internal control.

### Western blot

Hippocampal neurons were collected in the lysis buffer (50 mM Tris, 10 mM NaCl, 1% Nonidet P-40, 0.02% NaN<sub>3</sub>, protease inhibitor, and phosphatase inhibitor mixtures) and western blot performed as described (Peng *et al.* 2010). Primary antibodies included the following: anti-NogoR that recognizes both full-length and truncated soluble NgR1 (R&D Systems, Minneapolis, MN, USA), anti-Nogo-A (Santa Cruz Biotechnology, Santa Cruz, CA, USA) anti-PSD95 (UC Davis/NIH NeuroMab, Davis, CA, USA), anti-NR1 (GluN1; Millipore Corporation, Bedford, MA, USA), anti-NR2A (GluN2A; Millipore), anti-NR2B (GluN2B; UC Davis/NIH NeuroMab), anti-GluR1 (GluA1; Santa Cruz Biotechnology), anti-GluR2 (GluA2; UC Davis/NIH NeuroMab), anti-synapsin I (Millipore), anti-synaptophysin (1 : 1000; Millipore), anti-mTOR (Cell Signaling Technology, Beverly, MA, USA), anti-phospho-mTOR (Cell Signaling) followed by peroxidase-coupled secondary antibodies (Santa Cruz Biotechnology) and amplified using SuperSignal chemiluminescence reagent (Pierce, Rockford, IL, USA). The membranes were then stripped and reprobed with β-actin antibody (Santa Cruz Biotechnology). Quantitation of chemiluminescence emitted from each protein band was performed using ChemiDoc (Bio-Rad Laboratories, Hercules, CA, USA). No changes in β-actin levels in 20 µg of protein load were

observed in cell homogenates across different treatment groups and  $\beta$ -actin was used as internal control.

### Immunocytochemistry

Hippocampal neurons were fixed, blocked, and probed overnight with the following antibodies anti-Nogo-A (Santa Cruz Biotechnology), anti-NogoR (R&D), anti-NeuN (Millipore), anti-MAP2 (Millipore), anti-MAP2 (Sigma-Aldrich), anti-SMI31 (Covance, Princeton, NJ, USA), anti-synaptophysin (Millipore), anti-PSD95 (UC Davis/NIH NeuroMab), anti-GluR2 (GluA2; Santa Cruz), anti-NR2A (GluN2A; Millipore), vGluT1 (Santa Cruz Biotechnology), glial fibrillary acidic protein (GFAP) (Sigma) or myelin basic protein (MBP) (Millipore). The secondary antibodies were tagged-Alexa Fluor 594 or 488 (Invitrogen). Dendrites from randomly selected hippocampal neurons treated with Nogo-A siRNA or control siRNA were selected for quantitation of boutons. Three or four dendrites per each neuron were selected and measured in length starting at 10  $\mu\text{m}$  distance from the neuronal cell body to the endpoint. Total dendrite length analyzed was 600  $\mu\text{m}$  per treatment condition. The number of GluN2A and PSD95 immunofluorescent punctae along the dendrites stained with MAP2 antibody were counted, and the value represented as number of boutons per 10  $\mu\text{m}$  of dendrite length per treatment condition.

### Data analysis


The statistical significance of the difference between treated and control groups was determined by independent sample *t*-test except for data presented in Fig. 4a which was analyzed by one-way ANOVA using SPSS 12.0 for Windows (SPSS Inc. Armonk, NY, USA). Data are expressed as means  $\pm$  SE, with  $p < 0.05$  considered significant.

All the experiments were performed independently at least three times.

## Results


### Transcriptional control of Nogo-A by NgR1 in hippocampal neurons

To evaluate the regulation and function of neuronal Nogo-A in the absence of myelin inhibition we used primary postnatal hippocampal neurons and the cellular composition of the cultures was assessed by immunostaining. Approximately 80% of the neurons as determined by the neuronal marker NeuN express the vesicular glutamate transporter 1 consistent with a glutamatergic phenotype (Fig. 1a). Astrocyte visualized by glial fibrillary acidic protein (GFAP) staining accounted for less than 8% of the cells (Fig. 1b). No oligodendrocytes were detected using antibodies to myelin basic protein (MBP) or myelin associated glycoprotein (MAG). Hippocampal neurons (DIV15-17) have committed dendritic-axonal differentiation and elaborate extensive processes that form synaptic contacts as indicated by punctuate distribution of synaptophysin on dendrites (Fig. 1c). Nogo-A and NgR1 were both detected in neuronal cell bodies and dendrites indicated by co-localization with MAP-2 immunostaining, extending the length of the dendrites (Fig. 2a). Nogo-A and NgR1 were also found in axons as determined by co-localization with SMI31, a marker of phosphorylated neurofilaments and tau



**Fig. 1** Cellular composition of hippocampal cultures. P2 hippocampal neurons cultured in the presence of mitotic inhibitors as described in Methods. (a) Vesicular glutamate transporter 1 (red) protein expression was seen in 80% of neurons stained with NeuN antibody (green). Arrow shows neuron without vGluT1 staining in the field. Scale bar = 50  $\mu\text{m}$ . (b)

Less than 10% of all cells were astrocytes characterized by GFAP staining (green) and MAP2 (red). Scale bar = 100  $\mu\text{m}$ . (c) Hippocampal neurons (DIV15) have a mature phenotype with axon dendritic differentiation and widespread synaptic contacts. Immunostaining with MAP2 (red) and synaptophysin (green) as indicated. Scale bar = 20  $\mu\text{m}$ .



**Fig. 2** Localization of Nogo-A and NgR1 in hippocampal neurons. Hippocampal neurons (DIV15) show expression of Nogo-A and NgR1 in dendrites and axons. (a) Nogo-A and NgR1 immunostaining in cell body and dendrites but also extending beyond the limits

of MAP2 staining. Nogo-A (red) with MAP2 (green) and NgR (red) with MAP2 (green). Scale bar = 20  $\mu$ m. (b) Nogo-A and NgR (red) are seen in most axons stained with SMI31 (red). Scale bar = 20  $\mu$ m.


(Fig. 2b). Staining for NogoA along axons appeared more intense than for NgR1 but accurate quantitation cannot be made because of differences in antibody source and binding affinities, and because variation in staining intensity between different fibers, seen also with SMI31 antibody, may represent the number of axons forming the bundle. There was no detectable Nogo-A in astrocytes and there were no oligodendroglial cells in these hippocampal cultures (data not shown).

We exposed hippocampal neurons to Nogo-soluble receptor (NgSR) that recognizes the Nogo66 domain of Nogo-A, produced by release from HSV vector (QHNgSR) transfection of cortical neurons (Peng *et al.* 2010). NgSR treatment resulted in a significant down-regulation of Nogo-A mRNA and protein (Fig. 3a and b). To determine if the effect of NgSR was caused by capping of Nogo-A and internalization or by signal derived from receptor occupancy we used an RNAi approach. Treatment of hippocampal neurons with an NgR1-specific siRNA resulted in a 70% reduction in NgR1 protein as compared with neurons transfected with control siRNA (Fig. 4b). Reduction of NgR1 was accompanied by a


marked decrease in Nogo-A mRNA and protein (Fig. 4a and b). Taken together, these experiments indicate that expression of Nogo-A in hippocampal neurons is regulated, in part, by signals that derived from interactions of neuronal-expressed Nogo-A with NgR1.

#### Rho-ROCK and MAPK pathways converge in the regulation of neuronal Nogo-A expression

NgR1 signals through Rho-GTPase and Rho activated kinase (ROCK) to regulate actin polymerization, microtubule stabilization, and changes in transcriptional activity (Arimura *et al.* 2000; Fournier *et al.* 2001; Peng *et al.* 2010). To assess whether a similar signaling cascade might be involved in NgR1 regulation of Nogo-A expression in hippocampal neurons, we added the ROCK inhibitor Y-27632 for 24 h. ROCK inhibition resulted in a marked decrease in Nogo-A levels (Fig. 5a). ROCK inhibition also significantly lowered the basal activity of ERK1/2 in the treated neurons (Fig. 5b), suggesting that the MAPK pathway lies downstream of ROCK. Expression of Nogo-A mRNA and protein was also decreased by treatment of the hippocampal neurons with the



**Fig. 3** Soluble NgR1 reduces Nogo-A expression. Hippocampal neurons treated with or without NgSR released from QHNgSR for 24 h. NgSR decreases Nogo-A mRNA (a) and protein (b) as compared with control (QHGFP). Each sample was calculated as a ratio to β-actin and presented as percentage of control. \*\**p* < 0.01.




**Fig. 5** Inhibition of Rho-dependent kinase decreases Nogo-A levels and inhibits MAPK. Addition of 10 μM Y27632 for 24 h decreases Nogo-A (a) and pERK level (b). Each sample was calculated as a ratio to β-actin or total ERK and presented as percentage of control. \**p* < 0.05.

MEK inhibitor UO126 for 24 h (Fig. 6a and b) that inhibited phosphorylation of ERK (Fig. 6c) indicating that Rho-ROCK and MAPK signaling pathways converge to regulate neuronal Nogo-A expression.


**Regulation of NMDA and AMPA receptors by Nogo-A signaling**

To evaluate the possibility that Nogo-A-NgR1 interactions known to alter synaptic plasticity may regulate glutamate

receptor expression levels in differentiated postnatal hippocampal neurons, we designed an siRNA specific for Nogo-A and evaluated synaptic markers and glutamate receptor subunit proteins. Addition of Nogo-A siRNA to hippocampal neurons decreased Nogo-A protein without any change in the level of NgR1 (Fig. 7a), and resulted in increases in NMDA receptor GluN1, GluN2A and GluN2B subunit levels (Fig. 7b). The post-synaptic scaffolding protein PSD95, was increased in the Nogo-A siRNA-treated neurons



**Fig. 4** NgR1 knock-down causes a reduction of Nogo-A. Hippocampal neurons treated with control scrambled siRNA (csiRNA) or NgR1 siRNA for 72 h. NgR siRNA decreases NgR1 protein (b). Nogo-A mRNA (a) and protein (b) were decreased in NgR1 siRNA as compared with csiRNA. Each sample was calculated as a ratio to β-actin and presented as percentage of control. \**p* < 0.05.



**Fig. 6** Inhibition of MEK-MAPK reduces Nogo-A expression. Addition of 10  $\mu$ M UO126 for 24 h reduces Nogo-A mRNA (a) and protein (b). UO126 prevented the phosphorylation of ERK (c). Each sample was calculated as a ratio to  $\beta$ -actin or total ERK and presented as percentage of control. \* $p < 0.05$ .

(Fig. 7b). The ratio of GluN2A/GluN2B was not altered by knockdown of Nogo-A expression. Decreasing levels of Nogo-A by siRNA treatment of hippocampal neurons also resulted in an increase in AMPA receptor subunits GluA1 and GluA2 (Fig. 7b). By immunocytochemistry, representative subunits of AMPA and NMDA receptors as well as PSD95 showed punctuate staining along MAP2-positive dendrites (Fig. 7c and d) consistent with localization at synapses. Although we did not directly assess the relative levels of each subunit in different cellular compartments after siRNA treatment, the effect appeared to be most likely post-synaptic as knockdown of Nogo-A in hippocampal neurons by siRNA treatment did not result in any change in the amount of the SNARE proteins synapsin I and synaptophysin or the level of the vesicular glutamate transporter isoforms vGlut1 and vGlut2 (data not shown). This was supported by quantitative analysis of the number of GluN2A and PSD95 synaptic boutons on MAP2 stained dendrites that were significantly increased in Nogo-A siRNA treated neurons as compared with control siRNA (GluN2A per 10  $\mu$ m =  $3.12 \pm 0.13$  in

Nogo-A siRNA and  $2.63 \pm 0.15$  in control scrambled siRNA \* $p < 0.05$ ; PSD95 per 10  $\mu$ m =  $3.25 \pm 0.14$  in Nogo-A siRNA and  $2.46 \pm 0.05$  in control scrambled siRNA \* $p < 0.05$ ).

#### Rapamycin-sensitive mTOR pathway in the regulation of NMDA/AMPA by Nogo-A


mTOR and its downstream effectors eIF-4E-BP and eIF-4E, are localized in dendritic spines of hippocampal neurons, and play important roles in the development of late LTP (Vickers *et al.* 2005; Bekinschtein *et al.* 2007; Bramham and Wells 2007). mTOR activity measured by the amount of phospho-mTOR compared with total mTOR, was increased in hippocampal neuron cultures in which Nogo-A levels have been reduced by siRNA (Fig. 7e). Treatment of the neurons for 24 h with rapamycin blocked mTOR phosphorylation (Fig. 8a) and prevented the increase in NMDA and AMPA subunits and PSD95 protein levels resulting from knocking down Nogo-A by siRNA (Fig. 8b). Rapamycin treatment did not significantly change the levels of NMDA or AMPA receptor subunits in neurons treated with control siRNA (Fig. 8c). The results suggest that neuronal Nogo-A, regulation of the activity state of the rapamycin-sensitive mTOR-signaling pathway is important for the translation and/or maintenance of glutamate receptor protein subunits in hippocampal neurons.

#### Is regulation of NMDA and AMPA receptors by Nogo-A dependent on Ngr1?

NgR1 has been implicated in synaptic plasticity using transgenic models of NgR1 deletion and over-expression. To address if the regulation of NMDA and AMPA receptor subunit expression by Nogo-A was mediated through NgR1 signaling we used siRNA to decrease NgR1 levels in hippocampal neurons. We found an increase in NMDA and AMPA receptor subunits and PSD95 protein in neurons treated with NgR1 siRNA that was similar to the changes caused by Nogo-A knock down, but not in neurons treated with a scrambled sequence control siRNA (Fig. 9). As the down-regulation of NgR1 was accompanied by simultaneous decrease in Nogo-A levels (Fig. 4) the effects of NgR1 could not be fully dissociated from Nogo-A, but taken together these results suggest that decreased Nogo-A-NgR1 interaction as a result of reductions in either Nogo-A or NgR1 would favor enhanced glutamatergic transmission.


#### Discussion

The studies reported demonstrate four principal findings: (i) Nogo-A expression in neurons is under transcriptional control of NgR1; (ii) Nogo-A expression is regulated through Rho-ROCK and MEK-MAPK pathways; (iii) neuronal Nogo-A regulates NMDA and AMPA glutamate receptor subunit expression; (iv) Nogo-A regulation of NMDA and



**Fig. 7** Down-regulation of Nogo-A increases NMDA and AMPA receptor subunits and PSD95. Hippocampal neurons (DIV15) treated with control siRNA or Nogo-A siRNA. (a) Nogo-A siRNA significantly decreased Nogo-A without altering NogoR levels. (b) Nogo-A siRNA caused increase in GluN1, GluN2A and GluN2B, PSD95, GluR1 and GluR2. (c) Neurons treated with Nogo-A siRNA show den-

drogic synaptic immunolocalization of GluA2 (red) with PSD95 (green), of PSD95 (green) with MAP2 (red) and GluN2A (red) with MAP2 (green). (d) Similar neurons treated with control siRNA and stained as in panel c. Scale bar = 30  $\mu$ m. (e) Nogo-A siRNA caused increase in p-mTOR as compared with control siRNA. Each sample was calculated as a ratio to  $\beta$ -actin or total mTOR and presented as percentage of control. \* $p < 0.05$ .



**Fig. 8** Reversal of NMDA, AMPA subunits and PSD95 expression by rapamycin in Nogo-A knock-down. Addition of 100 nM rapamycin to hippocampal neurons transfected with Nogo-A siRNA prevented the phosphorylation of mTOR (a) and resulted in decreased GluN1, GluN2A, GluN2B, PSD95, GluA1, and GluA2 (b). Addition of

100 nM rapamycin transfected with control siRNA lowered p-mTOR below basal level but did not significantly change NMDA and AMPA subunits or PSD95 levels (c). Each sample was calculated as a ratio to  $\beta$ -actin or total mTOR and presented as percentage of control. \* $p < 0.05$ .


AMPA receptor subunits expression is effected through the rapamycin-sensitive mTOR pathway.

The assembly and maintenance of neuronal networks involves neurite growth, synapse formation and plasticity. Recent studies suggest that Nogo-A interaction with NgR1 plays an important role in many of these cellular processes. In postnatal hippocampal neurons *in vitro*, at a time when these cells have developed a large and distinct axonal and dendritic arborization with extensive axo-dendritic synaptic

contacts we found both Nogo-A and NgR1 in axons and dendrites, and that disengagement of NgR1 from Nogo-A, by either NgSR interference or by siRNA knockdown of NgR1, resulted in a reduction of Nogo-A mRNA and protein.

These results suggest the existence of a feedback loop in which interaction of NgR1 with one of its ligands regulates the level of Nogo-A. Ligand binding to NgR1 activates Rho-GTPases and downstream Rho-dependent kinase to mediate cellular responses responsible for growth cone collapse and





**Fig. 9** Down-regulation of NgR1 by siRNA increases NMDA and AMPA receptor subunits and PSD95. P2 hippocampal neurons cultured for 2 weeks and treated with csiRNA or NogoR1 siRNA for 72 h.

NgoR1 siRNA treatment results in increased GluN1, GluN2A, GluN2B, PSD95, GluA1 and GluA2. Each sample was calculated as a ratio to  $\beta$ -actin and presented as percentage of control. \* $p < 0.05$ .

axon growth inhibition. In hippocampal neurons, inhibition of ROCK activity diminished Nogo-A expression and reduced the basal state of ERK phosphorylation and inhibition of the MEK-MAPK signaling also resulted in a decrease in Nogo-A expression. These results suggest that, in neurons, ROCK and ERK pathways converge to regulate Nogo-A expression. Rho-ROCK signaling regulates transcription through the binding of serum response factor on serum response element (SRE). The serum response factor-SRE cassette utilizes as binding partners the transcription factors ternary complex factors, downstream of ERK, and the actin sensor megakaryocyte leukemia (MKL) factor, downstream from RhoA to regulate early and late gene transcription necessary for essential cellular responses including neurite growth, axonal regeneration and synapse formation (Knoll and Nordheim 2009). The regulation of Nogo-A by Rho-ROCK and MEK-MAPK pathways is consistent with the convergence of these signaling pathways on SRE. The observed regulation of Nogo-A by pathways downstream of NgR1 does not exclude the possibility of similar or other effects that may be triggered by other myelin inhibitors [OMgp, myelin associated glycoprotein (MAG)] through binding to NgR1.

In the hippocampal neuron culture system, down-regulation of Nogo-A resulted in increased expression of NMDA (GluN1, GluN2A and GluN2B) and AMPA (GluA1 and GluA2) receptor subunits and the post-synaptic anchoring protein PSD95. The effects of Nogo-A down-regulation appear to be post-synaptic, as there were no changes in pre-synaptic proteins (synapsin, synaptophysin vGlu1 or vGlu2), and there were no significant changes in neurite growth (Peng *et al.* 2010). These effects were mediated by

mTOR signaling, as blocking the mTOR pathway with rapamycin prevented the effects of knocking down Nogo-A on the levels of NMDA and AMPA receptors. mTOR is found in dendritic spines where it regulates the initiation of translation from local polyribosomal mRNAs to produce changes in synaptic function (Bourne *et al.* 2007) and where mTOR mediates translation of NMDA and AMPA receptors (Slipczuk *et al.* 2009). In dendritic spines, changes in synapse structure and function that occur dependent or independent of electrical activity, may persist for hours or days (Saneyoshi *et al.* 2010) These results suggest that enhanced mTOR activity by reduced Nogo-A signaling results in an increase in rapamycin-sensitive translation of NMDA and AMPA receptor subunits.

Dendritic spines are highly motile structures (Fischer *et al.* 2000; Yuste and Bonhoeffer 2001) and changes in spine morphology and development of LTP/LTD reflects multiple processes including protein synthesis from local mRNAs in dendrites, trafficking of synaptic proteins and incorporation of new neurotransmitter receptor subunits (Malenka and Bear 2004; Lin *et al.* 2005). As dendritic spines undergo actin-dependent shape changes regulated by glutamate neurotransmission, a two-step process has been proposed in which synaptic spines are formed in response to NMDA receptor activation and spines are subsequently stabilized by AMPA receptor-mediated transmission (Fischer *et al.* 2000); the enhanced translation of NMDA and AMPA demonstrated by suppression Nogo-A-NgR1 signaling could contribute to changes in spine morphology. Considered in light of studies demonstrating that Nogo-A is involved in altering conformation of membrane structures (Voeltz *et al.* 2006), the

observations that NgR1 knock-out mice display abnormal spines in apical dendrites of hippocampal CA1 neurons (Lee *et al.* 2008) and that neutralization of Nogo-A function alters dendritic and spine morphology in hippocampal pyramidal neurons (Zagrebelsky *et al.* 2010), our results suggest that neuronal Nogo-A-NgR1 signaling may be involved in the refinement of glutamatergic connections in hippocampal neurons. These data are complementary to published observations that describe modulatory effects of Nogo-A and NgR1 on synaptic plasticity: the observation of NgR1-dependent Nogo66 suppression of NMDA-mediated LTP (Raiker *et al.* 2010) and with the recent report that interfering with NgR1 or Nogo-A by function blocking antibodies increases LTP without affecting basal synaptic transmission, LTD or other measures of pre-synaptic function (Delekate *et al.* 2011).

## Acknowledgements

We acknowledge the technical support of Vikram Thakur for the purification of the HSV vectors used in the study. The work was supported by grants to MM and DJF from the National Institutes of Health and the Department of Veterans Affairs. The authors have no competing interests.

## References

- Aloy E. M., Weinmann O., Pot C., Kasper H., Dodd D. A., Rulicke T., Rossi F. and Schwab M. E. (2006) Synaptic destabilization by neuronal Nogo-A. *Brain Cell Biol.* **35**, 137–156.
- Arimura N., Inagaki N., Chihara K., Menager C., Nakamura N., Amano M., Iwamatsu A., Goshima Y. and Kaibuchi K. (2000) Phosphorylation of collapsin response mediator protein-2 by Rho-kinase. Evidence for two separate signaling pathways for growth cone collapse. *J. Biol. Chem.* **275**, 23973–23980.
- Bandtlow C. E., Dlaska M., Pirker S., Czech T., Baumgartner C. and Sperk G. (2004) Increased expression of Nogo-A in hippocampal neurons of patients with temporal lobe epilepsy. *Eur. J. Neurosci.* **20**, 195–206.
- Barrette B., Vallieres N., Dube M. and Lacroix S. (2007) Expression profile of receptors for myelin-associated inhibitors of axonal regeneration in the intact and injured mouse central nervous system. *Mol. Cell. Neurosci.* **34**, 519–538.
- Bekinschtein P., Katze C., Slipczuk L. N., Igaz L. M., Cammarota M., Izquierdo I. and Medina J. H. (2007) mTOR signaling in the hippocampus is necessary for memory formation. *Neurobiol. Learn. Mem.* **87**, 303–307.
- Bourne J. N., Sorra K. E., Hurlburt J. and Harris K. M. (2007) Polyribosomes are increased in spines of CA1 dendrites 2 h after the induction of LTP in mature rat hippocampal slices. *Hippocampus* **17**, 1–4.
- Bramham C. R. and Wells D. G. (2007) Dendritic mRNA: transport, translation and function. *Nat. Rev. Neurosci.* **8**, 776–789.
- Cheatwood J. L., Emerick A. J. and Kartje G. L. (2008a) Neuronal plasticity and functional recovery after ischemic stroke. *Top Stroke Rehabil.* **15**, 42–50.
- Cheatwood J. L., Emerick A. J., Schwab M. E. and Kartje G. L. (2008b) Nogo-A expression after focal ischemic stroke in the adult rat. *Stroke* **39**, 2091–2098.
- Chen M. S., Huber A. B., van der Haar M. E., Frank M., Schnell L., Spillmann A. A., Christ F. and Schwab M. E. (2000) Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature* **403**, 434–439.
- Craveiro L. M., Hakkoum D., Weinmann O., Montani L., Stoppini L. and Schwab M. E. (2008) Neutralization of the membrane protein Nogo-A enhances growth and reactive sprouting in established organotypic hippocampal slice cultures. *Eur. J. Neurosci.* **28**, 1808–1824.
- Davies S. P., Reddy H., Caivano M. and Cohen P. (2000) Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem. J.* **351**, 95–105.
- Delekate A., Zagrebelsky M., Kramer S., Schwab M. E. and Korte M. (2011) NogoA restricts synaptic plasticity in the adult hippocampus on a fast time scale. *Proc. Natl. Acad. Sci. USA* **108**, 2569–2574.
- Endo T., Spenger C., Tominaga T., Brene S. and Olson L. (2007) Cortical sensory map rearrangement after spinal cord injury: fMRI responses linked to Nogo signalling. *Brain* **130**, 2951–2961.
- Favata M. F., Horiuchi K. Y., Manos E. J. *et al.* (1998) Identification of a novel inhibitor of mitogen-activated protein kinase kinase. *J. Biol. Chem.* **273**, 18623–18632.
- Fischer M., Kaech S., Wagner U., Brinkhaus H. and Matus A. (2000) Glutamate receptors regulate actin-based plasticity in dendritic spines. *Nat. Neurosci.* **3**, 887–894.
- Fournier A. E., GrandPre T. and Strittmatter S. M. (2001) Identification of a receptor mediating Nogo-66 inhibition of axonal regeneration. *Nature* **409**, 341–346.
- Gil V., Nicolas O., Mingorance A., Urena J. M., Tang B. L., Hirata T., Saez-Valero J., Ferrer I., Soriano E. and del Rio J. A. (2006) Nogo-A expression in the human hippocampus in normal aging and in Alzheimer disease. *J. Neuropathol. Exp. Neurol.* **65**, 433–444.
- Gonzenbach R. R. and Schwab M. E. (2008) Disinhibition of neurite growth to repair the injured adult CNS: focusing on Nogo. *Cell. Mol. Life Sci.* **65**, 161–176.
- Huber A. B., Weinmann O., Brosamle C., Oertle T. and Schwab M. E. (2002) Patterns of Nogo mRNA and protein expression in the developing and adult rat and after CNS lesions. *J. Neurosci.* **22**, 3553–3567.
- Ishizaki T., Uehata M., Tamechika I., Keel J., Nonomura K., Maekawa M. and Narumiya S. (2000) Pharmacological properties of Y-27632, a specific inhibitor of rho-associated kinases. *Mol. Pharmacol.* **57**, 976–983.
- Josephson A., Trifunovski A., Scheele C., Widenfalk J., Wahlestedt C., Brene S., Olson L. and Spenger C. (2003) Activity-induced and developmental downregulation of the Nogo receptor. *Cell Tissue Res.* **311**, 333–342.
- Karlen A., Karlsson T. E., Mattsson A. *et al.* (2009) Nogo receptor 1 regulates formation of lasting memories. *Proc. Natl. Acad. Sci. USA* **106**, 20476–20481.
- Knoll B. and Nordheim A. (2009) Functional versatility of transcription factors in the nervous system: the SRF paradigm. *Trends Neurosci.* **32**, 432–442.
- Lee H., Raiker S. J., Venkatesh K., Geary R., Robak L. A., Zhang Y., Yeh H. H., Shrager P. and Giger R. J. (2008) Synaptic function for the Nogo-66 receptor NgR1: regulation of dendritic spine morphology and activity-dependent synaptic strength. *J. Neurosci.* **28**, 2753–2765.
- Lin B., Kramar E. A., Bi X., Brucher F. A., Gall C. M. and Lynch G. (2005) Theta stimulation polymerizes actin in dendritic spines of hippocampus. *J. Neurosci.* **25**, 2062–2069.

- Liu B. P., Cafferty W. B., Budel S. O. and Strittmatter S. M. (2006) Extracellular regulators of axonal growth in the adult central nervous system. *Philos. Trans. R Soc. Lond. B Biol. Sci.* **361**, 1593–1610.
- Malenka R. C. and Bear M. F. (2004) LTP and LTD: an embarrassment of riches. *Neuron* **44**, 5–21.
- McGee A. W., Yang Y., Fischer Q. S., Daw N. W. and Strittmatter S. M. (2005) Experience-driven plasticity of visual cortex limited by myelin and Nogo receptor. *Science* **309**, 2222–2226.
- Meier S., Brauer A. U., Heimrich B., Schwab M. E., Nitsch R. and Savaskan N. E. (2003) Molecular analysis of Nogo expression in the hippocampus during development and following lesion and seizure. *FASEB J.* **17**, 1153–1155.
- Montani L., Gerrits B., Gehrig P., Kempf A., Dimou L., Wollscheid B. and Schwab M. E. (2009) Neuronal Nogo-A modulates growth cone motility via Rho-GTP/LIMK1/cofilin in the unlesioned adult nervous system. *J. Biol. Chem.* **284**, 10793–10807.
- Narumiya S., Ishizaki T. and Uehata M. (2000) Use and properties of ROCK-specific inhibitor Y-27632. *Methods Enzymol.* **325**, 273–284.
- Nash M., Pribiag H., Fournier A. E. and Jacobson C. (2009) Central nervous system regeneration inhibitors and their intracellular substrates. *Mol. Neurobiol.* **40**, 224–235.
- Peng X., Zhou Z., Hu J., Fink D. J. and Mata M. (2010) Soluble Nogo receptor down-regulates expression of neuronal Nogo-A to enhance axonal regeneration. *J. Biol. Chem.* **285**, 2783–2795.
- Petrinovic M. M., Duncan C. S., Bourikas D., Weinman O., Montani L., Schroeter A., Maerki D., Sommer L., Stoekli E. T. and Schwab M. E. (2010) Neuronal Nogo-A regulates neurite fasciculation, branching and extension in the developing nervous system. *Development* **137**, 2539–2550.
- Raiker S. J., Lee H., Baldwin K. T., Duan Y., Shrager P. and Giger R. J. (2010) Oligodendrocyte-myelin glycoprotein and Nogo negatively regulate activity-dependent synaptic plasticity. *J. Neurosci.* **30**, 12432–12445.
- Saneyoshi T., Fortin D. A. and Soderling T. R. (2010) Regulation of spine and synapse formation by activity-dependent intracellular signaling pathways. *Curr. Opin. Neurobiol.* **20**, 108–115.
- Slipczuk L., Bekinschtein P., Katche C., Cammarota M., Izquierdo I. and Medina J. H. (2009) BDNF activates mTOR to regulate GluR1 expression required for memory formation. *PLoS ONE* **4**, e6007.
- Uehata M., Ishizaki T., Satoh H. *et al.* (1997) Calcium sensitization of smooth muscle mediated by a Rho-associated protein kinase in hypertension. *Nature* **389**, 990–994.
- Vickers C. A., Dickson K. S. and Wyllie D. J. (2005) Induction and maintenance of late-phase long-term potentiation in isolated dendrites of rat hippocampal CA1 pyramidal neurones. *J. Physiol.* **568**, 803–813.
- Voeltz G. K., Prinz W. A., Shibata Y., Rist J. M. and Rapoport T. A. (2006) A class of membrane proteins shaping the tubular endoplasmic reticulum. *Cell* **124**, 573–586.
- Wang X., Chun S. J., Treloar H., Vartanian T., Greer C. A. and Strittmatter S. M. (2002) Localization of Nogo-A and Nogo-66 receptor proteins at sites of axon-myelin and synaptic contact. *J. Neurosci.* **22**, 5505–5515.
- Yuste R. and Bonhoeffer T. (2001) Morphological changes in dendritic spines associated with long-term synaptic plasticity. *Annu. Rev. Neurosci.* **24**, 1071–1089.
- Zagrebelsky M., Schweigreiter R., Bandtlow C. E., Schwab M. E. and Korte M. (2010) Nogo-A stabilizes the architecture of hippocampal neurons. *J. Neurosci.* **30**, 13220–13234.