

# Structures and processes of care in ambulatory oncology settings and nurse-reported exposure to chemotherapy

Christopher R Friese,<sup>1</sup> Laurel Himes-Ferris,<sup>2</sup> Megan N Frasier,<sup>3</sup> Marjorie C McCullagh,<sup>4</sup> Jennifer J Griggs<sup>2,5</sup>

► An additional appendix is published online only. To view this file please visit the journal online (<http://dx.doi.org/10.1136/bmjqs-2011-000178>).

<sup>1</sup>Division of Nursing Business and Health Systems, University of Michigan School of Nursing, Ann Arbor, Michigan, USA

<sup>2</sup>University of Michigan School of Public Health, Ann Arbor, Michigan, USA

<sup>3</sup>University of Michigan College of Literature, Science, and Arts, Ann Arbor, Michigan, USA

<sup>4</sup>Division of Health Promotion and Risk Reduction, University of Michigan School of Nursing, Ann Arbor, Michigan, USA

<sup>5</sup>Division of Haematology/Oncology, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, Michigan, USA

## Correspondence to

Dr Christopher R Friese, Nursing Business and Health Systems, University of Michigan School of Nursing, 400 N. Ingalls 4162, Ann Arbor, MI 48109-5482, USA; [cfriese@umich.edu](mailto:cfriese@umich.edu)

Accepted 6 July 2011  
Published Online First  
16 August 2011

## ABSTRACT

**Purpose:** Oncology nurses in ambulatory settings are at increased risk for unintentional chemotherapy exposure due to the large volumes of agents delivered and the absence of regulatory enforcement. Given the limited data regarding the correlates of exposure, the authors sought to identify the relationship between the organisational structures and processes of care in ambulatory oncology settings associated with increased risk of unintentional chemotherapy.

**Methods:** Between April 2010 and June 2010, a state-wide sample of oncology nurses were surveyed who reported their employment outside of hospital inpatient units (n=1339). The survey examined the likelihood of self-reported exposure to chemotherapy as a function of perceived quality of the practice environment, nursing workload, and seven ambulatory chemotherapy administration safety standards.

**Results:** The response rate was 30.4%, with minimal demographic differences observed between respondents and non-respondents. The overall rate of exposure to the skin or eyes in the past year was 16.9%. In multivariable logistic regression models that controlled for demographic characteristics and clustering of nurses in practices, the likelihood of exposure decreased when nurses reported adequate staffing and resources (OR 0.35, 95% CI 0.17 to 0.73; p=0.001), and when nurses reported that chemotherapy doses were verified by two nurses frequently or very frequently (OR 0.17, 95% CI 0.05 to 0.59; p=0.001).

**Conclusions:** Oncology nurses in the ambulatory setting report substantial unintentional skin and eye exposure to chemotherapy. Ensuring adequate staffing and resources and adherence to recognised practice standards may protect oncology nurses from harm.

risks to patients and clinicians. In the United States, an estimated 23 million adult patient visits occur annually for chemotherapy. Of these visits, approximately 19 million (84%) are delivered in ambulatory settings, largely by nurses.<sup>1</sup> And despite numerous concerns for chemotherapy safety, less attention is paid to the risks that nurses face when they administer these agents and are unintentionally exposed because of a splash or spill. The antineoplastic drugs used for chemotherapy (eg, cyclophosphamide, ifosfamide, paclitaxel, methotrexate) confer significant health risks, such as immediate nervous system effects, acute and long-term reproductive effects, and subsequent risk of haematological malignancies.<sup>2 3</sup>

The absence of a consistent regulatory framework for chemotherapy delivery in ambulatory oncology settings exacerbates the potential risk to nurses. Apart from a recent Centres for Medicare and Medicaid Services statement<sup>4</sup> that a 'physician or non-physician practitioner's recurrent physical presence' is required during chemotherapy administration, Centres for Medicare and Medicaid Services performs little oversight of chemotherapy delivery. The National Institute for Occupational Safety and Health issued an alert on occupational exposure to antineoplastic and other hazardous drugs.<sup>5</sup> However, this alert is merely advisory and has no regulatory enforcement.

Chemotherapy administration processes vary across practices. In 2009, the Oncology Nursing Society (ONS) and American Society of Clinical Oncology (ASCO) acknowledged the potential problems imposed by practice variability and jointly issued voluntary safety standards for ambulatory chemotherapy

## INTRODUCTION

Chemotherapy administration is a high-priority target for quality improvement activities given high patient volumes and high

administration.<sup>6</sup> The adoption rate of these standards is unknown, and only 74 practices are certified currently by ASCO's Quality Oncology Practice Initiative for standard adherence.<sup>7</sup> The 17 standards assessed for Quality Oncology Practice Initiative certification do not address safe handling practices and related policies. Further, no reporting mechanism exists to track unintentional exposures to chemotherapy or the conditions under which the exposure occurred.

Noteworthy challenges to existing efforts to study unintentional exposure include measurement considerations and restricted sampling. Diverse measurement approaches and result interpretation vary across studies and hamper comparability. Studies conducted in the 1980s and 1990s examined both urine mutagenicity and drug levels excreted in urine.<sup>8–10</sup> More recently, biological measures have been correlated to healthcare workers' self reports.<sup>11</sup> Self report has been used successfully in assessing needle stick injuries to nurses.<sup>12</sup> A second limitation of the literature is the absence of multi-site, population-based samples. Because of its cost, biological sampling has been limited to convenience samples of single institutions, which limits generalisability to more diverse practices and participants.<sup>13 14</sup> Taken together, these limitations hamper examinations of variation in chemotherapy exposure across settings, and the correlates of increased exposure.

Published studies have not examined the relationship between chemotherapy exposure and the structures and processes employed by oncology practices. The decentralised nature of ambulatory chemotherapy administration hampers research efforts to study this problem. A better understanding of the organisational structures and processes of care that are associated with chemotherapy exposure to nurses in ambulatory oncology settings would inform policymakers, safety officials, managers, and clinicians on how best to reduce exposure risk and minimise harm to oncology nurses. In this context, we conducted a state-wide survey of oncology nurses employed outside of hospital inpatient units to quantify reported exposure and to identify organisational correlates of exposure.

## METHODS

### Study design and setting

Donabedian's Quality of Care Model<sup>15</sup> guided our study design. This model identified three components of quality: the organisational structures (eg, staffing, workplace context, leadership, management), the processes of care (eg, clinical and technical interventions provided), and the outcomes (eg, safety, satisfaction, and clinical endpoints). This report focuses on the organisational structures and processes of care in ambulatory

oncology settings, and their relationship to the safety outcome of nurse-reported chemotherapy exposure.

Completed between April and June, 2010, the Practice Environments of Oncology Nurses Study was a cross-sectional survey that examined nurse reports of the organisational structures, processes of care, and outcomes in ambulatory oncology settings; we have published our methodology previously.<sup>16</sup> Briefly, we used nursing registry data from one state in the southeast USA—updated biennially—to identify registered and licensed practical nurses who resided in the state and who reported a clinical specialty of oncology and a practice setting outside of hospital inpatient units (n=1339). We modified the Dillman survey method<sup>17</sup> by including a \$2 up-front incentive, a shorter interval for reminder notices, and randomised nurses to Internet versus paper survey completion. We observed no significant differences in response rates across arms.<sup>16</sup> We obtained human subjects approval from the University of Michigan's Institutional Review Board—Medicine.

### Measures

The internet and paper questionnaires had identical content. In addition to sociodemographic variables, the questionnaires included measures of organisational structure, processes of care, and perceived outcomes.

### Organisational structure

Nurses completed the Practice Environment Scale of the Nursing Work Index (PES-NWI).<sup>18</sup> The PES-NWI consists of 31 items that assess the presence of organisational features to support professional nursing practice and promote favourable patient outcomes. The original PES-NWI includes five subscales: nurse participation in hospital affairs; nursing foundations for quality of care; nurse manager leadership, ability, and support of nurses; staffing and resource adequacy; and collegial nurse—physician relationships. The instrument has published evidence of reliability (Cronbach  $\alpha$  values between 0.71 and 0.84 across subscales) and validity (higher PES-NWI scores for nurses employed in Magnet hospitals),<sup>19 20</sup> albeit in samples of inpatient nurses.

Using focus groups,<sup>21</sup> clinical nursing expert review, and cognitive interviews, we modified the items slightly for suitability to the outpatient environment. The revised PES-NWI had six subscales: the original five listed above plus the supportive relations with medical assistants subscale. The revised text of the revised items and corresponding subscales is available in the online appendix. In a previous study, we performed confirmatory factor analysis using structural equation modelling to examine model fit with revised items. In our final model specifying six constructs, a reduced set of 23 items achieved acceptable model fit, as reflected by

a comparative fit index of 0.95 and a root mean-square error of approximation of 0.057 (95% CI 0.049 to 0.064).<sup>22</sup> Cronbach  $\alpha$  coefficients for the subscales ranged from 0.80 to 0.90. Items were scored on a five-point Likert scale, ranging from 1=strongly disagree the characteristic was present in the workplace to 5=strongly agree the characteristic was present in the workplace. Subscale items were averaged for each nurse, where values above 3.0 suggested the presence of the characteristic in their workplace. Mean PES-NWI subscales were used in bivariate analysis, and later were dichotomised (>3.0 vs 3.0 and under) for use in multivariable models.

In addition to the revised PES-NWI subscales, we asked each participant an additional item: 'please describe the current practice environment for you as a nurse to delivery high-quality care'. Response options were favourable, mixed, or unfavourable.

To measure nursing workload, we used each nurse's report of the number of patients for whom they assumed the primary care on their last shift, and treated this measure as a continuous variable. This measure of nursing workload has been used in prior studies of hospital staffing and patient outcomes, and correlates well with both perceived staffing adequacy and administratively derived nurse staffing.<sup>19 23</sup>

### Processes of care

We developed seven process of care items from the 2009 ASCO/ONS standards for ambulatory chemotherapy administration.<sup>6</sup> Nurses rated the frequency that the process occurred on a five-point Likert scale (never to very frequently). Items included the presence of signed informed consent, documentation of treatment intent, and performance of pain assessments. In our exploratory data analysis, we identified only one process of care measure—the frequency that two nurses performed verification of the original chemotherapy order against the prepared dosage of chemotherapy—that was associated with the likelihood of chemotherapy exposure. We treated this dichotomously (verified chemotherapy order and dosage very frequently or frequently versus occasionally, rarely, very rarely, or never).

### Outcome

The primary outcome was nurse-reported exposure of a chemotherapeutic agent to their skin or to their eyes in the past year. We restricted our analyses to comparing nurses who reported zero versus one or more exposures in the past year (n=242).

### Nursing characteristics

We included the following nurse-reported covariates in our models: race (white or non-white), certification in oncology nursing (certified or not certified), and

education level (bachelor's degree or higher or less than a bachelor's degree).

### Analysis

First, we examined the response rate and non-response bias by comparing the demographic characteristics of our analytic sample from individuals who either did not respond at all to the survey, or did not answer the question regarding chemotherapy exposure. These demographic variables were provided on the entire sampling frame from the registry. Next, exposure to chemotherapy was treated as a binary variable (exposed vs not exposed), and bivariate analyses of the associations between categorical measures (perception of nurse practice environment, oncology certification and education level) were compared by exposure status using chi-square tests of independence. Differences in PES-NWI subscales (treated as continuous measures) and nursing workloads were conducted using two-sample t tests of mean differences.

Logistic regression, adjusted for clustering of nurses within practices (SAS GENMOD), was used to model odds of exposure among respondents. Six models were estimated. First, we estimated models to examine likelihood of chemotherapy exposure associated with perceived practice environment (model I), nursing workload (model II), and chemotherapy verification (model III). Two models were estimated for perceived practice environment (model IV) and chemotherapy verification (model V) that were adjusted for the following: nurse characteristics (race, education, experience as a nurse, and oncology certification status); and nursing workload. Model VI examined perceived practice environments and was adjusted for nurse characteristics, nurse workload, and chemotherapy verification. For our final model (VII), we used a backward selection process ( $p>0.20$ ) to retain the PES-NWI subscales related to chemotherapy exposure.<sup>24</sup> This model was adjusted for nursing workload, chemotherapy verification, and nurse characteristics. The overall perception of practice environment measure was excluded from this final analysis because the PES-NWI measures are considered components of the overall measure. All tests were two tailed and significance was established at a Cronbach  $\alpha$  of 0.05. All analyses were performed using SAS V.9.2 .

### Sensitivity analyses

We performed two sets of sensitivity analyses. Because of the relatively high rates of outcome and exposure, we replicated our analytic approach by specifying a Poisson distribution and calculating prevalence ratios that were adjusted for clustering.<sup>25</sup> Second, because missing data on PES-NWI items would exclude the respondent from the logistic regression models, we replicated our models

using the multiple imputation function in SAS V.9.2.<sup>26</sup> Specifically, we used a maximum likelihood function to replicate five complete datasets, and re-analyse the results from our final logistic regression model.

## RESULTS

### Response rate and analytic sample

Of the 402 nurses who responded to the survey (response rate of 30.5%),<sup>27</sup> this analysis was restricted to nurses who responded to the chemotherapy exposure item: 'In the past year, please indicate the number of times you have been exposed to a chemotherapeutic agent to your skin or eyes' (n=242). First, we examined demographic differences between our analytic sample and nurses who did not respond to the survey (n=937), or respond to the chemotherapy exposure question (n=160) using data provided by the registry on the entire sampling frame (table 1). Except for a higher proportion of nurses who did not respond employed outside of hospital ambulatory or physician practices, we observed no significant differences in observed demographics between our analytic sample and those who did not respond.

Next, we examined more detailed characteristics of our analytic sample (n=242). The individual characteristics of nurses who were exposed (race, oncology certification status, and education level) did not differ significantly from those who were not exposed (table 2). Though not significant, 23 or 62.2% of nurses who were exposed reported a favourable practice environment versus 142 or 74.7% of nurses who were not exposed to chemotherapy (p=0.12).

Compared with nurses who did not report exposure to chemotherapy within the last year, those who were

exposed had lower (worse) scores on several PES-NWI subscales. Mean scores on the participation in practice affairs and staffing and resource adequacy subscales differed by exposure status: 3.16 for nurses who were not exposed versus 2.73 for those who were exposed on a five-point scale (p< 0.01), and 3.61 for nurses who were not exposed versus 3.01 for those who were exposed (p<0.001). Compared with nurses who were exposed, the means of the remaining subscales (nurse manager leadership, ability, and support, collegial nurse–physician relations, nursing foundations for quality of care, and supportive medical assistant relations) were higher among those who reported no exposure, but the differences were not significant. Nurses who were exposed reported an average patient assignment of 11.1 patients per shift versus 8.43 patients per shift reported by those who were not exposed (p=0.02). The reported nurse workload across the sample ranged from 0 to 38 patients on the last shift. Chemotherapy orders were verified by two or more nurses on a frequent or very frequent basis 94.5% of the time for the entire sample, although only 82.9% of nurses exposed to chemotherapy indicated this was the case for them (compared with 96.9% of nurses not exposed to chemotherapy, p<0.01). Years employed as a nurse did not differ significantly by exposure status (p=0.58).

Multivariable models (table 3) were used to examine the relationships between chemotherapy exposure and organisational structures, and processes of care. Models I–III yielded significant relationships between reported chemotherapy exposure and practice environment, nursing workload, and chemotherapy verification, respectively, and were adjusted for nurse characteristics. In model IV, after adjusting for nurse characteristics and nursing workload, nurses who reported favourable

**Table 1** Demographic characteristics of the analytic sample and of nurses who did not respond

	Analytic sample (n=242) n (%)	Nurses in sampling frame who did not respond (n=1097)	p
Employed full time	146 (85.4)	913 (88.6)	0.25
Employed part time	25 (14.6)	118 (11.4)	
White	157 (91.8)	973 (94.4)	0.22
Non-white	14 (8.2)	58 (5.6)	
Women	166 (97.1)	994 (96.4)	0.65
Men	5 (2.9)	37 (3.6)	
Hospital outpatient	97 (56.7)	575 (55.8)	0.01
Physician practice	65 (38.0)	320 (31.0)	
Other practice setting	9 (5.3)	136 (13.2)	
Less than Bachelor's degree	86 (50.3)	521 (50.5)	1.0
Bachelor's degree or higher	85 (49.7)	510 (49.5)	
Resides in an MSA	129 (75.4)	819 (79.4)	0.26
Resides outside of an MSA	42 (24.6)	212 (20.6)	

Nurses in the sampling frame who did not respond include 160 who did not answer the chemotherapy exposure question, and 937 who did not return the questionnaire. Demographic characteristics from the state's nursing registry were not available for 254 nurses in the sampling frame. MSA, metropolitan statistical area.

**Table 2** Nurse-reported exposure to chemotherapy compared with work environment and individual level characteristics (n=242)

	Exposed to chemotherapy		p*
	Yes 41 (16.9%) Mean (SD)	No 201 (83.1%)	
<b>Structure</b>			
Revised PES-NWI subscales			
Nurse participation in practice affairs	2.73 (0.69)	3.16 (0.80)	<0.01
Nurse manager leadership and ability	3.37 (0.84)	3.57 (0.92)	0.20
Collegial nurse–physician relations	3.89 (0.63)	4.01 (0.73)	0.34
Staffing and resource adequacy	3.01 (1.00)	3.61 (0.91)	<0.01
Nursing foundations for quality of care	4.04 (0.54)	4.17 (0.60)	0.22
Supportive relations with medical assistants	3.41 (1.06)	3.62 (1.02)	0.25
Nursing workloads			
Number of patients primarily responsible for on last shift	11.11 (6.44)	8.43 (5.99)	0.02
	n (%)		p†
Perception of nurse practice environment			
Unfavourable or mixed	14 (37.8)	48 (25.3)	0.12
Favourable	23 (62.2)	142 (74.7)	
<b>Process</b>			
Chemotherapy orders and dosages are verified by two nurses			
Infrequently	7 (17.1)	6 (3.1)	<0.01
Frequently or very frequently	34 (82.9)	188 (96.9)	
<b>Nurse characteristics</b>			
<b>Race</b>			
Non-white	6 (14.6)	15 (7.5)	0.14
White	35 (85.4)	185 (92.5)	
<b>Oncology certification</b>			
Not certified	8 (19.5)	56 (27.9)	0.27
Certified	33 (80.5)	145 (72.1)	
<b>Education level</b>			
Less than a bachelor's degree	25 (61)	106 (52.7)	0.33
Bachelor's degree or higher	16 (39)	95 (47.3)	

\*Differences in exposure tested using two sample t tests.

†Differences in exposure tested using the chi-square test of independence. In cases of small cell sizes, Fisher's exact test was used. PES-NWI, Practice Environment Scale of the Nursing Work Index.

practice environments (as opposed to unfavourable or mixed environments) had a significantly lower likelihood for exposure (OR 0.44, 95% CI 0.21 to 0.92). In model V, after adjustment for nurse characteristics and nursing workloads, frequent or very frequent use of chemotherapy verification was associated with a significant reduction in exposure likelihood (OR 0.21, 95% CI 0.07 to 0.61). Model VI reports a significantly decreased likelihood of exposure for nurses who report favourable environments (as opposed to unfavourable/mixed environments), after adjusting for nurse characteristics, nursing workloads, and chemotherapy verification (OR 0.45, 95% CI 0.21 to 0.95).

One final model (model VII, right-sided panel of table 3) included specific PES-NWI subscales, nursing workloads, and verification of chemotherapy. Individual nurse characteristics were included in the model, but not displayed. Nurses were less likely to report chemotherapy exposure when they reported staffing and

resources were adequate (OR 0.35, 95% CI 0.17 to 0.73). Nurses who reported that chemotherapy orders and doses were consistently verified by two nurses had a significantly lower likelihood of chemotherapy exposure (OR 0.17, 95% CI 0.05 to 0.59).

### Sensitivity analysis

Our findings did not change when we specified a Poisson distribution and calculated prevalence ratios. Our results also did not change when the entire sample was used in a maximum-likelihood, five-dataset multiple imputation model to address missing data.

### DISCUSSION

In this state-wide sample of ambulatory oncology nurses, we identified a high rate of self-reported skin or eye exposure to chemotherapy (16.9%). While there is no target level of 'acceptable' exposure, unintentional occupational exposure to antineoplastics could be

**Table 3** Odds of exposure to chemotherapy, as estimated by structure and process variables, n=185\*

	OR (95% CI) adjusted for			
	Nurse characteristics	Nurse characteristics and nursing workload	Nurse characteristics, nursing workload, and chemotherapy verification	Nurse characteristics
Model	I	IV	VI	VII
Favourable practice environment	0.49 (0.24 to 1.00)	0.44 (0.21 to 0.92)	0.45 (0.21 to 0.95)	—
Model	II			
Nursing workload	1.07 (1.02 to 1.12)	—	—	1.06 (0.99 to 1.12)
Model	III	V		
Chemotherapy verification	0.22 (0.08 to 0.62)	0.21 (0.07 to 0.61)	—	0.17 (0.05 to 0.59)
Revised PES-NWI subscales				
Nurse participation in practice affairs (favourable vs unfavourable)				0.51 (0.24 to 1.06)
Staffing and resource adequacy (favourable vs unfavourable)				0.35 (0.17 to 0.73)

\*All models adjusted for clustering of nurses within practices using a generalised model function. PES-NWI, Practice Environment Scale of the Nursing Work Index.

considered a 'never' event, given the risks for harm. Both the structure and processes of ambulatory chemotherapy settings were associated with exposure. Favourable practice environments, lower nursing workloads, and adherence to two-nurse chemotherapy verification were protective. These relationships persisted after controlling for clustering of nurses in practices. Individual characteristics of nurses, such as years of experience, were not associated with exposure.

While it is not surprising that favourable practice environments and lower workloads for nurses were associated with reduced exposure, our finding of a relationship between chemotherapy exposure and dose verification is intriguing. Practices that adhere to this ASCO/ONS endorsed practice<sup>6</sup> likely are predisposed to a positive safety culture. Thus, the verification process may not directly influence exposure but rather may serve as a proxy for various processes to protect patients and nurses. Direct measures of safety behaviours<sup>28</sup> may clarify the mechanisms by which practices can protect nurses from potential harm.

Our study contributes to the literature in two important ways. First, it is one of the few to use a sampling frame of a diverse population of nurses employed outside of inpatient care units. The previous studies have focused on purposive samples from settings that have agreed to participate.<sup>8–11 13 14</sup> Facilities that participate in chemotherapy safety studies are likely to have heightened awareness of the risks and implement processes to protect employees. Second, we included and identified aspects of nurses' daily working conditions that are associated with increased exposure risk. The contribution of organisational structures and processes to exposure risk has not been reported previously.<sup>29</sup>

It is encouraging to note that the organisational characteristics associated with increased chemotherapy

exposure are modifiable by leadership. Practice managers can distribute nursing workloads more evenly, assure the availability of adequate time, space, and personnel for chemotherapy verification, and assure that requisite resources are available to administer chemotherapy in ways that minimise occupational exposure. Our findings also suggest that the current safety culture in ambulatory oncology practices could be strengthened to minimise risks to oncology nurses.

### Limitations

We did not measure nurses' use of safety devices, protective equipment, and adoption of protocols to minimise chemotherapy exposure. We also did not obtain information about the activities performed at the time of exposure (eg, mixing, administering, disconnecting, or disposal). These factors likely influence exposure risk and should be included in future studies. Second, our response rate, while within the range of recently published response rates of nursing personnel, raises concern for non-response bias. Except for practice setting, we observed no significant systematic differences between nurses who responded versus those who did not respond; this reduces concerns for bias.<sup>30</sup> Our non-response analysis suggests that nurses who did not respond were less likely to practice in hospital ambulatory or private practices, which are the target areas for our study. Third, our study did not include biological measures (eg, urine, blood, or surface swipe tests) to validate the self report of exposure. Historically, nurses have served as reliable informants on quality and safety phenomena.<sup>12</sup> In some cases, nurse reports of operational failures are lower than independent observation, which suggests that nurses may underreport exposure.<sup>31</sup> This would suggest that our exposure estimates are conservative. Finally, we interpret the

relationship between nursing workload and chemotherapy exposure with caution, as the reported workload of respondents ranged from 0 to 38 patients. However, our discussions with practicing oncology nurses confirm the wide variation in daily workloads.<sup>21</sup>

### Summary and practice implications

In this state-wide sample of oncology nurses practicing outside of hospital inpatient units, we observe high rates of nurse-reported chemotherapy exposure to their skin and eyes. Clinicians and managers can partner to manage workloads and implement steps to promote safe practices of hazardous drug administration. These include education, appropriate safety devices, personal protective equipment, and management support to adopt these methods by clinicians. Robust efforts to provide for the safety of patients receiving chemotherapy should be extended to the providers who administer this potentially harmful therapy.

**Acknowledgements** We thank the nurses who completed our questionnaires, and to Martha Polovich, RN, PhD, for content expertise.

**Funding** This research was supported by a Pathway to Independence Award (R00 NR 010750) from the National Institute of Nursing Research, National Institutes of Health, and in part from the University of Michigan Comprehensive Cancer Centre Support Grant (P30 CA46592).

**Competing interests** None.

**Ethics approval** University of Michigan—IRBMED.

**Provenance and peer review** Not commissioned; externally peer reviewed.

### REFERENCES

- Centres for Disease Control and Prevention. *Ambulatory Medical Care Utilization Estimates for 2007*. 2011. [http://www.cdc.gov/nchs/data/series/sr\\_13/sr13\\_169.pdf](http://www.cdc.gov/nchs/data/series/sr_13/sr13_169.pdf) (accessed 4 Aug 2011).
- Mason HJ, Blair S, Sams C, *et al*. Exposure to antineoplastic drugs in two UK hospital pharmacy units. *Ann Occup Hyg* 2005;49:603–10.
- Del Gaudio D, Menonna-Quinn D. Chemotherapy. Potential occupational hazards. *Am J Nurs* 1998;98:59–65.
- United States Department of Health and Human Services. *Center for Medicare and Medicaid Services: Medicare Program: hospital outpatient prospective payment system*. <http://edocket.access.gpo.gov/2010/pdf/2010-27926.pdf> (accessed 4 Apr 2011).
- National Institute for Occupational Safety and Health. *NIOSH alert: preventing occupational exposures to antineoplastic and other hazardous drugs in health care settings*. 2004. <http://www.cdc.gov/niosh/docs/2004-165/> (accessed 4 Apr 2011).
- Jacobson JO, Polovich M, McNiff KK, *et al*. American Society of Clinical Oncology/Oncology Nursing Society chemotherapy administration safety standards. *J Clin Oncol* 2009;27:5469–75.
- American Society of Clinical Oncology. *The Quality Oncology Practice Initiative: certification program*. 2010. <http://qopi.asco.org/certificationprocess> (accessed 4 Apr 2011).
- Benhamou S, Callais F, Sancho-Garnier H, *et al*. Mutagenicity in urine from nurses handling cytostatic agents. *Eur J Cancer Clin Oncol* 1986;22:1489–93.
- Newman MA, Valanis BG, Schoeny RS, *et al*. Urinary biological monitoring markers of anticancer drug exposure in oncology nurses. *Am J Public Health* 1994;84:852–5.
- Sessink PJ, Boer KA, Scheeffhals AP, *et al*. Occupational exposure to antineoplastic agents at several departments in a hospital. Environmental contamination and excretion of cyclophosphamide and ifosfamide in urine of exposed workers. *Int Arch Occup Environ Health* 1992;64:105–12.
- Connor TH, DeBord DG, Pretty JR, *et al*. Evaluation of antineoplastic drug exposure of health care workers at three university-based US cancer centers. *J Occup Environ Med* 2010;52:1019–27.
- Aiken LH, Sloane DM, Klocinski JL. Hospital nurses' occupational exposure to blood: prospective, retrospective, and institutional reports. *Am J Public Health* 1997;87:103–7.
- Castiglia L, Miraglia N, Pieri M, *et al*. Evaluation of occupational exposure to antineoplastic drugs in an Italian hospital oncological department. *J Occup Health* 2008;50:48–56.
- Turk M, Davas A, Ciceklioglu M, *et al*. Knowledge, attitude and safe behaviour of nurses handling cytotoxic anticancer drugs in Ege University Hospital. *Asian Pac J Cancer Prev* 2004;5:164–8.
- Donabedian A. Evaluating the quality of medical care. *Milbank Mem Fund Q* 1966;44:166–206.
- Friese CR, Lee CS, O'Brien S, *et al*. *Multi-mode and method experiment in a study of nurses*. Survey Practice, 2010. <http://surveypractice.org/2010/10/27/multi-mode-nurses-survey/> (accessed 20 Dec 2010).
- Dillman DA, Smyth J, Christian LM. *Internet, mail, and mixed-mode surveys: the tailored design method*. 3rd edn. Hoboken, NJ: John Wiley Co, 2009.
- Lake ET. Development of the practice environment scale of the Nursing Work Index. *Res Nurs Health* 2002;25:176–88.
- Friese CR, Lake ET, Aiken LH, *et al*. Hospital nurse practice environments and outcomes for surgical oncology patients. *Health Serv Res* 2008;43:1145–63.
- Lake ET. The nursing practice environment: measurement and evidence. *Med Care Res Rev* 2007;64(Suppl 2):104S–22S.
- Kamimura A, Schneider K, Lee CS, *et al*. Practice environments of nurses in ambulatory oncology settings: A thematic analysis. *Cancer Nurs*. Published Online First: 2 March 2011.
- Friese CR. Practice environments of nurses employed in ambulatory oncology settings: measure refinement. *Oncol Nurs Forum*. In press.
- Aiken LH, Clarke SP, Sloane DM, *et al*. Hospital nurse staffing and patient mortality, nurse burnout, and job dissatisfaction. *JAMA* 2002;288:1987–93.
- Vittinghoff E, Glidden D, Shiboski S, *et al*. *Regression methods in biostatistics: linear, logistic, survival, and repeated measures models*. New York: Springer, 2005.
- Zou G. A modified poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;159:702–6.
- Yuan Y. *Multiple imputation for missing data: concepts and new developments (Version 9.0)*. <http://support.sas.com/rnd/app/papers/multipleimputation.pdf> (accessed 4 Apr 2011).
- American Association for Public Opinion Research. *Standard definitions: final disposition of case codes and outcome rates for surveys*. 6th edn, 2009. [http://www.aapor.org/Standard\\_Definitions1.htm](http://www.aapor.org/Standard_Definitions1.htm) (accessed 10 Dec 2010).
- Vogus TJ, Sutcliffe KM. The Safety Organizing Scale: development and validation of a behavioral measure of safety culture in hospital nursing units. *Med Care* 2007;45:46–54.
- Connor TH, McDiarmid MA. Preventing occupational exposures to antineoplastic drugs in health care settings. *CA Cancer J Clin* 2006;56:354–65.
- Groves RM. *Survey non-response*. New York, NY: Wiley, 2002.
- Tucker AL, Spear SJ. Operational failures and interruptions in hospital nursing. *Health Serv Res* 2006;41:643–62.