

Continuing medical education activity in *American Journal of Hematology*

CME Editor: Ayalew Tefferi, MD
Author: Ryan A. Wilcox, MD, PhD

Article Title: Cutaneous B-cell lymphomas: 2013 update on diagnosis, risk-stratification, and management

If you wish to receive credit for this activity, please refer to the website: www.wileyhealthlearning.com

Accreditation and Designation Statement:

Blackwell Futura Media Services is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Blackwell Futura Media Services designates this journal-based CME for a maximum of 1 *AMA PRA Category 1 Credit*[™]. Physicians should only claim credit commensurate with the extent of their participation in the activity

Educational Objectives

Upon completion of this educational activity, participants will be better able to:

- To understand the clinical presentation, natural history, and therapeutic approach for the most common cutaneous B-cell lymphomas.

Activity Disclosures

No commercial support has been accepted related to the development or publication of this activity.

CME Editor – Ayalew Tefferi, MD has no conflicts of interest to disclose

Author – Ryan A. Wilcox, MD has no conflicts of interest to disclose

This activity underwent peer review in line with the standards of editorial integrity and publication ethics maintained by *American Journal of Hematology*. The peer reviewers have no conflicts of interest to disclose. The peer review process for *American Journal of Hematology* is single blinded. As such, the identities of the reviewers are not

disclosed in line with the standard accepted practices of medical journal peer review.

Conflicts of interest have been identified and resolved in accordance with Blackwell Futura Media Services's Policy on Activity Disclosure and Conflict of Interest. The primary resolution method used was peer review and review by a non-conflicted expert.

Instructions on Receiving Credit

This activity is intended for physicians. For information on applicability and acceptance of continuing medical education credit for this activity, please consult your professional licensing board.

This activity is designed to be completed within one hour; physicians should claim only those credits that reflect the time actually spent in the activity. To successfully earn credit, participants must complete the activity during the valid credit period, which is up to two years from initial publication. Additionally, up to 3 attempts and a score of 70% or better is needed to pass the post test.

Follow these steps to earn credit:

- Log on to www.wileyhealthlearning.com
- Read the target audience, educational objectives, and activity disclosures.
- Read the activity contents in print or online format.
- Reflect on the activity contents.
- Access the CME Exam, and choose the best answer to each question.
- Complete the required evaluation component of the activity.
- Claim your Certificate.

This activity will be available for CME credit for twelve months following its launch date. At that time, it will be reviewed and potentially updated and extended for an additional twelve months.

ANNUAL CLINICAL UPDATES IN HEMATOLOGICAL MALIGNANCIES:
A CONTINUING MEDICAL EDUCATION SERIES

Cutaneous B-cell lymphomas: 2013 update on diagnosis, risk-stratification, and management

Ryan A. Wilcox*

Disease overview: Approximately one-fourth of cutaneous lymphomas are B-cell derived and are generally classified into three distinct subgroups: primary cutaneous follicle-center lymphoma (PCFCL), primary cutaneous marginal zone lymphoma (PCMZL), and primary cutaneous diffuse large B-cell lymphoma, leg type (PCLBCL, LT).

Diagnosis: Diagnosis and disease classification is based on histologic review and immunohistochemical staining of an appropriate skin biopsy. Pathologic review and an appropriate staging evaluation are necessary to distinguish primary cutaneous B-cell lymphomas from systemic B-cell lymphomas with secondary skin involvement.

Risk-stratification: Disease histology remains the most important prognostic determinant. Both PCFCL and PCMZL are indolent lymphomas that infrequently disseminate to extracutaneous sites and are associated with an excellent long-term prognosis. In contrast, PCLBCL, LT is an aggressive lymphoma with an inferior prognosis.

Risk-adapted therapy: PCFCL and PCMZL patients with solitary or relatively few skin lesions may be affectively managed with local radiation therapy. Although single-agent rituximab may be employed for patients with more widespread skin involvement, multiagent chemotherapy is rarely appropriate. In contrast, management of patients with PCLBCL, LT is comparable to the management of patients with systemic DLBCL. *Am. J. Hematol.* 88:74–76, 2013. © 2012 Wiley Periodicals, Inc.

Disease Overview

Primary cutaneous lymphomas are a heterogeneous group of extranodal non-Hodgkin lymphomas, approximately 25% of which are B-cell derived and are classified into three major entities in the World Health Organization–European Organization for Research and Treatment of Cancer (EORTC) joint classification: primary cutaneous follicle-center lymphoma (PCFCL), primary cutaneous diffuse large B-cell lymphoma, leg type (PCLBCL, LT), and primary cutaneous marginal zone lymphoma (PCMZL) [1]. The incidence of cutaneous B-cell lymphomas (CBCL) has been increasing and is currently 3.1 per million persons, based on surveillance, epidemiology, and end results registry data, with the highest incidence rates being reported among males, non-Hispanic whites, and adults over the age of 50 [2].

Diagnosis

Diagnosis and classification of a CBCL requires an excisional or punch biopsy for careful morphologic and immunohistochemical analysis, and an appropriate staging evaluation to exclude systemic disease [3]. The use of appropriate immunohistochemical stains (e.g., CD5, cyclin D1) may also aid in distinguishing CBCL from secondary skin involvement by a systemic lymphoma.

PCFCL

PCFCL are commonly solitary plaques or tumors involving the trunk, particularly the head or scalp. Although grouped lesions may be observed, multifocal disease is less common. Histologically, PCFCL are characterized by a follicular, diffuse, or mixed growth pattern comprised of large centrocytes derived from germinal-center B-cells [1,4,5]. In contrast to systemic follicular lymphomas, the majority of PCFCL do not harbor the t(14;18) translocation involving the bcl-2 locus, and do not strongly express bcl-2 by immunohistochemistry, although weak expression may

be observed in a minority of cases [6–8]. These CBCL express bcl-6, variably express CD10, and are MUM-1/IRF-4 negative, consistent with their origin from germinal-center B cells.

PCLBCL, LT

In contrast to PCFCL, which is an indolent CBCL largely involving the head and trunk commonly affecting middle-aged adults, PCLBCL, LT commonly affects elderly females and presents with rapidly progressive tumors involving the lower legs. In contrast to PCFCL, these lymphomas are characterized by diffuse sheets of centroblasts that highly express bcl-2, likely due to gene amplification [9]. Most cases are MUM-1/IRF-4 and bcl-6 positive, CD10 negative, and have a gene expression profile resembling activated B cells [5].

PCMZL

Patients with PCMZL frequently present with multifocal patches, plaques or nodules involving the trunk and arms. Although an association with *Borrelia burgdorferi* has been observed in Europe, a similar association has not been observed in cases from the United States [10–13]. PCMZL are composed of a mixed infiltrate of small, marginal zone B cells, lymphoplasmacytic cells, plasma cells, and reactive

Division of Hematology/Oncology, University of Michigan Cancer Center, 1500 E. Medical Center Drive, Room 4310 CC, Ann Arbor, MI 48109-5948

Conflict of interest: Nothing to report

*Correspondence to: Ryan Wilcox, Division of Hematology/Oncology, University of Michigan Cancer Center, 1500 E. Medical Center Drive, Room 4310 CC, Ann Arbor, MI 48109-5948. E-mail: rywilcox@med.umich.edu

Received for publication 23 October 2012; Accepted 28 October 2012

Am. J. Hematol. 88:74–76, 2013.

Published online in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/ajh.23360

T cells. Marginal zone B cells characteristically express bcl-2, but lack bcl-6 or CD10 expression.

Risk-Stratification

The International Society for Cutaneous Lymphomas (ISCL) and EORTC recently proposed staging recommendations for cutaneous lymphomas other than mycosis fungoides and Sezary syndrome [3]. Staging should include a history, physical examination, appropriate laboratory studies (including lactate dehydrogenase), and imaging (either CT, PET, or increasingly PET/CT) of the chest, abdomen, pelvis, and neck (in cases with involvement of the head or neck). A bone marrow biopsy and aspirate should be performed in cases of PCLBCL, LT. Although the ISCL/EORTC does not endorse routine bone marrow examination in cases of PCFCL or PCMZL, approximately 10% of patients with PCFCL have bone marrow involvement [14]. Furthermore, bone marrow involvement was associated with significantly inferior disease-specific survival. Therefore, in the opinion of this author, bone marrow examination is justified in cases of PCFCL. While the TNM staging classification describes the extent of disease, staging in CBCL is of limited prognostic value, as the disease histology is the major determinant in risk-stratification. This is highlighted by a population-based study which identified histology and the site of skin involvement as important prognostic factors [15]. In contrast, the International Extranodal Lymphoma Study Group identified three independent prognostic factors (i.e., elevated LDH, >2 skin lesions, and nodular lesions) among patients with PCFCL and PCMZL. These factors were combined to form the cutaneous lymphoma international prognostic index (CLIPi). The absence of any adverse prognostic factor was associated with a 5-year progression-free survival of 91%. In contrast, the presence of two or three adverse prognostic factors was associated with a 5-year progression-free survival of 48%. As the vast majority of relapses were confined to the skin, the CLIPi was unable to risk-stratify patients by overall survival. The most important factor for risk-stratification among the CBCLs remains the histologic classification. Indolent CBCL (PCFCL and PCMZL) are associated with 5-year disease-specific survival $\geq 95\%$ [1]. Differences in growth pattern, the density of centroblasts, and cytogenetic findings do not appear to provide meaningful prognostic information. Bcl-2 expression among PCFCL with a diffuse large B-cell histology may be a notable exception [16]. In contrast, PCLBCL, LT is associated with a 5-year disease-specific survival of approximately 50% and with cytogenetic changes, including translocations involving c-myc, that confer a poor prognosis among systemic DLBCLs [1,17]. In contrast to patients presenting with only a single tumor, involvement of multiple sites, on one or both legs, is associated with a significantly inferior disease-specific survival [18].

Treatment

As no randomized controlled trials are available, treatment recommendations for CBCL are largely based on small retrospective studies and institutional experience. The EORTC and ISCL have published consensus treatment recommendations that are consistent with NCCN guidelines [19]. In most cases, optimal patient management requires a multidisciplinary approach, including dermatology, medical oncology, and radiation oncology.

PCFCL

For patients with solitary lesions, radiation therapy is safe and highly effective, with a complete remission rate approaching 100%. Radiation does not appear inferior to multiagent chemotherapy among patients with multiple lesions that can be included in multiple radiation fields

[20]. Although radiation therapy is generally recommended for patients with a solitary lesion, radiation therapy or observation (i.e., "watch and wait") are reasonable options for those patients with multiple lesions. Patients with more extensive skin involvement are effectively managed with single-agent rituximab [19]. Multiagent chemotherapy (e.g., R-CHOP) is rarely required in the management of PCFCL. Approximately one-third of patients may relapse following either radiation or single-agent rituximab, but relapses are usually confined to the skin and are approached in a manner similar to that described for the initial management of PCFCL.

PCMZL

Patients with PCMZL are approached in a manner analogous to that described in the initial management of PCFCL. Radiation therapy is associated with a similarly high response rate for patients with a single or few lesions [19]. Those with more widespread skin involvement may be observed. Once symptomatic, culprit lesions may be irradiated (or surgically excised). As for PCFCL, single-agent rituximab may be utilized in patients with symptomatic, widespread skin lesions. An initial trial of antibiotics for those with *B. burgdorferi*-associated PCMZL has been recommended [21], but is less relevant for North American patients.

PCLBCL, LT

As previously noted, the natural history of PCLBCL, LT more closely resembles that of systemic DLBCL. Therefore, R-CHOP (with or without radiation therapy) is utilized in these patients. Although few reports are available in the literature, the use of R-CHOP in these patients is associated with remission (and relapse) rates comparable to those reported for patients with high-risk systemic DLBCL [19].

References

1. Willemze R, Jaffe ES, Burg G, et al. WHO-EORTC classification for cutaneous lymphomas. *Blood* 2005;105:3768–3785.
2. Bradford PT, Devesa SS, Anderson WF, et al. Cutaneous lymphoma incidence patterns in the United States: A population-based study of 3884 cases. *Blood* 2009;113:5064–5073.
3. Kim YH, Willemze R, Pimpinelli N, et al. TNM classification system for primary cutaneous lymphomas other than mycosis fungoides and Sezary syndrome: A proposal of the International Society for Cutaneous Lymphomas (ISCL) and the Cutaneous Lymphoma Task Force of the European Organization of Research and Treatment of Cancer (EORTC). *Blood* 2007;110:479–484.
4. Gellrich S, Rutz S, Golembowski S, et al. Primary cutaneous follicle center cell lymphomas and large B cell lymphomas of the leg descend from germinal center cells. A single cell polymerase chain reaction analysis. *J Invest Dermatol* 2001;117:1512–1520.
5. Hoefnagel JJ, Dijkman R, Basso K, et al. Distinct types of primary cutaneous large B-cell lymphoma identified by gene expression profiling. *Blood* 2005;105:3671–3678.
6. Cerroni L, Kerl H. Immunoreactivity for bcl-2 protein in cutaneous lymphomas and lymphoid hyperplasias. *J Cutan Pathol* 1995;22:476–478.
7. Cerroni L, Volkenandt M, Rieger E, et al. bcl-2 protein expression and correlation with the interchromosomal 14;18 translocation in cutaneous lymphomas and pseudolymphomas. *J Invest Dermatol* 1994;102:231–235.
8. Cerroni L, Arzberger E, Putz B, et al. Primary cutaneous follicle center cell lymphoma with follicular growth pattern. *Blood* 2000;95:3922–3928.
9. Mao X, Lillington D, Child F, et al. Comparative genomic hybridization analysis of primary cutaneous B-cell lymphomas: Identification of common genomic alterations in disease pathogenesis. *Genes Chromosomes Cancer* 2002;35:144–155.
10. Cerroni L, Zochling N, Putz B, et al. Infection by *Borrelia burgdorferi* and cutaneous B-cell lymphoma. *J Cutan Pathol* 1997;24:457–461.
11. Goodlad JR, Davidson MM, Hollowood K, et al. *Borrelia burgdorferi*-associated cutaneous marginal zone lymphoma: A clinicopathological study of two cases illustrating the temporal progression of *B. burgdorferi*-associated B-cell proliferation in the skin. *Histopathology* 2000;37:501–508.
12. Goodlad JR, Davidson MM, Hollowood K, et al. Primary cutaneous B-cell lymphoma and *Borrelia burgdorferi* infection in patients from the Highlands of Scotland. *Am J Surg Pathol* 2000;24:1279–1285.
13. Wood GS, Kamath NV, Guitart J, et al. Absence of *Borrelia burgdorferi* DNA in cutaneous B-cell lymphomas from the United States. *J Cutan Pathol* 2001;28:502–507.

14. Senff NJ, Kluin-Nelemans HC, Willemze R. Results of bone marrow examination in 275 patients with histological features that suggest an indolent type of cutaneous B-cell lymphoma. *Br J Haematol* 2008;142:52–56.
15. Smith BD, Smith GL, Cooper DL, et al. The cutaneous B-cell lymphoma prognostic index: A novel prognostic index derived from a population-based registry. *J Clin Oncol* 2005;23:3390–3395.
16. Grange F, Petrella T, Beylot-Barry M, et al. Bcl-2 protein expression is the strongest independent prognostic factor of survival in primary cutaneous large B-cell lymphomas. *Blood* 2004;103:3662–3668.
17. Hallermann C, Kaune KM, Gesk S, et al. Molecular cytogenetic analysis of chromosomal breakpoints in the IGH, MYC, BCL6, and MALT1 gene loci in primary cutaneous B-cell lymphomas. *J Invest Dermatol* 2004;123:213–219.
18. Grange F, Bekkenk MW, Wechsler J, et al. Prognostic factors in primary cutaneous large B-cell lymphomas: A European multicenter study. *J Clin Oncol* 2001;19:3602–3610.
19. Senff NJ, Noordijk EM, et al. European organization for research and treatment of cancer and international society for cutaneous lymphoma consensus recommendations for the management of cutaneous B-cell lymphomas. *Blood* 2008;112:1600–1609.
20. Bekkenk MW, Vermeer MH, Geerts ML, et al. Treatment of multifocal primary cutaneous B-cell lymphoma: A clinical follow-up study of 29 patients. *J Clin Oncol* 1999;17:2471–2478.
21. Dreno B. Standard and new treatments in cutaneous B-cell lymphomas. *J Cutan Pathol* 2006;33 (Suppl 1):47–51.