

Particulate Matter Promotes In Vitro Receptor-Recognizable Low-Density Lipoprotein Oxidation and Dysfunction of Lipid Receptors

Natalia Manzano-León,¹ Jaime Mas-Oliva,² Laura Sevilla-Tapia,¹ Rocío Morales-Bárceñas,¹ Jesús Serrano,³ Marie S. O'Neill,⁴ Claudia M. García-Cuellar,¹ Raúl Quintana,¹ Inés Vázquez-López,¹ and Álvaro R. Osornio-Vargas^{1,5},

¹ Instituto Nacional de Cancerología, C.P. 14080 México, D.F., México

² Instituto de Fisiología Celular, Universidad Nacional Autónoma de México, C.P. 04510, México, D.F., México

³ Facultad de Ciencias, Universidad Nacional Autónoma de México, México, D.F., México

⁴ Departments of Environmental Health Sciences and Epidemiology, University of Michigan School of Public Health, Ann Arbor, MI 48109, USA

⁵ Department of Paediatrics, University of Alberta, Edmonton T6G 1C9, Canada; E-mail: osornio@ualberta.ca

Received 30 August 2012; accepted 7 October 2012

ABSTRACT: Particulate matter may promote cardiovascular disease, possibly as a consequence of its oxidative potential. Studies using susceptible animals indicate that particulate matter aggravates atherosclerosis by increasing lipid/macrophage content in plaques. Macrophage lipid uptake requires oxidized low-density lipoprotein and scavenger receptors; same receptors are involved in particulate matter uptake. We studied in vitro particulate matter potential to oxidize low-density lipoproteins and subsequent cell uptake through scavenger receptors. Particulate matter-induced low-density lipoproteins oxidation was evaluated by the thiobarbituric acid assay. Binding/internalization was tested in wild type and scavenger receptor-transfected Chinese hamster ovary cells, and in RAW264.7 cells using fluorescently labeled low-density lipoproteins. Dose-dependent binding/internalization only occurred in scavenger receptor-transfected Chinese hamster ovary cells and RAW264.7 cells. Competition binding/internalization using particles showed that particulate matter induced decreased binding (~50%) and internalization (~70%) of particle-oxidized low-density lipoproteins and native low-density lipoproteins. Results indicate that particulate matter was capable of oxidizing low-density lipoproteins, favoring macrophage internalization, and

also altered scavenger and low-density lipoproteins receptor function. © 2012 Wiley Periodicals, Inc. *J Biochem Mol Toxicol* 27:69–76, 2013; View this article online at wileyonlinelibrary.com. DOI 10:1002/jbt.21452

KEYWORDS: Particulate Matter; Low-Density Lipoprotein; Scavenger Receptor; Macrophages; Oxidative Potential

INTRODUCTION

Exposure to particulate matter (PM) in polluted cities represents a significant risk to human health [1]. PM is a complex, dynamic mixture of materials with toxic potential that depends on size and composition [2, 3]. Several studies have linked cardiovascular mortality and complications of preexisting atherosclerotic disease after exposure to PM present in polluted air [4–6].

The mechanisms by which exposure to PM complicates a preexisting cardiovascular condition have not been fully elucidated yet. However, experimental studies in humans and animals have provided a wealth of evidence on how PM exposure can influence relevant cardiovascular processes including increased blood pressure, increased blood viscosity, and altered fibrinogen levels [7]. Arrhythmias, changes in electrocardiographic parameters, and increased vasoconstriction have been also identified as complications in elderly subjects [8] and healthy people occupationally exposed to different types of particles [9, 10]. Subjects

Correspondence to: Álvaro R. Osornio-Vargas.

Contract Grant Sponsor: Consejo Nacional de Ciencia y Tecnología.

Contract Grant Number: CONACyT-M43138.

Contract Grant Sponsor: NIH/NIEHS.

Contract Grant Numbers: R01 ES016932-01 and R01 ES017022-

01.

© 2012 Wiley Periodicals, Inc.

implanted with defibrillators show an increase in the frequency of defibrillator interventions and an increase in the level of serum C-reactive protein in response to PM exposure [7, 11, 12].

Several studies in humans have also shown a relationship between carotid intimal-medial thickness and exposure to PM less than 2.5 μm in aerodynamic diameter ($\text{PM}_{2.5}$), in conjunction with atherosclerosis development and coronary artery calcification [13, 14]. Suwa et al. [15] observed advanced coronary artery and aorta atherosclerotic plaques in Watanabe heritable hyperlipidemic rabbits exposed to PM less than 10 μm in aerodynamic diameter (PM_{10}) compared to unexposed control animals of the same strain. Atherosclerotic plaques in exposed animals had a larger number of cells containing an increased amount of intracellular lipids, resulting in a greater quantity of total lipids in aortic lesions. Moreover, Araujo et al. [16] studied the proatherogenic effects of ultrafine particles and $\text{PM}_{2.5}$ in genetically susceptible mice (apolipoprotein $\text{E}^{-/-}$), which after exposure to PM presented larger atherosclerotic lesions than control animals. Likewise, apolipoprotein $\text{E}^{-/-}$ mice exposed to $\text{PM}_{2.5}$ showed increased content of lipids, cholesterol, and macrophages in atherosclerotic plaques, along with a reduced response to arterial dilatation. Also apolipoprotein $\text{E}^{-/-}$ mice fed an enriched fat diet exposed to PM presented an increased development of atherosclerotic plaques in comparison to animals receiving a balanced diet [17]. Finally, PM causes oxidative stress in cultured cells and cell-free systems [18–21].

The central objective of our research was to explore *in vitro* the capacity of PM to oxidize low-density lipoprotein (LDL) and to study the possible consequences of PM-induced LDL oxidation in the binding/internalization of PM-oxidized LDL (PM-ox-LDL) through the scavenger receptor (SR). SRs are involved with the development of foam cells by inducing internalization and accumulation of a large amount of ox-LDL within macrophages [22]. Since PM is internalized by macrophages through SR [23], the same receptor involved in ox-LDL uptake, we performed several assays to explore PM-ox-LDL and PM competition for the same receptor. Finally, to evaluate the effect of PM on a different group of receptors, we explored PM and native-LDL (nat-LDL) interaction with the LDL receptor (LDL-R).

MATERIAL AND METHODS

PM Sampling

$\text{PM}_{2.5}$ and PM_{10} were collected in the northern (industrial) zone of Mexico City, using high-volume samplers (1.13 m^3/min) (GMW model 1200, VCF HVPM10;

Sierra Andersen, Smyrna, GA) on nitrocellulose membranes (Sartorius, Goettingen, Germany). Integrated 24-h samples were collected 5 days a week. PM was recovered from the membranes and handled as previously reported [24]. The chemical characterization of samples from the industrial zone of Mexico City has been reported elsewhere [2, 25].

Low-Density Lipoproteins

Human LDL (density = 1.019–1.063 g/mL) were isolated from the plasma of three randomly selected healthy blood bank donors who were in a fasting state by sequential density ultracentrifugation in potassium bromide at 4°C as previously described [26]. The experiments were performed separately on the LDL from each of the three donors. Lipoproteins were dialyzed first against 0.15 M NaCl and 0.01% ethylene diamine tetraacetic acid (EDTA) and then against phosphate buffered saline (PBS) to eliminate EDTA. LDL concentration is given in terms of its protein content determined by the bicinchoninic acid method (Pierce, Rockford, IL).

LDL Oxidation by $\text{PM}_{2.5}$ and PM_{10}

Isolated LDL were incubated with increasing doses of $\text{PM}_{2.5}$ or PM_{10} (20, 40, and 80 $\mu\text{g}/\text{mg}$ LDL) with and without H_2O_2 (1 mM) over 24 h at 37°C, to evaluate oxidation by the thiobarbituric acid method (TBA) using a commercially available kit and following the manufacturer instructions (Northwest, Vancouver, WA, Canada). This method is based on the reaction of malondialdehyde with TBA, forming a complex that absorbs at 532 nm.

The PM dose range used in this study corresponds to concentration commonly reported in the literature as having significant *in vitro* cellular effects. They are also in a range relevant in magnitude to the exposure of an average person living in Mexico City at hot spots on airway bifurcations [27].

LDL Acetylation

Acetylated-LDL (ac-LDL) was prepared by LDL treatment with acetic anhydride and sodium acetate [28]. Briefly, an equal volume of saturated solution of sodium acetate was added to LDL with continuous stirring at a temperature of 4°C. Then, adequate amount of acetic anhydride was added to have a total mass of acetic anhydride equal to 1.5 times the mass of protein used and incubated for 1 h. Ac-LDL were dialyzed

against 0.15 M NaCl, centrifuged at 10,000 revolutions per minute (rpm) for 30 min and filtered.

LDL Fluorescent Labeling

LDL were labeled fluorescently with 1,1'-dioctadecyl-3,3,3',3'-tetramethylindocarbocyanine perchlorate (DiI) (Molecular Probes, Eugene, OR) as previously described [29]. LDL-DiI was dialyzed against 0.15 M NaCl and 0.01% EDTA as well as PBS to eliminate EDTA.

Cell Culture

We used Chinese hamster ovary (CHO), SR-transfected CHO (CHO-SR), and RAW264.7 cells. CHO and RAW cells were obtained from the American Type Culture Collection. CHO cells were used as a control due to the lack of scavenger receptor A (SR-A). CHO-SR were stably transfected with bovine SR-A (cells were kindly donated by Dr. Robert E. Pitas, Gladstone Institute of Cardiovascular Disease, University of California, San Francisco, CA). Cells were cultured in 10% fetal bovine serum-DMEM (Dulbecco's modified Eagle's Media)/F12 containing penicillin (50 U/mL)/streptomycin (50 µg/mL) for CHO, or geneticin (400 µg/mL) for CHO-SR. RAW cells (murine monocyte/macrophage) were used to compare SR activity in a naturally expressing SR cell line. Cells were grown in DMEM supplemented with 10% fetal bovine serum and penicillin (50 U/mL)/streptomycin (50 µg/mL) at 37°C in a 5% CO₂/95% air atmosphere.

PM-ox-LDL-DiI Binding/Internalization Assays

LDL-DiI were incubated with 80 µg PM₁₀/mg for 24 h at 37°C to induce oxidation, centrifuged at 5,000 rpm for 5 min to eliminate PM, and quantified once again. Confluent cells were incubated for 1 h in serum-free medium, and then in the presence of PM-ox-LDL-DiI (2.5, 5, 10, 20, and 40 µg/mL) for 4 h at 4°C (binding assay) or 37°C (internalization assay). Parallel assays were performed in the presence of 10 times the amount of nat-LDL and H₂O₂-ox-LDL to measure nonspecific binding internalization. Cells were washed five times with PBS-bovine serum albumin (2 mg/mL), recovered and analyzed by flow cytometry (FACScalibur instrument, Becton-Dickinson, NJ). To obtain specific binding/internalization data, fluorescence intensity values in the presence of an excess of lipoproteins (nat-LDL and H₂O₂-ox-LDL) were subtracted from total fluorescence intensity values to present final results.

Experiments were carried out using LDL from three different donors, and results are expressed as relative units where basal cell fluorescence corresponds to one relative unit. Relative units were used to normalize for the fluorescence variability observed between LDL-DiI samples obtained from different donors. Fluorescence variability of samples varied from 30 to 70 fluorescence units in the binding assays and between 170 and 280 fluorescence units in internalization assays, when a dose of 40 µg/mL of PM-ox-LDL-DiI was used in CHO-SR cells.

PM/LDL Competition Assays

Confluent CHO-SR were exposed to 0, 20, 40, and 80 µg/cm² of PM_{2.5} or PM₁₀, and to 20 µg/mL of PM-ox-LDL-DiI or nat-LDL-DiI for 4 h at 4 or 37°C. Parallel assays were performed using an excess (10 times the amount) of nat-LDL and H₂O₂-ox-LDL as indicated above. Cells were washed, and samples were analyzed by flow cytometry. RAW cells were only tested using 40 µg/cm² of PM₁₀ and 20 µg/mL of PM-ox-LDL-DiI or nat-LDL-DiI.

Cell Viability

Cell viability was assessed in parallel under the same experimental conditions described above by trypan-blue exclusion using a Neubauer camera. Trypan blue was added to cell suspensions at a final concentration of 0.8 mg/mL and incubated for 1 min at room temperature. The differentiation among the viable and nonviable cells was done through blue cell coloration under optical microscopy.

PM/LDL Interactions

We also evaluated possible PM-LDL (nat-LDL and PM-ox-LDL) binding interactions under cell-free conditions, using the same PM and LDL concentrations and conditions described in the binding/internalization experiments, except for the absence of cells in the assays. Samples were centrifuged and analyzed by fluorometry. We found that only ~10% of PM-ox-LDL-DiI or nat-LDL-DiI total fluorescence was retained by PM. This indicates that "unspecific" PM binding to LDL introduced an experimental error no larger than 10%.

Statistical Analysis

All experiments were done in triplicate, using plasma from three independent donors. Graphics were created to show the average of the three independent

experiments \pm standard deviation (SD). Data were analyzed by one-way analysis of variance (ANOVA) for competition assays and by factorial ANOVA for ox-LDL-binding/internalization assays, the last to evaluate the effect of ox-LDL concentration, cell line and the interaction between both factors (ox-LDL concentration \times cell line). When necessary, variables were transformed to homogenize the variances. Both ANOVA procedures were followed by multiple comparison tests (Scheffe test) and specifically for factorial ANOVA, orthogonal contrasts were performed to determine differences between cell lines according to the ox-LDL concentration. Statistical analysis was performed using SPSS version 13.0 for Windows. The results were considered significant when $p < 0.05$.

RESULTS

LDL exposure to three increasing concentrations of PM_{2.5} and PM₁₀ (0, 20, 40, and 80 $\mu\text{g}/\text{mg}$ LDL) for 24 h in the absence of H₂O₂ resulted in a concentration-related LDL oxidation. Exposures in the presence of H₂O₂ resulted in a corresponding increased oxidation (Figure 1). All further experimentation used PM₁₀ + H₂O₂ induced LDL oxidation, due to limited accessibility to PM_{2.5} samples.

Cell viability evaluation in cells (CHO, CHO-SR, and RAW) exposed to PM-ox-LDL and/or PM using the same experimental conditions used for binding/internalization and competition assays never resulted in viability decrements larger than 10% (data

not shown). Results obtained from each experimental condition were adjusted according to viability.

PM-ox-LDL-binding/internalization assays in CHO cells showed a relatively low dose-related basal activity. In the case of CHO-SR cells, a higher dose-related PM-ox-LDL binding/internalization was observed ($p < 0.05$) (Figure 2).

Experimentation was also done using ac-LDL to investigate whether similar results would be obtained in our cell system when using traditionally modified LDL. The ac-LDL yielded similar results to the ones obtained with PM-ox-LDL using CHO-SR and CHO cells (data not shown).

PM-ox-LDL-binding/internalization competition assays in the presence of PM_{2.5} or PM₁₀ (20, 40, and 80 $\mu\text{g}/\text{cm}^2$) resulted in a PM concentration-related decreased binding (53%–55% at 80 $\mu\text{g}/\text{cm}^2$) (Figures 3A and 3C) and internalization (73%–78% at 80 $\mu\text{g}/\text{cm}^2$) ($p < 0.05$) (Figures 3B and 3D).

PM/LDL-R competition binding/internalization assays performed in CHO-SR cells showed that PM_{2.5} and PM₁₀ also interfered in a dose-related manner with nat-LDL binding (62% at 80 $\mu\text{g}/\text{cm}^2$) (Figure 4A) and internalization (86% at 80 $\mu\text{g}/\text{cm}^2$) ($p < 0.05$) (Figure 4B).

When RAW cells binding/internalization assays were performed, PM-ox-LDL binding was as low as observed in CHO cells, but internalization was as high as in CHO-SR cells ($p < 0.05$). Competition experiments showed that PM₁₀ (40 $\mu\text{g}/\text{cm}^2$) induced decreased binding/internalization of PM-ox-LDL (22% and 32%, respectively; $p < 0.05$).




FIGURE 1. LDL oxidation by PM. LDL were exposed to increasing concentrations of PM_{2.5} or PM₁₀ (20, 40, and 80 $\mu\text{g}/\text{mg}$ LDL) during 24 h at 37°C with and without H₂O₂, and LDL oxidation was observed. Thiobarbituric acid reactive substances (TBARS). Mean of three independent assays \pm standard deviation (SD).




FIGURE 2. PM-ox-LDL binding/internalization by CHO and CHO-SR cells. CHO and CHO-SR cells were incubated with 2.5–40 µg/mL PM-ox-LDL-DiI for 4 h at 4°C (binding) or 37°C (internalization). LDL-DiI were previously oxidized with 80 µg PM₁₀ for 24 h. Cells were recovered and evaluated by flow cytometry. Mean of three independent assays ± SD. * *p* < 0.05 vs. similar concentration in CHO cells.




FIGURE 3. Competition of PM-ox-LDL and PM for SR-binding/internalization. Cultures of CHO-SR cells were exposed to 20, 40, and 80 µg/cm² of PM_{2.5} (A, B) or PM₁₀ (C, D), and 20 µg/mL PM-ox-LDL-DiI for 4 h at 4°C (A, C) or 37°C (B, D). Competition assays in the presence of PM_{2.5} or PM₁₀ resulted in decreased ox-LDL binding (53%–55%) and internalization (73%–78%) at 80 µg/cm². Mean ± SD of three independent experiments. * *p* < 0.05 vs. control.

Binding/internalization decrements were also observed in RAW cells when exposed to PM₁₀ (40 µg/cm²) and nat-LDL (45% and 55%, respectively; *p* < 0.05) (data not shown).

DISCUSSION

Evidence suggests that the mechanism by which PM exerts some of its adverse effects is through




FIGURE 4. Competition of nat-LDL and PM for LDL-R binding/internalization. CHO-SR cells were exposed to 20, 40, and 80 $\mu\text{g}/\text{cm}^2$ of $\text{PM}_{2.5}$ and 20 $\mu\text{g}/\text{mL}$ of native-LDL-DiI for 4 h at 4°C (A) or 37°C (B). PM interfered with nat-LDL binding (62%) and internalization (86%) at 80 $\mu\text{g}/\text{cm}^2$ with CHO-SR cells. Mean of three independent experiments \pm SD. * $p < 0.05$ vs. control.

oxidative potential [18–21]. Several studies support the existence of a relation between atherosclerosis and exposure to PM in both humans [13, 14] and animals [15–17], without fully understanding of the mechanisms involved. Results presented in this paper support that PM is capable of oxidizing LDL. Both PM fractions, PM_{10} and $\text{PM}_{2.5}$, showed similar LDL-oxidative potentials in spite of differences in chemical composition [2]. PM-induced LDL oxidation seems to be mediated by metallic and nonmetallic PM constituents, since oxidation was induced in the presence/absence of hydrogen peroxide. Therefore, our experimental data support the hypothesis that PM could have the potential to aggravate atherosclerosis by promoting the oxidation of LDL and subsequent uptake by macrophages.

We report that PM is capable of competing for SR and the LDL-receptor blocking natural ligands binding. Although it is known that various types of particles (e.g., diesel, PM_{1648}) [30–33] bind to SR, no one has previously reported that PM also alters LDL receptor functionality. This is an indication that PM could affect cell receptors with potential detrimental effects on cell function; the most obvious being the endocytic capacity of macrophages and a consequent altered primary immune response [34, 35]. Since SR is also present in other cell types like endothelial cells [36], it would be interesting to explore PM interactions with other receptors in various cell types.

The exact nature of PM/SR interactions and the mechanisms involved in the observed changes in receptor functionality remains to be described. These interactions may be related to receptor downregulation [37], direct physical PM/receptor interaction, or secondary to PM-related oxidative potential, as described under different conditions [38–41].

Since in this report we only studied PM-ox-LDL internalization through the type A scavenger receptor, studies evaluating the participation of other receptors involved in the atherosclerotic process, such as macrophage receptor with collagenous structure (MARCO), type B scavenger receptor, or lectin-type oxidized LDL receptor 1 (LOX-1), are still needed [42–44].

In spite of a growing body of research, mechanisms for PM-related cardiovascular effects are still not completely understood. For instance, we do not know whether PM can reach the bloodstream and directly affect vascular or cardiac cells, or if the effects observed *in vivo* are the result of inflammatory responses taking place in the respiratory system and then affecting the cardiovascular system by nervous or chemical mediators [45–50]. While it is possible that PM-induced systemic inflammation promotes the oxidation of LDL, our current study explored LDL oxidation in the event that PM or some of its compounds could enter the bloodstream and reach LDL.

In conclusion, we demonstrated that PM can oxidize LDL in a SR recognizable form. Although biological and epidemiological evidence relating PM exposure and atherosclerosis exists, the present work provides additional mechanistic evidence toward the elucidation of this phenomenon. We also observed that PM is able to induce changes in cell surface lipid receptors' functionality. This observation requires more studies since serious health implications would be related to these changes.

CONFLICT OF INTEREST STATEMENT

The authors declare that there are no conflicts of interest.

ACKNOWLEDGMENTS

This project was supported Consejo Nacional de Ciencia y Tecnología (CONACyT-M43138), Fogarty International Research Collaboration Award (FIRCA) and NIH/NIEHS grants R01 ES016932-01 and R01 ES017022-01. We thank the Gobierno de la Ciudad de México, Red Automática de Monitoreo Atmosférico (RAMA), Armando Retama and Rafael Ramos for making air sampling possible. A fellowship from Instituto de Ciencia y Tecnología del Distrito Federal (ICyTDF) has been granted to N. Manzano-León. We also thank Geraldine Flores-Rojas, Salvador Damián-Zamacona, and Nalleli Reyes-Martínez for technical assistance.

REFERENCES

- Samet J, Krewski D. Health effects associated with exposure to ambient air pollution. *J Toxicol Environ Health A* 2007;70:227–242.
- Osornio-Vargas AR, Bonner JC, Alfaro-Moreno E, Martínez L, García-Cuellar C, Ponce de León-Rosales S, Miranda J, Rosas I. Proinflammatory and cytotoxic effects of Mexico City air pollution particulate matter in vitro are dependent on particle size and composition. *Environ Health Perspect* 2003;111:1289–1293.
- Pérez L, Medina-Ramón M, Künzli N, Alastuey A, Pérez N, García R, Tobias A, Querol X, Sunyer J. Size fractionate particulate matter, vehicle traffic, and case-specific daily mortality in Barcelona, Spain. *Environ Sci Technol* 2009;43:4707–4714.
- Braga AL, Zanobetti A, Schwartz J. The lag structure between particulate air pollution and respiratory and cardiovascular deaths in 10 US cities. *J Occup Environ Med* 2001;43:927–933.
- Kang JY. Cardiovascular toxicity of particulate air pollution. *Cardiovasc Toxicol* 2002;2:233–236.
- Zanobetti A, Schwartz J, Dockery DW. Airborne particles are a risk factor for hospital admissions for heart and lung disease. *Environ Health Perspect* 2000;108:1071–1077.
- Peters A, Doring A, Wichmann H, Koenig W. Increased plasma viscosity during an air pollution episode: a link to mortality?. *Lancet* 1997;349:1582–1587.
- Gold DR, Litonjua A, Schwartz J, Lovett E, Larson A, Nearing B, Allen G, Verrier M, Cherry R, Verrier R. Ambient pollution and heart rate variability. *Circulation* 2000;101:1267–1273.
- Brook RD, Brook JR, Urch B, Vincent R, Rajagopalan S, Silverman F. Inhalation of fine particulate air pollution and ozone causes acute arterial vasoconstriction in healthy adults. *Circulation* 2002;105:1534–1536.
- Magari SR, Hauser R, Schwartz J, Williams PL, Smith TJ, Christiani DC. Association of heart rate variability with occupational and environmental exposure to particulate air pollution. *Circulation* 2001;104:986–991.
- Peters A, Liu E, Verrier RL, Schwartz J, Gold DR, Mittleman M, Baliff J, Oh JA, Allen G, Monohan K, Dockery DW. Air pollution and incidence of cardiac arrhythmia. *Epidemiology* 2000;11:11–17.
- Pope CA III, Hansen ML, Long RW, Nielsen KR, Eatough NL, Wilson WE, Eatough DJ. Ambient particulate air pollution, heart rate variability, and blood markers of inflammation in a panel of elderly subjects. *Environ Health Perspect* 2004;112:339–345.
- Diez Roux AV, Auchincloss AH, Franklin TG, Raghunathan T, Barr RG, Kaufman J. Long-term exposure to ambient particulate matter and prevalence of subclinical atherosclerosis in the Multi-Ethnic study of atherosclerosis. *Am J Epidemiol* 2008;167:667–675.
- Künzli N, Jerrett M, Mack WJ, Beckerman B, LeBree L, Gilliland F, Thomas D, Peters J, Hodis HN. Ambient air pollution and atherosclerosis in Los Angeles. *Environ Health Perspect* 2004;113:201–206.
- Suwa T, Hogg JC, Quinlan KB, Ohgami A, Vincent R, van Eeden SF. Particulate air pollution induces progression of atherosclerosis. *J Am Coll Cardiol* 2002;39:935–942.
- Araujo JA, Barajas B, Kleinman M, Wang X, Bennett BJ, Gong KW, Navab M, Harkema J, Sioutas C, Lulis AJ, Nel AE. Ambient particulate pollutants in the ultrafine range promote early atherosclerosis and systemic oxidative stress. *Circ Res* 2008;102:589–596.
- Sun Q, Wang A, Jin X, Natanzon A, Duquaine D, Brook RD, Aguinaldo JG, Fayad ZA, Fuster V, Lippmann M, Chen LC, Rajagopalan S. Long-term air pollution exposure and acceleration of atherosclerosis and vascular inflammation in an animal model. *JAMA* 2005;294:3003–3010.
- Chirino YI, Sánchez-Pérez Y, Osornio-Vargas AR, Morales-Bárceñas R, Gutiérrez-Ruiz MC, Segura-García Y, Rosas I, Pedraza-Chaverri J, García-Cuellar CM. PM₁₀ impairs the antioxidant defense system and exacerbates oxidative stress driven cell death. *Toxicol Lett* 2010;193:209–216.
- García-Cuellar C, Alfaro-Moreno E, Martínez-Romero F, Ponce de León Rosales S, Rosas I, Pérez-Cárdenas E, Osornio-Vargas AR. DNA damage induced by PM₁₀ from different zones of Mexico City. *Ann Occup Hyg* 2002;46(Suppl 1):425–428.
- Knaapen AM, Schins RPF, Steinfartz Y, Höhr D, Duneemann L, Borm P. Ambient particulate matter induces oxidative DNA damage in lung epithelial cells. *Inhal Toxicol* 2000;12(Suppl 3):125–132.
- Wessels A, Birmili W, Albrecht C, Hellack B, Jermann E, Wick G, Harrison RM, Schins RP. Oxidant generation and toxicity of size-fractionated ambient particles in human lung epithelial cells. *Environ Sci Technol* 2010;44:3539–3545.
- Steinberg D. Arterial metabolism of lipoproteins in relation to atherogenesis. *Ann NY Acad Sci* 1990;598:125–135.
- Obot CJ, Morandi MT, Beebe TP, Hamilton RF, Holian A. Surface components of airborne particulate matter induce macrophage apoptosis through scavenger receptors. *Toxicol Appl Pharmacol* 2002;184:98–106.
- Alfaro-Moreno E, Torres V, Miranda J, Martínez L, García-Cuellar C, Nawrot TS, Vanaudenaerde B, Hoet P, Ramírez-López P, Rosas I, Nemery B, Osornio-Vargas AR. Induction of IL-6 and inhibition of IL-8 secretion in the human airway cell line Calu-3 by urban particulate matter collected with a modified method of PM sampling. *Environ Res* 2009;109:528–535.
- Rosas-Pérez I, Serrano J, Alfaro-Moreno E, Baumgardner D, García-Cuellar C, Miranda-Martín del Campo J, Raga GB, Castillejos M, Drucker-Colín R, Osornio-Vargas AR. Relations between PM₁₀ composition and cell toxicity:

- A multivariate and graphical approach. *Chemosphere* 2007;67:1218–1228.
26. Pitas RE, Innerarity TL, Mahley RW. Cell surface receptor binding of phospholipid protein complexes containing different ratios of receptor-active and inactive E apoprotein. *J Biol Chem* 1980;255:5454–5460.
 27. Alfaro-Moreno E, García-Cuellar C, De Vizcaya Ruiz A, Rojas-Bracho L, Osornio-Vargas AR. Cellular Mechanism behind Particulate Matter Air Pollution Related Health Effects. In: Gurjar BR, Molina LT, Ojha CSP, editors. *Air pollution: Health & environmental impacts*. Boca Raton, FL: CRC Press (Taylor & Francis); 2010. pp 249–274.
 28. Basu SK, Goldstein JL, Anderson RG, Brown MS. Degradation of cationized low density lipoprotein and regulation of cholesterol metabolism in homozygous familial hypercholesterolemia fibroblast. *Proc Natl Acad Sci USA* 1976;73:3178–3182.
 29. Innerarity TL, Pitas RE, Mahley RW. Lipoprotein-receptor interactions. *Methods Enzymol* 1986;129:542–565.
 30. Beamer CA, Holian A. Scavenger receptor class A type I/II (CD204) null mice fail to develop fibrosis following silica exposure. *Am J Physiol Lung Cell Mol Physiol*. 2005;289:L186–L195.
 31. Palecanda A, Paulauskis J, Al-Mutairi E, Imrich A, Quin G, Suzuki H, Kodama T, Tryggvason K, Koziel H, Kobzik L. Role of the scavenger receptor MARCO in alveolar macrophage binding of unopsonized environmental particles. *J Exp Med* 1999;189:1497–1506.
 32. Resnick D, Freedman NJ, Xu S, Krieger M. Secreted extracellular domains of macrophage scavenger receptors form elongated trimers which specifically bind crocidolite asbestos. *J Biol Chem* 1993;268:3538–3545.
 33. Hamilton RF Jr, Pfau JC, Marshall GD, Holian A. Silica and PM1648 modify human alveolar macrophage antigen-presenting cell activity in vitro. *J Environ Pathol Toxicol Oncol* 2001;20(Suppl 1):75–84.
 34. Arredouani MS, Palecanda A, Koziel H, Huang YC, Imrich A, Sulahian TH, Ning YY, Yang Z, Pikkarainen T, Sankala M, Vargas SO, Takeya M, Tryggvason K, Kobzik L. MARCO is the major binding receptor for unopsonized particles and bacteria on human alveolar macrophages. *J Immunol* 2005;175:6058–6064.
 35. Monn C, Fendt R, Koller T. Ambient PM₁₀ extracts inhibit phagocytosis of defined inert model particles by alveolar macrophages. *Inhal Toxicol* 2002;14:369–385.
 36. Murphy JE, Tedbury PR, Homer-Vanniasakam S, Walker JH, Ponnambalam S. Biochemistry and cell biology of mammalian scavenger receptors. *Atherosclerosis* 2005;182:1–15.
 37. Lo YH, Pan MH, Li S, Yen JH, Kou MC, Ho CT, Wu MJ. Nobiletin metabolite, 3',4'-dihydroxy-5,6,7,8-tetramethoxyflavone, inhibits LDL oxidation and down-regulates scavenger receptor expression and activity in THP-1 cells. *Biochim Biophys Acta* 2010;1801:114–126.
 38. Aguilar-Gaytán R, Mas-Oliva J. Oxidative stress impairs endocytosis of the scavenger receptor class A. *Biochem Biophys Res Commun* 2003;305:510–517.
 39. Guo MY, Satoh K, Qi B, Narita T, Katsumata-Kato O, Matsuki-Fukushima M, Fujita-Yoshigaki J, Sugiya H. Thiol-oxidation reduces the release of amylase induced by β -adrenergic receptor activation in rat parotid acinar cells. *Biomed Res* 2010;31:293–299.
 40. Manzano-León N, Delgado-Coello B, Guaderrama-Díaz M, Mas-Oliva J. Beta-adaptin: key molecule for microglial scavenger receptor function under oxidative stress. *Biochem Biophys Res Commun* 2006;351:588–594.
 41. Yu WF, Nordberg A, Ravid R, Guan ZZ. Correlation of oxidative stress and the loss of the nicotinic receptor alpha4 subunit in the temporal cortex of patients with Alzheimer's disease. *Neurosci Lett* 2003;338:13–16.
 42. Catanese MT, Graziani R, von Hahn T, Moreau M, Huby T, Paonessa G, Santini C, Luzzago A, Rice CM, Cortese R, Vitelli A, Nicosia A. High-avidity monoclonal antibodies against the human scavenger class B type I receptor efficiently block hepatitis C virus infection in the presence of high-density lipoprotein. *J Virol* 2007;81:8063–8071.
 43. Lund AK, Lucero J, Harman M, Madden MC, McDonald JD, Seagrave JC, Campen MJ. The oxidized low-density lipoprotein receptor mediates vascular effects of inhaled vehicle emissions. *Am J Respir Crit Care Med* 2011;184:82–91.
 44. Thakur SA, Beamer CA, Migliaccio CT, Holian A. Critical role of MARCO in crystalline silica-induced pulmonary inflammation. *Toxicol Sci* 2009;108:462–471.
 45. Donaldson K, Stone V, Seaton A, MacNee W. Ambient particle inhalation and the cardiovascular system: potential mechanisms. *Environ. Health Perspect* 2001;109(Suppl 4):523–527.
 46. Furuyama A, Kanno S, Kobayashi T, Hirano S. Extrapulmonary translocation of intratracheally instilled fine and ultrafine particles via direct and alveolar macrophage-associated routes. *Arch Toxicol* 2009;83:429–437.
 47. Kato T, Yashiro T, Murata Y, Herbert DC, Oshikawa K, Bando M, Ohno S, Sugiyama Y. Evidence that exogenous substances can be phagocytized by alveolar epithelial cells and transported into blood capillaries. *Cell Tissue Res* 2003;311:47–51.
 48. Nemmar A, Vanbilloen H, Hoylaerts MF, Hoet PHM, Verbruggen A, Nemery B. Passage of intratracheally instilled ultrafine particles from the lung into the systemic circulation in hamster. *Am J Respir Crit Care Med* 2001;164:1665–1668.
 49. Oberdörster G, Sharp Z, Atudorei V, Elder A, Gelein R, Lunts A, Kreyling W, Cox C. Extrapulmonary translocation of ultrafine carbon particles following whole-body inhalation exposure of rats. *J Toxicol Environ Health A* 2002;65:1531–1543.
 50. van Eeden SF, Tan WC, Suwa T, Mukae H, Terashima T, Fujii T, Qui D, Vincent R, Hogg JC. Cytokines involved in the systemic inflammatory response induced by exposure to particulate matter air pollutants (PM₁₀). *Am J Respir Crit Care Med* 2001;164:826–830.