



# A spatial time-to-event approach for estimating associations between air pollution and preterm birth

Howard H. Chang,

*Emory University, Atlanta, USA*

Brian J. Reich

*North Carolina State University, Raleigh, USA*

and Marie Lynn Miranda

*University of Michigan, Ann Arbor, USA*

[Received June 2011. Revised March 2012]

**Summary.** The paper describes a Bayesian spatial discrete time survival model to estimate the effect of air pollution on the risk of preterm birth. The standard approach treats prematurity as a binary outcome and cannot effectively examine time varying exposures during pregnancy. Time varying exposures can arise either in short-term lagged exposures due to seasonality in air pollution or long-term cumulative exposures due to changes in length of exposure. Our model addresses this challenge by viewing gestational age as time-to-event data where each pregnancy becomes at risk at a prespecified time (e.g. the 28th week). The pregnancy is then followed until either a birth occurs before the 37th week (preterm), or it reaches the 37th week, and a full-term birth is expected. The model also includes a flexible spatially varying baseline hazard function to control for unmeasured spatial confounders and to borrow information across areal units. The approach proposed is applied to geocoded birth records in Mecklenburg County, North Carolina, for the period 2001–2005. We examine the risk of preterm birth that is associated with total cumulative and 4-week lagged exposure to ambient fine particulate matter.

**Keywords:** Air pollution; Fine particulate matter; Preterm birth; Reproductive epidemiology; Spatial survival data

## 1. Introduction

Preterm birth, which is defined as gestational age at delivery of less than 37 weeks, is linked to significant neonatal morbidity and mortality (Lorenz *et al.*, 1998; Goldenberg *et al.*, 2008; Saigal and Doyle, 2008), long-term health and developmental problems (Swamy *et al.*, 2008; Moster *et al.*, 2008) and medical costs (Institute of Medicine, 2006). There is a growing interest in examining the association between environmental exposures during pregnancy and adverse birth outcomes. Population studies have found consistent positive associations between ambient air pollution levels and low birth weight; however, epidemiological evidence remains mixed for preterm birth (Šràm *et al.*, 2005; Bosetti *et al.*, 2010).

Studies of air pollution and preterm birth utilize large birth record databases that provide extensive information on individual live births. For each birth, average exposures to air

*Address for correspondence:* Howard H. Chang, Department of Biostatistics and Bioinformatics, Emory University Rollins School of Public Health, 1518 Clifton Road North East, Mailstop 1518-002-3AA, Atlanta, GA 30322, USA.

E-mail: howard.chang@emory.edu

pollutants over specific susceptible pregnancy windows are then derived from air quality measurements. The most common analytic approach is carried out via logistic regression where preterm *versus* full-term births are treated as binary outcomes (Wilhelm and Ritz, 2005; Huynh *et al.*, 2006; Leem *et al.*, 2006; Ritz *et al.*, 2007; Brauer *et al.*, 2008). This approach is most suitable to examine exposure metrics that do not vary during pregnancy, such as the first 6 weeks since conception, or the first and second trimester (Woodruff *et al.*, 2009). However, many long-term and short-term exposure windows are time varying because ambient pollution levels exhibit strong seasonality.

Consider using the average air pollution level over the *entire* pregnancy to estimate the long-term effect of air pollution on preterm birth. In a logistic regression model, bias in risk estimates may arise with this overall exposure metric because the lengths of exposure differ between preterm and full-term births. For pregnancies that are conceived in the winter months, preterm births are more likely to experience lower average exposure than full-term births. This is because full-term births have a longer exposure window extending into the summer months when ambient pollution concentrations are typically higher. However, for pregnancies that are conceived in the summer, preterm births experience higher exposure levels compared with full-term births. The direction and magnitude of the bias therefore depend on the seasonality in both air pollution and the number of on-going pregnancies in the population (Darrow *et al.*, 2009b). This challenge is also present in estimating the effect of air pollution during the third trimester (27th week till birth).

Estimating the short-term effects of air pollution on preterm birth during late pregnancy is also problematic by using logistic regression. A common approach is to capture late pregnancy exposure with a window before delivery (e.g. 4 weeks or 6 weeks before birth). However, this exposure metric does not coincide with the period when a full-term birth is at risk of being preterm. For example, consider a 40-week full-term pregnancy that experienced high exposure during the month before birth. This pregnancy will contribute to a protective effect of air pollution even though it cannot be preterm after week 37. Using only the weeks before birth also discards data from earlier weeks that are informative about the short-term effect.

The main contribution of this paper is to describe a model for preterm birth that addresses the above challenges in estimating the effects of long-term and short-term exposures that are time varying during pregnancy. This is accomplished by viewing gestational age as time-to-event (survival) data where each pregnancy enters the risk set at a prespecified time (e.g. the 28th week). The pregnancy is then followed until either a birth occurs before the 37th week (preterm), or it reaches the 37th week and a full-term birth is expected. Therefore, we align the data such that pregnancies are compared with each other only during a window at risk of being preterm (i.e. 28th–37th week). This allows us to examine

- (a) long-term effects by using a time varying *cumulative* average instead of an average over the entire pregnancy and
- (b) short-term effects by using a time varying *lagged* average instead of a fixed period defined before delivery.

The risk estimates from the time-to-event approach have similar interpretations to that obtained from a time series analysis stratified by gestational week (Darrow *et al.*, 2009a). In a time series analysis, the outcome of interest is the daily number of preterm births aggregated over a geographic region. The corresponding daily exposure metric is obtained by averaging air pollution exposures across all on-going pregnancies on each day. The time series design also overcomes the challenges in defining time varying exposures because the aggregate exposure is allowed to vary between days. The time series design was originally motivated by the issue of

unmeasured confounders. In contrast, our proposed approach utilizes the full spatial and temporal contrast in air pollution levels and can control for individual level covariates. Moreover, a time series analysis has limited power to detect long-term effects because considerable temporal variation in the exposure is removed when controlling for seasonality in preterm births.

In studies that rely on birth certificate data, the issue of residual confounding due to unmeasured risk factors is well recognized because the health outcome is compared across space and time (Northam and Knapp, 2009; Strickland *et al.*, 2009). Examples of known risk factors for preterm birth that are typically not available from birth certificates include the mother's socio-economic status, maternal body mass index, level of stress and anxiety, amount of physical work, the quality of the built environment and infections status. Often these factors may also, at least partially, determine the amount of personal exposure to air pollution due to outdoor sources. We control for unmeasured spatial confounders by including a flexible baseline hazard model that is spatially varying. This approach also allows us to borrow information across spatial units to estimate the baseline hazards at locations with small numbers of births.

Spatially referenced survival models often account for association between nearby regions by including a random effect (frailty) for the region of residence in the linear predictor, and smoothing the frailties by using a Gaussian spatial model (Banerjee *et al.*, 2003). Although this approach allows different regions to have different baseline risks, it assumes that the general shape of the survival curve is the same for each region after accounting for the spatial frailty term. For example, this does not allow for some regions to have elevated risk very early in pregnancy but low risk late in the pregnancy. For continuous survival data, one generalization of the frailty model is to use an accelerated failure time model, i.e. suitably transform the survival times and model the transformed responses by using linear regression while allowing the mean and the entire shape of the residual distribution to vary spatially. Many models for the spatially varying residual density exist; see for example Griffin and Steel (2006), or Reich and Fuentes (2007). In contrast, with the model for continuous survival data that was described above, the North Carolina birth certificate records gestational age as the number of completed weeks. Therefore, we propose a simpler model for discrete time survival data.

We apply the proposed model to estimate the total cumulative and 4-week lag effects of ambient particulate matter that is less than  $2.5 \mu\text{m}$  in diameter ( $\text{PM}_{2.5}$ ). We treat ambient  $\text{PM}_{2.5}$ -concentration as a surrogate measure for personal exposure to fine particulate matter from outdoor sources. The  $\text{PM}_{2.5}$ -mass represents a chemically diverse mixture of solids and liquids that arise from combustion processes. Exposure to  $\text{PM}_{2.5}$ -pollution has been associated with numerous health outcomes including mortality, emergency department visits and hospital admissions (Pope and Dockery, 2006). The biological mechanisms by which particulate matter might affect preterm birth focus on initiation of the inflammation pathway (Kannan *et al.*, 2006).

The remainder of this paper is organized as follows. Section 2 gives the general modelling framework of our discrete time spatial survival model for preterm birth. Section 3 describes the air pollution data and a data set of geocoded births from Mecklenburg County, North Carolina, for the period 2001–2005. Because past studies predominantly use logistic regression to analyse preterm birth, Section 4 describes a simulation study that compares the model proposed and the standard approach that treats prematurity as a binary outcome. We highlight the potential bias in risk estimates when surrogate time invariant exposures (e.g. pregnancy average or 4 weeks before birth) are used instead of time varying exposures (e.g. cumulative average or 4-week lag). Results from the health analysis are given in Section 5. Finally, discussion and future work appear in Section 6.

## 2. Spatial time-to-event model for preterm birth

For pregnancy  $i$ , we observe the follow-up time  $t_i$ , an indicator of whether the pregnancy was censored  $c_i$ , the spatial location  $s_i$  and a vector of  $p$  potentially time-dependent covariates  $\mathbf{X}_i(t) = (X_{1i}(t), \dots, X_{pi}(t))'$ . We assume a discrete domain for spatial locations. For example, in our application  $s_i$  represents census tracts in Mecklenburg County. We also assume a discrete domain for event times because gestational age is typically recorded as the number of completed weeks. We define gestational weeks 28–36 as the at-risk period for a pregnancy being preterm. For preterm births with gestational age less than 37 weeks,  $t_i$  represents the completed week of gestation and  $c_i = 0$ . Full-term births of at least 37 weeks are censored at week 36 ( $t_i = 36$  and  $c_i = 1$ ) because they are no longer at risk of being preterm. Therefore, under this modelling framework (which is illustrated in Fig. 1),

- (a) no censoring occurs between gestational weeks 28–35,
- (b) all preterm births experienced an event and
- (c) all full-term births are censored.


The model is defined through the event hazard rate,  $\pi\{\mathbf{X}_i(t), s_i, t\} \in [0, 1]$ , which is the probability of birth for pregnancy  $i$  at week  $t$  given that pregnancy  $i$  reaches week  $t$ . This implies the survival probability

$$P(t_i > m) = \prod_{t=1}^m [1 - \pi\{\mathbf{X}_i(t), s_i, t\}], \tag{1}$$

i.e. the usual life table model. We model the discrete event hazard rate by using spatial probit regression:

$$\pi\{\mathbf{X}_i(t), s_i = s, t\} = \Phi\{\beta_0(s, t) + \mathbf{X}_i(t)' \boldsymbol{\beta}\}, \tag{2}$$

where  $\Phi$  is the standard normal distribution function and  $\boldsymbol{\beta}$  is the vector of regression coefficients. The probit link is chosen to facilitate Bayesian inference by using Markov chain Monte Carlo sampling and other link functions may be considered, e.g. a logistic link (Holmes and Held, 2006). Parameter  $\beta_0(s, t)$  determines the baseline risks (the event rate for a subject with  $X_{1i}(t) = \dots = X_{pi}(t) = 0$ ). Because  $\beta_0(s, t)$  varies with both space and time, this model spans the entire class of baseline models on this discrete domain of  $s_i \in \{1, \dots, S\}$  and  $t \in \{28, \dots, 36\}$ , where  $S$  is the total number of discrete spatial units in the study region.



**Fig. 1.** A time-to-event approach for preterm birth and air pollution: a preterm birth and a full-term birth are shown with pregnancywide cumulative exposure and 4-week lagged exposure given at week 30 (●, conception date; ×, birth date; ○, censored)

Let  $\beta_0(s, t) = \eta + \mu(s) + \gamma(t) + \theta(s, t)$ , where  $\eta$  is the overall average,  $\mu(s)$  is a spatial effect,  $\gamma(t)$  is a temporal effect and  $\theta(s, t)$  is the space–time interaction. Since the spatial terms  $\boldsymbol{\mu} = (\mu(1), \dots, \mu(S))'$  are discrete areal units, we model them by using the conditionally auto-regressive (CAR) model (Besag, 1974). This spatial model is specified through spatial adjacencies. Let  $s \sim s'$  indicate that regions  $s$  and  $s'$  are spatial neighbours and  $m_s$  be the number of spatial neighbours of region  $s$ . The CAR model is often defined through the full conditional distribution of  $\mu(s)$  given  $\boldsymbol{\mu}$  at all other locations. The full conditional distribution is Gaussian with

$$E\{\mu(s)|\boldsymbol{\mu}(s'), s' \neq s\} = \rho_\mu \sum_{s' \sim s} \mu(s')/m_s, \tag{3}$$

$$V\{\mu(s)|\boldsymbol{\mu}(s'), s' \neq s\} = \sigma_\mu^2/m_s. \tag{4}$$

The full conditional mean is proportional to the average of the spatial neighbours, where  $\rho_\mu \in [0, 1]$  controls the degree of spatial association, and the variance is controlled by  $\sigma_\mu^2 > 0$ . The joint model for the vector  $\boldsymbol{\mu}$  is multivariate normal with mean 0 and covariance  $\sigma_\mu^2(M_S - \rho_\mu C_S)^{-1}$ , where the  $(s, s')$  element of  $C_S$  is  $C_S(s, s') = I(s \sim s')$  and  $M_S$  is diagonal with diagonal elements  $\sum_{s' \neq s} C_S(s, s') = m_s$ . We denote this model as  $\boldsymbol{\mu} \sim \text{CAR}(\rho_\mu, \sigma_\mu^2, C_S)$ .

The temporal effects  $\boldsymbol{\gamma} = (\gamma(28), \dots, \gamma(36))'$  control the temporal average baseline hazard function. The vector  $\boldsymbol{\gamma}$  has a lag 1 auto-regressive model which can be written  $\boldsymbol{\gamma} \sim \text{CAR}(\rho_\gamma, \sigma_\gamma^2, C_T)$ , where  $C_T$  is the  $9 \times 9$  temporal adjacency matrix with  $(t, t')$  element equal to  $I(|t - t'| = 1)$ . The spatiotemporal random effects have the dynamic spatial model (Banerjee *et al.*, 2003)

$$\theta(s, t) = \rho_\theta \theta(s, t - 1) + \delta(s, t), \quad t = 29, \dots, 36, \tag{5}$$

where  $\rho_\theta \in (0, 1)$  and  $\boldsymbol{\delta}_t = (\delta(1, t), \dots, \delta(S, t))' \sim \text{CAR}(\rho_\delta, \sigma_\delta^2, C_S)$ . For identification purposes, we fix  $\theta(s, 28) = 0$  for all  $s$ .

The above baseline hazard function model has several special cases. If  $\theta(s, t) \equiv 0$ , then  $\beta_0(s, t) = \eta + \mu(s) + \gamma(t)$ , and the baseline risk function varies spatially only through the spatial frailties  $\mu(s)$ . Therefore the shape of the risk function for all regions is constant and controlled by  $\boldsymbol{\gamma}$ . If  $\rho_\mu = \rho_\delta = 0$ , then the baseline risk functions are exchangeable across locations, and hierarchically centred on  $\eta + \boldsymbol{\gamma}$ .

Inference is carried out in a Bayesian framework by specifying priors for the model parameters. Parameter  $\eta$  and each component of  $\boldsymbol{\beta}$  are assigned  $N(0, 100^2)$ . The variances  $\sigma_\mu^2, \sigma_\gamma^2$  and  $\theta_\theta^2 \sim \text{gamma}(a_1, b_1)$ . Following Kelsall *et al.* (1999), we take  $a_1 = 0.5$  and  $b_1 = 0.005$ . The CAR association parameters  $\rho_\mu, \rho_\gamma, \rho_\theta, \rho_\delta \sim \text{beta}(a_2, b_2)$ . We discretize the prior to 1000 equally spaced points spanning  $[0, 1]$  and, to give an uninformative prior, we take  $a_2 = b_2 = 1$ .

For each pregnancy, we augment the data  $(c_i, t_i)$  to  $(Y_i(28), \dots, Y_i(t_i))$ , where  $Y_i(t) = 0$  for  $t < t_i$  and  $Y_i(t_i) = 1 - c_i$ . Therefore, at each time point during the pregnancy,  $Y_i(t)$  indicates whether a preterm birth occurred. The model for pregnancy  $i$  can then be written as  $Y_i(t) \sim \text{Bernoulli}[\Phi\{\beta_0(s, t) + \mathbf{X}_i(t)' \boldsymbol{\beta}\}]$ , independent across time. The Bernoulli model for  $Y_i(t)$  is equivalent to the model  $Y_i(t) = I\{Z_i(t) > 0\}$ , where  $Z_i(t)$  is a latent variable with  $Z_i(t) \sim N\{\beta_0(s, t) + \mathbf{X}_i(t)' \boldsymbol{\beta}, 1\}$ .

After introducing the latent variables  $Z_i(t)$ , the model is entirely conjugate, and we used Gibbs sampling (Casella and George, 1992) to analyse the posterior distributions of all unknown parameters. All analysis was carried out in R 2.8.0 (R Development Core Team, 2009). We generated 20 000 samples and discarded the first 5000 samples as burn-in. Convergence was monitored by using trace plots and auto-correlation plots for several representative parameters. In the on-line supplementary material, we describe the Markov chain Monte Carlo algorithm in detail and provide the R code for fitting the spatial survival model.

### 3. Health and exposure data

Birth data for Mecklenburg County were obtained from the North Carolina detailed birth record database. Mecklenburg County is the most populous county in North Carolina and contains the city Charlotte. We included all pregnancies that were conceived from the period 2001–2005 using the clinical estimate of gestation in the birth record to back-calculate the date of conception. We restricted the analysis to singleton live births with birth weight 400 g or more and no congenital anomalies. We further restricted the data set to those mothers aged 15–44 years who self-declared as non-Hispanic white, non-Hispanic black or Hispanic.

Daily PM<sub>2.5</sub>-data were obtained from the statistically fused air quality database ([http://www.epa.gov/esd/land-sci/lcb/lcb\\_sfads.html](http://www.epa.gov/esd/land-sci/lcb/lcb_sfads.html)). The database is a recent product from the US Environmental Protection Agency that provides predicted daily PM<sub>2.5</sub>-concentration averaged over contiguous 12 km × 12 km grid cells. We chose this data set because monitors in the air quality system network typically measure PM<sub>2.5</sub> only every third or sixth day. The database predictions are based on a Bayesian space–time hierarchical model (McMillan *et al.*, 2009) that combines

- (a) PM<sub>2.5</sub>-data from the air quality system network and
- (b) outputs from the models-3–community multiscale air quality model (Byun and Schere, 2006), which is an air quality model that simulates the complex interactions between weather and air pollutants on the basis of atmospheric chemistry and physics. Although this model provides higher spatial and temporal resolution compared with the air quality system network, its output is known to exhibit bias, particularly for capturing short-term variation between days (Mebust *et al.*, 2003). The statistically fused air quality database attempts to adjust the bias in the community multiscale air quality model by using the observed PM<sub>2.5</sub>-concentrations from the air quality system network.

Maternal residential addresses at the time of delivery were geocoded to the street block level by using ArcGIS 9.3 software (Esri, Redlands, California). We used 2006 topologically integrated geographic encoding and referencing street data from the US Census Bureau as the spatial reference file. The geocoding success rate was 97.1%, owing to invalid, missing or unmatched addresses. Using the latitude and longitude co-ordinates that were delivered by the geocoding process, we linked each pregnancy in space and time to one of the statistically fused air quality grid cells overlapping Mecklenburg County.

### 4. Simulation study

This section describes a simulation study to compare the approach proposed *versus* viewing prematurity as a binary outcome. Reproducing the PM<sub>2.5</sub> exposure levels in Mecklenburg County, we generated 1000 replicates of simulated exposures and gestational age for births that were conceived in the year 2001 as follows. Let  $\hat{X}(c)$  denote the average PM<sub>2.5</sub>-level during the week leading up to day  $c$ . For each pregnancy  $i$  conceived on day  $c$ , we generated its weekly PM<sub>2.5</sub> exposure series  $X_{ij}$  for gestational week  $j = 28, \dots, 42$  as

$$X_{ij} = \hat{X}(c + j) + \varepsilon_{ij}, \quad \varepsilon_{ij} \sim N(0, \sigma^2).$$

The above exposure model assumes that pregnancies conceived on the same day share mean exposure time series, and parameter  $\sigma^2$  controls the between-pregnancies variation on a particular day. The total sample size was 10588 births with a median of 30 conceptions per day.

**Table 1.** Simulation study results: bias and coverage probability of a 95% confidence interval based on 1000 simulated replicate data sets

Relative risk	Bias ( $\times 100$ )				Coverage probability			
	Cumulative		4-week lag		Cumulative		4-week lag	
	Survival	Probit	Survival	Probit	Survival	Probit	Survival	Probit
1.00	-0.01	0.23	0.00	0.34	0.96	0.94	0.96	0.90
1.01	0.02	0.42	-0.01	0.37	0.95	0.91	0.95	0.87
1.02	-0.01	0.51	0.00	0.42	0.95	0.90	0.95	0.85
1.03	-0.03	0.60	0.02	0.47	0.94	0.89	0.95	0.84
1.04	0.01	0.78	0.01	0.50	0.95	0.85	0.96	0.81
1.05	-0.01	0.91	0.02	0.53	0.95	0.80	0.95	0.79

We estimated  $\hat{X}(c)$  and  $\sigma^2 = 0.30$  on the basis of the actual exposure series in our study. Weekly  $PM_{2.5}$ -averages show strong temporal correlation with a lag 1 auto-correlation of 0.94. Using  $X_{ij}$  we constructed total cumulative and 4-week lagged exposure  $X_i(t)$  for each pregnancy.

The gestational age  $t_i$  for pregnancy  $i$  was generated with probabilities

$$P(t_i = t) = h_i(t) \prod_{t' < t} \{1 - h_i(t')\}, \quad t = 28, \dots, 42,$$

where  $h_i(t) = \Phi\{\hat{h}_0(t) + \beta X_i(t)\}$  for  $t \leq 36$  and  $h_i(t) = h_0(t)$  for  $t > 36$ . In other words, we generated  $Y_i(t)$  for  $t = 28, \dots, 42$  and took  $t_i = \min\{t | Y_i(t) = 1\}$ . We estimated  $\hat{h}_0(t)$  from the data and allowed the hazard ratio  $\beta$  for  $PM_{2.5}$  to vary in the simulation. Here we do not consider spatial variation in baseline risks.

In the time-to-event approach, we modelled  $Y_i(t)$ , which is an indicator of whether a birth occurred in week  $t = 28, \dots, t_i$ , by using a discrete time survival model  $\Phi\{Y_i(t) = 1\} = \beta_0(t) + \beta_1 X_i(t)$ . We also modelled the occurrence of a preterm birth via probit regression as  $\Phi\{P(t_i < 37)\} = \beta_0 + \beta_1 X_i$ . In the time-to-event approach,  $X_i(t)$  represents time varying cumulative and 4-week lagged exposure. In the probit regression,  $X_i$  represents the surrogate measures of using the average  $PM_{2.5}$ -level for the entire pregnancy and the 4-week exposure before delivery.

Table 1 gives the bias and 95% confidence interval coverage probability for various values of approximate relative risk ( $1.927\beta$ ) per interquartile range of  $PM_{2.5}$ -exposures. The root-mean-squared error and average confidence interval length are given in the on-line supplementary materials, section 3. We found that the survival model consistently outperforms the probit regression based on the exposure levels and variations in our study population. Also, when treating prematurity as a binary outcome, the surrogate time invariant metrics led to a positive bias in the risk estimates. The bias can be attributed to the seasonality in conceptions and  $PM_{2.5}$ -levels. Specifically, the largest number of conceptions occurred in May 2001. Among this birth cohort, full-term births experience lower  $PM_{2.5}$ -levels later in the pregnancy which coincides with the winter months. Therefore, in the simulation full-term births are more likely to have lower average exposures across the entire pregnancy and the 4 weeks before birth, even though they were not at risk of being preterm past the 37th week.

## 5. Analysis of North Carolina preterm birth data

### 5.1. Health model for preterm birth and $PM_{2.5}$

We examined the effects of average  $PM_{2.5}$ -levels over two time varying exposure windows. Given a pregnancy-completed gestational week  $t$ , we considered the fixed length short-term exposure of 4-week lag (4 weeks leading up to the date that week  $t$  was completed). We also considered the long-term cumulative exposure of conception till week  $t$  where the exposure window length varies with gestation age.

We controlled for the following time-independent variables: maternal age (15–19, 20–24, 25–29, 30–34, 35–39 and 40–44 years), maternal education (less than 9, 9–11, 12, 13–15 and more than 15 years), race or ethnicity (non-Hispanic white, non-Hispanic black and Hispanic), tobacco use during pregnancy (yes or no), marital status (married or unmarried), first born (yes or no), infant sex (male or female) and percentage population below poverty of each census tract obtained from the 2000 US census. This choice of covariates as potential confounders was based on a previous study of air pollution and birth weight in the same study population (Gray *et al.*, 2009). To control for unmeasured time varying confounders, we included the season of conception (winter, December–February; spring, March–May; summer, June–August; autumn, September–November) and indicators for conception year. We also calculated a 1-week lagged average temperature for gestational weeks 28–36. We modelled the short-term effect of temperature as a smooth function via natural cubic splines with 4 degrees of freedom.

We considered three models for the baseline risk: non-spatial with  $\beta_0(s, t) = \gamma(t)$ , spatial frailty with  $\beta_0(s, t) = \mu(s) + \gamma(t)$  and space–time interaction  $\beta_0(s, t) = \mu(s) + \gamma(t) + \theta(s, t)$ . Here  $\beta_0(s, t)$  represents the baseline prevalence of preterm birth at tract  $s$  among pregnancies that reached gestational week  $t$ . We assume that the effects of all other covariates are constant in space and time.

Finally, we discuss the interpretation of the regression coefficients  $\beta$  where the probit link makes interpretation difficult. However, for small probabilities, the Gaussian distribution function can be approximated with an exponential function leading to an approximate relative risk interpretation. Specifically, for  $z \in (-3, -1)$ , and thus  $\Phi(z) \in (0.001, 0.159)$ ,  $\Phi(z) \approx \exp(0.136 + 1.927z)$ . This approximation is quite accurate; over this range of  $z$ ,  $\exp(0.136 + 1.927z)$  explains over 99.7% of the variation in  $\Phi(z)$ . Therefore, we present the posteriors of  $\beta^* = 1.927\beta$  and refer to  $\exp(\beta^*)$  as the approximate relative risk of preterm birth due to a unit increase in the covariates.

### 5.2. Results

Our study included a total of 55647 geocoded births (7.7% preterm) representing all 144 census tracts in Mecklenburg County, North Carolina. In the study population, the average  $PM_{2.5}$ -level across the entire pregnancy had a mean of  $15.5 \mu\text{g m}^{-3}$  and an interquartile range of  $1.37 \mu\text{g m}^{-3}$ . The average  $PM_{2.5}$ -level across a 4-week window had a mean of  $15.5 \mu\text{g m}^{-3}$  and an interquartile range of  $4.56 \mu\text{g m}^{-3}$ .

Table 2 gives the posterior means and 95% posterior intervals of the coefficients, in terms of approximate relative risk ( $1.927\beta_j$ ). Estimates are from a model that includes a space–time interaction baseline hazard and average  $PM_{2.5}$ -levels over the entire pregnancy. Higher risks of preterm birth were observed for older, unmarried, non-Hispanic black mothers and among those who reported tobacco use. Mothers with more than 15 years of education were at a reduced risk of preterm birth compared with mothers with 12 years of education. First-born babies and those that were conceived during the summer months were also more likely to be preterm. We did not find an acute effect of temperature during late pregnancy. Also, census tracts with higher



**Table 2.** Posterior mean and 95% posterior interval for the relative increase in preterm birth risk associated with various factors†

<i>Covariate</i>	<i>Estimate (95% posterior interval)</i>
Male	1.00 (0.96, 1.05)
Tobacco	1.34 (1.23, 1.46)
Unmarried	1.19 (1.12, 1.27)
Firstborn	1.21 (1.15, 1.27)
Tract level % poverty (×10)	1.05 (1.01, 1.08)
<i>Ethnicity</i>	
Non-Hispanic white	Reference
Non-Hispanic black	1.40 (1.31, 1.49)
Hispanic	0.99 (0.91, 1.08)
<i>Mother's education (years)</i>	
< 9	0.99 (0.88, 1.09)
9–11	1.09 (1.00, 1.17)
12	Reference
13–15	1.00 (0.93, 1.07)
> 15	0.83 (0.78, 0.90)
<i>Mother's age (years)</i>	
Age 15–19	0.95 (0.86, 1.03)
Age 20–24	0.94 (0.88, 1.00)
Age 25–29	Reference
Age 30–34	1.14 (1.08, 1.21)
Age 35–39	1.31 (1.21, 1.42)
Age 40–44	1.47 (1.27, 1.67)
<i>Conception season</i>	
June–August	Reference
September–November	0.92 (0.84, 1.01)
December–February	0.92 (0.83, 1.01)
March–May	0.98 (0.92, 1.05)

†Estimates are from a model that includes space–time interaction baseline hazards and average PM<sub>2.5</sub>-levels over the entire pregnancy.

proportions of families below the federal poverty line were associated with higher rates of preterm birth. These results are consistent with findings from previous studies (Wilhelm and Ritz, 2005).

Table 3 gives the estimates and 95% probability intervals of the PM<sub>2.5</sub>-coefficients under various baseline risk models. The estimates are presented as approximate relative risk per interquartile range. We found a consistent positive association between average total cumulative PM<sub>2.5</sub>-exposure and the risk of preterm birth. Specifically, controlling for a tract-specific baseline hazard model (space–time interaction), an interquartile range (1.73 μg m<sup>-3</sup>) increase was associated with a 7.3% (95% posterior interval 2.5, 11.7) increase in the risk of preterm birth. The magnitude of our risk estimate is consistent with previous studies using average exposure across the entire pregnancy (Brauer *et al.*, 2008). The deviance information criterion, effective degrees of freedom, posterior predictive loss and estimates for the CAR parameters are given in the on-line supplementary materials, section 2.

We did not find a statistically significant association between a 4-week lagged exposure and preterm birth. The PM<sub>2.5</sub>-level in Mecklenburg County is below the national ambient air quality standards and short-term exposure may not be sufficiently high to induce a shift in gestation.

**Table 3.** Posterior mean and 95% posterior interval of the PM<sub>2.5</sub>-coefficients under various baseline risk models†

Baseline hazard	Cumulative estimate (95% posterior interval)	4-week lag estimate (95% posterior interval)
Non-spatial	1.067 (1.020, 1.116)	1.023 (0.961, 1.087)
Spatial frailty	1.069 (1.020, 1.248)	1.017 (0.956, 1.074)
Space–time interaction	1.073 (1.025, 1.117)	1.031 (0.977, 1.088)

†The estimates are presented as approximate relative risks ( $1.927\beta_j$ ) per inter-quartile range ( $1.37 \mu\text{g m}^{-3}$  for total cumulative and  $4.56 \mu\text{g m}^{-3}$  for 4-week lag).



**Fig. 2.** Baseline tract-specific rates of (a) preterm births (less than 37 gestational weeks) (□, [6.8, 7.25]%; □, [7.25, 7.45]%; ■, [7.45, 7.55]%; ■, [7.55, 8]%) and very preterm birth (less than 34 gestational weeks) (□, [1.45, 1.55]%; □, [1.55, 1.65]%; ■, [1.65, 1.75]%; ■, [1.75, 1.85]%) baseline hazard rates are calculated at the average value of each covariate across all areas; estimates are from a model that includes space–time interaction baseline hazards and average PM<sub>2.5</sub>-levels over the entire pregnancy

Several studies have reported evidence linking short-term exposure to PM<sub>2.5</sub> and preterm births in urban communities with higher levels of PM<sub>2.5</sub> such as Atlanta, Georgia (Darrow *et al.*, 2009a), California (Huynh *et al.*, 2006) and Pennsylvania (Sagiv *et al.*, 2005).

To visualize the spatial variation in baseline risk across counties, Fig. 2 plots the tract-specific baseline rates of preterm birth and very preterm birth (less than 34 gestational weeks). The hazard rates are centred at the average value of each covariate across the study population. Tract-specific baseline hazards for the individual gestational week are given in Fig. 1 in the on-line supplementary material. The rates are categorized into four groups indicated by different shadings by a *k*-means algorithm that minimizes within-group variation. This analysis is not

intended to suggest that it is the geography itself that is driving these differences in preterm birth rates. Rather there is a spatially patterned latent variable which we cannot account for when relying solely on birth certificate data. We found relatively small spatial variation at the census tract level in our study region. We also found that the differences in deviance information criterion values are extremely small and found no statistically significant differences in baseline risks across tracts. We note that our objective is not to identify a model that best predicts the occurrence of preterm birth, but to assess the robustness of risk estimates under different ways to control for unmeasured spatial confounders. The ability to model baseline hazard functions flexibly may be crucial in other settings such as a county level analysis across the entire state of North Carolina.

### 6. Discussion

We present a model for preterm birth by viewing gestational age as discrete time survival data. The Bayesian modelling framework also incorporates a flexible spatial baseline hazard function. The approach proposed can examine *both* long-term and short-term environmental exposures, such as ambient air pollution, that are potentially time varying during pregnancy. Bosetti *et al.* (2010) noted that previous studies often did not report results for all exposure metrics, resulting in the possibility of selective reporting and difficulty in synthesizing findings. Although we report only the total cumulative and 4-week lagged exposure metrics to demonstrate our approach, additional time varying metrics such as the third trimester, 6-week lag and 1-week lag can be examined.

Several extensions of our spatial survival model are possible. For example, it would be straightforward to incorporate non-stationarity by allowing  $\sigma_\mu^2$ ,  $\sigma_\gamma^2$  or  $\sigma_\delta^2$  to vary with space or time. The variances could be modelled as independent draws from a common prior, or as a log-Gaussian process to encourage the variability to change smoothly over space or time. Also, we have centred all random effects at the constant  $\eta$ . A constant baseline mortality rate is similar to an exponential distribution. Centring on other parametric distributions is also possible; for example, replacing  $\eta$  with

$$g(t|\eta_1, \eta_2) = (\eta_1/\eta_2)(t/\eta_2)^{\eta_1-1}$$

would approximate a Weibull distribution. Also, about two-thirds of preterm births had low birth weight (less than 2500 g) and another model extension is to consider joint modelling of gestational age and the risk of low birth weight.

Each regression coefficient can also be modelled by using the spatiotemporal CAR model following the model for  $\beta_0(s, t)$ , i.e.

$$\beta_j(s, t) = \eta_j + \mu_j(s) + \gamma_j(t) + \theta_j(s, t),$$

where  $\mu_j \sim \text{CAR}(\rho_{\mu_j}, \sigma_{\mu_j}^2, C_S)$ ,  $\gamma_j \sim \text{CAR}(\rho_{\gamma_j}, \sigma_{\gamma_j}^2, C_T)$ ,  $\theta_j(s, t) = \rho_{\theta_j} \theta_j(s, t - 1) + \delta_j(s, t)$  and  $\delta_{jt} \sim \text{CAR}(\rho_{\delta_j}, \sigma_{\delta_j}^2, C_S)$ . In full generality, this allows the effect of the  $j$ th covariate (e.g. exposure to PM<sub>2.5</sub>-pollution) to vary either by spatial location or gestational age. Posterior inference is also straightforward via Gibbs sampling. One potential future analysis is to examine whether air pollution effects vary across spatial unit and whether the spatial variation in risks is associated with spatial variation in population characteristic.

There are additional challenges that are common in the analysis of preterm birth and air pollution that our model does not consider and warrant further investigation. The first challenge arises from assigning PM<sub>2.5</sub>-exposure to each individual pregnancy and the associated potential measurement error. We used the statistically fused air quality data set to avoid missing daily

observations in calculating average exposure. However, we ignored the spatial change of support problem by assigning the average exposure over a 12 km × 12 km grid cell to the point level (geocoded residence of the mother). A significant subset of mothers also moves during pregnancy (Canfield *et al.*, 2006). The second challenge concerns outcome misclassification, particularly around pregnancies with gestational age near the 37th-week cut-off. The gestational length for each pregnancy was clinically estimated by physicians and the measurement error could differ on the basis of when prenatal care was initiated, as well as whether ultrasound fetal diagnostics were part of the routine prenatal care. For example, using Illinois data from 1989–1991, Mustafa and David (2001) found that the concordance between gestational age obtained by using reported last menstrual period and clinical estimates are 78% for the 1-week difference and 87% for 2-week differences. Ananth (2007) also found that the rates of preterm birth based on clinical estimates were lower relative to that based on last menstrual periods. Finally, there is also a growing interest in differentiating spontaneous and medically indicated preterm births (Savitz *et al.*, 2005) where the effects of air pollution may be heterogeneous across clinical subtypes and severity of preterm births.

## Acknowledgements

The research is supported by grant DMS-0635449 from the National Science Foundation and grant RD-83329301-4 from the US Environmental Protection Agency.

## References

- Ananth, C. V. (2007) Menstrual versus clinical estimate of gestational age dating in the United States: temporal trends and variability in indices of perinatal outcomes. *Paed. Perinat. Epidemiol.*, **21**, 20–30.
- Banerjee, S., Carlin, B. and Gelfand, A. (2003) Hierarchical modeling and analysis for spatial data. Boca Raton: Chapman and Hall.
- Besag, J. (1974) Spatial interaction and the statistical analysis of lattice systems (with discussion). *J. R. Statist. Soc. B*, **36**, 192–236.
- Bosetti, C., Nieuwenhuijsen, M. J., Gallus, S., Cipriani, S., La Vecchia, C. and Parazzini, F. (2010) Ambient particulate matter and preterm birth or birth weight: a review of the literature. *Arch. Toxicol.*, **84**, 447–460.
- Brauer, M., Lencar, C., Tamburic, L., Koehoorn, M., Demers, P. and Karr, C. (2008) A cohort study of traffic-related air pollution impacts on birth outcomes. *Environ. Health Perspect.*, **116**, 680–686.
- Byun, D. J. and Schere, K. L. (2006) Review of the governing equations, computational algorithms, and other components of the Model-3 Community Multiscale Air Quality (CMAQ) modeling system. *Appl. Mech. Rev.*, **59**, 51–77.
- Canfield, M. A., Ramadhani, T. A., Langlois, P. H. and Waller, D. K. (2006) Residential mobility patterns and exposure misclassification in epidemiologic studies of birth defects. *J. Expos. Sci. Environ. Epidemiol.*, **16**, 538–543.
- Casella, G. and George, E. I. (1992) Explaining the Gibbs sampler. *Am. Statist.*, **6**, 167–174.
- Darrow, L. A., Klein, M., Flanders, W. D., Waller, L. A., Correa, A., Marcus, M., Mulholland, J. A., Russell, A. G. and Tolbert, P. E. (2009a) Ambient air pollution and preterm birth: a time-series analysis. *Epidemiology*, **20**, 689–698.
- Darrow, L. A., Strickland, M. J., Klein, M., Waller, L. A., Flanders, W. D., Correa, A., Marcus, M. and Tolbert, P. E. (2009b) Seasonality of birth and implications for temporal studies of preterm birth. *Epidemiology*, **20**, 699–706.
- Goldenberg, R. L., Culhane, J. F., Iams, J. D. and Romero, R. (2008) Epidemiology and causes of preterm birth. *Lancet*, **371**, 75–89.
- Gray, S. C., Edwards, S. E. and Miranda, M. L. (2009) Assessing exposure metrics for PM and birth weight models. *J. Expos. Sci. Environ. Epidemiol.*, **20**, 469–477.
- Griffin, J. E. and Steel, M. F. J. (2006) Order-based dependent Dirichlet processes. *J. Am. Statist. Ass.*, **101**, 179–194.
- Holmes, C. C. and Held, L. (2006) Bayesian auxiliary variable models for binary and multinomial regression. *Bayes Anal.*, **1**, 145–168.
- Huynh, M., Woodruff, T. J., Parker, J. D. and Schoendorf, K. C. (2006) Relationships between air pollution and preterm birth in California. *Paed. Perinat. Epidemiol.*, **20**, 454–461.
- Institute of Medicine (2006) *Preterm Birth: Causes, Consequences, and Prevention*. Washington DC: National Academies Press.

- Kannan, S., Misra, D. P., Dvonch, J. T. and Krishnakumar, A. (2006) Exposures to airborne particulate matter and adverse perinatal outcomes: a biologically plausible mechanistic framework for exploring potential effect modification by nutrition. *Environ. Hlth Perspect.*, **114**, 1636–1642.
- Kelsall, J. E., Wakefield, J. C., Bernardo, J. M., Berger, J. O., Dawid, A. P. and Smith, A. F. M. (eds) (1999) Comment on Bayesian model for spatially correlated disease and exposure data. In *Bayesian Statistics 6*. Oxford: Oxford University Press.
- Leem, J. H., Kaplan, B. M., Shim, Y. K., Pohl, H. R., Gotway, C. A., Bullard, S. M., Rogers, J. F., Smith, M. M. and Tylenda, C. A. (2006) Exposures to air pollutants during pregnancy and preterm delivery. *Environ. Hlth Perspect.*, **114**, 905–910.
- Lorenz, J., Wooliever, D., Jetton, J. and Paneth, N. (1998) A quantitative review of mortality and developmental disability in extremely premature newborns. *Arch. Ped. Adolesc. Med.*, **152**, 425–435.
- McMillan, N. J., Holland, D. M., Morara, M. and Feng, J. (2009) Combining numerical model output and particulate data using Bayesian space-time modeling. *Environmetrics*, **21**, 48–65.
- Mebust, M. R., Eder, B. K., Binkowski, F. S. and Roselle, S. J. (2003) Models-3 Community Multiscale Air 392 Quality (CMAQ) model aerosol component 2 model evaluation. *J. Geophys. Res.*, **108**, 4184–4202.
- Moster, D., Lie, R. T. and Markestad, T. (2008) Long-term medical and social consequences of preterm birth. *New Engl. J. Med.*, **359**, 262–273.
- Mustafa, G. and David, R. (2001) Comparative accuracy of clinical estimate versus menstrual gestational age in computerized birth certificates. *Publ. Hlth Rep.*, **116**, 15–21.
- Northam, S. and Knapp, T. R. (2009) The reliability and validity of birth certificates. *J. Obstetr. Gyn. Neonl Nurs.*, **35**, 3–12.
- Pope III, C. A. and Dockery, D. W. (2006) Health effects of fine particulate air pollution: lines that connect. *J. Air Waste Mangmnt Ass.*, **56**, 709–742.
- R Development Core Team (2009) *R: a Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Reich, B. J. and Fuentes, M. (2007) A multivariate semiparametric Bayesian spatial modeling framework for hurricane surface wind fields. *Ann. Appl. Statist.*, **1**, 249–264.
- Ritz, B., Wilhelm, M., Hoggatt, K. J. and Ghosh, J. C. (2007) Ambient air pollution and preterm birth in the environment and pregnancy outcome study at the University of California, Los Angeles. *Am. J. Epidem.*, **166**, 1045–1052.
- Sagiv, S., Mendola, P., Loomis, D., Herring, A. H., Neas, L. M., Savitz, D. A. and Poole, C. (2005) A time series analysis of air pollution and preterm birth in Pennsylvania. *Environ. Hlth Perspect.*, **113**, 602–606.
- Saigal, S. and Doyle, L. W. (2008) An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet*, **371**, 261–269.
- Savitz, D. A., Dole, N., Herring, A., Kaczor, D., Murphy, J., Siega-Riz, Jr, A. M. and MacDonald, T. L. (2005) Should spontaneous and medically indicated preterm births be separated for study aetiology? *Paed. Perntl Epidem.*, **19**, 97–105.
- Šrám, R. J., Binkova, B., Dejmek, J. and Bobak, M. (2005) Ambient air pollution and pregnancy outcomes: a review of the literature. *Environ. Hlth Perspect.*, **113**, 375–382.
- Strickland, M. J., Klein, M., Darrow, L. A., Flanders, W. D., Correa, A., Marcus, M. and Tolbert, P. E. (2009) The issue of confounding in epidemiological studies of ambient air pollution and pregnancy outcomes. *J. Epidem. Commty Hlth*, **63**, 500–504.
- Swamy, G. K., Østbye, T. and Skjærven, R. (2008) Association of preterm birth with long-term survival, reproduction, and next-generation preterm birth. *J. Am. Med. Ass.*, **299**, 1429–1436.
- Wilhelm, M. and Ritz, B. (2005) Local variations in CO and particulate air pollution and adverse birth outcomes in Los Angeles County, California, USA. *Environ. Hlth Perspect.*, **113**, 1212–1221.
- Woodruff, T. J., Parker, J. D., Darrow, L. A., Slama, R., Bell, M., Choi, H., Glinianaia, S., Hoggatt, K., Karr, C., Lobdell, D. T. and Wilhelm, M. (2009) Methodological issues in studies of air pollution and reproductive health. *Environ. Res.*, **109**, 311–320.

#### Supporting information

Additional 'supporting information' may be found in the on-line version of this article:

'Supplementary materials for A spatial time-to-event approach for estimating associations between air pollution and preterm birth'.

Please note: Wiley-Blackwell are not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the author for correspondence for the article.