


Survival Analysis of Study Participants with serum acylcarnitine data that received either L-Carnitine (18g) or Placebo in the RACE Clinical Trial (<https://clinicaltrials.gov/ct2/show/NCT01665092>)



Hazard Ratio (Mantel-Haenszel)	1.134	
Ratio (and its reciprocal)	0.7299 to 1.763	
95% CI of ratio		
	Placebo-treated	L-Carnitine (18g)-treated
Number of rows	170	170
# of blank lines	100	70
# rows with impossible data	0	0
# censored subjects	33	53
# deaths/events	37	47

Number of subjects at risk at various times:		
Survival Days	Placebo-treated	Carnitine (18g)-treated
0	70	100
0.08		100
0.16		99
0.2		98
0.21	70	
0.36		97
1	69	96
2	63	86
3	60	85
4	57	83
5	56	81
6	55	79
7	53	76
8	52	
9	50	74
10		72
11	49	
13	48	
14	47	
15		71
16	46	68
17		65
18		64
21	45	63
22	43	62
23		61
25		60
27		59
28	42	58
29	41	
30	40	
36	39	57
52		56
54		55
56	37	
66	36	
72		54
83	35	
84	34	
108		53
148		52
166	33	
190		51
227	32	
242	31	
269	30	
365	29	50